

Evaluation of crop yield and fruit quality in organic apple production

Gonda, I., Rakonczás, N. & Holb, I.J.

Debrecen University, Centre of Agricultural Sciences, Department of Horticulture and Plant Biotechnology
H-4015 Debrecen, Böszörményi str 138., Hungary

Summary: In this study, crop yield and fruit quality parameters (soluble solid concentration, sugar, total acid, and vitamin C content) of 15 apple cultivars including old, resistant and currently grown cultivars. The highest fruit yield among all 15 cultivars was obtained by Jonagold with 15.5 kg/tree, while the lowest yield was measured on the resistant cultivar Reanda with 8.1 kg/tree. There were no significant differences among the cultivars in fruit diameter ranging from 70 mm in the case of cv. Húsvéti rozmaring to 82 mm for cv. Mutsu. Variations in soluble solids concentration of the cultivars varied between 15.5% for cv. Téli Banán and 19.2% for cv. Renora. Sugar content values showed a similar tendency to soluble solids content. The highest and lowest total sugar content was observed on cvs. Jonagold and Renora and Retina with 17% and 12.4%, respectively. Total acid content values ranged between 0.18% and 0.53% for cvs. Jonagold and Remo, respectively. Values of vitamin C content for the 15 apple cultivars varied between 2 mg% and 4 mg% for cv. Retina and cv. Mutsu, respectively.

Key words: yield, fruit diameter, soluble solid concentration, sugar content, total acid, vitamin C, organic apple, resistant apple cultivar

Introduction

Sustainable and environmentally-benign production including organic apple production is a major principle of today's fruit growing sector (Gonda, 1993, 2000). In organic production, only organic manures can be applied, while in the integrated production artificial fertilizers can also be used, with respect to environmental and health protection limitations (Soltész & Szabó, 1997). In the pest management used in organic orchards, copper- and sulphur-based and natural products can be applied, while in the integrated systems, the use of numerous synthetic, environmentally-friendly products is permitted (Soltész & Szabó, 1997; Holb & Heijne, 2001).

Increasing interest of consumers toward organic fruits has an increasing impact on the growers also, who show higher affinity to organic apple production. However, some parts of organic production technology, such as nutrient supply, phytotechnical operations and plant protection, are a great challenge to most growers (Gonda, 1993, 2000). Especially plant protection can be a difficult task to be solved in organic apple production (Holb, 2000). In several cases, costs of organic apple production are higher compared to integrated or conventional production, and therefore, economic aspects of organic production cannot be neglected (Ellis et al. 1998). However, priority is given to 'organic fruits' as harmful pesticide residues of fruits are less compared with traditional apple production. Soltész (1998) and Gonda et al (2000) presented class intervals of inner content parameters of some apple cultivars (vitamin C, soluble solid content, sugar, and

acid content). Inner content parameters of fruits in organic vs. integrated or conventional apple production investigated by some researchers. Ruger (1984) and K et al. (1990) showed no significant differences of content of fruit between conventional and organic production. Gonda et al. (2000) were not able to prove differences in vitamin C, soluble solid content, sugar, and acid content of organic apple fruits compared to integrated ones. The study on crop yield and fruit quality parameters has not been conducted in organic apple production on a large number of apple cultivars.

The aim of the present study was to determine crop yield and fruit quality parameters (soluble solid content, sugar, total acid, and vitamin C content) of 15 apple cultivars including old, resistant and currently grown cultivars.

Materials and methods

Orchard site

Data assessments were made in an apple collection at Debrecen-Pallag, Debrecen University Department of Horticulture and Plant Biotechnology orchard was planted in 1997, on M26 rootstock, with a spacing of 4 x 1.5m. In one plot, there were 7 trees of one cultivar. Plots were arranged in a randomized block system in 3 replicates in the orchard. The orchard was treated according to the Hungarian Organic Guidelines (Anon, 1997). Orchard soil type was brown

soil. Orchard was not irrigated. Nutrient supply was maintained with synthetic fertilizers in the integrated orchard part. Stable manure, 30 t ha⁻¹ was applied to the soil in 2001 and 2003 in the orchard. The orchard consisted of 39 apple cultivars. Fifteen cultivars (six resistant ones: cv. Retina, cv. Rewena, cv. Remo, cv. Reanda, cv. Renora, and cv. Liberty; six currently grown ones: cv. Gala Must, cv. Jonagold, cv. Jonica, cv. Idared, cv. Red Elstar, and cv. Mutsu; and three old ones: cv. Húsvéti rozmaring, cv. Téli banán, and cv. Batul) were selected for evaluating crop yield and four fruit quality parameters of harvested fruits.

Yield

Fruits of each cultivar were harvested at the technological maturity phase in 2001, 2002 and 2004. The largest fruit diameter of each cultivar was measured with a vernier caliper then total weight of fruits per tree was measured with a commercial scale. Both measures (fruit diameter and weight) were averaged to get mean fruit diameter and yield values per tree for each cultivar.

Fruit quality parameters

After measuring fruit diameter and weight, 5 fruits from 4 trees of each cultivar were randomly selected for fruit quality measures. Fruit samples were used to measure soluble solids concentration, total sugar, total acidic and vitamin C content for each cultivar. Soluble solids concentration was determined on a freshly squeezed juice sample from each fruit using a refractometer calibrated prior to use with a 10% (weight:volume) sucrose solution. Total sugar, total acidic and vitamin C contents were determined by standard Hungarian Fruit Quality Evaluation Methods.

Statistical analysis

Analyses of variance were used to separate means of fruit yield and fruit




Figure 1 Mean yield (kg/tree) of 15 selected apple cultivars in an organic apple orchard (Debrecen-Pallag, Hungary, 2001, 2002 and 2004).

(%)
Total
sugar

Fig
(Del




Figure 2 Mean fruit diameter (mm) of 15 selected apple cultivars in an organic apple orchard (Debrecen-Pallag, Hungary, 2001, 2002 and 2004).

(%)
Total
acid

Fig
Pall




Figure 3 Mean soluble solids concentration (%) of 15 selected apple cultivars in an organic apple orchard (Debrecen-Pallag, Hungary, 2003).

(%)
Vitamin C
content

Fi
(D




Figure 4 Mean total sugar content (%) of 15 selected apple cultivars in an organic apple orchard (Debrecen-Pallag, Hungary, 2001, 2002, 2004).




Figure 5 Mean total acid content (%) of 15 selected apple cultivars in an organic apple orchard (Debrecen-Pallag, Hungary, 2001, 2002 and 2004).




Figure 6 Mean vitamin C content (mg%) on 15 selected apple cultivars in an organic apple orchard (Debrecen-Pallag, Hungary, 2001, 2002 and 2004).

quality parameters for the cultivars. Significant differences were determined at $P = 0.05$.

Results and discussion

Yield and fruit diameter

In the average of three years, highest fruit yield among all cultivars was obtained on cv. Jonagold with 15.5 kg/tree, while the lowest yield was measured on the resistant cultivar Reanda with 8.1 kg/tree (Figure 1). The highest yield among the resistant and old cultivars was measured on cv. Liberty and cv. Baul with 12 kg/tree and 10.8 kg/tree respectively. Cvs. Húsvéti rozmaring and Gala Must gave the lowest yield among the old and currently grown cultivars with 10.5 kg/tree and 10.8 kg/tree, respectively.

There were no significant differences among the cultivars in fruit diameter ranging from 70 mm in the case of cv. Húsvéti rozmaring to 82 mm for cv. Mutsu (Figure 2).

Fruit quality parameters

Soluble solids concentration

Values of soluble solids concentration of the cultivars varied between 15.5% for cv. Téli Banán and 19.2% for cv. Renora (Figure 3). Among the currently grown cultivars, cv. Jonagold and cv. Mutsu had the highest (19%) and lowest (16.5%) soluble solids content, respectively. The lowest soluble solids content were measured on cv. Retina and Téli Banán among the resistant and old cultivars, respectively. Fruits with higher soluble solids concentration contain relatively less water, and they are more compact. Fruits having these properties can be stored better.

Total sugar content

Sugar content values showed a similar tendency to soluble solids content (Figure 4). The highest and lowest total sugar content

observed on cvs. Jonagold and Renora and cv. Retina with 17% and 12.4%, respectively. Among the old cultivars, the highest and lowest sugar contents were obtained for cvs. Batul and Téli Banán, respectively. The lowest sugar content among the currently grown cultivars was measured in cv. Idared, its total sugar content was similar to that of cv. Batul, which had the highest sugar content among the old cultivars.

Total acid content

Total acid content values ranged between 0.18% and 0.53% for cvs. Jonagold and Remo, respectively (Figure 5). The highest values were similar for all three groups of the cultivars. Among the currently grown and old cultivars, the highest values were obtained for cv. Red Elstar (0.52%) and cv. Batul (0.51%) respectively. Apple fruits with higher acid content maintain their freshness and good texture for a longer time and decay more slowly during storage.

Vitamin C content

Values of vitamin C content for the 15 apple cultivars varied between 2 mg% and 4 mg% for cv. Retina and cv. Mutsu, respectively (Figure 6). The largest vitamin C content among the resistant and the old cultivars was detected in cvs. Remo and Húsvéti rozsmaring, respectively. The lowest vitamin C contents were measured in cv. Jonica (2.8 mg%) among the currently grown cultivars and cv. Batul (2.2 mg%) among the old cultivars.

As a summary, it can be stated that inner content values of the examined cultivars were within the optimal values given by Soltész (1998). Similarly to a previous study of Gonda et al. (2000) higher vitamin C, soluble solid content, sugar, and acid content of organic apple fruits were found compared with fruits from integrated orchards.

References

- Ellis, M. A., Ferree, D. C., & Madden, L. V. (1998): Effects of an apple scab-resistant cultivar on use patterns of inorganic and organic fungicides and economics of disease control. *Plant Disease* 82: 428–433.
- Gonda, I. (1993): A fitotechnikai műveletek szerepe az alma integrált termesztéstechnológiájában. *Integrált termesztés a kertészetben* 14: 72–78.
- Gonda, I. (2000): Minőségi almatermesztés, 1st Ed.; Primom Vállalkozáselnökítő Alapítvány: Nyíregyháza, Hungary, 378 pp.
- Gonda, I., Holb, I.J., Bitskey, K. 2000: Rate of scab infection and quality parameters of apple fruit in organic and integrated production systems. *International Journal of Horticultural Science* 6 (4): 63–67.
- Holb, I. (2000): Az alma ventúriás varasodásának mértéke integrált és organikus védekezési programokban. *Kertgazdaság* 32(2): 25–35.
- Holb, I.J., & Heijne, B. (2001): Evaluating primary scab control in organic apple production. *European Journal of Horticultural Science* 66 (5): 254–261.
- Keipert, K., Wedler, A., & Overbeck, G. (1990): Alternativer Anbau von Äpfeln und Gemüse. (Alternative cultivation of apples and vegetables.) Schriftenreihe-der-Landwirtschaftskammer-Rheinland. 66: 1–195.
- Ruger, H. (1984): Vorläufige Ergebnisse aus biologisch-dynamischen Apfelanbauversuchen. (Preliminary results of tests on the bio-dynamic method of apple growing.) *Obstbau*. 9 (3): 110–114.
- Soltész, M. (1998): A fajták megválasztása. In: Soltész M. (eds.): *Gyümölcsfajtaismeret és -használat*. Mezőgazda, Budapest. p. 119–154.
- Soltész, M., & Szabó, T. (1997): Alma. In: Soltész M. (eds.): *Gyümölcsfajtaismeret és -használat*. Mezőgazda, Budapest. p. 428–437.