

On the Partial Approximation of Sets

Zoltán Csajbók

University of Debrecen, Faculty of Health

Abstract 1 According to Pawlak's classic rough set theory the vagueness of a subset of a finite universe U is defined by the difference of its upper and lower approximations with respect to an equivalence relation on U . A natural way of the generalization of this idea is that the equivalence relation is replaced by either any other type of binary relations on U or an arbitrary covering of the universe. In this paper, our starting point will be an *arbitrary* family of subsets of an *arbitrary* universe U . Within this framework, we shall investigate a possible generalization of Pawlak's idea. Both Pawlak's rough set theory and our approach can extensively be applied in medical informatics.

Keywords: vagueness, rough set theory, partial approximation of sets, medical informatics

Lektor: Dr. Mihálydeák Tamás, egyetemi tanár, Debreceni Egyetem

1. Introduction

The rough set theory was introduced by the Polish mathematician Z. Pawlak in the early 1980s [15, 16]. Let U be a *finite* set of distinguishable objects which is called the universe of discourse, and $\varepsilon \subseteq U \times U$ be an equivalence relation on U . The elements of partition generated by ε are called ε -elementary sets. An ε -elementary set can be viewed as a set of indiscernible objects characterized by the same available information about them [18, 20]. Any union of ε -elementary sets is referred to as *definable* set. An arbitrary subset $X \subseteq U$ may not necessarily be

a union of some ε -elementary sets. However, it can be naturally approximated by two sets, by the union of ε -elementary sets that are subsets of X , called the lower ε -approximation of X , and by the union of ε -elementary sets that have a nonempty intersection with X , called the upper ε -approximation of X .

The basic idea of Pawlak's rough set theory is that the vagueness [12, 17, 19, 20] of a set is described by the difference of its upper and lower ε -approximations called the ε -boundary of the set. A set is *rough* if its ε -boundary is nonempty.

Using partitions, however, is a strict requirement. Moreover, in practice, there are attributes which do not characterize all members of an observed collection of objects [6, 13].

A natural way of the generalization of Pawlak's idea is that the equivalence relation is replaced by any other type of binary relations on U [10, 11, 24]. Another generalization is the assumption that the starting point is an arbitrary covering of the universe [1, 23, 26, 27]. In this paper, our starting point will be an *arbitrary* family of subsets of an *arbitrary* universe U . We will not assume whether this family of sets covers the universe or the universe is finite.

The paper is organized as follows. In Section 2 we summarize the basic notations used throughout the paper. Section 3 presents the basic concepts and their properties of the classical Pawlak's rough set theory. Only those facts which are important from the point of view of the generalization will be considered. The major contributions of this paper are covered in Section 4 which summarizes the basic principles of the partial approximation of sets.

2. Basic Notations

Let U be any set. The powerset of U is 2^U . If $\mathfrak{A} \subseteq 2^U$, then the union of \mathfrak{A} is $\bigcup \mathfrak{A} = \{x \mid \exists A \in \mathfrak{A} (x \in A)\}$, the intersection of \mathfrak{A} is $\bigcap \mathfrak{A} = \{x \mid \forall A \in \mathfrak{A} (x \in A)\}$. If \mathfrak{A} is an empty family of sets we define $\bigcup \emptyset = \emptyset$ and $\bigcap \emptyset = U$.

If ϵ is an arbitrary binary relation on U , let $[x]_\epsilon$ denote the ϵ -related elements to x , i.e., $[x]_\epsilon = \{y \in U \mid (x, y) \in \epsilon\}$. They are called ϵ -elementary sets, and the family of $[x]_\epsilon$ is denoted by U/ϵ .

Let X and Y be nonempty sets and $f : X \rightarrow Y$ be a map. If $\text{dom} f = X$ then f is *total*, if $\text{dom} f \subsetneq X$ then f is *partial*. If f is a partial map, then $\text{dom} f = \emptyset$ is allowed. For the purpose of simplicity we will talk about partial maps without direct references to their partial properties. However, statements with respect to partial maps always concern their restrictions to their domains.

A nonempty set P together with a partial order \leq on P is called a *poset*, in symbols (P, \leq) . Let (P, \leq_P) and (Q, \leq_Q) be two posets. A map $f : P \rightarrow Q$ is *monotone* if $x \leq_P y \Rightarrow f(x) \leq_Q f(y)$, *order-embedding* if $x \leq_P y \Leftrightarrow f(x) \leq_Q f(y)$, and *order-isomorphism* if f is an order-embedding onto Q . In general, monotone maps are many-to-one correspondences. An order embedding is always monotone and injective. Hence, f is an order-isomorphism if and only if f is a bijection, and both f and f^{-1} are monotone.

3. Basics of Pawlak's Rough Set Theory

Definition 3.1. The pair (U, ε) where U is a finite universe of discourse and ε is an equivalence relation on U is called *Pawlak's approximation space*.

Equivalence classes generated by the equivalence relation ε are ε -*elementary sets*. A subset $X \subseteq U$ is ε -*definable*, if it is a union of ε -elementary sets, otherwise X is ε -*undefinable*. By definition, the empty set is considered to be an ε -definable set.

Let $\mathfrak{D}_{U/\varepsilon}$ denote the family of ε -definable subsets of U .

Remark 3.2. The idea of approximation *space* is a bit younger than Pawlak's initial works. For an evolutionary survey of approximation spaces, see [20].

The following statement is elementary, however, in the context of Pawlak's rough set theory it is an important fact. For the sake of simple reference, it is formulated in a lemma. It follows just from the fact that the partition U/ε generated by ε consists of nonempty pairwise disjoint subsets of U .

Lemma 3.3. $\forall \mathfrak{X} \in 2^{U/\varepsilon} \forall X \in U/\varepsilon (X \subseteq \bigcup \mathfrak{X} \Leftrightarrow X \in \mathfrak{X})$.

Clearly, by Definition 3.1, $\mathfrak{D}_{U/\varepsilon}$ contains the empty set and is closed under complementation and unions. According to Lemma 3.3, it is also closed under intersections, i.e., $\mathfrak{D}_{U/\varepsilon}$ is a σ -algebra with basis U/ε .

Proposition 3.4 ([3], Theorem 8) Let (U, ε) be a Pawlak's approximation space.

Then the map $u_\varepsilon : 2^{U/\varepsilon} \rightarrow \mathfrak{D}_{U/\varepsilon}, \mathfrak{X} \mapsto \bigcup \mathfrak{X}$ is an order isomorphism between $(2^{U/\varepsilon}, \subseteq)$ and $(\mathfrak{D}_{U/\varepsilon}, \subseteq)$.

Corollary 3.5. Any ε -definable subset $D \in \mathfrak{D}_{U/\varepsilon}$ of U can be written uniquely in the following form:

$$D = \bigcup \mathfrak{X}, \text{ where } \mathfrak{X} = \{X \mid X \in U/\varepsilon, X \subseteq D\} \in 2^{U/\varepsilon},$$

that is, there is no other $\mathfrak{X}' \in 2^{U/\varepsilon}$ satisfying $D = \bigcup \mathfrak{X}'$.

Proof. Since $D \in \mathfrak{D}_{U/\varepsilon}$, thus $D = \bigcup \mathfrak{X}$ immediately holds by Lemma 3.3. However, u_ε is a bijection, so $u_\varepsilon^{-1}(D) \in 2^{U/\varepsilon}$ is unique and $u_\varepsilon^{-1}(D) = \mathfrak{X}$. \square

In Pawlak's approximation spaces, lower and upper approximations of $X \in 2^U$ can be defined in three equivalent forms [21, 22, 25].

Definition 3.6. Let (U, ε) be a Pawlak's approximation space and $X \in 2^U$ be any subset of U .

The lower ε -approximation of X is

$$\underline{\varepsilon}(X) = \{x \in U \mid [x]_{\varepsilon} \subseteq X\}, \quad (3.1a)$$

$$= \bigcup \{Y \mid Y \in U/\varepsilon, Y \subseteq X\} \quad (3.1b)$$

$$= \bigcup \{Y \mid Y \in \mathfrak{D}_{U/\varepsilon}, Y \subseteq X\}, \quad (3.1c)$$

the upper ε -approximation of X is

$$\bar{\varepsilon}(X) = \{x \in U \mid [x]_{\varepsilon} \cap X \neq \emptyset\}, \quad (3.2a)$$

$$= \bigcup \{Y \mid Y \in U/\varepsilon, Y \cap X \neq \emptyset\}, \quad (3.2b)$$

$$= \bigcap \{Y \mid Y \in \mathfrak{D}_{U/\varepsilon}, X \subseteq Y\}. \quad (3.2c)$$

It follows just from the definitions that $\underline{\varepsilon}(X), \bar{\varepsilon}(X) \in \mathfrak{D}_{U/\varepsilon}$, in addition the maps $\underline{\varepsilon}, \bar{\varepsilon} : 2^U \rightarrow \mathfrak{D}_{U/\varepsilon}$ are total and many-to-one.

Proposition 3.7 Let (U, ε) be a Pawlak's approximation space and $X \in 2^U$ be a subset of U . The sets $\underline{\varepsilon}(X), \bar{\varepsilon}(X)$ can be written uniquely in the following forms:

$$\underline{\varepsilon}(X) = \bigcup \mathfrak{X}, \text{ where } \mathfrak{X} = \{Y \mid Y \in U/\varepsilon, Y \subseteq X\} \in 2^{U/\varepsilon},$$

$$\bar{\varepsilon}(X) = \bigcup \bar{\mathfrak{X}}, \text{ where } \bar{\mathfrak{X}} = \{Y \mid Y \in U/\varepsilon, Y \cap X \neq \emptyset\} \in 2^{U/\varepsilon},$$

that is, there are no other $\mathfrak{X}', \bar{\mathfrak{X}}' \in 2^{U/\varepsilon}$ satisfying $\underline{\varepsilon}(X) = \bigcup \mathfrak{X}'$ and $\bar{\varepsilon}(X) = \bigcup \bar{\mathfrak{X}}'$.

Proof. According to Definition 3.6 (3.1b), (3.2b), we only have to prove the uniqueness.

$\underline{\varepsilon}(X), \bar{\varepsilon}(X) \in \mathfrak{D}_{U/\varepsilon}$, and so, by Proposition 3.4, $u_{\varepsilon}^{-1}(\underline{\varepsilon}(X))$ and $u_{\varepsilon}^{-1}(\bar{\varepsilon}(X))$ are unique. Hence, by Lemma 3.3, we get

$$\begin{aligned} u_{\varepsilon}^{-1}(\underline{\varepsilon}(X)) &= \{Y \mid Y \in U/\varepsilon, Y \subseteq \underline{\varepsilon}(X)\} \\ &= \{Y \mid Y \in U/\varepsilon, Y \subseteq \bigcup \{Y' \mid Y' \in U/\varepsilon, Y' \subseteq X\}\} \\ &= \{Y \mid Y \in U/\varepsilon, Y \subseteq \{Y' \mid Y' \in U/\varepsilon, Y' \subseteq X\}\} \\ &= \{Y \mid Y \in U/\varepsilon, Y \subseteq X\} = \mathfrak{X}. \\ u_{\varepsilon}^{-1}(\bar{\varepsilon}(X)) &= \{Y \mid Y \in U/\varepsilon, Y \subseteq \bar{\varepsilon}(X)\} \\ &= \{Y \mid Y \in U/\varepsilon, Y \subseteq \bigcup \{Y' \mid Y' \in U/\varepsilon, Y' \cap X \neq \emptyset\}\} \\ &= \{Y \mid Y \in U/\varepsilon, Y \subseteq \bigcup \{Y' \mid Y' \in U/\varepsilon, Y' \cap X \neq \emptyset\}\} \\ &= \{Y \mid Y \in U/\varepsilon, Y \cap X \neq \emptyset\} = \bar{\mathfrak{X}}. \end{aligned}$$

□

Basic properties of lower and upper ε -approximations can be found, e.g. in [10, 16]. Here we cite only a few among them which will be important in the following.

Proposition 3.8 ([16], Proposition 2.1, point a)) Let (U, ε) be a Pawlak's approximation space. Then $X \in \mathfrak{D}_{U/\varepsilon}$ if and only if $\underline{\varepsilon}(X) = \bar{\varepsilon}(X)$.

Proposition 3.9 ([16], Proposition 2.2, points 1, 9, 10) Let (U, ε) be a Pawlak's approximation space. Then

$$\forall X \in 2^U (\underline{\varepsilon}(X) \subseteq X \subseteq \bar{\varepsilon}(X)),$$

that is, the maps $\underline{\varepsilon}$ and $\bar{\varepsilon}$ are contractive and extensive, respectively.

Corollary 3.10. $\underline{\varepsilon}(X) = X$ if and only if $X = \bar{\varepsilon}(X)$.

Proof. Since $\underline{\varepsilon}(X) \in \mathfrak{D}_{U/\varepsilon}$ ($\bar{\varepsilon}(X) \in \mathfrak{D}_{U/\varepsilon}$), then $X = \underline{\varepsilon}(X) \in \mathfrak{D}_{U/\varepsilon}$ ($X = \bar{\varepsilon}(X) \in \mathfrak{D}_{U/\varepsilon}$), and so, by Proposition 3.8, $X = \underline{\varepsilon}(X) = \bar{\varepsilon}(X)$ ($X = \bar{\varepsilon}(X) = \underline{\varepsilon}(X)$). □

Definition 3.11. Let (U, ε) be Pawlak's approximation space and $X \subseteq U$. The ε -boundary of X is

$$B_{\varepsilon}(X) = \bar{\varepsilon}(X) \setminus \underline{\varepsilon}(X).$$

X is ε -crisp, if $B_{\varepsilon}(X) = \emptyset$, otherwise X is ε -rough.

Proposition 3.12 Let (U, ε) be Pawlak's approximation space and $X \subseteq U$.

1. X is ε -crisp if and only if X is ε -definable.
2. X is ε -rough if and only if X is ε -undefinable.

Proof. (1) (\Rightarrow) X is ε -crisp $\Leftrightarrow B_{\varepsilon}(X) = \bar{\varepsilon}(X) \setminus \underline{\varepsilon}(X) = \emptyset \Leftrightarrow \bar{\varepsilon}(X) \subseteq \underline{\varepsilon}(X)$. Proposition 3.9 implies $\underline{\varepsilon}(X) \subseteq \bar{\varepsilon}(X)$, and so $\underline{\varepsilon}(X) = \bar{\varepsilon}(X)$. According to Proposition 3.8, $\underline{\varepsilon}(X) = \bar{\varepsilon}(X) \Leftrightarrow X \in \mathfrak{D}_{U/\varepsilon}$.

(\Leftarrow) Since $X \in \mathfrak{D}_{U/\varepsilon} \Leftrightarrow \underline{\varepsilon}(X) = \bar{\varepsilon}(X)$, so $B_{\varepsilon}(X) = \bar{\varepsilon}(X) \setminus \underline{\varepsilon}(X) = \emptyset$ trivially satisfies.

- (2) It is the contrapositive version of (1). □

As a consequence of Proposition 3.12, the notions ' ε -crisp' and ' ε -definable' are synonymous to each other, and so are ' ε -rough' and ' ε -undefinable'.

4. Partial Approximation of Sets

Let U be any nonempty set called the *universe of discourse*.

Definition 4.1. Let $\mathfrak{B} \subseteq 2^U$ be a nonempty family of nonempty subsets of U called the *base system*. Its elements are the \mathfrak{B} -sets.

The family of sets $\mathfrak{D} \subseteq 2^U$ is \mathfrak{B} -definable, if its elements are \mathfrak{B} -sets, otherwise \mathfrak{D} is \mathfrak{B} -undefinable.

A nonempty subset $X \in 2^U$ is \mathfrak{B} -definable, if there exists a \mathfrak{B} -definable family of sets \mathfrak{D} such that $X = \bigcup \mathfrak{D}$, otherwise X is \mathfrak{B} -undefinable. The empty set is considered to be a \mathfrak{B} -definable set.

Let $\mathfrak{D}_{\mathfrak{B}}$ denote the family of \mathfrak{B} -definable sets of U .

Definition 4.2. Let $\mathfrak{B} \subseteq 2^U$ be a base system and X be any subset of U .

The *weak lower \mathfrak{B} -approximation* of X is

$$\mathfrak{C}_{\mathfrak{B}}^b(X) = \bigcup \{Y \mid Y \in \mathfrak{B}, Y \subseteq X\}, \quad (4.1)$$

and the *weak upper \mathfrak{B} -approximation* of X is

$$\mathfrak{C}_{\mathfrak{B}}^{\sharp}(X) = \bigcup \{Y \mid Y \in \mathfrak{B}, Y \cap X \neq \emptyset\}. \quad (4.2)$$

Clearly, $\mathfrak{C}_{\mathfrak{B}}^b(X), \mathfrak{C}_{\mathfrak{B}}^{\sharp}(X) \in \mathfrak{D}_{\mathfrak{B}}$, and the maps $\mathfrak{C}_{\mathfrak{B}}^b, \mathfrak{C}_{\mathfrak{B}}^{\sharp}$ are total, onto, and, in general, many-to-one. Furthermore, both of them are monotone.

Proposition 4.3 ([3], Theorem 17) Let the fixed base system $\mathfrak{B} \subseteq 2^U$ be given.

1. $\forall X \in 2^U (\mathfrak{C}_{\mathfrak{B}}^b(X) \subseteq \mathfrak{C}_{\mathfrak{B}}^{\sharp}(X))$.
2. $\forall X \in 2^U (\mathfrak{C}_{\mathfrak{B}}^b(X) \subseteq X)$ —that is, $\mathfrak{C}_{\mathfrak{B}}^b$ is contractive.
3. $\forall X \in 2^U (X \subseteq \mathfrak{C}_{\mathfrak{B}}^{\sharp}(X))$ if and only if $\bigcup \mathfrak{B} = U$ —that is, $\mathfrak{C}_{\mathfrak{B}}^{\sharp}$ is extensive if and only if \mathfrak{B} covers the universe.

Proposition 4.4 ([3], Theorem 19) Let $\mathfrak{B} \subseteq 2^U$ be a base system. Then

1. $X \in \mathfrak{D}_{\mathfrak{B}}$ if and only if $\mathfrak{C}_{\mathfrak{B}}^b(X) = X$.
2. $X \notin \mathfrak{D}_{\mathfrak{B}}$ if and only if $\mathfrak{C}_{\mathfrak{B}}^b(X) \neq X$.

Unlike Pawlak's approximation spaces (cf. Proposition 3.8), by Proposition 4.4, the \mathfrak{B} -definable property is generally not equivalent to $\mathfrak{C}_{\mathfrak{B}}^b(X) = \mathfrak{C}_{\mathfrak{B}}^{\sharp}(X)$.

Definition 4.5. Let $\mathfrak{B} \subseteq 2^U$ be a base system and X be any subset of U .

The \mathfrak{B} -boundary of X is

$$\mathfrak{N}_{\mathfrak{B}}(X) = \mathfrak{C}_{\mathfrak{B}}^{\sharp}(X) \setminus \mathfrak{C}_{\mathfrak{B}}^b(X).$$

X is \mathfrak{B} -approximatable if $X \subseteq \mathfrak{C}_{\mathfrak{B}}^{\sharp}(X)$, otherwise it is said that X has a \mathfrak{B} -approximation gap.

Provided that $X \in 2^U$ is \mathfrak{B} -approximatable, X is \mathfrak{B} -crisp, if $\mathfrak{N}_{\mathfrak{B}}(X) = \emptyset$, otherwise is \mathfrak{B} -rough.

In general, $\mathfrak{N}_{\mathfrak{B}}(X) \notin \mathfrak{D}_{\mathfrak{B}}$, i.e., \mathfrak{B} -boundaries are usually \mathfrak{B} -undefinable.

A \mathfrak{B} -approximation gap calls our attention that the available knowledge about the system encoded in \mathfrak{B} is not enough to approximate X . However, it may be natural or not.

According to Proposition 4.4, point (1), X is \mathfrak{B} -definable if and only if $X = \mathfrak{C}_{\mathfrak{B}}^b(X)$. If $X = \mathfrak{C}_{\mathfrak{B}}^b(X)$, then $X = \mathfrak{C}_{\mathfrak{B}}^b(X) \subseteq \mathfrak{C}_{\mathfrak{B}}^{\sharp}(X)$. However, it can easily be seen that $X = \mathfrak{C}_{\mathfrak{B}}^b(X)$ generally does not imply $X = \mathfrak{C}_{\mathfrak{B}}^{\sharp}(X)$. Hence, the notion ' \mathfrak{B} -definable' does not imply the notion ' \mathfrak{B} -crisp'. Thus, unlike Pawlak's approximation spaces (cf. Proposition 3.12), the notions ' \mathfrak{B} -crisp' and ' \mathfrak{B} -definable' are not synonymous to each other.

Possible interpretations of lower and upper \mathfrak{B} -approximations are the following [16, 18]:

- $\mathfrak{C}_{\mathfrak{B}}^b(X)$ is the set of all elements in U which can be *certainly* classified in a way that they belong to X with respect to \mathfrak{B} (\mathfrak{B} -positive region of X).
- $\mathfrak{C}_{\mathfrak{B}}^{\sharp}(X)$ is the set of all elements in U which can be *possibly* classified in a way that they belong to X with respect to \mathfrak{B} .
- $U \setminus \mathfrak{C}_{\mathfrak{B}}^{\sharp}(X)$ is the set of all elements in U which can be *certainly* classified in a way that they *do not belong* to X with respect to \mathfrak{B} (\mathfrak{B} -negative region of X).
- The elements in $\mathfrak{C}_{\mathfrak{B}}^{\sharp}(X) \setminus \mathfrak{C}_{\mathfrak{B}}^b(X)$ are *abstained* because they cannot be uniquely classified either as belonging to X or as not belonging to X with respect to \mathfrak{B} (\mathfrak{B} -borderline region of X).

Notice that if $\bigcup \mathfrak{B} \neq U$, then $\forall X \subseteq U \setminus \bigcup \mathfrak{B} \forall B \in \mathfrak{B} (X \cap B = \emptyset)$. Consequently, for all these subsets $\mathfrak{C}_{\mathfrak{B}}^{\sharp}(X) = \bigcup \emptyset = \emptyset$, i.e., the empty set is the weak upper \mathfrak{B} -approximation of certain nonempty subsets of U .

Such cases may be excluded by a partial map, a so-called *strong upper \mathfrak{B} -approximation*. For more details, see [2, 3, 4, 5].

5. Conclusions and Future Work

In this paper we have presented a generalization of the rough set theory. Most notions of Pawlak's classical approximation spaces constitute compound ones which, however, split in two or more parts in our approach. This new approach helps us to understand the state of the compound nature of these notions and to specify their constituents.

The next most important task is to work out a partial approximative information system model by analogy with Pawlak's one [16] on which practical implementations of our theoretical model can be built up.

Finally, it must be mentioned that both Pawlak's rough set theory and our approach can extensively be applied in medical informatics, see, e.g., [7, 8, 9, 14].

References

- [1] BONIKOWSKI, Z., BRYNIARSKI, E., AND WYBRANIEC-SKARDOWSKA, U. Extensions and intentions in the rough set theory. *Information Sciences* 107, 1-4 (1998), 149-167.
- [2] CSAJBÓK, Z. Partial approximative set theory. In *Programs, Proofs, Processes, 6th Conference on Computability in Europe (CiE 2010), Ponta Delgada (Azores), Portugal, June 30 - July 4, 2010, Abstract and Handout Booklet (2010)*, F. Ferreira, H. Guerra, E. Mayordomo, and J. Rasga, Eds., Centre for Applied Mathematics and Information Technology, Dept. of Mathematics, University of Azores, pp. 113-122.
- [3] CSAJBÓK, Z. Partial approximative set theory: A generalization of the rough set theory. In *Proceedings of the International Conference of Soft Computing and Pattern Recognition (SoCPaR 2010), December 7-10, 2010, Cergy Pontoise/Paris, France (2010)*, T. Martin, A.K. Muda, A. Abraham, H. Prade, A. Laurent, D. Laurent, and V. Sans, Eds., IEEE, pp. 51-56. DOI:10.1109/SOCPAR.2010.5686424. Also published in IEEE Xplore Digital Library.
- [4] CSAJBÓK, Z. A security model for personal information security management based on partial approximative set theory. In *Proceedings of the International Multiconference on Computer Science and Information Technology (IMCSIT 2010), October 18-20, 2010., Wista, Poland (Katowice, Poland - Los Alamitos, USA, 2010)*, M. Ganzha and M. Paprzycki, Eds., vol. 5, Polskie Towarzystwo Informatyczne - IEEE Computer Society Press, pp. 839-845. Also published in IEEE Xplore Digital Library.
- [5] CSAJBÓK, Z. Partial approximative set theory: A view from Galois connections. In *Proceedings of the 8th International Conference on Applied Informatics (ICAI 2010), January 27-30, 2010, Eger, Hungary (in press)*, Eszterházy Károly College, Eger.
- [6] CSAJBÓK, Z., AND MIHÁLYDEÁK, T. A general tool-based approximation framework based on partial approximation of sets. In *13th International Conference on Rough Sets, Fuzzy Sets and Granular Computing (RSFDGrC-2011), June 25-27, 2011, Moscow, Russia (2011)*, S.O. Kuznetsov et al., Eds., vol. 6743, Lecture Notes in Computer Science - Lecture Notes in Artificial Intelligence, Springer-Verlag, Berlin Heidelberg, pp. 52-59.
- [7] HASSANIEN, A. E., ABRAHAM, A., PETERS, J. F., AND SCHAEFER, G. Rough sets in medical informatics applications. In *Applications of Soft Computing: From Theory to Praxis (Berlin, Heidelberg 2009)*, J. Mehnen, M. K. A. Saad, and A. Tiwari, Eds., vol. 58 of *Advances in Intelligent and Soft Computing (AISC)*, Springer-Verlag, pp. 23-30.
- [8] HASSANIEN, A. E., ABRAHAM, A., PETERS, J. F., SCHAEFER, G., AND HENRY, C. Rough sets and near sets in medical imaging: a review. *IEEE Trans. on Information Technology in Biomedicine* 13, 6 (2009), 955-968.
- [9] HIRANO, S., AND TSUMOTO, S. Rough representation of a region of interest in medical images. *International Journal of Approximate Reasoning* 40, 1-2 (2005), 23-34.
- [10] JÄRVINEN, J. Lattice theory for rough sets. In *Transactions on Rough Sets VI: Commemorating the Life and Work of Zdzisław Pawlak, Part I*, J. F. Peters, A. Skowron, I. Düntsch, J. W. Grzymała-Busse, E. Orlowska, and L. Polkowski, Eds., vol. 4374 of *Lecture Notes in Computer Science*. Springer-Verlag, Berlin, Heidelberg, 2007, pp. 400-498.
- [11] JÄRVINEN, J., RADELECZKI, S., AND VERES, L. Rough sets determined by quasiorders. *Order* 26, 4 (2009), 337-355.
- [12] KEEFE, R. *Theories of Vagueness*. Cambridge Studies in Philosophy. Cambridge University Press, Cambridge, UK, 2000.
- [13] MIHÁLYDEÁK, T. On tarskian models of general type-theoretical languages. In *Proceedings of the 7th Panhellenic Logic Symposium (Patras, 2009)*, C. Drossos, P. Peppas, and C. Tsinakis, Eds., Patras University Press, pp. 127-131.
- [14] ØHRN, E., AND ROWL, T. Rough sets: A knowledge discovery technique for multifactorial medical outcomes. *American Journal of Physical Medicine and Rehabilitation* 79 (2000), 100-108.
- [15] PAWLAK, Z. Rough sets. *International Journal of Information and Computer Science* 11, 5 (1982), 341-356.
- [16] PAWLAK, Z. *Rough Sets: Theoretical Aspects of Reasoning about Data*. Kluwer Academic Publishers, Dordrecht, 1991.
- [17] PAWLAK, Z., POLKOWSKI, L., AND SKOWRON, A. Rough sets: An approach to vagueness. In *Encyclopedia of Database Technologies and Applications*, L. C. Rivero, J. Doorn, and V. Ferraggine, Eds. Idea Group Inc., Hershey, PA, 2005, pp. 575-580.

- [18] PAWLAK, Z., AND SKOWRON, A. Rudiments of rough sets. *Information Sciences* 177, 1 (2007), 3–27.
- [19] RUSSELL, B. Vagueness. *Australasian Journal of Philosophy and Psychology* 1 (1923), 84–92.
- [20] SKOWRON, A. Vague concepts: A rough-set approach. In *Current Issues in Data and Knowledge Engineering: Proceedings of EUROFUSE 2004, The Eighth Meeting of the EURO Working Group on Fuzzy Sets, Workshop on Data and Knowledge Engineering, September 22-25, 2004, Warszawa, Poland*, B. De Baets, R. De Caluwe, G. De Tré, J. Fodor, J. Kacprzyk, and S. Zadrożny, Eds. Akademicka Oficyna Wydawnicza EXIT, Warszawa, 2004, pp. 480–493.
- [21] XU, F., YAO, Y., AND MIAO, D. Rough set approximations in formal concept analysis and knowledge spaces. In *ISMIS (2008)*, A. An, S. Matwin, Z. W. Ras, and D. Slezak, Eds., vol. 4994 of *Lecture Notes in Computer Science*, Springer, pp. 319–328.
- [22] YAO, Y., AND CHEN, Y. Rough set approximations in formal concept analysis. In *Transactions on Rough Sets V (2006)*, J. F. Peters and A. Skowron, Eds., vol. 4100 of *Lecture Notes in Computer Science*, Springer, pp. 285–305.
- [23] YAO, Y. Y. Two views of the theory of rough sets in finite universes. *International Journal of Approximation Reasoning* 15, 4 (1996), 291–317.
- [24] YAO, Y. Y. Constructive and algebraic methods of the theory of rough sets. *Information Sciences* 109, 1–4 (1998), 21–47.
- [25] YAO, Y. Y. On generalizing rough set theory. In *Proceedings of the 9th International Conference Rough Sets, Fuzzy Sets, Data Mining, and Granular Computing (RSFDGrC 2003)* (2003), LNAI 2639, Springer-Verlag, pp. 44–51.
- [26] ZHU, W. Topological approaches to covering rough sets. *Information Sciences* 177, 6 (2007), 1499–1508.
- [27] ZHU, W. Relationship between generalized rough sets based on binary relation and covering. *Inf. Sci.* 179, 3 (2009), 210–225.

... • ...

Zoltán Csajbók University of Debrecen, Faculty of Health, Nyíregyháza, 4400, Sóstói út 2-4.

Szerzőink

Balatoni Ildikó - a DE OEC Minőségügyi Központjának vezetője, 2006-tól stratégiai igazgatóhelyettese. A SOTE-n szerzett általános orvosi diplomát, majd az ELTE-n jogi szakokleveles orvos végzettséget. 2010. óta a Debreceni Egyetem Esélyegyenlőségi Bizottságának elnöke.

Bánki M. Csaba - orvos, pszichiáter, farmakológus, karunk részállású egyetemi tanára. A „Sántha Kálmán” Szakkórház főorvosa. A Magyar Tudományos Akadémia doktora minősítéssel rendelkezik.

Csajbók Zoltán: karunk főállású oktatója. A debreceni Kossuth Lajos Tudományegyetem Természettudományi Karán szerzett okleveles matematikus és matematika szakos középiskolai tanár diplomát. Alapító tagja az ISACA Magyarországi Tagszervezetének, majd 1994 és 2000 között elnökségi tagja. 2011-ben az International Rough Set Society tagjává választották. Jelenleg Ph.D. doktorjelölt, és doktori disszertációján dolgozik.

Diószeghy Péter: A Jósa András Oktatókórház Nonprofit Kft. Neurológiai Osztályának osztályvezető főorvosa. Neurológus, pszichiáter, klinikai neurofiziológiai szakvizsgával rendelkezik. Az orvostudomány kandidátusa, habilitációval rendelkezik.

Ferenczi Gábor: karunk főállású oktatója. Szociális munkás képesítést nyert a DOTE Egészségügyi Főiskolán, majd a KLTE-n szerzett szociálpolitikusi diplomát. Jelenleg a Debreceni Református Hittudományi Egyetem doktori iskoljának hallgatója.

Gaál Zsolt: A Jósa András Oktatókórház Nonprofit Kft. IV. Belgyógyászatának osztályvezető főorvosa. Belgyógyász, endokrinológus, klinikofarmakológus, foglalkozás-egészségügyi szakvizsgával, diabetológus minősítéssel rendelkezik.

Husztai Éva: karunk főállású oktatója, a Debreceni Egyetem Egészségügyi Főiskolai Karon szerzett általános szociális munkás végzettséget, majd szociológusi diplomát az ELTE-n. Az ELTE Társadalomtudományi Kar Szociológia Doktori Iskola, Szociológia program doktorjelöltje. A disszertáció témája: A társas kapcsolatok hálózata.