

# Renal control of disease tolerance to malaria

Susana Ramos<sup>a</sup>, Ana Rita Carlos<sup>a,1</sup>, Balamurugan Sundaram<sup>a,1,2</sup>, Viktoria Jeney<sup>b,1</sup>, Ana Ribeiro<sup>a,3</sup>, Raffaella Gozzelino<sup>a,4</sup>, Claudia Bank<sup>a</sup>, Erida Gjini<sup>a</sup>, Faouzi Braza<sup>a</sup>, Rui Martins<sup>a</sup>, Temitope Wilson Ademolue<sup>a</sup>, Birte Blankenhaus<sup>a,5</sup>, Zélia Gouveia<sup>a,6</sup>, Pedro Faisca<sup>a</sup>, Damian Trujillo<sup>c</sup>, Sílvia Cardoso<sup>a</sup>, Sofia Rebelo<sup>a</sup>, Laura del Barrio<sup>a</sup>, Abolfazl Zarjou<sup>d</sup>, Subhashini Bolisetty<sup>d</sup>, Anupam Agarwal<sup>d,e</sup>, and Miguel P. Soares<sup>a,7</sup>

<sup>a</sup>Instituto Gulbenkian de Ciência, 2780-156 Oeiras, Portugal; <sup>b</sup>Faculty of Medicine, Department of Internal Medicine, University of Debrecen, H-4032, Debrecen, Hungary; <sup>c</sup>Department of Microbiology and Immunology, Stanford University, Stanford, CA 94305-5124; <sup>d</sup>Department of Medicine, Division of Nephrology, University of Alabama, Birmingham, AL 35294; and <sup>e</sup>Birmingham Veterans Administration Medical Center, Birmingham, AL 35294

Edited by Peter Agre, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, and approved February 8, 2019 (received for review January 3, 2019)

**Malaria, the disease caused by *Plasmodium* spp. infection, remains a major global cause of morbidity and mortality. Host protection from malaria relies on immune-driven resistance mechanisms that kill *Plasmodium*. However, these mechanisms are not sufficient per se to avoid the development of severe forms of disease. This is accomplished instead via the establishment of disease tolerance to malaria, a defense strategy that does not target *Plasmodium* directly. Here we demonstrate that the establishment of disease tolerance to malaria relies on a tissue damage-control mechanism that operates specifically in renal proximal tubule epithelial cells (RPTEC). This protective response relies on the induction of heme oxygenase-1 (*HMOX1*; HO-1) and ferritin H chain (*FTH*) via a mechanism that involves the transcription-factor nuclear-factor E2-related factor-2 (*NRF2*). As it accumulates in plasma and urine during the blood stage of *Plasmodium* infection, labile heme is detoxified in RPTEC by HO-1 and FTH, preventing the development of acute kidney injury, a clinical hallmark of severe malaria.**

infection | malaria | disease tolerance | heme | kidney

**D**isease tolerance is an evolutionarily conserved defense strategy against infection, first described as a central component of plant immunity (1). Over the past decade it became apparent that this defense strategy is also operational in animals, including mammals where it confers protection against malaria (2, 3).

The blood stage of *Plasmodium* spp. infection is characterized by the invasion of host red blood cells (RBC), in which this protozoan parasite proliferates extensively, consuming up to 60–80% of the RBC hemoglobin (HB) content (4). *Plasmodium* spp. do not express a *HMOX1* ortholog gene (5) and cannot catalyze the extraction of Fe from heme, acquiring Fe via heme auto-oxidation while also polymerizing labile heme into redox-inert hemozoin and avoiding its cytolytic effects (6).

Once the physical integrity of infected RBC becomes compromised, the remaining RBC HB content is released into plasma, where extracellular  $\alpha\beta_2$  HB tetramers disassemble into  $\alpha\beta$  dimers that undergo auto-oxidation, eventually releasing their noncovalently bound heme (7). As it accumulates in plasma, labile heme is loosely bound to plasma acceptor proteins, macromolecules, or low-molecular-weight ligands that fail, however, to control its redox activity (8). A fraction of the labile heme in plasma becomes bioavailable, acting in a pathogenic manner and compromising the establishment of disease tolerance to malaria (2, 7, 9).

Heme accumulation in plasma and urine of malaria patients is associated with the development of acute kidney injury (AKI), a clinical hallmark of severe malaria (10–12). Similarly, heme accumulation in plasma, as a consequence of rhabdomyolysis, is also associated with the development of AKI (13). While heme participates in the pathogenesis of AKI associated with rhabdomyolysis, whether this is the case for severe malaria has not been established.

We have previously shown that heme detoxification by the stress-responsive enzyme HO-1 is a limiting factor in the establishment of disease tolerance to malaria (2, 7). In a similar manner, heme detoxification by HO-1 also prevents the development of AKI following rhabdomyolysis (13). This protective effect requires that the Fe extracted from heme is neutralized by

the ferroxidase active FTH component of the ferritin complex (14), establishing disease tolerance to malaria (9) and preventing development of AKI following rhabdomyolysis (14). Here we asked whether heme catabolism by HO-1 and Fe sequestration by FTH act locally in the kidney to prevent the development of AKI and establish disease tolerance to malaria.

## Results

Malaria is associated with HO-1 induction in renal proximal tubule epithelial cells (RPTEC). Consistent with heme accumulation in plasma and urine of individuals developing severe forms of malaria (9, 15), *Plasmodium chabaudi chabaudi* (Pcc)-infected C57BL/6 mice accumulated heme in plasma and urine (Fig. 1A). Approximately 15% of the heme in plasma and 6% in urine were bioavailable (Fig. 1B), as assessed using a cellular-based heme-reporter assay (SI Appendix, Fig. S1 A and B). Heme and labile heme concentration in urine was higher than in plasma, suggesting that labile heme is actively excreted into urine.

## Significance

**Malaria, the disease caused by *Plasmodium* spp. infection, remains a major global cause of morbidity and mortality, claiming the lives of over  $\sim 4.5 \times 10^5$  individuals per year. Paradoxically, however, up to 98% of infected individuals survive the infection, establishing disease tolerance to malaria. We found that this host defense strategy, which does not target *Plasmodium* directly, relies on the capacity of renal proximal tubule epithelial cells to detoxify labile heme, a pathologic by-product of hemolysis that accumulates in plasma and urine during the blood stage of infection. This defense strategy prevents the onset of acute kidney injury, a clinical hallmark of severe malaria.**

Author contributions: S. Ramos, A.R.C., V.J., R.G., and M.P.S. designed research; S. Ramos, A.R.C., B.S., V.J., A.R., R.G., C.B., E.G., F.B., R.M., T.W.A., B.B., Z.G., P.F., D.T., S.C., S. Rebelo, and L.d.B. performed research; A.Z., S.B., and A.A. contributed new reagents/analytic tools; S. Ramos, A.R.C., B.S., V.J., A.R., R.G., C.B., E.G., F.B., R.M., T.W.A., B.B., Z.G., P.F., D.T., S.C., S. Rebelo, and M.P.S. analyzed data; and S. Ramos and M.P.S. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

This open access article is distributed under Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND).

<sup>1</sup>A.R.C., B.S., and V.J. contributed equally to this work.

<sup>2</sup>Present address: Institute of Molecular Medicine, Department of Molecular Medicine II, Medical Faculty, Heinrich-Heine-University, 40225 Dusseldorf, Germany.

<sup>3</sup>Present address: Department of Animal and Avian Sciences, University of Maryland at College Park, MD 20742-2311.


<sup>4</sup>Present address: Chronic Diseases Research Center, NOVA Medical School, 1150-082 Lisbon, Portugal.

<sup>5</sup>Present address: Instituto de Medicina Molecular, Faculty of Medicine, University of Lisbon, 1649-028 Lisbon, Portugal.

<sup>6</sup>Present address: Institute Curie, UMR144, 75000 Paris, France.

<sup>7</sup>To whom correspondence should be addressed. Email: mpssoares@igc.gulbenkian.pt.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1822024116/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1822024116/-DCSupplemental).



**Fig. 1.** HO-1 expression is induced in RPTEC during *Plasmodium* infection. (A) Heme and (B) bioavailable heme concentration (mean  $\pm$  SD) in plasma and urine of C57BL/6 mice, not infected (NI;  $n = 7$ ) or 7 d after *Pcc* infection ( $n = 8$ ). Data are from one experiment. (C) *Hmox1* normalized to *Arbp0* mRNA (mean  $\pm$  SD) in brain (B), liver (Li), spleen (S), kidney (K), muscle (M), lung (Lu), and heart (H) of C57BL/6 mice, not infected (NI;  $n = 3$ ) or 7 d after *Pcc* infection ( $n = 6$ ). Data are from one experiment. (D) HO-1 and Gapdh protein expression detected by Western blot in the brain (B), liver (Li), spleen (S), kidney (K), muscle (M), lung (Lu), and heart (H) of C57BL/6 mice, not infected (NI) or 7 d after *Pcc* infection. Data are representative of four mice per group in one experiment. (E) Densitometry analysis (mean  $\pm$  SD) of proteins in the same experiment as D. #: protein levels below detection limit for the exposure time analyzed. (F) HO-1 and Gapdh protein detected by Western blot in total kidney (T), renal cortex (C), and renal medulla (M) of C57BL/6 mice, not infected (NI) or 7 d after *Pcc* infection. Data are representative of four mice per group in one experiment. (G) Kidney immunostaining in C57BL/6 mice, not infected (NI) or 7 d after *Pcc* infection. Gamma glutamyl transferase 1 (Ggt1; red) was used as a RPTEC marker. Image is representative of three mice per group in one experiment. (Scale bar: 1,000  $\mu$ m.) (H) Kidney immunostaining, as in G. DAPI (blue) was used to counterstain DNA. Arrowheads highlight HO-1 (cyan) in Ggt1<sup>+</sup> RPTEC. Images are representative of five random fields from four to six mice per group in one experiment. (Scale bar: 50  $\mu$ m.) *P* values in A and B, comparing log-transformed heme concentration values, were determined using two-way ANOVA with Tukey's multiple comparisons tests, and in C and E using Mann–Whitney *U* test. NS: not significant ( $P > 0.05$ ); \* $P < 0.05$ ; \*\*\* $P < 0.001$ .

Consistent with our previous findings (9, 16), *Hmox1* mRNA (Fig. 1C) and HO-1 protein (Fig. 1D and E) were induced in the liver of *Pcc*-infected mice. *Hmox1* mRNA and HO-1 protein were also induced in other organs, including in the kidneys (Fig. 1C–E and *SI Appendix*, Fig. S2A and B). Expression of HO-1 in the kidney was induced in the cortex and in the medulla (Fig. 1F and G and *SI Appendix*, Fig. S2A), predominantly in RPTEC (Fig. 1H and *SI Appendix*, Fig. S2B). This suggests that, as it accumulates in plasma and urine during the blood stage of *Plasmodium* infection, labile heme is up taken by RPTEC, where it is catabolized by HO-1.

HO-1 expression in RPTEC is essential to establish disease tolerance to malaria. To determine whether heme catabolism in RPTEC is involved in the establishment of disease tolerance to malaria, we generated *Hmox1*<sup>PepckΔ/Δ</sup> mice, in which *Hmox1* is deleted specifically in RPTEC (17) (*SI Appendix*, Fig. S2C). *Pcc*-infected *Hmox1*<sup>PepckΔ/Δ</sup> mice succumbed to *Pcc* infection, compared with control *Hmox1*<sup>fl/fl</sup> mice that survived and resolved the infection (Fig. 2A). Mortality of *Hmox1*<sup>PepckΔ/Δ</sup> mice was not associated with changes in parasitemia, compared with control *Hmox1*<sup>fl/fl</sup> mice (Fig. 2A). This suggests that heme catabolism by HO-1 in RPTEC is essential to establish disease tolerance to malaria.

Lethality of *Pcc*-infected *Hmox1*<sup>PepckΔ/Δ</sup> mice was associated with the development of AKI, as illustrated by the accumulation

of Lipocalin 2 (LCN2) in urine and blood urea nitrogen (BUN) (Fig. 2B). AKI in *Pcc*-infected *Hmox1*<sup>PepckΔ/Δ</sup> mice was characterized histologically by widespread acute tubular necrosis and HB casts affecting up to 15% of the renal cortex, compared with sparse tubular single-cell death and HB casts in control *Pcc*-infected *Hmox1*<sup>fl/fl</sup> mice (Fig. 2C). This suggests that during the blood stage of *Plasmodium* infection, the induction of heme catabolism by HO-1 in RPTEC is essential to prevent the development of AKI, a clinical hallmark of severe malaria (10–12).

The extent of liver damage, another clinical hallmark of severe malaria captured in *Pcc*-infected mice (9, 16), was similar in infected *Hmox1*<sup>PepckΔ/Δ</sup> vs. *Hmox1*<sup>fl/fl</sup> mice, as assessed serologically (*SI Appendix*, Fig. S2D). This suggests that heme catabolism in RPTEC does not counter the development of liver damage.

The complete blood count profile of *Pcc*-infected *Hmox1*<sup>PepckΔ/Δ</sup> and *Hmox1*<sup>fl/fl</sup> mice was indistinguishable (*SI Appendix*, Table S1), suggesting that HO-1 expression in RPTEC does not impact on the development of hemolytic anemia during *Plasmodium* infection. Blood neutrophilia was also similar in *Pcc*-infected *Hmox1*<sup>PepckΔ/Δ</sup> vs. *Hmox1*<sup>fl/fl</sup> mice (*SI Appendix*, Table S1), suggesting that HO-1 expression in RPTEC does not impact on neutrophil activation.


Disease tolerance can be inferred from the analysis of “disease trajectories” established by the temporal relationship of specific host homeostatic variables versus pathogen load (18). The disease trajectories of *Pcc*-infected *Hmox1*<sup>PepckΔ/Δ</sup> mice were distinct from those of *Pcc*-infected control *Hmox1*<sup>fl/fl</sup> mice, when analyzing body temperature vs. body weight vs. pathogen load (Fig. 2D and E). Differences among genotypes were statistically significant as early as 7 d after infection, before the onset of lethality in *Hmox1*<sup>PepckΔ/Δ</sup> mice (Fig. 2D and E). This suggests that heme catabolism by HO-1 in RPTEC is required to maintain vital homeostatic parameters, such as body temperature and body weight, within a dynamic range compatible with survival from *Plasmodium* infection.

HO-1 expression in cell compartments other than RPTEC is not essential to establish disease tolerance to malaria. Ubiquitous deletion of *Hmox1* in *Hmox1*<sup>R26Δ/Δ</sup> mice (*SI Appendix*, Fig. S3A) recapitulated the deletion of *Hmox1* specifically in RPTEC, impairing survival to *Pcc* infection under pathogen loads similar to control *Hmox1*<sup>fl/fl</sup> mice (*SI Appendix*, Fig. S3B). The disease trajectories established by *Pcc*-infected were also distinct from those of *Pcc*-infected control *Hmox1*<sup>fl/fl</sup> mice (*SI Appendix*, Fig. S3C and D). We then asked whether *Hmox1* deletion in cell compartments—where heme catabolism by HO-1 would be expected to act in a protective manner—impaired disease tolerance to *Pcc* infection. However, *Hmox1* deletion specifically in myeloid cells of *Hmox1*<sup>LysMΔ/Δ</sup> mice (19), macrophages and dendritic cells of *Hmox1*<sup>Cd11cΔ/Δ</sup> mice, vascular endothelial and hematopoietic cells of *Hmox1*<sup>Tie2Δ/Δ</sup> mice, hepatocytes and hematopoietic cells of *Hmox1*<sup>Mx1Δ/Δ</sup> mice, or neuronal cells of *Hmox1*<sup>NestinΔ/Δ</sup> mice failed to compromise disease tolerance to *Pcc* infection (*SI Appendix*, Fig. S4A–F). This reinforces the notion that heme catabolism by HO-1 in RPTEC is a limiting factor in the establishment of disease tolerance to malaria.

We then questioned whether HO-1 expression in RPTEC is also essential to establish disease tolerance to other infectious hemolytic conditions and tested this using an experimental model of hemolytic *Escherichia coli* (clinical isolate strain CFT073) infection (20). The lethality of *E. coli* CFT073 infection was similar in *Hmox1*<sup>PepckΔ/Δ</sup> vs. control *Hmox1*<sup>fl/fl</sup> mice (*SI Appendix*, Fig. S5A), indicating that heme catabolism by HO-1 in RPTEC is not essential to establish disease tolerance to hemolytic bacterial infections.

Malaria is associated with the induction of FTH in RPTEC. The Fe released via heme catabolism by HO-1 induces post-transcriptionally the expression of FTH (21). Consistent with this notion and in keeping with our previous findings (9), Fth expression was induced in the liver of *Pcc*-infected C57BL/6 mice,





**Fig. 2.** HO-1 expression in RPTEC establishes disease tolerance to malaria. (A) Survival and percentage of iRBC (mean ± SD) in *Pcc*-infected *Hmxo1<sup>fl/fl</sup>* (n = 7) and *Hmxo1<sup>PepckΔ/Δ</sup>* (n = 12) mice. Data from four independent experiments with similar trend. (B) Lcn2 concentration in urine and BUN (mean ± SD) in noninfected (NI) and *Pcc*-infected *Hmxo1<sup>fl/fl</sup>* (NI: 5–11; *Pcc*: n = 7–12) and *Hmxo1<sup>PepckΔ/Δ</sup>* (NI: 5–11; *Pcc*: n = 7–12) mice. Data from two experiments with a similar trend. (C) Kidney H&E staining in noninfected and 7 d after *Pcc* infection. Representative of four to five mice per genotype in two experiments. (Scale bar: 50 μm.) Arrowhead indicates HB casts, and dashed lines outline tubular necrosis. (D) Individual disease trajectories of the same mice as in A. Circles represent median of disease trajectories, and corresponding numbers are days after infection (day 0). Portion of the disease trajectories in darker color indicates days before onset of mortality in *Hmxo1<sup>PepckΔ/Δ</sup>* mice. (E) The 3D plot of disease trajectories from the same mice as in D. Statistical differences between genotypes became significant ( $P = 0.008$ ; Mann–Whitney  $U$  test) on day 7 before onset of mortality in *Hmxo1<sup>PepckΔ/Δ</sup>* mice.  $P$  values in A were determined using the log-rank (Mantel–Cox) test and in B using two-way ANOVA with Tukey’s multiple comparison test. NS, not significant ( $P > 0.05$ ); \*\*\* $P < 0.001$ .

as assessed by Western blotting (Fig. 3A and B). This was also the case in the kidney (Fig. 3A and B), where we had previously failed to detect Fth induction in response to *Pcc* infection (9), most likely owing to technical reasons. Induction of Fth expression in the kidney occurred both in the cortex and the medulla (Fig. 3C and D and *SI Appendix*, Fig. S6A), predominantly in RPTEC (Fig. 3E and *SI Appendix*, Fig. S6B).

FTH expression in RPTEC is essential to establish disease tolerance to malaria. To assess whether FTH expression in RPTEC contributes to the establishment of disease tolerance to malaria, we generated *Fth<sup>PepckΔ/Δ</sup>* mice, in which *Fth* is deleted specifically in RPTEC (*SI Appendix*, Fig. S7A) (14). More than 75% of *Fth<sup>PepckΔ/Δ</sup>* mice succumbed to *Pcc* infection, as compared to 13.3% of control *Pcc*-infected *Fth<sup>fl/fl</sup>* mice (Fig. 4A). This was not associated with changes in pathogen load (Fig. 4A), suggesting that Fe storage by Fth in RPTEC is essential to establish disease tolerance to *Pcc* infection. Of note, *Hmxo1* expression was highly induced in the kidneys of *Pcc*-infected *Fth<sup>PepckΔ/Δ</sup>* vs. *Fth<sup>fl/fl</sup>* mice (*SI Appendix*, Fig. S7B), demonstrating that heme catabolism by HO-1 per se fails to establish disease tolerance to malaria, if not coupled to Fe storage by Fth. Moreover, when considering *Hmxo1* induction as a biomarker of cell stress, it becomes apparent that, in the absence of Fth, heme catabolism by HO-1 is deleterious in RPTEC.

*Pcc*-infected *Fth<sup>PepckΔ/Δ</sup>* mice developed AKI, as illustrated by Lcn2, BUN, and cystatin C accumulation in plasma (Fig. 4B and *SI Appendix*, Fig. S7C), compared with control *Fth<sup>fl/fl</sup>* mice. AKI in *Pcc*-infected *Fth<sup>PepckΔ/Δ</sup>* mice was associated with acute tubular necrosis and intraluminal HB casts, affecting up to 40% of the renal cortex, compared with control *Pcc*-infected *Fth<sup>fl/fl</sup>* mice that developed only discrete and rare tubular epithelial cell necrosis or HB casts in the renal cortex (Fig. 4C). Expression of *Lcn2* mRNA was also higher in the kidneys of *Pcc*-infected *Fth<sup>PepckΔ/Δ</sup>* vs. *Fth<sup>fl/fl</sup>* mice (*SI Appendix*, Fig. S7D).

The disease trajectories established by the temporal relationship of body temperature vs. body weight vs. pathogen load showed significant differences in *Pcc*-infected *Fth<sup>PepckΔ/Δ</sup>* vs. *Fth<sup>fl/fl</sup>* mice (Fig. 4D and E), including before the onset of lethality. This suggests that FTH expression in RPTEC is essential to maintaining vital homeostatic factors such as body temperature and body weight within a range compatible with host survival.


We have previously shown that composite *Fth* deletion in *Fth<sup>Mx1Δ/Δ</sup>* mice compromises disease tolerance to *Pcc* infection (9). This suggests that Fe storage by FTH, in cellular compartments other than RPTEC, may contribute to disease tolerance to malaria. However, *Fth* deletion in myeloid cells of *Fth<sup>LysMΔ/Δ</sup>* mice, hepatocytes of *Fth<sup>AlbΔ/Δ</sup>* mice, or vascular endothelial cells of *Fth<sup>Cdh5Δ/Δ</sup>* mice (*SI Appendix*, Fig. S8A–E) failed to compromise disease tolerance to *Pcc* infection. This suggests that deletion of *Fth* in the kidney (22) contributes critically to our previous observation that *Fth<sup>Mx1Δ/Δ</sup>* mice fail to establish disease tolerance to malaria (9).

FTH expression in RPTEC does not regulate local immune responses. We asked whether FTH expression in RPTEC modulates kidney immunopathology, as assessed by renal leukocyte infiltration. Neutrophils, inflammatory monocytes, and CD4<sup>+</sup> as well as CD8<sup>+</sup> T cells accumulated to a similar extent in the kidneys of *Pcc*-infected *Fth<sup>PepckΔ/Δ</sup>* vs. *Fth<sup>fl/fl</sup>* mice (*SI Appendix*, Fig. S9A–F). The same was true for T cell proliferation, as monitored by Ki67 expression (*SI Appendix*, Fig. S9G–J). This suggests that FTH expression in RPTEC does not impact on the extent of kidney leukocyte infiltration.

FTH expression by RPTEC is not essential to establish disease tolerance to hemolytic bacterial infections. Mortality from *E. coli* CFT073 infection in *Fth<sup>PepckΔ/Δ</sup>* was similar to that in *Fth<sup>fl/fl</sup>* mice (*SI Appendix*, Fig. S10A). This suggests that, similarly to HO-1, FTH expression in RPTEC is not required per se to provide a survival advantage against infection by this hemolytic bacterium.

FTH is cytoprotective in RPTEC. We reasoned that FTH might protect RPTEC from the cytotoxic effects of labile heme, as it accumulates in plasma and urine during the blood stage of *Plasmodium* infection (Fig. 1A). In keeping with previous findings in other parenchyma cells (16, 23, 24), heme enhanced the cytotoxic effect of hydrogen peroxide ( $H_2O_2$ ) on human RPTEC in vitro, compared with exposure to heme or  $H_2O_2$  alone (Fig. 5A). Transduction of human RPTEC in vitro with a recombinant adenovirus (Rec.Ad.) encoding FTH (Fig. 5B) was cytoprotective against heme plus  $H_2O_2$ , compared with control cells transduced with a Rec.Ad. encoding  $\beta$ -galactosidase (*LacZ*) (Fig. 5A). Human RPTEC were not protected against heme plus  $H_2O_2$  when transduced with a Rec.Ad. encoding a FTH mutant (FTH<sup>m</sup>) lacking ferroxidase activity (Fig. 5A and B) (9, 25), suggesting that the ferroxidase activity of FTH is required to counter the cytotoxic effects of labile heme and  $H_2O_2$  in human RPTEC.

We then tested whether the cytoprotective effect of FTH is mediated via a mechanism that prevents Fe from catalyzing the production of reactive oxygen species (ROS). Human RPTEC accumulated high levels of ROS when exposed in vitro to heme plus  $H_2O_2$ , compared with heme or  $H_2O_2$  alone (Fig. 5C). This is in keeping with previous observations that heme promotes ROS generation in parenchyma cells exposed to a variety of inflammatory agonists (23, 24). Transduction of human RPTEC with an FTH, but not an FTH<sup>m</sup> Rec.Ad., suppressed ROS accumulation, compared with control cells transduced with a *LacZ* Rec.Ad. (Fig. 5D and E). This further supports the interpretation that the ferroxidase activity



**Fig. 3.** FTH expression is induced in RPTEC during *Plasmodium* infection. (A) Fth and Gapdh protein expression in brain (B), liver (Li), spleen (S), kidney (K), muscle (M), lung (Lu), and heart (H) of C57BL/6 mice, noninfected (NI) or 7 d after *Pcc* infection. Western blot representative of nine mice, from two independent experiments with the same trend. (B) Densitometry analysis (mean  $\pm$  SD) of proteins shown in A.  $n = 9$  mice per group from two experiments. (C) Fth and Gapdh protein levels in total kidney (T), renal cortex (C), and renal medulla (M) of C57BL/6 mice, not infected (NI) or 7 d after *Pcc* infection. Data are representative of four mice per group from one experiment. (D) Kidney immunostaining in C57BL/6 mice, noninfected (NI) or 7 d after *Pcc* infection. Gamma glutamyl transferase 1 (Ggt1; red) was used as a RPTEC marker. Image is representative of three mice per group in one experiment. (Scale bar: 1,000  $\mu$ m.) (E) Kidney immunostaining, as in D. DAPI (blue) was used to counterstain DNA. Arrowheads highlight Fth (cyan) in Ggt1<sup>+</sup> RPTEC. Images are representative of five random fields from four to six mice per group in one experiment. (Scale bar: 50  $\mu$ m.)  $P$  values in B determined using Mann-Whitney  $U$  test. NS, not significant ( $P > 0.05$ ); \*\* $P < 0.01$ ; \*\*\* $P < 0.001$ .

of FTH is critical to limit the pro-oxidant and cytotoxic effects of intracellular labile Fe, generated via heme catabolism by HO-1 in RPTEC. Presumably, this explains why, in the absence of FTH, HO-1 fails to confer disease tolerance to malaria.

The transcription factor NRF2 controls the establishment of disease tolerance to malaria. Upon activation NRF2 can induce *Hmox1* (26) and *Fth* (27), and therefore we asked whether NRF2 is required to support the induction of these genes in the kidneys of *Pcc*-infected mice, preventing the development of AKI and establishing disease tolerance. Expression of NAD(P)H quinone dehydrogenase 1 (Nqo1), a prototypical NRF2-responsive gene (28), was induced in the kidneys of *Pcc*-infected wild type (*Nrf2*<sup>+/+</sup>) but not in *Pcc*-infected *Nrf2*-deficient (*Nrf2*<sup>-/-</sup>) mice (Fig. 6A). This suggests that NRF2 is activated in the kidneys in response to *Pcc* infection. Similar, although less pronounced, the expression of *Hmox1*, *Fth*, and Ferroportin-1 (Fpn1; solute carrier family 40 member 1; Slc40a1) was also induced in the kidney during *Pcc* infection in *Nrf2*<sup>+/+</sup> but to a lesser extent in *Nrf2*<sup>-/-</sup> mice (Fig. 6A). This suggests that NRF2 activation in response to *Plasmodium* infection supports the induction of HO-1, FTH, and FPN1 in the kidney.

About 80% of *Nrf2*<sup>-/-</sup> mice succumbed to *Pcc* infection, compared with 20% of control *Pcc*-infected *Nrf2*<sup>+/+</sup> mice (Fig. 6B), with similar pathogen loads (Fig. 6C). *Pcc*-infected *Nrf2*<sup>-/-</sup> mice developed AKI, as illustrated by a higher BUN (Fig. 6D), and extensive acute tubular necrosis (Fig. 6E), compared with *Nrf2*<sup>+/+</sup> mice. This suggests that NRF2 acts upstream of HO-1, FTH, and FPN1 to promote the establishment of disease tolerance to malaria.


Given the involvement of other transcription factors, such as the heat shock factor 1 (HSF1) (29) and hypoxia inducible factor 1 alpha (HIF1 $\alpha$ ) (30), in the regulation of *HMOX-1* expression, we addressed whether these transcription factors were also involved in the establishment of disease tolerance to *Pcc* infection. To test this hypothesis, we generated *Hsf1*<sup>R26 $\Delta$ /Δ</sup> and *Hif1 $\alpha$* <sup>R26 $\Delta$ /Δ</sup> mice, in which *Hsf1* or *Hif1 $\alpha$* , respectively, are globally deleted in an inducible manner (SI Appendix, Fig. S11 A and B). Lethality and parasitemias of *Pcc*-infected *Hsf1*<sup>R26 $\Delta$ /Δ</sup> and *Hif1 $\alpha$* <sup>R26 $\Delta$ /Δ</sup> mice were similar to

control *Hsf1*<sup>fl/fl</sup> and *Hif1 $\alpha$* <sup>fl/fl</sup> mice, respectively (SI Appendix, Fig. S11 C–F). This suggests that neither HSF1 nor HIF1 $\alpha$  are involved in the establishment of disease tolerance to *Pcc* infection.

## Discussion

Most of the Fe contained in the prosthetic heme groups of HB must be continuously recycled, redirected toward heme biosynthesis, and incorporated into nascent HB during erythropoiesis (31). Hemophagocytic macrophages are at a center stage in this Fe-recycling process, engulfing an estimated  $2 \times 10^6$  senescent RBC and processing  $2 \times 10^{15}$  heme molecules per second in healthy humans (31, 32). The pathologic outcomes associated with disruption of this Fe-recycling process during hemolytic conditions are countered by the recruitment of circulating monocytes that differentiate into tissue resident Fe-recycling hemophagocytic macrophages (33). When the recruitment and differentiation of these macrophages is impaired, hemolytic conditions become associated with the development of AKI (33). Given that AKI is a hallmark of severe malaria (10–12), it is reasonable to speculate that impairment of hemophagocytic macrophage function (34) contributes to the pathogenesis of malaria-associated AKI, but this remains to be formally established.

Presumably, defective hemophagocytic macrophage function associated with malaria (34) also favors the accumulation of damaged RBC in the circulation, leading to intravascular hemolysis



**Fig. 4.** FTH expression in RPTEC is essential to establish disease tolerance to malaria. (A) Survival and percentage of iRBC (mean  $\pm$  SD) of *Pcc*-infected *Fth*<sup>Pepck $\Delta$ /Δ</sup> ( $n = 23$ ) and *Fth*<sup>fl/fl</sup> ( $n = 30$ ) mice. Data are from six independent experiments with a similar trend. (B) Lcn2 plasma concentration and BUN (mean  $\pm$  SD) in noninfected and *Pcc*-infected *Fth*<sup>Pepck $\Delta$ /Δ</sup> (NI:  $n = 5$ –10; *Pcc*:  $n = 7$ ) and *Fth*<sup>fl/fl</sup> (NI:  $n = 5$ ; *Pcc*:  $n = 7$ ) mice, 7 d after infection. Data are from three experiments. (C) H&E staining of kidney sections from noninfected (NI) and *Pcc*-infected *Fth*<sup>Pepck $\Delta$ /Δ</sup> and *Fth*<sup>fl/fl</sup> mice, 9 d after infection. Arrowheads indicate HB casts, and dashed lines outline tubular necrosis. Images are representative of two to seven mice per genotype in four experiments. (Scale bar: 50  $\mu$ m.) (D) Individual disease trajectories of *Pcc*-infected *Fth*<sup>Pepck $\Delta$ /Δ</sup> ( $n = 17$ ) and *Fth*<sup>fl/fl</sup> ( $n = 10$ ) mice from three independent experiments with a similar trend. Circles represent median of disease trajectories, and corresponding numbers indicate days after infection (day 0). Portion of disease trajectories in darker color indicates days before onset of mortality in *Fth*<sup>Pepck $\Delta$ /Δ</sup> mice. (E) The 3D plot of disease trajectories from the same mice as in D. Statistical differences between genotypes became significant on day 6 after *Pcc* infection ( $P = 0.00007$ ; Mann-Whitney  $U$  test) before onset of mortality of *Fth*<sup>Pepck $\Delta$ /Δ</sup> mice.  $P$  values in A were determined using a log-rank (Mantel-Cox) test and in B using two-way ANOVA with Tukey's multiple comparison test. NS, not significant; \* $P < 0.05$ ; \*\* $P < 0.01$ ; \*\*\* $P < 0.001$ .





RPTEC is essential to prevent acute tubular necrosis underlying the development of AKI.

As suggested by previous studies (18), the establishment of disease tolerance to *Pcc* infection can be inferred from the analyses of disease trajectories (Fig. 2 *D* and *E* and *SI Appendix*, Figs. S3 and Fig. 4 *D* and *E*). Accordingly, *Hmox1* or *Fth* deletion in RPTEC gave rise to disease trajectories with a sharp decline of body temperature, occurring irrespective of pathogen load (Figs. 2 *D* and *E* and 4 *D* and *E*). One possible explanation for why the expression of these genes in the kidney controls body temperature likely relates to a cross-talk between Fe and glucose metabolism in which intracellular Fe accumulation inhibits endogenous glucose production via gluconeogenesis (25). Of note, kidney gluconeogenesis is essential to control of glycemia in response to different forms of stress (39), and glucose is a major source of energy controlling thermoregulation and the establishment of disease tolerance to *Plasmodium* infection (40). We speculate that expression of HO-1 and FTH in RPTEC might regulate kidney glucose production in a manner that impacts on thermoregulation and the establishment of disease tolerance to *Plasmodium* infection, which, however, remains to be established.

In conclusion, this study demonstrates the existence of a previously unsuspected tissue damage-control mechanism that operates specifically in the kidneys to establish disease tolerance to *Plasmodium* infection. We propose that targeting components of this defense mechanism may be of therapeutic value in the treatment of severe malaria without the selection of drug resistance in *Plasmodium* spp.

## Materials and Methods

**Plasmodium Infections.** Mice were bred and maintained under specific pathogen-free (SPF) conditions at the Instituto Gulbenkian de Ciência (IGC). Experimental protocols were approved by the Ethics Committee of the IGC

(A008.2010 and A009.2011) and the Portuguese National Entity (Direcção Geral de Alimentação e Veterinária; 008959 and 018071). Experimental procedures were performed according to the Portuguese (Decreto-Lei 113/2013) and European (Directive 2010/63/EU) legislations. Mice were infected by i.p. inoculation of blood isolated from mice infected with a *Pcc* AS strain [ $2 \times 10^6$  infected red blood cells (iRBC)]. Mice were monitored daily for parasitemia, weight, temperature, RBC number, and survival.

**Serology.** Mice were killed and plasma was obtained at the indicated time points after *Plasmodium* infection. BUN and aspartate aminotransferase (AST) were quantified using quantitative colorimetric determination kits (Quantichrom and EnzyChrom; Bioassay Systems) and Lcn2 and Cystatin C by ELISA, as detailed in *SI Appendix*.

**Adenoviral Transduction in Human RPTEC.** Human RPTEC (ScienCell Research Laboratories) were transduced (50% confluence) with recombinant adenovirus encoding FTH, ferroxidase-deleted FTH, and LacZ, as described (9, 25). Briefly, RPTEC were exposed to Rec.Ad. (10 pfu/cell, 6 h) and, within 48–72 h, exposed to hemin (5  $\mu$ M) and/or H<sub>2</sub>O<sub>2</sub> (50  $\mu$ M) and monitored for survival and ROS production, as detailed in *SI Appendix*.

For detailed materials and methods, see *SI Appendix*.

**ACKNOWLEDGMENTS.** This work was funded by Fundação para a Ciência e Tecnologia Grants PTDC/SAU-TOX/116627/2010, HMSP-ICT/0018/2011, and LISBOA-01-0145-FEDER-029411 (to M.P.S.); SFRH/BPD/101608/2014 (to A.R.C.); and SFRH/BD/51877/2012 (to A.R.). S. Rebelo and T.W.A. are supported by Fundação Calouste Gulbenkian Grant 217/BD/17; B.S., S.C., L.d.B., R.G., and S. Ramos by European Union 7th Framework Grant ERC-2011-AdG 294709 (to M.P.S.); V.J. by Hungarian National Research, Development, and Innovation Office Grant K116024; F.B. by Marie Skłodowska-Curie Research Fellowship 707998; R.M. by European Molecular Biology Organization (EMBO) Long-Term Fellowship ALTF290-2017; A.A. by NIH Grants R01 DK059600 and P30 DK079337; and S.B. by NIH Grant K01 DK103931.

- Medzhitov R, Schneider DS, Soares MP (2012) Disease tolerance as a defense strategy. *Science* 335:936–941.
- Ferreira A, et al. (2011) Sickle hemoglobin confers tolerance to Plasmodium infection. *Cell* 145:398–409.
- Råberg L, Sim D, Read AF (2007) Disentangling genetic variation for resistance and tolerance to infectious diseases in animals. *Science* 318:812–814.
- Francis SE, Sullivan DJ, Jr, Goldberg DE (1997) Hemoglobin metabolism in the malaria parasite *Plasmodium falciparum*. *Annu Rev Microbiol* 51:97–123.
- Sigala PA, Crowley JR, Hsieh S, Henderson JP, Goldberg DE (2012) Direct tests of enzymatic heme degradation by the malaria parasite *Plasmodium falciparum*. *J Biol Chem* 287:37793–37807.
- Orjih AU, Banyal HS, Chevli R, Fitch CD (1981) Hemin lyses malaria parasites. *Science* 214:667–669.
- Pamplona A, et al. (2007) Heme oxygenase-1 and carbon monoxide suppress the pathogenesis of experimental cerebral malaria. *Nat Med* 13:703–710.
- Gouveia Z, et al. (2017) Characterization of plasma labile heme in hemolytic conditions. *FEBS J* 284:3278–3301.
- Gozzelino R, et al. (2012) Metabolic adaptation to tissue iron overload confers tolerance to malaria. *Cell Host Microbe* 12:693–704.
- Marsh K, et al. (1995) Indicators of life-threatening malaria in African children. *N Engl J Med* 332:1399–1404.
- Sypniewska P, et al. (2017) Clinical and laboratory predictors of death in African children with features of severe malaria: A systematic review and meta-analysis. *BMC Med* 15:147.
- Cruz LAB, Barral-Netto M, Andrade BB (2018) Distinct inflammatory profile underlies pathological increases in creatinine levels associated with *Plasmodium vivax* malaria clinical severity. *PLoS Negl Trop Dis* 12:e0006306.
- Nath KA, et al. (1992) Induction of heme oxygenase is a rapid, protective response in rhabdomyolysis in the rat. *J Clin Invest* 90:267–270.
- Zarjou A, et al. (2013) Proximal tubule H-ferritin mediates iron trafficking in acute kidney injury. *J Clin Invest* 123:4423–4434.
- Dalko E, et al. (2015) Multifaceted role of heme during severe *Plasmodium falciparum* infections in India. *Infect Immun* 83:3793–3799.
- Seixas E, et al. (2009) Heme oxygenase-1 affords protection against noncerebral forms of severe malaria. *Proc Natl Acad Sci USA* 106:15837–15842.
- Higgins DF, et al. (2007) Hypoxia promotes fibrogenesis in vivo via HIF-1 stimulation of epithelial-to-mesenchymal transition. *J Clin Invest* 117:3810–3820.
- Schneider DS (2011) Tracing personalized health curves during infections. *PLoS Biol* 9:e1001158.
- Parada E, et al. (2013) The microglial  $\alpha 7$ -acetylcholine nicotinic receptor is a key element in promoting neuroprotection by inducing heme oxygenase-1 via nuclear factor erythroid-2-related factor 2. *Antioxid Redox Signal* 19:1135–1148.
- Martins R, et al. (2016) Heme drives hemolysis-induced susceptibility to infection via disruption of phagocyte functions. *Nat Immunol* 17:1361–1372.
- Eisenstein RS, Garcia-Mayol D, Pettingell W, Munro HN (1991) Regulation of ferritin and heme oxygenase synthesis in rat fibroblasts by different forms of iron. *Proc Natl Acad Sci USA* 88:688–692.
- Darshan D, Vanoaica L, Richman L, Beermann F, Kühn LC (2009) Conditional deletion of ferritin H in mice induces loss of iron storage and liver damage. *Hepatology* 50:852–860.
- Balla G, et al. (1992) Ferritin: A cytoprotective antioxidant strategem of endothelium. *J Biol Chem* 267:18148–18153.
- Larsen R, et al. (2010) A central role for free heme in the pathogenesis of severe sepsis. *Sci Transl Med* 2:51ra71.
- Weis S, et al. (2017) Metabolic adaptation establishes disease tolerance to sepsis. *Cell* 169:1263–1275.e14.
- Alam J, et al. (1999) Nrf2, a Cap'n'Collar transcription factor, regulates induction of the heme oxygenase-1 gene. *J Biol Chem* 274:26071–26078.
- Pietsch EC, Chan JY, Torti FM, Torti SV (2003) Nrf2 mediates the induction of ferritin H in response to xenobiotics and cancer chemopreventive dithiolethiones. *J Biol Chem* 278:2361–2369.
- Suzuki T, Motohashi H, Yamamoto M (2013) Toward clinical application of the Keap1-Nrf2 pathway. *Trends Pharmacol Sci* 34:340–346.
- Alam J, Cook JL (2007) How many transcription factors does it take to turn on the heme oxygenase-1 gene? *Am J Respir Cell Mol Biol* 36:166–174.
- Lee PJ, et al. (1997) Hypoxia-inducible factor-1 mediates transcriptional activation of the heme oxygenase-1 gene in response to hypoxia. *J Biol Chem* 272:5375–5381.
- Muckenthaler MU, Rivella S, Hentze MW, Galy B (2017) A red carpet for iron metabolism. *Cell* 168:344–361.
- Soares MP, Hamza I (2016) Macrophages and iron metabolism. *Immunity* 44:492–504.
- Theurl I, et al. (2016) On-demand erythrocyte disposal and iron recycling requires transient macrophages in the liver. *Nat Med* 22:945–951.
- Wu J, et al. (2014) Strain-specific innate immune signaling pathways determine malaria parasitemia dynamics and host mortality. *Proc Natl Acad Sci USA* 111:E511–E520.
- Fervenza FC, et al. (2008) Induction of heme oxygenase-1 and ferritin in the kidney in warm antibody hemolytic anemia. *Am J Kidney Dis* 52:972–977.
- Nath KA, et al. (2001) Heme protein-induced chronic renal inflammation: Suppressive effect of induced heme oxygenase-1. *Kidney Int* 59:106–117.
- Poss KD, Toneyawa S (1997) Heme oxygenase-1 is required for mammalian iron reutilization. *Proc Natl Acad Sci USA* 94:10919–10924.
- Yachie A, et al. (1999) Oxidative stress causes enhanced endothelial cell injury in human heme oxygenase-1 deficiency. *J Clin Invest* 103:129–135.
- Soty M, Gautier-Stein A, Rajas F, Mithieux G (2017) Gut-brain glucose signaling in energy homeostasis. *Cell Metab* 25:1231–1242.
- Cummock K, et al. (2018) Host energy source is important for disease tolerance to malaria. *Curr Biol* 28:1635–1642.e3.