

See discussions, stats, and author profiles for this publication at: <http://www.researchgate.net/publication/229424075>

Myrmica salina (Hymenoptera: Formicidae) as a host of Maculineaalcon (Lepidoptera: Lycaenidae)

ARTICLE · JANUARY 2005

CITATIONS

11

READS

27

1 AUTHOR:



András Tartally

University of Debrecen

52 PUBLICATIONS 237 CITATIONS

SEE PROFILE

***Myrmica salina* (Hymenoptera: Formicidae) as a Host of *Maculineaalcon* (Lepidoptera: Lycaenidae)**

by

András Tartally

ABSTRACT

Numerous overwintered *Maculineaalcon* (Denis & Schiffermüller 1775) larvae and a single pupa were found in nests of *Myrmica salina* Ruzsky 1905 in the Szatmár lowland, NE Hungary. This ant species appears to be the primary host of *M.alcon* at this site. *M. salina* has previously been recorded as a host of *Maculinea teleius* (Bergsträsser 1779) at this, and one other site. Nearby, other local populations of *M.alcon* use *Myrmica scabrinodis* Nylander 1846 as a host, so it appears that this particular *M.alcon* population has become locally adapted to use *M. salina* as a host ant because of the secondary salinization of this site.

Keywords: *Myrmica salina*, *Maculineaalcon*, host specificity, myrmecophily, new host species, local adaptation, Hungary.

INTRODUCTION

Larvae of *Maculinea* van Eecke 1915 are obligate parasites of *Myrmica* Latreille 1804 colonies during most of their development. The host ant species has been shown to be crucial for the protection of these endangered butterflies (see e.g. Elmes *et al.* 1998). *Myrmica rubra* (Linnaeus 1758), *M. ruginodis* Nylander 1846 and *M. scabrinodis* Nylander 1846 are the known hosts of *Maculineaalcon* (Denis & Schiffermüller 1775) in western Europe, where different populations of this butterfly have evolved to use these three hosts in varied proportions (Thomas *et al.* 1989; Elmes *et al.* 1994, 1998; Als *et al.* 2002). In contrast, only *M. scabrinodis* was considered a host of *M.alcon* in central Europe (Sielezniew & Stankiewicz 2002, Höttinger *et al.* 2003, Tartally & Csősz 2004). Sielezniew & Stankiewicz (2004) recently found two localities in Poland where both *M. scabrinodis* and *M. vandeli* Bondroit 1920 are the hosts of *M.alcon*, increasing the global number of known *Myrmica* host species (see: Als *et al.* 2004: Supplementary Table 10) to four.

METHODS

During 2002 and 2004 I repeatedly searched a known *M. alcon* site in a slightly salinized marshy meadow near the village of Fülesd (Szatmár lowland, NE Hungary; N 48.05°, E 22.75°) for *Myrmica* nests. Each nest (2 *M. gallienii* Bondroit 1919, 1 *M. ruginodis*, 9 *M. salina* Ruzsky 1905 and 4 *M. scabrinodis* nests; determination according to Seifert 1988) situated within 2 m of the initial host plant *Gentiana pneumonanthe* L. was investigated (see: Als *et al.* 2002) in a 20 x 50 m area. Further nests were not opened in order to minimize the disturbance to this small, protected and endangered site.

RESULTS

On 7 May 2004 two *M. salina* nests were found to contain 61 and 5 *M. alcon* larvae respectively. The nests were marked and the site was revisited on 29 June 2004 (the flying of *M. alcon* started three weeks later, Varga personal communication). At this time, the ants had vacated these disturbed nests, presumably taking the *M. alcon* larvae with them. However, a single *M. alcon* pupa and 39 prepupal larvae (which is a huge number; see: Elmes *et al.* 1994, Thomas & Elmes 1998, Als *et al.* 2002, Höttinger *et al.* 2003, Tartally & Csősz 2004) were found in another *M. salina* nest, about 25 m from the nest which contained 61 *M. alcon* larvae in May. An additional three prepupal *M. alcon* larvae were also found in another *M. salina* nest. No *M. alcon* larvae or pupae were found in the nests of other *Myrmica* species on this site. Voucher samples (revised by Csősz) are stored in the authors collection and in the Hymenoptera Collection of the Hungarian Natural History Museum/Budapest.

DISCUSSION

These results do not exclude the possibility that *M. alcon* may use other species of *Myrmica* as hosts at Fülesd because the number of *M. scabrinodis*, *M. gallienii* and *M. ruginodis* nests examined was small. However, they do show that *M. salina* is an important host ant of *M. alcon* at this site, as had already been suggested by adoption experiments (Tartally & Csősz 2004, Tartally unpublished work). Presumably, *M. scabrinodis* was the only host ant of *M. alcon* from Hungary recorded in the published literature (Tartally & Csősz 2004) and *M. salina* has never previously been implicated as a host of *M. alcon* in Hungary (Tartally unpublished work) or in Transylvania, W. Romania (Csősz personal communication). Moreover, prepupal *M. alcon* larvae were observed exclusively in *M. scabrinodis* nests at a near lying

lowland site in NE Hungary (about 32 km from Fülesd, at Hetefejércse, Bereg lowland; N 48.15, E 22.45; Tartally unpublished work). These observations suggest that *M. alcon* has locally adapted to use *M. salina* as its primary host ant at Fülesd. This is likely to be a relatively recent phenomenon, since one part of the site at Fülesd was used as a rice field during the 1950s, which may have resulted in the secondary salinization of the soil. *M. salina* is characteristic for habitats with a particularly high salinity (Seifert 1988).

Sielezniew & Stankiewicz (2004) hypothesized that *M. vandeli* (probably a temporary social parasite of *M. scabrinodis* nests, Elmes *et al.* 2003) has a similar chemical profile to *M. scabrinodis*, allowing *M. alcon* larvae to easily develop in the nests of *M. vandeli* at two Polish sites. It would also be interesting to compare the chemical signatures of *M. salina* with those of *M. scabrinodis* and *M. alcon* larvae from Fülesd.

Nests of *M. salina* from the Fülesd site and at another Hungarian site were found to have *Maculinea teleius* (Bergsträsser 1779) larvae (Tartally & Csősz 2005). These were the first records for *M. salina* as a host ant of any *Maculinea* species (see: Als *et al.* 2004: Supplementary Table 10, but see: Tartally & Csősz 2004). My observations in 2004, of a single caterpillar being found with *M. salina* on 7 May and another on 29 June, give additional support for the use of *M. salina* by *M. teleius* at Fülesd. I have never found larvae of *M. teleius* and *M. alcon* together in the same *M. salina* nest.

Under laboratory conditions a *Maculinea rebeli* (Hirschke 1904) larva from Hungary has also been raised to pupation in a *M. salina* colony (Tartally 2004). This is not surprising considering the overall genetic similarity of *M. alcon* and *M. rebeli* (Als *et al.* 2004, Bereczki *et al.* 2005), although these butterflies often have different host ant species in the same region (see: Thomas *et al.* 1989; Elmes *et al.* 1994, 1998; Als *et al.* 2002; Sielezniew & Stankiewicz 2002; Höttinger *et al.* 2003; Steiner *et al.* 2003; Sielezniew & Stankiewicz 2004; Tartally & Csősz 2004). This larva pupated only 35 days after adoption, but this laboratory result has not been confirmed by field records yet (Tartally 2005).

ACKNOWLEDGMENTS

This study would not be possible without the pioneering work of the late Walter Wettstein, who made, during his tragically short life, several important ornithological and entomological discoveries at the Upper Tisza region in NE Hungary.

I would like to thank E. Tóth, Z. Varga, G.W. Elmes, D.R. Nash, J.A. Thomas and S. Csősz for their help and for their critical comments.

Research has been funded by the EC within the RTD project "MacMan" (EVK2-CT-2001-00126).

REFERENCES

- Als, T.D., D.R. Nash & J.J. Boomsma 2002. Geographical variation in host-ant specificity of the parasitic butterfly *Maculinea alcon* in Denmark. *Ecological Entomology*, 27: 403-414.
- Als, T.D., R. Vila, N.P. Kandul, D.R. Nash, S.-H. Yen, Y.-F. Hsu, A.A. Mignault, J.J. Boomsma & N.E. Pierce 2004. The evolution of alternative parasitic life histories in Large Blue butterflies. *Nature*, 432: 386-390.
- Bereczki, J., K. Pecsénye, L. Peregovits & Z. Varga 2005. Pattern of genetic differentiation in the *Maculinea alcon* species group (Lepidoptera, Lycaenidae) in Central Europe. *Journal of Zoological Systematics and Evolutionary Research*, (in press).
- Elmes, G.W., A.G. Radchenko & J.A. Thomas 2003. First records of *Myrmica vandeli* Bondroit (Hymenoptera, Formicidae) for Britain. *British Journal of Entomology and Natural History*, 16: 145-152.
- Elmes, G.W., J.A. Thomas, O. Hammarstedt, M.L. Munguira, J. Martin & J. van der Made 1994. Differences in host-ant specificity between Spanish, Dutch and Swedish populations of the endangered butterfly, *Maculinea alcon* (Denis et Schiff.) (Lepidoptera). *Memorabilia Zoologica*, 48: 55-68.
- Elmes, G.W., J.A. Thomas, J.C. Wardlaw, M. Hochberg, R.T. Clarke & D.J. Simcox 1998. The ecology of *Myrmica* ants in relation to the conservation of *Maculinea* butterflies. *Journal of Insect Conservation*, 2: 67-78.
- Höttinger, H., B.C. Schlick-Steiner & F.M. Steiner 2003. The Alcon blue *Maculinea alcon* (Lepidoptera: Lycaenidae) in eastern Austria: endangering and conservation measures. *Ecology (Bratislava)*, 22: 107-118.
- Seifert, B. 1988. A taxonomic revision of the *Myrmica* species of Europe, Asia Minor and Caucasia (Hymenoptera, Formicidae). *Abhandlungen und Berichte des Naturkundemuseums Görlitz*, 62: 1-75.
- Sielezniew, M. & A. Stankiewicz 2002. First data on host-ant specificity of parasitic butterfly *Maculinea alcon* (Den. & Schiff.) (Lepidoptera: Lycaenidae) in Poland and eastern Europe. *Fragmenta Faunistica*, 45: 123-130.
- Sielezniew, M. & A.M. Stankiewicz 2004. Simultaneous exploitation of *Myrmica vandeli* and *M. scabrinodis* (Hymenoptera: Formicidae) colonies by the endangered myrmecophilous butterfly *Maculinea alcon* (Lepidoptera: Lycaenidae). *European Journal of Entomology*, 101: 693-696.
- Steiner, F.M., M. Sielezniew, B.C. Schlick-Steiner, H. Höttinger, A. Stankiewicz & A. Górnicki 2003. Host specificity revisited: New data on *Myrmica* host ants of the Lycaenid butterfly *Maculinea rebeli*. *Journal of Insect Conservation*, 7: 1-6.
- Tartally, A. 2005. Accelerated development of *Maculinea rebeli* larvae under artificial conditions. *Nota Lepidopterologica*, 27: (in press).

- Tartally, A. & S. Csősz 2004. Adatok a magyarországi *Maculinea* fajok (Lepidoptera: Lycaenidae) hangyagazdáiról (Data on the ant hosts of the *Maculinea* butterflies (Lepidoptera: Lycaenidae) of Hungary). Természetvédelmi Közlemények, 11: 309-317.
- Thomas, J.A. & G.W. Elmes 1998. Higher productivity at the cost of increased host-specificity when *Maculinea* butterfly larvae exploit ant colonies through trophallaxis rather than by predation. Ecological Entomology, 23: 457-464.
- Thomas, J.A., G.W. Elmes, J.C. Wardlaw & M. Woyciechowski 1989. Host specificity among *Maculinea* butterflies in *Myrmica* ant nests. Oecologia, 79: 452-457.

