

Review

The Neglected Sibling: NLRP2 Inflammasome in the Nervous System

László Ducza^{##}, Botond Gaál[#]

Department of Anatomy, Histology and Embryology, Faculty of Medicine, University of Debrecen, Hungary, Hungary

[Received August 1, 2023; Revised September 25, 2023; Accepted September 26, 2023]

ABSTRACT: While classical NOD-like receptor pyrin domain containing protein 1 (NLRP1) and NLRP3 inflammasomal proteins have been extensively investigated, the contribution of NLRP2 is still ill-defined in the nervous system. Given the putative significance of NLRP2 in orchestrating neuroinflammation, further inquiry is needed to gain a better understanding of its connectome, hence its specific targeting may hold a promising therapeutic implication. Therefore, bioinformatical approach for extracting information, specifically in the context of neuropathologies, is also undoubtedly preferred. To the best of our knowledge, there is no review study selectively targeting only NLRP2. Increasing, but still fragmentary evidence should encourage researchers to thoroughly investigate this inflammasome in various animal- and human models. Taken together, herein we aimed to review the current literature focusing on the role of NLRP2 inflammasome in the nervous system and more importantly, we provide an algorithm-based protein network of human NLRP2 for elucidating potentially valuable molecular partnerships that can be the beginning of a new discourse and future therapeutic considerations.

Key words: inflammasome, NLRP2, neurological disorders, connectome, STRING

1. Introduction

Acute inflammation is a physiological defense mechanism in response to pathogens and cell damages; chronic inflammation, however, is considered as a dysregulated maladaptive clinical phenomenon without any recuperative benefits [1]. Chronic inflammation has been associated with many neurological disorders, [2] thus studying inflammasomes is of particular importance.

Classically, the currently known canonical inflammasomes (NLRP1, NLRP2, NLRP3, NLRP6, NLRP7, NLRP9, NLRP12, absent in melanoma 2 (AIM2) and pyrin inflammasomes) recruit caspase-1 enzyme, cleaving the zymogen interleukin-1 β (IL-1 β), IL-18 or IL-37 to induce lytic pyroptotic cell death and subsequent inflammatory downstream signaling [3-7].

Inflammasome machinery was first characterized in macrophage lineage cells by Martinon et al [8] in 2002, functioning as a caspase-activator normally inactive

intracellular protein platform releasing mature proinflammatory mediators upon activation by microbial antigens and foreign and/or host-derived danger factors. Since then inflammasomes have been identified in several cell types, they were found partake in the activation of innate immune system in numerous organs including the central nervous system (CNS) as well. Innate immunity is the major core component of neuroinflammation regarding both causative and consequential viewpoints of neuropathologies [9, 10, 11]. Nowadays, it is undoubtedly accepted that neuroinflammatory- then neurodegenerative states are the aftermaths of abnormal protein aggregations such as amyloid- β , α -synuclein or prions that may directly induce inflammasome- assembling [12].

NLRP2 (alternate names: NALP2, PYPAF2, NBS1, PAN1, CLR19.9) inflammasome is an intracellular multimer protein signaling hub assembled by three main elements: (i) a sensory component, termed NLRP, involved in the recognition of Pathogen-Associated

*Correspondence should be addressed to: Dr. László Ducza, Faculty of Medicine, University of Debrecen, Hungary, Hungary. Email: ducza.laszlo@anat.med.unideb.hu.

Copyright: © 2023 Ducza L & Gaal B. This is an open-access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Molecular patterns (PAMPs) and Damage-Associated Molecular Patterns (DAMPs), (ii) an adaptor unit, termed Apoptosis-associated Speck-like protein containing a caspase-activation and recruitment domain (ASC, CARD) as well as the (iii) effector caspase-1 enzyme [3, 8-16]. NLRP2 activation recruits „function to find” (FIIND) and a CARD-containing protein Cardinal that interacts with caspase-1 [15]. Of note, NLRP2 was earlier shown to govern inflammasome signaling by inhibiting nuclear factor kappa B (NF- κ B) transcription factor [17, 18]. Recent data have indicated that NLRP2 robustly increases the amount of NF- κ B regulated cytokines in cystinosis [19]. Moreover, several lines of evidence support the

critical regulator role of NLRP2 in the reproductive system, hence *NLRP2* gene was reported as one of the mammalian maternal effect genes associated with murine embryogenesis, age-related maternal fertility and idiopathic recurrent miscarriage, respectively [20-22]. Besides, NLRP2 expression was also proven to associate with arsenic-induced skin lesion, chromosomal damage, and respiratory disorders as well [23]. Intriguingly, despite the aforementioned past research on non-neural tissues, our gap of knowledge regarding NLRP2 and its connectome in neuropathological context has still yet to be improved (Fig. 1, Fig. 2).




Figure 1. Litmaps visualisation of original research articles targeting NLRP2 inflammasome in the nervous system Articles are shown in display mode “Momentum”. Calculation is based on “Cited By” count, weighed by recency.

Although our comprehension of the role of certain inflammasomes in CNS disorders has substantially improved throughout the past two decades, the causes of multiple number of NLRP2-related neurological disorders are indispensable to be further researched. Therefore, first, our study focused on the past experiments related to NLRP2 inflammasome in human and rodent models by providing a state-of-the-art of the literature. Secondly, a publicly available cutting-edge biomedical database called Search Tool for the Retrieval of Interacting Genes/Proteins (STRING, ver.11.5; string-db.org) [24] was employed to envisage potential protein interactions of NLRP2.

Apart from a few exceptions, currently there are future directions neither in NLRP2 research pipelines nor in therapeutic avenues. In this overview, following clarification of the main aspects of NLRP2 research, we aimed to describe a set of proteins (connectome) that had been already identified in a broad spectrum of brain disorders, but still none of their actions were directly linked with NLRP2. Eventually, we found 15 proteins that may associate with NLRP2, representing promising therapeutic targets and signaling pathways in NLRP2 associated neurological disorders.




Figure 2. Number of publications and citations related to NLRP2 inflammasome in the nervous system (2005-2023) based on Web of Science Core Collection (WOSCC).

2. Cellular distribution of major inflammasomes in the nervous system

CNS parenchyma has a remarkably complex architecture represented by diverse and highly interconnected cell types comprising neurons and glial cells (mainly astrocytes, microglia, oligodendrocytes) that collaborate to provide ideal circumstances for proper neuronal development and function. Adding another depth of complexity, in the vast majority of brain disorders not exclusively neurons, but glial- and even peripheral immune cells (granulocytes, monocytes, and lymphocytes) are recruited as well, all endowed with great level of heterogeneity including dynamically changing phenotypes upon noxious stimuli. These cells can vary spatiotemporally in their morphological and ultrastructural features as well as their genetic and protein signature. Complex neuron-glia-immune cell interactions even aggravate the determination of causative relationships and identification of primary disease vs. secondary “reactive” states [25, 26, 27]. Expression of inflammasomal NLRPs have been meticulously investigated in the nervous system. Main expression patterns are as follows: NLRP1 and NLRP3: primarily in microglial cells [28, 29], NLRP2 and NLRP3: in astrocytes [7, 30-35], NLRP2: dorsal root ganglia (DRG) cells and neural stem cells [36, 37], AIM2 and NLRP1: in neurons [38, 39]. Recently, oligodendrocytes were also shown to induce NLRP3 activation upon glycolytic stress in Alzheimer’s disease (AD) [40]. Of these types, the distribution of NLRP1 and NLRP3 inflammasomes is the most abundant in the CNS, they play a critical role in

neuroinflammation as well as wide range of neurological abnormalities like ischaemic stroke, traumatic brain- and spinal cord injury, neurodegenerative diseases, epilepsy or variety of brain infections [11, 41-44].

3. NLRP2 expression in the nervous system

3.1 Human NLRP2

For many years it remained elusive whether NLRP2 could be validated as a functional inflammasome in the human nervous system. The pioneering milestone was achieved by Julia Minkiewicz et al. in 2013 [7] by verifying the significance of NLRP2 inflammasome in human cortical astrocyte cultures. They proposed that NLRP2 interacted with ionotropic purinergic receptor X7 (P2X7) receptor and pannexin 1 (PNX1), hence mimicking inflammation with ATP stimulation rapidly resulted in caspase-1 activation and subsequent IL-1 β release. Conversely, pharmacological blockade of astrocytic PNX1 and P2X7 receptor with probenecid and Brilliant Blue G (BBG) diminished the ATP-induced NLRP2 activation. In addition, siRNA-based silencing of *NLRP2* was also found to disrupt caspase-1 activation, emphasizing its pivotal role in astrocyte immune responses. *NLRP2* gene dysregulation was shown to be involved in human fetal brain development indicating a robust difference between cases of bipolar disorder and healthy individuals [45, 46]. In line with this, Truvé et al. [37] in 2020 found considerable increase of NLRP2 expression in neural stem- and mature cells of patients suffering from bipolar disorder.

3.2 Rodent NLRP2

To date, still more NLRP2 related experimental data are accessible from rodent models. Several rodent models are continuously used in drug development pipelines, but regrettably the limitations of these concepts are major hindrances to translation attempts from bench to bedside. Few years previously Sun et al. in 2016 [32] have already determined NLRP2 distribution at specific areas of the murine CNS. Low constitutive NLRP2 levels were observed in the cortex, hippocampus and striatum, which were significantly elevated following occlusion of middle cerebral artery mimicking ischemic stroke. Similar results were obtained in primary astrocyte cultures upon oxygen-glucose deprivation. These findings were further corroborated by Cheon et al. in 2018 [33], who searched the interaction of apoptosis signal-regulating kinase 1 (ASK1) protein with astroglial NLRP2 inflammasome after ischemic stroke. ASK1 inhibition effectively downregulated NLRP2, as well as proinflammatory cytokine release in murine model and cultured astrocytes. In former studies, NLRP1 and NLRP3 inflammasomes were recognised in neuropathic models and spinal injury [47-50], but only scanty data were provided about NLRP2 protein following peripheral inflammation. Matsuoka and his colleagues in their study [36] characterised NLRP2 inflammasome in NeuN- and peripherin positive DRG cells of C-fibers implying its significant role in nociceptive processing. They found that both siRNA gene silencing of NLRP2 expression and pharmacological blockade of caspase-1 prevented nociceptive hypersensitivity, measured in two peripheral inflammatory models using either complete Freund adjuvant (CFA)- or ceramide. Their data did not support possible activation of matrix metalloproteinase enzyme 9 or other inflammasome types except NLRP2, therefore they hypothesized that caspase-1 enzyme in DRG neurons worked presumably in NLRP2-dependent manner. Enigmatically, inhibition of caspase-1 and siRNA silencing of NLRP2 attenuated mechanical, but not thermal hypersensitivity upon inflammation. This finding can be explained by the low NLRP2 level in thermal sensor transient receptor potential vanilloid subtype-1 positive neurons.

Our workgroup also investigated the putative changes of NLRP2 expression in CFA- induced inflammatory pain model within the spinal dorsal horn [31]. Although, NLRP1, NLRP2 and NLRP3 inflammasomes have been all detected in CNS [46, 51], our results showed that dorsal horn astrocytes mainly expressed NLRP2, colocalisation was entirely lacking between NLRP1 and astrocytes, we did not investigate DRG cells. With protein blot analysis we also verified a robust increase of NLRP2 protein in spinal cord tissue lysates upon CFA

injection. Not unexpectedly, the role of astroglial NLRP2 has also been reported in chronic mild stress induced depression [34], where they introduced an intraperitoneal injection of tryptophane derived metabolite kynurenine (Kyn), which upregulated NLRP2 in hippocampal primary astrocytes cultures. Simultaneously, Kyn treatment aided nuclear translocation of NF κ B to promote *NLRP2* transcription, eventually leading to inflammasome activation. These findings also illustrated that Kyn elicited depressive behaviour was eliminated following astrocytic *NLRP2* knockdown, proposing a crucial role for this inflammasome in depressive behaviours.

Recently, the role of NLRP2 inflammasome has been elucidated in murine AD [35]. In this study, glucagon-like peptide-1 (GLP-1) receptor agonist exenatide attenuated neuroinflammation in piriform cortex, resulting in cognitive improvement of 5x FAD transgenic mice. Additionally, upon exendine-4 treatment of cultured astrocytes, lower amyloid β_{1-42} level was detected confirming the reduced inflammation and oxidative stress presumably via downregulation of NLRP2 inflammasome.

3.3 Controversial findings of NLRP2 expression

Taken together, in agreement of results obtained from human and rodent models, expression of NLRP2 inflammasome is relatively scanty, therefore it is less studied in CNS compared with NLRP1 and NLRP3. Furthermore, distribution of NLRP2 appears to be more scattered, hence it has also been detected in different types of astrocytes (such as primary astrocyte cultures or astrocytes of spinal dorsal horn tissue), and also in neurons (such as DRG cells or neural stem cells).

However, the question arises, what the reason could be for the simultaneous duplicated expression in dorsal horn astrocytes and DRG cells during CFA induced inflammatory pain. Unfortunately, although there is limited information from experimental approaches, we hypothesize that NLRP2 function may span multiple cell types and induce/influence signalings that require further investigation including data mining techniques.

4. Human NLRP2 connectome with STRING database

The complex nature of cells derives from a wide variety of protein-protein interactions. Knowledge of major interactions shows an increasing trend, but novel interactions should also be explored. Meanwhile the information is frequently divided across databases with differing profiles, STRING [24] is a freely accessible online biological web resource that aims to collect protein-protein associations taken from many organisms

disorders at least in one article), are introduced in the order of their interaction score (chapter 4.14-4.15). We excluded those items that were either experimentally investigated with NLRP2 protein in the nervous system or reviewed in chapter 1-3.

The detailed principles of how scoring is precisely calculated are legible on the string.db website. Briefly,

STRING determines the numerical value of the interaction score by combining the probability values from all of the evidence channels and makes correction for the probability of randomly occurring interactions applying a freely available python script (combine_subscores.py).

Table 1. List of direct interactions of NLRP2.

	Node protein hits	Full name	Interaction score
1	CASP1	Caspase-1	0.862
2	PYCARD	Apoptosis-associated speck-like protein containing a CARD	0.825
3	CASP5	Caspase-5	0.811
4	MEFV	Pyrin	0.798
5	CARD8	Caspase recruitment domain-containing protein 8	0.771
6	SLA	Src-like-adapter; Adapter protein	0.752
7	KHDC3L	KHDC3-like protein; KH domain containing 3 like, subcortical maternal complex member	0.739
8	BECN1	Beclin-1	0.664
9	AIM2	Interferon-inducible protein AIM2	0.645
10	P2RX7	P2X purinoceptor 7; Receptor	0.638
11	CARD16	Caspase recruitment domain-containing protein 16	0.629
12	PANX1	Pannexin-1	0.600
13	IL18	Interleukin-18	0.597
14	SLA2	Src-like-adapter 2; Adapter protein	0.590
15	PAGE4	P antigen family member 4	0.575
16	EIF1AX	Eukaryotic translation initiation factor 1A	0.567
17	IL1B	Interleukin-1 beta	0.563
18	ZFP57	Krab domain-containing zinc finger protein	0.563
19	VGLL1	Transcription cofactor vestigial-like protein 1	0.562
20	CHUK	Inhibitor of nuclear factor kappa-B kinase subunit alpha	0.560
21	NLRP5	NACHT, LRR and PYD domains-containing protein 5	0.556
22	PCDH11X	Protocadherin-11 X-linked; Potential calcium-dependent cell-adhesion protein	0.547
23	GYG2	Glycogenin-2	0.515
24	OOEP	Oocyte-expressed protein homolog	0.514
25	MRE11A	Double-strand break repair protein MRE11	0.490
26	TLE6	Transducin-like enhancer protein 6	0.487
27	SUGT1	SGT1 homolog, MIS12 kinetochore complex assembly cochaperone	0.485
28	RIPK2	Receptor-interacting serine/threonine-protein kinase 2	0.473
29	SPON2	Spondin-2	0.471
30	IKBKG	NF-kappa-B essential modulator	0.470
31	EPS15	Epidermal growth factor receptor substrate 15	0.468
32	NEUROD1	Neurogenic differentiation factor 1	0.466
33	CAPN6	Calpain-6	0.462
34	PRKX	cAMP-dependent protein kinase catalytic subunit PRKX	0.456
35	TXLNG	Gamma-taxilin	0.455
36	DLGAP2	Disks large-associated protein 2	0.449
37	APLNR	Apelin receptor	0.445
38	SLC29A1	Solute carrier family 2	0.444
39	CARD18	Caspase recruitment domain-containing protein 18	0.437
40	AS3MT	Arsenite methyltransferase	0.423
41	CCDC50	Coiled-coil domain-containing protein 50	0.421
42	TBL1X	F-box-like/WD repeat-containing protein TBL1X	0.418
43	PADI6	Protein-arginine deiminase type-6	0.418
44	FCAR	Immunoglobulin alpha Fc receptor	0.408
45	RCC1	Regulator of chromosome condensation	0.400

4.1 Src-like-adapter proteins (SLA proteins)

SLA proteins are signal transducing proteins received their name after the protooncogen Src family tyrosine

kinases (STK). Unlike Src kinases, SLA proteins do not have tyrosine kinase activity, yet they contribute to multiple cellular processes. Specifically, the SLA protein 1 (SLA1) was recognised earlier in proteosomic

degradation of redundant N-methyl-D-aspartate (NMDA) receptors preventing excitotoxicity [52]. Furthermore, STKs were also investigated because of their role in central sensitisation, hence intrathecal injection of STK inhibitors mitigated mechanical allodynia in chronic pain models [53]. In addition, STK inhibitor PP2 could efficiently downregulate lipopolysaccharide (LPS)-induced microglial activation and production of neuroinflammatory factors via NF- κ B inhibition in MPTP-induced murine model of Parkinson's disease (PD) [54, 55].

4.2 Beclin 1 (BECN1)

BECN1 is a multifaceted protein, having a well-established role in an apoptosis-related multimer complex, which is essential for nucleation of autophagic vesicles [56-58]. Convincing data earlier confirmed the contribution of BECN1 to neuroinflammation [59], associated with extracellular increase of A β plaques as well as microglia-derived proinflammatory cytokines. Similarly, disruption of autophagy via decrease of BECN1 ultimately leads to exaggerated microglial release of IL-1 β and IL-18. In patients suffering from Alzheimer's disease (AD) microglial cells exhibited strongly reduced level of BECN1 [60, 61]. In addition, beneficial effect of BECN1 overexpression was unveiled in murine huntingtin (HTT) model of Huntington's disease (HD). They concluded that in early disease stage BECN1 administration restored network impairments occurred accumulation of mutant HTT aggregates [62].

4.3 Protocadherin-11X (PCDH11X)

Protocadherins (PCDHs), as the most versatile cadherin proteins, are involved in several neural events such as synaptogenesis, dendrite- and axon growth. They are classified into three main α , β , and γ gene clusters with approximately 60 genes extended with 10 non-clustered δ genes [63]. It is a well-known fact that *PCDH* gene dysregulations may result in neurological disorders, especially HD and AD. In knock-in murine model of HD, RNA-seq experiments performed by Langfelder et al. [64] determined 4 gene modules associated with 37 clustered *PCDH* genes together with non-clustered δ members *PCDH20* and *PCDH11X/Y*. Despite the fact that Carrasquillo and his group [65] earlier reported 4 single nucleotide polymorphisms (SNPs) of *PCDH11X* (SNPs: rs5984894, rs2573905, rs5941047, rs4568761) with hypothesized role in late-stage AD, later works with thorough genome analysis revealed no correlation between these SNPs and AD [66, 67].

4.4 Glycogenin-2 (GYG-2)

Glycogen is a primarily used glucose reservoir that provides rapidly accessible energy, participating in blood sugar control as well as muscle contraction. Glycogenesis requires the cooperation of both GYG glycosyltransferase and glycogen synthase (GS) enzymes. GS is allosterically activated by glucose-6-phosphate, but inhibited by glycogen synthase kinase (GSK) phosphorylation [68, 69]. Defects of glycogen metabolism, specifically the function of GYG, may lead to a genetically heterogeneous and rapidly progressing neurodegenerative disorder called Leigh syndrome that displays severe bilateral pathological alterations in cerebellum, spinal cord and brainstem. *In vitro* experiments earlier unveiled that human hemizygous missense mutation of *GYG-2* gene impaired self-glycosylation leading to disruption of glycogenesis [70]. Another glycogenesis related abnormality, the Lafora disease is characterised by aberrant intracellular glycogen deposits in both neurons and astrocytes [71]. Recently, ubiquitous serine/threonine GSK-3 enzyme has become an important player in neuronal metabolism. GSK-3 acts as the main negative regulator of GS, its pathological role has been already confirmed in AD. GSK-3 targets include A β and tau hallmark proteins of senile plaques and neurofibrillary tangles [72], confirmed by the significant increase of GSK-3 level in postmortem brain of AD patients [73]. Moreover, recent findings have demonstrated that lysophosphatidic acid, an identified GSK-3 activator, showed association with AD biomarkers A β , total tau and phospho-tau [74].

4.5 Meiotic recombination 11A (MRE 11A)

Undoubtedly, genome maintenance is vital in every cell, by preventing potential DNA damage via sophisticated DNA repair mechanisms [75]. Considering that neurons are post-mitotic cells producing high number of free radicals through their active metabolism, unlike dividing cells, they can not use the machinery of homologous recombination to repair double-strand break (DSB) in an error-free manner. Thus, they rather recruit error-prone repair methods like nonhomologous end joining [76]. Of the several types of DNA damages, DSBs are regarded the most deleterious for neurons, worsening their function and survival. Owing to the fact that no complementary strand template exists for repair, genome instability, and ultimately age-related cognitive disorders and neurodegeneration may develop [77, 78]. Given its key determinant role in MRE11-RAD50-NBS1 (MRN) damage sensor complex, MRE11 orchestrates ATM- or ATR-mediated DNA damage responses upon DSBs [79]. MRE11 nuclease defects were found to be associated with

human pathologies such as progressive myoclonic ataxia [80], Nijmegen breakage syndrome-like disorder [81] and Ataxia-Telangiectasia-like disorder [82]. Furthermore, role of MRE11 was also revealed in microcephaly, cognitive disabilities and cerebellar degeneration [83], but so far only one article reported substantially reduced level of MRE11 and other members of MRN complex in AD patients [84].

4.6 Suppressor of G2 allele of Skp1 (SUGT1)

SUGT1 is a major component of the CBF3 kinetochore as well as SCF ubiquitin ligase complexes [85, 86]. SUGT1 binds heat shock protein 90 (Hsp90) via CHORD-Sgt1 domain, therefore it is also considered as a member of chaperone complexes [87, 88]. Additionally, SUGT1 was found to be associated with innate immunity through Nod1 stabilization [89]. SUGT1 is abundantly expressed in hippocampal and cortical neurons as well as in Purkinje cells and cerebellar glial cells. Unsurprisingly, the significance of SUGT1 was earlier described in neurodegenerative diseases including AD and PD respectively [90, 91]. In AD brain, fewer SUGT1 immunopositive neurons were found compared with control [92], but in contrast, *SUGT1* mRNA level was robustly enhanced in fronto-temporal cortex of human PD [91].

4.7 Epidermal growth factor receptor substrate 15 (EPS15)

EPS15 tyrosine kinase substrate was first identified thirty years ago as the binding partner of AP-2 complex member alpha-adaptin, playing a crucial role in clathrin-mediated endocytosis as well as vesicular transport [93, 94]. Given its localisation in coated pits, it was proven that EPS15 contributed to internalization of the epidermal growth factor receptor (EGFR). For instance, antibody microinjection against EPS15 or using *EPS15* knockdown animals inhibited clathrin-mediated endocytosis of EGFR [95-97]. EPS15 participates in the EGF-AKT pathway mediated pro-survival signaling that is highly reduced in PD patients. In accordance with this, striatal dopaminergic neurodegeneration can be prevented upon EGFR activation in PD animal model [98]. Nevertheless, even if EGF-EGFR binding promotes AKT pathway, it also stimulates ubiquitin-mediated receptor endocytosis and degradation. In doing so, ubiquitinated EGFRs, guided by EPS15, are trafficked directly to proteosomal degradation. However, EPS15-EGFR interaction can be inhibited by E3 ubiquitin-protein ligase called Parkin that aids ubiquitination of EPS15 while preventing receptor internalisation [99, 100].

4.8 Neurogenic differentiation factor 1 (NEUROD1)

It has been well documented that basic Helix-loop-Helix (bHLH) transcription superfamily, especially proneural bHLH subgroup, termed NEUROD family, is a critical determinant of neuronal differentiation in spinal cord, brainstem, and cerebral cortex [101, 102]. The family consists of four members: NEUROD1, NEUROD2, NEUROD4, NEUROD6 that show overlapping, but not identical expression pattern in the cerebral cortex. In addition, they are substantially expressed not only in prenatal, but in adult cortex, cerebellum, and hippocampus as well [103, 104, 105].

In pathological context, increased level of hippocampal *NEUROD1* has been detected in murine model of environmental stress normalised by antidepressant agomelatine [106]. Rubio-Cabezas et al. [107] identified the first human mutations of *NEUROD1* gene in the form of two homozygous frameshift mutations found in patients with neonatal diabetes and neurological disorders including cerebellar hypoplasia, visual impairment, and sensorineural deafness. *NEUROD1* gene ablation produced murine epilepsy due to excessive neurodegeneration of hippocampal dentate granule cells [108], moreover its reduced level was also detected in R6/2 murine model of HD [109]. Despite its significance, *NEUROD1* expression has been debated in AD patients based on RNA-seq results [110] from APPxPS1 murine model. However, the overexpression of NEUROD1 greatly raised dendritic spine density, improving the spatial memory in hippocampus [111]. Nowadays, novel therapeutic approaches attempt to either block ubiquitin-dependent proteosomal degradation of NEUROD1 to prevent neuronal loss or apply induced pluripotent stem cells-derived hippocampal spheroids [112, 113]. Besides, manipulation of *NEUROD1* gene in AD and ischemic stroke efficiently produced cortical neurons from *in vivo* reprogrammed reactive glial cells [114, 115].

4.9 Calpain-6 (CAPN6, CANPX, calpamodulin)

CAPN is a Ca²⁺ regulated intracellular non-lysosomal cysteine protease that cleaves numerous substrates resulting in either degradation or modulation of cell responses. To this date, 16 *CAPN* genes have been identified including *CAPN6* gene that belongs to the non-classic (atypical) *CAPN* genes [116, 117]. Despite the high level of *CAPN6* mRNA during embryogenesis, especially in the first brancial arch, skeletal muscle, bronchial epithelium, kidney- and lung sacs, its expression was robustly attenuated after birth [118-121]. In the last decade CAPN6 has attracted the neuroscientists' attention, as its overexpression was detected in prion disease-affected sheep brain tissues as

well as ischemia induced white matter injury [122, 123]. CAPN6 is yet to be investigated in diseases like AD and PD, nevertheless calpains have already been linked to neurodegenerative hallmark proteins such as A β plaque and tau hyperphosphorylation in AD. Notably, calpain contributes to reactive astrocyte transformation, since its pharmacological blockade abrogates the neuroinflammatory activity of both astrocyte and microglial cells meanwhile improving cognitive decline in the 3xTg AD model [124, 125].

4.10 γ -taxilin (TXLNG, γ -TXLN)

Members of the taxilin family were first recognised as interactive partners of syntaxins involved in intracellular vesicular transport. The family is composed of α -, β -, and γ -TXLNs sharing a coiled-coil domain at their C-terminal. Meanwhile α - and γ -TXLNs are ubiquitously expressed, β -TXLN is less abundant in skeletal- and heart muscle tissues [126-129]. During cell interphase γ -TXLN has a centrosomal localisation with reduced expression at G2/M transition owing to proteosomal degradation. Recently, γ -TXLN has been proposed to control Never in mitosis A-related kinase A2 mediated centrosome disjunction [130].

It has already been shown that inhibition of ER stress sensor γ -TXLN via GSK-3 signaling activates unfolded protein production leading to apoptosis. Moreover γ -TXLN ablation may induce tau hyperphosphorylation in AD patients [131]. In the past few years, even if large body of evidence [132, 133] supported physiological relevance of the taxilin family, further research on the role of family member γ -TXLN in neuropathology would be of great value.

4.11 Disks large-associated protein 2 (DLGAP2)

As a specialised matrix, postsynaptic density (PSD) plays a role in mediating excitatory neurotransmission at the postsynaptic terminal. Discs large scaffold proteins (DLGAPs) of PSD are required for proper function of both ionotropic and metabotropic glutamate receptors contributing to synaptic scaling [134-137]. Regardless of the fact that DLGAP1-4 is ubiquitously expressed in CNS, higher protein levels were found within the brain and DLGAP2 showed abundant expression in cortex, hippocampus, olfactory bulb and striatum, but not in cerebellum or thalamus [138-141]. Evidence is accumulating for its involvement in autism, hence *DLGAP2* knockout mice exhibited aggressivity, social dominance and aberrant synaptic dysfunction in orbitofrontal cortex [142]. Epigenetic studies also implied that *DLGAP2* contributed to other neuropsychiatric disorders like schizophrenia and post-traumatic stress syndrome [143, 144]. Unsurprisingly, cross-species

analyses verified *DLGAP2* as a major regulator in cognitive decline of AD patients, its low cortical expression was associated with neurodegenerative hallmarks [145].

4.12 Apelin receptor (APLNR)

APLNR is a G protein-coupled receptor, highly resembling the structure of angiotensin II receptor, providing a potentially therapeutic target in cardiovascular diseases. Its bioactive ligand apelin was reported to play a crucial role in myriad of physiological actions such as cardiac contractility, neoangiogenesis as well as immunoregulation, nitric oxide-induced vasorelaxation, fluid- and bone homeostasis, human immunodeficiency virus-1 infection, neuroendocrine stress, glucose metabolism, appetite, and drink behavior, respectively [146-155].

Compelling evidence emphasizes the involvement of apelin in neuropathologies as well. For instance, apelin-13 was identified to exert neuroprotective effects on cortical neurons via inhibiting serum deprivation-induced oxidative stress and apoptosis [156]. Apelin also downregulates NMDA receptor-mediated excitotoxicity in rat hippocampus [157]. Recently, apelin-13 has been shown to disrupt A β - induced memory deficit in A β ₂₅₋₃₅-treated animals [158]. Based on earlier observations of Agostinho, Cunha, and Oliveira [159], exacerbated neuroinflammation, characterised by glial-derived proinflammatory cytokines, facilitates neurodegenerative progression in AD patients' brain. In this context, apelin-13 certainly to be neuroprotective by alleviating ischemic stroke as well as the effects of proinflammatory cytokines [160]. Furthermore, insulin-like growth factor 1 (IGF-1), which exerts inhibitory effect on α -Syn aggregation in rodent PD, was found to be upregulated by apelin [161]. It has recently been reported in SH-SY5Y cell model of PD that apelin-13 attenuates selective 6-OHDA induced impairment of dopaminergic neurons [162, 163]. Additionally, apelin also ameliorates NLRP3 inflammasome resulting in dampened systemic inflammation [164]. Apelin has also been targeted in HD, since PolyQ-HTT, a pathological form of HTT that perturbs postsynaptic plasticity and cytoskeleton organisation, [165] is inhibited by apelin via PI3K/AKT and MAPK/ERK signaling [166]. Aforementioned research by Yang [163] has already demonstrated that apelin-IGF interaction may have significance in PD, corroborated by recent findings identifying IGF in clearance of HTT aggregates in HD [167].

4.13 Solute carrier family 29A1 (SLC29A1, ENT1)

SLC is a membrane transporter superfamily with more than 400 members and 65 families mediating translocation of versatile substances across biological membranes, such as electrolytes, lipids, sugars, amino acids, neurotransmitters or drugs [168, 169]. It is worth mentioning that SLCs have been considered significant pharmacological targets, thus developing new agents related to SLCs may aid their bioavailability in drug interactions- and disposition as well as toxicity [170-172]. SLC29 contains four human intrinsic energy independent membrane proteins termed Equilibrative Nucleoside Transporters (hENT1-4). ENTs regulate facilitative bidirectional nucleoside transport showing capability for transporting both pyrimidines and purines [173, 174].

Significance of SLC29A1 (ENT1) adenosine transporter has been highlighted in 2017 when Yu-Han Kao et al. [175] shed light on dysregulation of adenosine metabolism in HD. Striatal ENT1 expression was robustly enhanced in R6/2 and Hhd^{150Q} models and reduced following either intrastratial injection of ENT1 inhibitor JMF1907 or genetic suppression leading to higher survival rate of R6/2 mice. Later, others [176] found that application of adenosine analogue ENT1 inhibitor J4 was promising in preventing spatial memory decline in murine APP/PS1 model of AD. Moreover, chronic administration of J4 reversed the impaired basic synaptic signaling, long-term potentiation, excitatory synaptic expression and signaling of neuronal plasticity such as GSK-3 β or PKA. Recently, J4 has been tested in murine model of tauopathy, and was found to block tau hyperphosphorylation while improving memory, mitochondrial dysfunction, and synaptic loss [177]. Collectively, the evidence opens new prospects for neuropathological studies with adenosine dysregulation in mind.

4.14 Arsenite methyltransferase (AS3MT)

Inorganic arsenic (iAs) has been found in highest amount in the air as arsenic trioxide (As₂O₃), but it may be present in food, water or soil, predominantly in the form of either AsO₃ or AsO₂ (arsenite) [178]. Regrettably, an increasing number of people suffer from the consequences of iAs exposure to drinking water. It is particularly a concern in Bangladesh and Eastern parts of India, where arsenic inflicted crisis is a natural calamity affecting approximately 43 million people [179]. Apart from this, the most frequent causes of poisoning include occupational contamination, moonshine alcohol or malevolent delivery [180].

The role of as has robustly emerged in numerous deleterious neurological symptoms such as memory

deficit, Guillain–Barre-like neuropathy, encephalopathy, peripheral neuropathy, PD and AD. However, to date, particularly in terms of mechanisms underlying arsenic neurotoxicity, no remarkable success has been achieved. A growing body of experimental evidence proposes that free radicals provoke neuroinflammation, mitochondrial dysregulation, ER stress, apoptosis and protein homeostasis that may be implicated in AD [181-183]. Oxidative stress, mostly incriminated in arsenic-induced neuronal damage, substantially elevates the level of reactive oxygen species while reducing expression of superoxide dismutase and glutathione (GSH) [184, 185]. AS3MT contributes to iAs processing in the presence of GSH, thus GSH deficit may prevent detoxification resulting in AD [186]. It is nowadays considered trivial that neuroinflammation is involved in AD pathology [187]. As also appears to activate microglia cells that secrete proinflammatory cytokines [188, 189]. Besides, several data showed toxic effects of as on brain mitochondria [190, 191]. As has also been found to induce apoptosis of cerebellar neurons via activation of JNK and p38MAPK signaling [192]. Recent evidence shows that as upregulates apoptotic Bax and decreases antiapoptotic Bcl-2 factor [193]. Arsenic-mediated posttranslational modifications and impairment of ubiquitination may also damage proteostasis via accumulation of abnormal hallmark protein aggregates [194, 195].

4.15 Coiled-coil domain containing protein 50 (CCDC50)

Autophagy is a phylogenetically conserved lysosomal degradation program that coordinates lysosomal fusion with autophagosomes [196]. Mutations or deficiencies associated with autophagy highlight their significance in exacerbated inflammatory disorders [197]. CCDC50, also termed as Ymer, functions as an ubiquitous autophagy receptor that suppresses inflammatory responses by disrupting NLRP3 inflammasome [198, 199]. Previous findings have reported that CCDC50 also prevents ligand-induced downregulation of EGFR as well as regulate NF- κ B, Fas and interferon signaling. Thus, dysregulation of CCDC50 may play a pivotal role in clinical implications such as viral infection and autoinflammatory disorders including systematic lupus erythematosus [200].

Albeit insufficient data are yet available on CCDC50 in the context of neurodegeneration, genome wide association studies showed that SNPs of 3q28 locus were correlated with tau/ptau levels of cerebrospinal fluid and a broad range of AD phenotypes. However, no significant SNPs were found in *CCDC50* and other genes such as *GEMC1*, *OSTN*, *SNAR1*, *IL6RAP* or *UTS2D* from the same locus, they are substantially expressed in brain tissue and contribute to synaptogenesis [201].

5. Discussion and future prospects

To date, neurodegenerative disorders lack effective therapeutic agents or programs to prevent, efficiently influence or at least decelerate disease progrediation. Moreover, the rise in absolute numbers of people inflicted far and wide proposes that management of main neurological disorders is globally inadequate. These people pose a significant socioeconomic burden owing to disability, illness and premature death [202, 203].

Unquestionably, inflammasome signaling is affected in several neurodegenerative disorders including AD, PD or HD that attract more increasing attention in studies investigating their place in clinical practice. Specifically, the NLRP3 inflammasome has been meticulously investigated in this context [204, 205], but recently, role of other inflammasomes as pathological drivers in brain diseases was also conceptualised by Anna Chiarini et al

[46]. This attempt markedly indicates that there is an urgent need to extend our comprehension regarding neurodegenerative mechanisms associated with inflammasomes other than NLRP3.

First, in this study we attempted to encompass the literature focusing on the expression and distribution of NLRP2 inflammasome in human and rodent neuropathological disorders (Fig. 4). Taken together, in these models major NLRP2 expression was upregulated in primary astrocyte cultures [7, 32, 33, 34, 35], as well as in astrocytes of rodent spinal dorsal horn, respectively [31]. Beyond astrocytes, the role of NLRP2 was also verified in NeuN- and peripherin positive mechanical, but not thermal sensor subsets of DRG cells by Matsuoka et al [36]. Furthermore, tissue distribution of NLRP2 protein was observed not only in spinal cord, adult cortex, hippocampus or striatum [31, 32], but in neural stem cells as well [37].




Figure 4. State-of-the-art in relation to NLRP2 protein in neuropathologies. ASC: Apoptosis-associated Speck-like protein containing a caspase-activation and recruitment domain, ASK1: Apoptosis signal-regulating kinase 1, BBG: Brilliant Blue G, GLP-1: Glucagon-like peptide-1, NFκB: Nuclear factor kappa B, P2X4, P2X7- P2X purinoreceptor X4,7, PNX1-Pannexin-1, ROS-reactive oxygen species.

Regarding activatory stimuli, particular attention is given to purinergic signaling in human NLRP2 activation, hence ATP is a well-known and adeptly characterised

DAMP that facilitates inflammasome activation upon trauma [206]. Indeed, exogenous stimulation with ATP activates NLRP2 inflammasome resulting in caspase-1

mediated production of mature IL-1 β [7]. According to all indications, ATP acts on P2X7 receptor that cooperates with PNX1, hence application of P2X7 receptor inhibitor BBG and PNX inhibitor probenecid diminishes NLRP2 activation. Nevertheless, it is important to underline that P2X4 receptor has been found to be functionally coupled with P2X7 receptor and pannexin-1 in NLRP3 inflammasome of gingival epithelial cells [207, 208]. Overall, these observations are in accordance with findings in which heme activates NLRP3 inflammasome in macrophages via P2X7 and P2X4 signaling during kidney inflammation [209]. We postulate a similar case with NLRP2 inflammasome, however there are no confirming experimental data. In our earlier article we hypothesized that overexpression of astroglial NLRP2 in spinal dorsal horn may be associated with significant P2X4 upregulation upon intraplantar CFA injection [31, 210]. Other triggering stimulus of NLRP2 activation is ischemic stroke, evoked by occlusion of middle cerebral artery or oxygen-glucose deprivation in astrocyte cultures [32]. Unsurprisingly, mounting evidence suggests that astrocyte-mediated inflammation may be potentially involved in the pathogenesis of mental disorders such as depression [211, 212]. In their recent study, Zhang et al [34] found that tryptophan metabolite Kyn, a specific biomarker of depressive behaviours, upregulated NLRP2 in astrocytes, supported by others also reporting significantly elevated levels of proinflammatory cytokines [213]. In bipolar disorder, neuroinflammatory biomarkers of cerebrospinal fluid were found to associate with cognitive decline. Moreover, cognitive impairment is increasingly recognised in people suffering from bipolar disorder, suggesting a link between neuroinflammation, neurodegenerative states and mood abnormalities [214], which may be significantly modulated by NLRP2 inflammasome as well [37, 45, 46].

Secondly, the present study collects the connectome of NLRP2, envisaged with STRING platform. The obtained set of proteins were further filtered to 15 possible protein interactors involved in brain disorders; their pathophysiological roles are discussed in details in chapter 4.1-4.15, TABLE S1. Many of these potential partnerships have not been earlier experimentally investigated. Besides signaling of NLRP2 inflammasome is also lacking from the collection of the Kyoto Encyclopedia of Genes and Genomes (KEGG) database Supplementary Figure 1.

Concluding the literature and STRING based matches we hypothesize that the filtered NLRP2 connectome - influences a plethora of molecular processes in neurons, such as glutamatergic excitotoxicity, apoptosis/ survival signaling and neuroinflammation.

Glutamatergic excitotoxicity - The integrity of glutamatergic signaling is essential for preserving

neuronal homeostasis and evading neurodegeneration. SLA1 prevents glutamatergic excitotoxicity by degrading excessive NMDA receptors [52]. As a synergist, apelin also disrupts NMDA receptor-mediated excitotoxicity in rat hippocampus through survival kinases AKT and Raf/ERK-1/2 [157]. Furthermore, the role of DLGAPs is to anchor glutamate receptors in the postsynaptic membrane and to link them with other proteins including other glutamate receptors, signaling- and cytoskeletal factors while regulating both ionotropic and metabotropic glutamate receptors via synaptic scaling [134-137]. Low *DLGAP2* expression was detected in age-related cognitive decline and AD [145], which may relate to other findings, such as low spine density or reduced excitatory postsynaptic current, revealed in orbitofrontal cortex of *DLGAP2* knockout mice [140, 142]. Directed expression bHLH transcription factor *NEUROD1* stimulates neuronal maturation as well as ameliorates deficits of dendritic spine density in hippocampal neurons of APPxPS1 model of AD [111]. Besides, ENT1 inhibitor J4 mitigates the damages of long-term potentiation, excitatory synaptic expression as well as GSK-3 β or PKA signaling of neuronal plasticity and alleviates memory deficits in APP/PS1 model of AD [176].

Apoptosis/survival signaling- It is of particular importance to gain a better understanding of the signaling of neuronal apoptosis/survival in neurodegenerative states. Impairment of autophagy via reduction of BECN1 level results in overproduction of microglial IL-1 β and IL-18 proinflammatory cytokines in AD patients. Deficits of BECN1 mediated phagocytosis causes dysfunctional recruitment of phagocytic receptors that may associate with extracellular accumulation of A β plaques and other cellular debris. Recently, in early phase of HD BECN1 administration has successfully cleared mutant HTT accumulation and reversed progrediation [59-62]. Apelin exerts positive impacts on redox homeostasis and prevents mitochondrial cytochrome c release and caspase-3 activation in cultured murine cortical neurons [157].

Furthermore, EPS15 participates in the EGF-AKT pathway mediated pro-survival cell signaling reduced in dopaminergic neurodegeneration [98, 99]. CCDC50 autophagy receptor inhibits inflammatory responses by disrupting NLRP3 [197, 198] and prevents EGFR downregulation as well as regulate NF- κ B, Fas and interferon signaling. As exposure, in contrast, triggers apoptosis of cerebellar neurons via activation of JNK and p38MAPK signaling [182] as well as upregulates Bax and decreases Bcl-2 factor [192]. A rescue factor SUGT1, functioning as a chaperone protein, was also recognised to counteract pathological aggregation of α -synuclein and neurotoxicity in PD. *SUGT1* mRNA was highly elevated in fronto-temporal cortex in human PD [90, 91]. Its significance has also been earlier described in

AD, hence decreased SUGT1 immunopositivity was found in degenerating neurons of AD patients [92]

Dysregulation of aforementioned GSK-3 signaling, inhibited by PDK1/AKT kinase [215, 216], contributes to the hyperphosphorylation of tau protein as well as A β -induced cell death in AD pathogenesis. Abundant GSK-3 level was found in postmortem brain of AD patients, moreover large body of evidence support that GSK-3 activator lysophosphatidic acid associates with AD biomarkers A β , total tau and phospho-tau [72, 74]. In contrast, GSK-3 inhibits ER stress sensor γ -TXLN, promoting apoptosis- and autophagy. In addition, blockade of γ -TXLN alone may lead to tau hyperphosphorylation in AD [131].

Neuroinflammation- Several key mechanisms have been identified in neurodegenerative diseases as summarized above, also including neuroinflammation, which evokes a great challenge to clinical practice. Neuroinflammation has been recognised in dementia and is typically linked to cognitive decline with elevated levels of proinflammatory markers (IL-1, IL-6, IL-8, C-reactive protein) in patients suffering from dementia [217-219]. However, recent data suggest that inflammatory proteins may express both pro- and anti-inflammatory actions making the interpretation even more difficult in complex neurodegenerative states [220]. In fact, the versatility of postmortem samples as well as challenges of appropriate resolution in detection of cytokines may all lead to controversial conclusions [221, 222]. Indeed, neuroinflammation is still regarded as one of the crucial molecular processes of dementia, but we still do not know whether this mechanism is consequential or causative, in regards of neurodegenerative progression. Of note, it receives growing evidence that neuroinflammation may appear as an early temporal red flag event that may link to other mechanisms contributing to neuropathologies [223-225].

In fact, neuroinflammation and generally the therapeutic potential of targeting inflammasomes has been increasingly recognised in neurodegenerative conditions since 2013, when Heneka et al. [226] proved the significance of NLRP3 inflammasome with *NLRP3* -/- mice in AD. Since then, many efforts have been made to seek selective and potent NLRP3 inhibitors, because the currently US Food and Drug Administration (FDA) approved inhibitors of multiple inflammatory diseases include only canakinumab, anakinra and riloncept. Nevertheless, these inhibitors do not cross efficiently the blood-brain barrier and lack proper pharmacokinetic properties [227, 228].

A great achievement was reached when a potent diarylsulfonylurea compound MCC950 (also termed as CRID3, CP-456773), endowed with NLRP3 selectivity, showed therapeutic improvement in several preclinical

models such as experimental autoimmune encephalomyelitis, such as AD and PD [229-232], nevertheless it failed in clinical trials because of off-target toxicity.

In the search for new inhibitors Stavudine (d4T), acting as an inhibitor of nucleoside reverse transcriptase, has been published recently to downregulate NLRP3 activation in AD [233]. Furthermore, Gastaldi et al. [234] by applying the pharmacophore-hybridization method synthesised several benzo[d]imidazole-2-one derivatives to test their inhibitory effect on NLRP3 evoked pyroptosis and IL-1 β production.

Epigenetic research has also recently provided ammunition against inflammasomes. Three molecular events have come into focus: CpG DNA methylation, posttranslational modification of histones as well as noncoding RNA expression. From this point forward microRNAs may be promising therapeutic targets. Currently, as NLRP3 represents the most investigated inflammasome till date, testing of miR-30e and miR-7 have already started against this inflammasome type in PD [235, 236, 237].

However, it is increasingly becoming clear that NLRP3 is not the only inflammasome involved in neurodegenerative states. Notably, Kaushal et al. [238] has already reported elevated mRNA level of *NLRP1* and highlighted the causative role of NLRP1-caspase 1-caspase-6 signaling in the accumulation of A β ₄₂ deposits in AD.

For clarity, we need to examine those common reasons why proper therapeutic options targeting human inflammasomes including NLRP2 are still limited till date:

a) There are theories and evidence that inflammasomes can functionally take over each other's roles, albeit these mostly unrevealed interplays among canonical and/or non-canonical inflammasomes need to be further investigated. Denes et al. [29] showed in their rodent model that NLRC4 (NLR family, CARD domain containing 4) and AIM2 contribute a lot to the pathogenesis of acute ischemic brain injury in the presence of pharmacologically blocked NLRP3.

b) Unfortunately, it has been confirmed that the genetic signature as well as organs of rodents and humans differ greatly. This inter-species discrepancy also applies to the architecture of CNS, as in the most used rodent models usually only the structurally less complex olfactory cortex was put under scrutiny. Furthermore, even at a cellular level many morphofunctional aspects differ between humans and rodents [46, 239], and if that is not enough of a problem specifically their inflammasome regulation is also markedly distinct [240, 241]. Thus, even though we have a lot of animal experiments available from animals' models, low

efficiency rate of rodent-human translation has been delaying the development of breakthrough drugs for brain disorders for decades.

Currently, there is only one promising “game changer” in this struggle we can carry, the Lecanemab, IgG1 monoclonal antibody, which has been recently proven to reduce amyloidosis and cognitive deterioration in early-stage AD patients compared with placebo group, but we cannot forget about its unpleasant side effects such as brain swelling or hemorrhage [242].

c) Above that, even myriad of inhibitors is mentioned with potential therapeutic benefits against inflammasomes like NLRP3 [46, 227, 228, 229], these inhibitions mostly result from blocking related signaling pathways (such as dopamine receptor [243], adiponectin receptor [244], estrogen receptor [245], Angiotensin II receptor [246]) rather than direct inhibitory strategies. Thus, the same may be the case with NLRP2, if the major signaling pathways are precisely revealed. Data mining techniques like STRING can help us to target directly the potentially NLRP2-associated protein candidates and test them experimentally.

Nevertheless, it has been suggested to reduce the function of not only NLRP3 but all canonical inflammasomes by inhibiting the ASC protein with a newly developed small molecule inhibitor MM01 [247].

d) Finally, despite accumulating experimental evidence, unfortunately, few experts in the field still take seriously the idea that the rodent and human NLRP2 inflammasome should be addressed more seriously.

6. Conclusions

Taken together, based on STRING analysis human NLRP2 inflammasome may cooperate with proteins that are related to excitotoxicity, cellular apoptosis/survival and neuroinflammation. Thus, this filtered connectome in the nervous system requires an experimental approach both in rodent and human models. In the future, complete understanding of inflammasome signaling including NLRP2 mediated processes may open up new prospects for efficient inhibitors and/or therapeutic agents, which are currently lacking from our armamentarium.

Supplementary Materials

The Supplementary data can be found online at: www.aginganddisease.org/EN/10.14336/AD.2021.0926.

Author Contributions

Conceptualization: LD Manuscript writing, data visualisation, editing and supervision: LD, BG. All

authors have read and agreed to the published version of the manuscript.

Acknowledgements

This project was supported by the Department of Anatomy, Histology and Embryology, University of Debrecen, Hungary and received no other external funding.

Data Availability Statement

The data sets and materials supporting the conclusions of this study are included within the article.

Competing interests

The authors declare no competing interest.

References

- [1] Yi YS (2006). Functional crosstalk between non-canonical caspase-11 and canonical NLRP3 inflammasomes during infection-mediated inflammation. *Immunology*, 159(2):142-155.
- [2] Singh N, Baby D, Rajguru JP, Patil PB, Thakkannavar SS, Pujari VB (2019). Inflammation and cancer. *Ann Afr Med*, 18(3):121-126.
- [3] Schroder K, Tschopp J (2002). The inflammasomes. *Cell*, 140(6):821-32.
- [4] Ross C, Chan AH, von Pein JB, Maddugoda MP, Boucher D, Schroder K (2022). Inflammatory Caspases: Toward a Unified Model for Caspase Activation by Inflammasomes. *Annu Rev Immunol*, 40:249-269.
- [5] Kelley N, Jeltama D, Duan Y, He Y (2019). The NLRP3 Inflammasome: An Overview of Mechanisms of Activation and Regulation. *Int J Mol Sci*, 20(13):3328.
- [6] Akbal A, Dernst A, Lovotti M, Mangan MSJ, McManus RM, Latz E (2022). How location and cellular signaling combine to activate the NLRP3 inflammasome. *Cell Mol Immunol*, 19(11):1201-1214.
- [7] Minkiewicz J, de Rivero Vaccari JP, Keane RW (2013). Human astrocytes express a novel NLRP2 inflammasome. *Glia*, 61(7):1113-21.
- [8] Martinon F, Burns K, Tschopp J (2002). The inflammasome: a molecular platform triggering activation of inflammatory caspases and processing of proIL-beta. *Mol Cell*, 10(2):417-26.
- [9] Bergsbaken T, Fink SL, Cookson BT (2009). Pyroptosis: host cell death and inflammation. *Nat Rev Microbiol*, 7(2):99-109.
- [10] Heneka MT, Kummer MP, Latz E (2014). Innate immune activation in neurodegenerative disease. *Nat Rev Immunol*, 14(7):463-77.
- [11] Mamik MK, Power C (2017). Inflammasomes in neurological diseases: emerging pathogenic and therapeutic concepts. *Brain*, 140(9):2273-2285.

- [12] Voet S, Srinivasan S, Lamkanfi M, van Loo G (2019). Inflammasomes in neuroinflammatory and neurodegenerative diseases. *EMBO Mol Med*, 11(6):e10248.
- [13] de Rivero Vaccari JP, Dietrich WD, Keane RW (2014). Activation and regulation of cellular inflammasomes: gaps in our knowledge for central nervous system injury. *J Cereb Blood Flow Metab*, 34(3):369-75.
- [14] Xue Y, Enosi Tuipulotu D, Tan WH, Kay C, Man SM (2019). Emerging Activators and Regulators of Inflammasomes and Pyroptosis. *Trends Immunol*, 40(11):1035-1052.
- [15] Sonnessa M, Cioffi A, Brunetti O, Silvestris N, Zito FA, Saponaro C, et al. (2020). NLRP3 Inflammasome From Bench to Bedside: New Perspectives for Triple Negative Breast Cancer. *Front Oncol*, 10:1587.
- [16] Khare S, Luc N, Dorfleutner A, Stehlik C (2010). Inflammasomes and their activation. *Crit Rev Immunol*, 30(5):463-87.
- [17] Bruey JM, Bruey-Sedano N, Newman R, Chandler S, Stehlik C, Reed JC (2004). PAN1/NALP2/PYPAF2, an inducible inflammatory mediator that regulates NF-kappaB and caspase-1 activation in macrophages. *J Biol Chem*, 279(50):51897-907.
- [18] Fontalba A, Gutierrez O, Fernandez-Luna JL (2007). NLRP2, an inhibitor of the NF-kappaB pathway, is transcriptionally activated by NF-kappaB and exhibits a nonfunctional allelic variant. *J Immunol*, 179(12):8519-24.
- [19] Rossi MN, Pascarella A, Licursi V, Caiello I, Taranta A, Rega LR, et al. (2019). NLRP2 Regulates Proinflammatory and Antiapoptotic Responses in Proximal Tubular Epithelial Cells. *Front Cell Dev Biol*, 7:252.
- [20] Peng H, Chang B, Lu C, Su J, Wu Y, Lv P et al. (2012). *Nlrp2*, a maternal effect gene required for early embryonic development in the mouse. *PLoS One*, 7(1):e30344.
- [21] Kuchmiy AA, D'Hont J, Hochepped T, Lamkanfi M (2016). NLRP2 controls age-associated maternal fertility. *J Exp Med*, 213(13):2851-2860.
- [22] Huang JY, Su M, Lin SH, Kuo PL (2012). A genetic association study of NLRP2 and NLRP7 genes in idiopathic recurrent miscarriage. *Hum Reprod*, 28(4):1127-34.
- [23] Bhattacharjee P, Das N, Chatterjee D, Banerjee A, Das JK, Basu S, et al. (2013). Association of NALP2 polymorphism with arsenic induced skin lesions and other health effects. *Mutat Res*, 755(1):1-5.
- [24] Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Huerta-Cepas J, et al. (2019). STRING v11: protein-protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res*, 47(D1):D607-D613.
- [25] Tremblay MÈ (2020). A Diversity of Cell Types, Subtypes and Phenotypes in the Central Nervous System: The Importance of Studying Their Complex Relationships. *Front Cell Neurosci*, 14:628347.
- [26] Butt A, Verkhratsky A (2012). Neuroglia: Realising their true potential. *Brain Neurosci Adv*, 2:2398212818817495.
- [27] Dumas AA, Prinz M (2020). The myeloid side of the CNS. *Brain Pat*, 30(6):1158.
- [28] Cho MH, Cho K, Kang HJ, Jeon EY, Kim HS, Kwon HJ, et al. (2012). Autophagy in microglia degrades extracellular β -amyloid fibrils and regulates the NLRP3 inflammasome. *Autophagy*, 10(10):1761-75.
- [29] Denes A, Coutts G, Lénárt N, Cruickshank SM, Pelegrin P, Skinner J, et al. (2015). AIM2 and NLRC4 inflammasomes contribute with ASC to acute brain injury independently of NLRP3. *Proc Natl Acad Sci USA*, 112(13):4050-5.
- [30] Lu M, Sun XL, Qiao C, Liu Y, Ding JH, Hu G (2014). Uncoupling protein 2 deficiency aggravates astrocytic endoplasmic reticulum stress and nod-like receptor protein 3 inflammasome activation. *Neurobiol Aging*, 35(2):421-30.
- [31] Ducza L, Szücs P, Hegedűs K, Bakk E, Gajtkó A, Wéber I, et al. (2021). NLRP2 Is Overexpressed in Spinal Astrocytes at the Peak of Mechanical Pain Sensitivity during Complete Freund Adjuvant-Induced Persistent Pain. *Int J Mol Sci*, 22(21):11408.
- [32] Sun X, Song X, Zhang L, Sun J, Wei X, Meng L, et al. (2016). NLRP2 is highly expressed in a mouse model of ischemic stroke. *Biochem Biophys Res Commun*, 479(4):656-662.
- [33] Cheon SY, Kim EJ, Kim SY, Kim JM, Kam EH, Park JK, et al. (2018). Apoptosis Signal-regulating Kinase 1 Silencing on Astroglial Inflammasomes in an Experimental Model of Ischemic Stroke. *Neuroscience*, 390:218-230.
- [34] Zhang Q, Sun Y, He Z, Xu Y, Li X, Ding J, et al. (2020). Kynurenine regulates NLRP2 inflammasome in astrocytes and its implications in depression. *Brain Behav Immun*, 88:471-481.
- [35] Zhang M, Wu Y, Gao R, Chen X, Chen R, Chen Z (2022). Glucagon-like peptide-1 analogs mitigate neuroinflammation in Alzheimer's disease by suppressing NLRP2 activation in astrocytes. *Mol Cell Endocrinol*, 542:111529.
- [36] Matsuoka Y, Yamashita A, Matsuda M, Kawai K, Sawa T, Amaya F (2022). NLRP2 inflammasome in dorsal root ganglion as a novel molecular platform that produces inflammatory pain hypersensitivity. *Pain*, 160(9):2149-2160.
- [37] Truvé K, Parris TZ, Vizlin-Hodzic D, Salmela S, Berger E, Ågren H et al. (2020). Identification of candidate genetic variants and altered protein expression in neural stem and mature neural cells support altered microtubule function to be an essential component in bipolar disorder. *Transl Psychiatry*, 10(1):390.
- [38] Adamczak SE, de Rivero Vaccari JP, Dale G, Brand FJ 3rd, Nonner D, Bullock MR, et al. (2014). Pyroptotic neuronal cell death mediated by the AIM2 inflammasome. *J Cereb Blood Flow Metab*, 34(4):621-9.
- [39] Tan MS, Tan L, Jiang T, Zhu XC, Wang HF, Jia CD, et al. (2014). Amyloid- β induces NLRP1-dependent

- neuronal pyroptosis in models of Alzheimer's disease. *Cell Death Dis*, 5(8):e1382.
- [40] Zhang X, Wang R, Hu D, Sun X, Fujioka H, Lundberg K et al. (2020). Oligodendroglial glycolytic stress triggers inflammasome activation and neuropathology in Alzheimer's disease. *Sci Adv*, 6(49):eabb8680.
- [41] Yap JKY, Pickard BS, Chan EWL, Gan SY (2019). The Role of Neuronal NLRP1 Inflammasome in Alzheimer's Disease: Bringing Neurons into the Neuroinflammation Game. *Mol Neurobiol*, 56(11):7741-7753.
- [42] Mi L, Min X, Chai Y, Zhang J, Chen X (2022). NLRP1 Inflammasomes: A Potential Target for the Treatment of Several Types of Brain Injury. *Front Immunol*, 13:863774.
- [43] Saresella M, La Rosa F, Piancone F, Zoppis M, Marventano I, Calabrese E, et al. (2016). The NLRP3 and NLRP1 inflammasomes are activated in Alzheimer's disease. *Mol Neurodegener*, 11:23.
- [44] Song L, Pei L, Yao S, Wu Y, Shang Y (2017). NLRP3 Inflammasome in Neurological Diseases, from Functions to Therapies. *Front Cell Neurosci*, 11:63.
- [45] Vizlin-Hodzic D, Zhai Q, Illes S, Södersten K, Truvé K, Parris TZ, et al. (2017). Early onset of inflammation during ontogeny of bipolar disorder: the NLRP2 inflammasome gene distinctly differentiates between patients and healthy controls in the transition between iPS cell and neural stem cell stages. *Transl Psychiatry*, 7(1):e1010.
- [46] Chiarini A, Armato U, Gui L, Dal Prà I (2022). "Other Than NLRP3" Inflammasomes: Multiple Roles in Brain Disease. *Neuroscientist*, 11:10738584221106114.
- [47] Li Q, Tian Y, Wang ZF, Liu SB, Mi WL, Ma HJ, et al. (2013). Involvement of the spinal NALP1 inflammasome in neuropathic pain and aspirin-triggered-15-epi-lipoxin A4 induced analgesia. *Neuroscience*, 254:230-40.
- [48] Liu S, Li Q, Zhang MT, Mao-Ying QL, Hu LY, Wu GC, et al. (2016). Curcumin ameliorates neuropathic pain by down-regulating spinal IL-1 β via suppressing astroglial NALP1 inflammasome and JAK2-STAT3 signalling. *Sci Rep*, 6:28956.
- [49] He W, Long T, Pan Q, Zhang S, Zhang Y, Zhang D, et al. (2019). Microglial NLRP3 inflammasome activation mediates IL-1 β release and contributes to central sensitization in a recurrent nitroglycerin-induced migraine model. *J Neuroinflammation*, 16(1):78.
- [50] Grace PM, Strand KA, Galer EL, Urban DJ, Wang X, Baratta MV, et al. (2016). Morphine paradoxically prolongs neuropathic pain in rats by amplifying spinal NLRP3 inflammasome activation. *Proc Natl Acad Sci USA*, 113(24):E3441-50.
- [51] Chiarini A, Gui L, Viviani C, Armato U, Dal Prà I (2023). NLRP3 Inflammasome's Activation in Acute and Chronic Brain Diseases-An Update on Pathogenetic Mechanisms and Therapeutic Perspectives with Respect to Other Inflammasomes. *Biomedicines*, 11(4):999.
- [52] Marton N, Baricza E, Érsek B, Buzás EI, Nagy G (2015). The Emerging and Diverse Roles of Src-Like Adaptor Proteins in Health and Disease. *Mediators Inflamm*, 2015:952536.
- [53] Ge MM, Zhou YQ, Tian XB, Manyande A, Tian YK, Ye DW et al. (2020). Src-family protein tyrosine kinases: A promising target for treating chronic pain. *Biomed Pharmacother*, 125:110017.
- [54] Yang H, Wang L, Zang C, Wang Y, Shang J, Zhang Z, et al. (2020). Src Inhibition Attenuates Neuroinflammation and Protects Dopaminergic Neurons in Parkinson's Disease Models. *Front Neurosci*, 14:45.
- [55] Giraud F, Pereira E, Anizon F, Moreau P (2021). Recent Advances in Pain Management: Relevant Protein Kinases and Their Inhibitors. *Molecules*, 26(9):2696.
- [56] Tran S, Fairlie WD, Lee EF (2021). BECLIN1: Protein Structure, Function and Regulation. *Cells*, 10(6):1522.
- [57] Kang R, Zeh HJ, Lotze MT, Tang D (2011). The Beclin 1 network regulates autophagy and apoptosis. *Cell Death Differ*, 18(4):571-80.
- [58] Zhang D, Wang W, Sun X, Xu D, Wang C, Zhang Q, et al. (2016). AMPK regulates autophagy by phosphorylating BECN1 at threonine 388. *Autophagy*, 12(9):1447-59.
- [59] Lucin KM, O'Brien CE, Bieri G, Czirr E, Mosher KI, Abbey RJ, et al. (2013). Microglial beclin 1 regulates retromer trafficking and phagocytosis and is impaired in Alzheimer's disease. *Neuron*, 79(5):873-86.
- [60] Heppner FL, Ransohoff RM, Becher B (2015). Immune attack: the role of inflammation in Alzheimer disease. *Nat Rev Neurosci*, 16(6):358-72.
- [61] Houtman J, Freitag K, Gimber N, Schmoranz J, Heppner FL, Jendrach M (2019). Beclin1-driven autophagy modulates the inflammatory response of microglia via NLRP3. *EMBO J*, 38(4):e99430.
- [62] Brattås PL, Hersbach BA, Madsen S, Petri R, Jakobsson J, Pircs K (2021). Impact of differential and time-dependent autophagy activation on therapeutic efficacy in a model of Huntington disease. *Autophagy*, 17(6):1316-1329.
- [63] Peek SL, Mah KM, Weiner JA (2017). Regulation of neural circuit formation by protocadherins. *Cell Mol Life Sci*, 74(22):4133-4157.
- [64] Langfelder P, Cattle JP, Chatzopoulou D, Wang N, Gao F, Al-Ramahi (2016). Integrated genomics and proteomics define huntingtin CAG length-dependent networks in mice. *Nat Neurosci*, 19(4):623-33.
- [65] Carrasquillo MM, Zou F, Pankratz VS, Wilcox SL, Ma L, Walker LP (2019). Genetic variation in PCDH11X is associated with susceptibility to late-onset Alzheimer's disease. *Nat Genet*, 41(2):192-8.
- [66] Beecham GW, Naj AC, Gilbert JR, Haines JL, Buxbaum JD, Pericak-Vance MA (2010). PCDH11X variation is not associated with late-onset Alzheimer disease susceptibility. *Psychiatr Genet*, 20(6):321-4.
- [67] Miar A, Alvarez V, Corao AI, Alonso B, Díaz M, Menéndez M (2011). Lack of association between protocadherin 11-X/Y (PCDH11X and PCDH11Y) polymorphisms and late onset Alzheimer's disease. *Brain Res*, 1383:252-6.
- [68] Marr L, Biswas D, Daly LA, Browning C, Vial SCM, Maskell DP (2022). Mechanism of glycogen synthase inactivation and interaction with glycogenin. *Nat Commun*, 13(1):3372.

- [69] Fastman NM, Liu Y, Ramanan V, Merritt H, Ambing E, DePaoli-Roach AA (2022). The structural mechanism of human glycogen synthesis by the GYS1-GYG1 complex. *Cell Rep*, 40(1):111041.
- [70] Imagawa E, Osaka H, Yamashita A, Shiina M, Takahashi E, Sugie H (2014). A hemizygous GYG2 mutation and Leigh syndrome: a possible link? *Hum Genet*, 133(2):225-34.
- [71] Duran J, Hervera A, Markussen KH, Varea O, López-Soldado I, Sun RC, et al. (2021). Astrocytic glycogen accumulation drives the pathophysiology of neurodegeneration in Lafora disease. *Brain*, 144(8):2349-2360.
- [72] Sayas CL, Ávila Jb (2021). GSK-3 and Tau: A Key Duet in Alzheimer's Disease. *Cells*, 10(4):721.
- [73] Leroy K, Yilmaz Z, Brion JP (2007). Increased level of active GSK-3beta in Alzheimer's disease and accumulation in argyrophilic grains and in neurones at different stages of neurofibrillary degeneration. *Neuropathol Appl Neurobiol*, 33(1):43-55.
- [74] Ahmad S, Orellana A, Kohler I, Frölich L, de Rojas I, Gil S (2020). Association of lysophosphatidic acids with cerebrospinal fluid biomarkers and progression to Alzheimer's disease. *Alzheimers Res Ther*, 12(1):124.
- [75] Madabhushi R, Pan L, Tsai LH (2014). DNA damage and its links to neurodegeneration. *Neuron*, 83(2):266-282.
- [76] Kannan A, Bhatia K, Branzei D, Gangwani L (2018). Combined deficiency of Senataxin and DNA-PKcs causes DNA damage accumulation and neurodegeneration in spinal muscular atrophy. *Nucleic Acids Res*, 46(16):8326-8346.
- [77] Suberbielle E, Sanchez PE, Kravitz AV, Wang X, Ho K, Eilertson K (2013). Physiologic brain activity causes DNA double-strand breaks in neurons, with exacerbation by amyloid- β . *Nat Neurosci*, 16(5):613-21.
- [78] Thadathil N, Hori R, Xiao J, Khan MM (2019). DNA double-strand breaks: a potential therapeutic target for neurodegenerative diseases. *Chromosome Res*, 27(4):345-364.
- [79] Lu R, Zhang H, Jiang YN, Wang ZQ, Sun L, Zhou ZW (2021). Post-Translational Modification of MRE11: Its Implication in DDR and Diseases. *Genes (Basel)*, 12(8):1158.
- [80] Miyamoto R, Morino H, Yoshizawa A, Miyazaki Y, Maruyama H, Murakami N, et al. (2014). Exome sequencing reveals a novel MRE11 mutation in a patient with progressive myoclonic ataxia. *J Neurol Sci*, 337(1-2):219-23.
- [81] Matsumoto Y, Miyamoto T, Sakamoto H, Izumi H, Nakazawa Y, Ogi T (2011). Two unrelated patients with MRE11A mutations and Nijmegen breakage syndrome-like severe microcephaly. *DNA Repair (Amst)*, 10(3):314-21.
- [82] Sedghi M, Salari M, Moslemi AR, Kariminejad A, Davis M, Goullée H, et al. (2018). Ataxia-telangiectasia-like disorder in a family deficient for MRE11A, caused by a *MRE11* variant. *Neurol Genet*, 4(6):e295.
- [83] Ding M, Qing X, Zhang G, Baade-Büttner C, Gruber R, Lu H (2022). The Essential DNA Damage Response Complex MRN Is Dispensable for the Survival and Function of Purkinje Neurons. *Front Aging Neurosci*, 13:786199.
- [84] Jacobsen E, Beach T, Shen Y, Li R, Chang Y (2004). Deficiency of the Mre11 DNA repair complex in Alzheimer's disease brains. *Brain Res Mol Brain Res*, 128(1):1-7.
- [85] Kitagawa K, Skowrya D, Elledge SJ, Harper JW, Hieter P (1999). SGT1 encodes an essential component of the yeast kinetochore assembly pathway and a novel subunit of the SCF ubiquitin ligase complex. *Mol Cell*, 4(1):21-33.
- [86] Willhoft O, Kerr R, Patel D, Zhang W, Al-Jassar C, Daviter T (2017). The crystal structure of the Sgt1-Skp1 complex: the link between Hsp90 and both SCF E3 ubiquitin ligases and kinetochores. *Sci Rep*, 7:41626.
- [87] Lee YT, Jacob J, Michowski W, Nowotny M, Kuznicki J, Chazin WJ (2004). Human Sgt1 binds HSP90 through the CHORD-Sgt1 domain and not the tetratricopeptide repeat domain. *J Biol Chem*, 279(16):16511-7.
- [88] Eisele F, Eisele-Bürger AM, Hao X, Berglund LL, Höög JL, Liu B (2021). An Hsp90 co-chaperone links protein folding and degradation and is part of a conserved protein quality control. *Cell Rep*, 35(13):109328.
- [89] da Silva Correia J, Miranda Y, Leonard N, Ulevitch R (2007). SGT1 is essential for Nod1 activation. *Proc Natl Acad Sci USA*, 104(16):6764-9.
- [90] Bohush A, Góral A, Sierant M, Nawrot B, Leśniak W, Filipek A (2021). Sgt1 Regulates α -Synuclein Subcellular Localization and Expression of Parkinson's Disease Related Genes, *PINK1* and *PARK9*. *Biomolecules*, 11(11):1675.
- [91] Bohush A, Niewiadomska G, Weis S, Filipek A (2019). HSP90 and Its Novel Co-Chaperones, SGT1 and CHP-1, in Brain of Patients with Parkinson's Disease and Dementia with Lewy Bodies. *J Parkinsons Dis*, 9(1):97-107.
- [92] Spiechowicz M, Bernstein HG, Dobrowolny H, Leśniak W, Mawrin C, Bogerts B, et al. (2006). Density of Sgt1-immunopositive neurons is decreased in the cerebral cortex of Alzheimer's disease brain. *Neurochem Int*, 49(5):487-93.
- [93] Fazioli F, Minichiello L, Matoskova B, Wong WT, Di Fiore PP (1993). eps15, a novel tyrosine kinase substrate, exhibits transforming activity. *Mol Cell Biol*, 13(9):5814-28.
- [94] Benmerah A, Gagnon J, Bègue B, Mégarbané B, Dautry-Varsat A, Cerf-Bensussan N (1995). The tyrosine kinase substrate eps15 is constitutively associated with the plasma membrane adaptor AP-2. *J Cell Biol*, 131(6 Pt 2):1831-8.
- [95] Carbone R, Fré S, Iannolo G, Belleudi F, Mancini P, Pelicci PG, et al. (1997). eps15 and eps15R are essential components of the endocytic pathway. *Cancer Res*, 57(24):5498-504.
- [96] Huang F, Khvorova A, Marshall W, Sorkin A (2004). Analysis of clathrin-mediated endocytosis of epidermal growth factor receptor by RNA interference. *J Biol Chem*, 279(16):16657-61.

- [97] van Bergen En Henegouwen PM (2009). Eps15: a multifunctional adaptor protein regulating intracellular trafficking. *Cell Commun Signal*, 7:24.
- [98] Iwakura Y, Piao YS, Mizuno M, Takei N, Kakita A, Takahashi H, et al. (2005). Influences of dopaminergic lesion on epidermal growth factor-ErbB signals in Parkinson's disease and its model: neurotrophic implication in nigrostriatal neurons. *J Neurochem*, 93(4):974-83.
- [99] Atkin G, Paulson H (2014). Ubiquitin pathways in neurodegenerative disease. *Front Mol Neurosci*, 7:63.
- [100] Conway JA, Kinsman G, Kramer ER (2022). The Role of NEDD4 E3 Ubiquitin-Protein Ligases in Parkinson's Disease. *Genes (Basel)*, 13(3):513.
- [101] Dokucu ME, Zipursky SL, Cagan RL (1996). Atonal, rough and the resolution of proneural clusters in the developing *Drosophila* retina. *Development*, 122(12):4139-47.
- [102] Sommer L, Ma Q, Anderson DJ (1996). neurogenins, a novel family of atonal-related bHLH transcription factors, are putative mammalian neuronal determination genes that reveal progenitor cell heterogeneity in the developing CNS and PNS. *Mol Cell Neurosci*, 8(4):221-41.
- [103] Bertrand N, Castro DS, Guillemot F (2002). Proneural genes and the specification of neural cell types. *Nat Rev Neurosci*, 3(7):517-30.
- [104] Tutukova S, Tarabykin V, Hernandez-Miranda LR (2002). The Role of NeuroD Genes in Brain Development, Function, and Disease. *Front Mol Neurosci*, 14:662774.
- [105] Schwab MH, Druffel-Augustin S, Gass P, Jung M, Klugmann M, Bartholomae A, et al. (1998). Neuronal basic helix-loop-helix proteins (NEX, neuroD, NDRF): spatiotemporal expression and targeted disruption of the NEX gene in transgenic mice. *J Neurosci*, 18(4):1408-18.
- [106] Boulle F, Massart R, Stragier E, Païzani E, Zaidan L, Marday S, et al. (2014). Hippocampal and behavioral dysfunctions in a mouse model of environmental stress: normalization by agomelatine. *Transl Psychiatry*, 4(11):e485.
- [107] Rubio-Cabezas O, Minton JA, Kantor I, Williams D, Ellard S, Hattersley AT (2010). Homozygous mutations in *NEUROD1* are responsible for a novel syndrome of permanent neonatal diabetes and neurological abnormalities. *Diabetes*, 59(9):2326-31.
- [108] Liu MH, Li W, Zheng JJ, Xu YG, He Q, Chen G (2020). Differential neuronal reprogramming induced by *NeuroD1* from astrocytes in grey matter *versus* white matter. *Neural Regen Res*, 15(2):342-351.
- [109] Fedele V, Roybon L, Nordström U, Li JY, Brundin P (2011). Neurogenesis in the R6/2 mouse model of Huntington's disease is impaired at the level of *NeuroD1*. *Neuroscience*, 173:76-81.
- [110] Satoh J, Yamamoto Y, Asahina N, Kitano S, Kino Y (2014). RNA-Seq data mining: downregulation of *NeuroD6* serves as a possible biomarker for Alzheimer's disease brains. *Dis Markers*, 2014:123165.
- [111] Richetin K, Leclerc C, Toni N, Gallopin T, Pech S, Roybon L, et al. (2015). Genetic manipulation of adult-born hippocampal neurons rescues memory in a mouse model of Alzheimer's disease. *Brain*, 138(Pt 2):440-55.
- [112] Lee TY, Cho IS, Bashyal N, Naya FJ, Tsai MJ, Yoon JS, et al. (2020). ERK Regulates *NeuroD1*-mediated Neurite Outgrowth via Proteasomal Degradation. *Exp Neurobiol*, 29(3):189-206.
- [113] Pomeschik Y, Klementieva O, Gil J, Martinsson I, Hansen MG, de Vries T, et al. (2020). Human iPSC-Derived Hippocampal Spheroids: An Innovative Tool for Stratifying Alzheimer Disease Patient-Specific Cellular Phenotypes and Developing Therapies. *Stem Cell Reports*, 15(1):256.
- [114] Guo Z, Zhang L, Wu Z, Chen Y, Wang F, Chen G (2014). In vivo direct reprogramming of reactive glial cells into functional neurons after brain injury and in an Alzheimer's disease model. *Cell Stem Cell*, 14(2):188-202.
- [115] Ge LJ, Yang FH, Li W, Wang T, Lin Y, Feng J, et al. (2020). *In vivo* Neuroregeneration to Treat Ischemic Stroke Through *NeuroD1* AAV-Based Gene Therapy in Adult Non-human Primates. *Front Cell Dev Biol*, 8:590008.
- [116] Sorimachi H, Hata S, Ono Y (2011). Calpain chronicle--an enzyme family under multidisciplinary characterization. *Proc Jpn Acad Ser B Phys Biol Sci*, 87(6):287-327.
- [117] Chen L, Xiao D, Tang F, Gao H, Li X (2020). CAPN6 in disease: An emerging therapeutic target (Review). *Int J Mol Med*, 46(5):1644-1652.
- [118] Dear N, Matena K, Vingron M, Boehm T (1997). A new subfamily of vertebrate calpains lacking a calmodulin-like domain: implications for calpain regulation and evolution. *Genomics*, 45(1):175-84.
- [119] Dear TN, Boehm T (1999). Diverse mRNA expression patterns of the mouse calpain genes *Capn5*, *Capn6* and *Capn11* during development. *Mech Dev*, 89(1-2):201-9.
- [120] Moretti D, Del Bello B, Allavena G, Maellaro E (2014). Calpains and cancer: friends or enemies? *Arch Biochem Biophys*, 564:26-36.
- [121] Ono Y, Ojima K, Shinkai-Ouchi F, Hata S, Sorimachi H (2016). An eccentric calpain, *CAPN3/p94/calpain-3*. *Biochimie*, 122:169-87.
- [122] Filali H, Vidal E, Bolea R, Márquez M, Marco P, Vargas A, et al. (2013). Gene and protein patterns of potential prion-related markers in the central nervous system of clinical and preclinical infected sheep. *Vet Res*, 44(1):14.
- [123] Su X, Xiao D, Huang L, Li S, Ying J, Tong Y, et al. (2019). MicroRNA Alteration in Developing Rat Oligodendrocyte Precursor Cells Induced by Hypoxia-Ischemia. *J Neuropathol Exp Neurol*, 78(10):900-909.
- [124] Medeiros R, Kitazawa M, Chabrier MA, Cheng D, Baglietto-Vargas D, Kling A, et al. (2012). Calpain inhibitor A-705253 mitigates Alzheimer's disease-like pathology and cognitive decline in aged 3xTgAD mice. *Am J Pathol*, 181(2):616-25.
- [125] Mahaman YAR, Huang F, Kessete Afewerky H, Maibouge TMS, Ghose B, Wang X (2019). Involvement

- of calpain in the neuropathogenesis of Alzheimer's disease. *Med Res Rev*, 39(2):608-630.
- [126] Nogami S, Satoh S, Nakano M, Shimizu H, Fukushima H, Maruyama A, et al. (2003). Taxilin; a novel syntaxin-binding protein that is involved in Ca²⁺-dependent exocytosis in neuroendocrine cells. *Genes Cells*, 8(1):17-28.
- [127] Nogami S, Satoh S, Tanaka-Nakadate S, Yoshida K, Nakano M, Terano A, et al. (2004). Identification and characterization of taxilin isoforms. *Biochem Biophys Res Commun*, 319(3):936-43.
- [128] Horii Y, Sakane H, Nogami S, Ohtomo N, Tomiya T, Shirataki H (2014). Expression of α -taxilin in the murine gastrointestinal tract: potential implication in cell proliferation. *Histochem Cell Biol*, 141(2):165-80.
- [129] Sakane H, Makiyama T, Nogami S, Horii Y, Akasaki K, Shirataki H (2016). β -Taxilin participates in differentiation of C2C12 myoblasts into myotubes. *Exp Cell Res*, 345(2):230-8.
- [130] Makiyama T, Higashi S, Sakane H, Nogami S, Shirataki H (2018). γ -Taxilin temporally regulates centrosome disjunction in a Nek2A-dependent manner. *Exp Cell Res*, 362(2):412-423.
- [131] Hotokezaka Y, Katayama I, van Leyen K, Nakamura T (2015). GSK-3 β -dependent downregulation of γ -taxilin and α NAC merge to regulate ER stress responses. *Cell Death Dis*, 6(4):e1719.
- [132] Ma D, Wang F, Wang R, Hu Y, Chen Z, Huang N, et al. (2022). α -/ γ -Taxilin are required for centriolar subdistal appendage assembly and microtubule organization. *Elife*, 11:e73252.
- [133] Higashi S, Makiyama T, Sakane H, Nogami S, Shirataki H (2022). Regulation of Hook1-mediated endosomal sorting of clathrin-independent cargo by γ -taxilin. *J Cell Sci*, 135(1):jcs258849.
- [134] Kornau HC, Schenker LT, Kennedy MB, Seeburg PH (1995). Domain interaction between NMDA receptor subunits and the postsynaptic density protein PSD-95. *Science*, 269(5231):1737-40.
- [135] Naisbitt S, Kim E, Tu JC, Xiao B, Sala C, Valtschanoff J, et al. (1999). Shank, a novel family of postsynaptic density proteins that binds to the NMDA receptor/PSD-95/GKAP complex and cortactin. *Neuron*, 23(3):569-82.
- [136] Chen L, Chetkovich DM, Petralia RS, Sweeney NT, Kawasaki Y, Wenthold RJ (2000). Stargazin regulates synaptic targeting of AMPA receptors by two distinct mechanisms. *Nature*, 408(6815):936-43.
- [137] Steiner P, Higley MJ, Xu W, Czervionke BL, Malenka RC, Sabatini BL (2008). Destabilization of the postsynaptic density by PSD-95 serine 73 phosphorylation inhibits spine growth and synaptic plasticity. *Neuron*, 60(5):788-802.
- [138] Naisbitt S, Kim E, Weinberg RJ, Rao A, Yang FC, Craig AM (1997). Characterization of guanylate kinase-associated protein, a postsynaptic density protein at excitatory synapses that interacts directly with postsynaptic density-95/synapse-associated protein 90. *J Neurosci*, 17(15):5687-96.
- [139] Welch JM, Wang D, Feng G (2004). Differential mRNA expression and protein localization of the SAP90/PSD-95-associated proteins (SAPAPs) in the nervous system of the mouse. *J Comp Neurol*, 472(1):24-39.
- [140] Yao I, Iida J, Nishimura W, Hata Y (2003). Synaptic localization of SAPAP1, a synaptic membrane-associated protein. *Genes Cells*, 8(2):121-9.
- [141] Rasmussen AH, Rasmussen HB, Silahatoglu A (2017). The DLGAP family: neuronal expression, function and role in brain disorders. *Mol Brain*, 10(1):43.
- [142] Jiang-Xie LF, Liao HM, Chen CH, Chen YT, Ho SY, Lu DH (2014). Autism-associated gene *Dlgap2* mutant mice demonstrate exacerbated aggressive behaviors and orbitofrontal cortex deficits. *Mol Autism*, 5:32.
- [143] Li JM, Lu CL, Cheng MC, Luu SU, Hsu SH, Hu TM (2014). Role of the *DLGAP2* gene encoding the SAP90/PSD-95-associated protein 2 in schizophrenia. *PLoS One*, 9(1):e85373.
- [144] Gilbertson MW, Shenton ME, Ciszewski A, Kasai K, Lasko NB, Orr SP, et al. (2002). Smaller hippocampal volume predicts pathological vulnerability to psychological trauma. *Nat Neurosci*, 5(11):1242-7.
- [145] Ouellette AR, Neuner SM, Dumitrescu L, Anderson LC, Gatti DM, Mahoney ER, et al. (2020). Cross-Species Analyses Identify *Dlgap2* as a Regulator of Age-Related Cognitive Decline and Alzheimer's Dementia. *Cell Rep*, 32(9):108091.
- [146] O'Dowd BF, Heiber M, Chan A, Heng HH, Tsui LC, Kennedy JL, et al. (1993). A human gene that shows identity with the gene encoding the angiotensin receptor is located on chromosome 11. *Gene*, 136(1-2):355-60.
- [147] Hu G, Wang Z, Zhang R, Sun W, Chen X (2021). The Role of Apelin/Apelin Receptor in Energy Metabolism and Water Homeostasis: A Comprehensive Narrative Review. *Front Physiol*, 12:632886.
- [148] Luo H, Han L, Xu J (2021). Apelin/APJ system: A novel promising target for neurodegenerative diseases. *J Cell Physiol*, 235(2):638-657.
- [149] Pi J, Cheng Y, Sun H, Chen X, Zhuang T, Liu J, et al. (2017). *Apln-CreERT:mT/mG* reporter mice as a tool for sprouting angiogenesis study. *BMC Ophthalmol*, 17(1):163.
- [150] Mughal A, O'Rourke ST (2018). Vascular effects of apelin: Mechanisms and therapeutic potential. *Pharmacol Ther*, 190:139-147.
- [151] Tatemoto K, Takayama K, Zou MX, Kumaki I, Zhang W, Kumano K, et al. (2001). The novel peptide apelin lowers blood pressure via a nitric oxide-dependent mechanism. *Regul Pept*, 99(2-3):87-92.
- [152] O'Carroll AM, Lolait SJ, Harris LE, Pope GR (2013). The apelin receptor APJ: journey from an orphan to a multifaceted regulator of homeostasis. *J Endocrinol*, 219(1):R13-35.
- [153] Kazemi F, Zahediasl S (2018). Effects of exercise training on adipose tissue apelin expression in streptozotocin-nicotinamide induced diabetic rats. *Gene*, 662:97-102.
- [154] Ashley E, Chun HJ, Quertermous T (2016). Opposing cardiovascular roles for the angiotensin and apelin signaling pathways. *J Mol Cell Cardiol*, 41(5):778-81.
- [155] Taheri S, Murphy K, Cohen M, Sujkovic E, Kennedy A, Dhillon W, et al. (2002). The effects of centrally

- administered apelin-13 on food intake, water intake and pituitary hormone release in rats. *Biochem Biophys Res Commun*, 291(5):1208-12.
- [156] Zeng XJ, Yu SP, Zhang L, Wei L (2010). Neuroprotective effect of the endogenous neural peptide apelin in cultured mouse cortical neurons. *Exp Cell Res*, 316(11):1773-83.
- [157] O'Donnell LA, Agrawal A, Sabnekar P, Dichter MA, Lynch DR, Kolson DL (2007). Apelin, an endogenous neuronal peptide, protects hippocampal neurons against excitotoxic injury. *J Neurochem*, 102(6):1905-1917.
- [158] Aminyavari S, Zahmatkesh M, Farahmandfar M, Khodaghali F, Dargahi L, Zarrindast MR (2019). Protective role of Apelin-13 on amyloid β 25-35-induced memory deficit; Involvement of autophagy and apoptosis process. *Prog Neuropsychopharmacol Biol Psychiatry*, 89:322-334.
- [159] Agostinho P, Cunha RA, Oliveira C (2010). Neuroinflammation, oxidative stress and the pathogenesis of Alzheimer's disease. *Curr Pharm Des*, 16(25):2766-78.
- [160] Xin Q, Cheng B, Pan Y, Liu H, Yang C, Chen J (2015): Neuroprotective effects of apelin-13 on experimental ischemic stroke through suppression of inflammation. *Peptides*, 63:55-62.
- [161] Yang L, Wang H, Liu L, Xie A (2018). The Role of Insulin/IGF-1/PI3K/Akt/GSK3 β Signaling in Parkinson's Disease Dementia. *Front Neurosci*, 12:73.
- [162] Haghparast E, Esmaeili-Mahani S, Abbasnejad M, Sheibani V (2018). Apelin-13 ameliorates cognitive impairments in 6-hydroxydopamine-induced substantia nigra lesion in rats. *Neuropeptides*, 68:28-35.
- [163] Poursmaeili-Babaki E, Esmaeili-Mahani S, Abbasnejad M, Ravan H (2018). Protective Effect of Neuropeptide Apelin-13 on 6-Hydroxydopamine-Induced Neurotoxicity in SH-SY5Y Dopaminergic Cells: Involvement of Its Antioxidant and Antiapoptotic Properties. *Rejuvenation Res*, 21(2):162-167.
- [164] Chi Y, Chai J, Xu C, Luo H, Zhang Q (2015). Apelin inhibits the activation of the nucleotide-binding domain and the leucine-rich, repeat-containing family, pyrin-containing 3 (NLRP3) inflammasome and ameliorates insulin resistance in severely burned rats. *Surgery*, 157(6):1142-52.
- [165] Fernández-Nogales M, Santos-Galindo M, Hernández IH, Cabrera JR, Lucas JJ (2016). Faulty splicing and cytoskeleton abnormalities in Huntington's disease. *Brain Pathol*, 26(6):772-778.
- [166] Li Y, Bai YJ, Jiang YR, Yu WZ, Shi X, Chen L, et al. (2018). Apelin-13 Is an Early Promoter of Cytoskeleton and Tight Junction in Diabetic Macular Edema via PI-3K/Akt and MAPK/Erk Signaling Pathways. *Biomed Res Int*, 2018:3242574.
- [167] Bryan MR, Bowman AB (2017). Manganese and the Insulin-IGF Signaling Network in Huntington's Disease and Other Neurodegenerative Disorders. *Adv Neurobiol*, 18:113-142.
- [168] Zhang Y, Zhang Y, Sun K, Meng Z, Chen L (2019). The SLC transporter in nutrient and metabolic sensing, regulation, and drug development. *J Mol Cell Biol*, 11(1):1-13.
- [169] César-Razquin A, Snijder B, Frappier-Brinton T, Isserlin R, Gyimesi G, Bai X (2015). A Call for Systematic Research on Solute Carriers. *Cell*, 162(3):478-87.
- [170] Brzica H, Abdullahi W, Ibbotson K, Ronaldson PT (2017). Role of Transporters in Central Nervous System Drug Delivery and Blood-Brain Barrier Protection: Relevance to Treatment of Stroke. *J Cent Nerv Syst Dis*, 9:1179573517693802.
- [171] Türková A, Zdrzil B (2019). Current Advances in Studying Clinically Relevant Transporters of the Solute Carrier (SLC) Family by Connecting Computational Modeling and Data Science. *Comput Struct Biotechnol J*, 17:390-405.
- [172] Hu C, Tao L, Cao X, Chen L (2020). The solute carrier transporters and the brain: Physiological and pharmacological implications. *Asian J Pharm Sci*, 15(2):131-144.
- [173] Young JD, Yao SY, Baldwin JM, Cass CE, Baldwin SA (2013). The human concentrative and equilibrative nucleoside transporter families, SLC28 and SLC29. *Mol Aspects Med*, 34(2-3):529-47.
- [174] Pastor-Anglada M, Urtasun N, Pérez-Torras S (2018). Intestinal Nucleoside Transporters: Function, Expression, and Regulation. *Compr Physiol*, 8(3):1003-1017.
- [175] Kao YH, Lin MS, Chen CM, Wu YR, Chen HM, Lai HL (2017). Targeting ENT1 and adenosine tone for the treatment of Huntington's disease. *Hum Mol Genet*, 26(3):467-478.
- [176] Lee CC, Chang CP, Lin CJ, Lai HL, Kao YH, Cheng SJ, et al. (2018). Adenosine Augmentation Evoked by an ENT1 Inhibitor Improves Memory Impairment and Neuronal Plasticity in the APP/PS1 Mouse Model of Alzheimer's Disease. *Mol Neurobiol*, 55(12):8936-8952.
- [177] Chang CP, Chang YG, Chuang PY, Nguyen TNA, Wu KC, Chou FY, et al. (2021). Equilibrative nucleoside transporter 1 inhibition rescues energy dysfunction and pathology in a model of tauopathy. *Acta Neuropathol Commun*, 9(1):112.
- [178] Chung JY, Yu SD, Hong YS (2014). Environmental source of arsenic exposure. *J Prev Med Public Health*, 47(5):253-7.
- [179] Ahmad SA, Khan MH, Haque M (2018). Arsenic contamination in groundwater in Bangladesh: implications and challenges for healthcare policy. *Risk Manag Healthc Policy*, 11:251-261.
- [180] Emadi A, Gore SD (2010). Arsenic trioxide - An old drug rediscovered. *Blood Rev*, 24(4-5):191-9.
- [181] Singh AP, Goel RK, Kaur T (2011). Mechanisms pertaining to arsenic toxicity. *Toxicol Int*, 18(2):87-93.
- [182] Rahman MA, Hannan MA, Uddin MJ, Rahman MS, Rashid MM, Kim B (2021). Exposure to Environmental Arsenic and Emerging Risk of Alzheimer's Disease: Perspective Mechanisms, Management Strategy, and Future Directions. *Toxics*, 9(8):188.
- [183] Chandravanshi LP, Gupta R, Shukla RK (2018). Developmental Neurotoxicity of Arsenic: Involvement

- of Oxidative Stress and Mitochondrial Functions. *Biol Trace Elem Res*, 186(1):185-198.
- [184] Hannan MA, Dash R, Sohag AAM, Haque MN, Moon IS (2020). Neuroprotection Against Oxidative Stress: Phytochemicals Targeting TrkB Signaling and the Nrf2-ARE Antioxidant System. *Front Mol Neurosci*, 13:116.
- [185] Dwivedi N, Flora SJ (2011). Concomitant exposure to arsenic and organophosphates on tissue oxidative stress in rats. *Food Chem Toxicol*, 49(5):1152-9.
- [186] Dash R, Mitra S, Ali MC, Oktaviani DF, Hannan MA, Choi SM, et al. (2021). Phytosterols: Targeting Neuroinflammation in Neurodegeneration. *Curr Pharm Des*, 27(3):383-401.
- [187] Wisessaowapak C, Visitnonthachai D, Watcharasi P, Satayavivad J (2021). Prolonged arsenic exposure increases tau phosphorylation in differentiated SH-SY5Y cells: The contribution of GSK3 and ERK1/2. *Environ Toxicol Pharmacol*, 84:103626.
- [188] Medda N, Patra R, Ghosh TK, Maiti S (2020). Neurotoxic Mechanism of Arsenic: Synergistic Effect of Mitochondrial Instability, Oxidative Stress, and Hormonal-Neurotransmitter Impairment. *Biol Trace Elem Res*, 198(1):8-15.
- [189] Prakash C, Soni M, Kumar V (2016). Mitochondrial oxidative stress and dysfunction in arsenic neurotoxicity: A review. *J Appl Toxicol*, 36(2):179-88.
- [190] Prakash C, Soni M, Kumar V (2015). Biochemical and Molecular Alterations Following Arsenic-Induced Oxidative Stress and Mitochondrial Dysfunction in Rat Brain. *Biol Trace Elem Res*, 167(1):121-9.
- [191] Rahman MA, Rahman MS, Uddin MJ, Mamun-Or-Rashid ANM, Pang MG, Rhim H (2020). Emerging risk of environmental factors: insight mechanisms of Alzheimer's diseases. *Environ Sci Pollut Res Int*, 27(36):44659-44672.
- [192] Zhang W, Cui X, Gao Y, Sun L, Wang J, Yang Y, et al. (2019). Role of pigment epithelium-derived factor (PEDF) on arsenic-induced neuronal apoptosis. *Chemosphere*, 215:925-931.
- [193] Dash R, Jahan I, Ali MC, Mitra S, Munni YA, Timalina B, et al. (2021). Potential roles of natural products in the targeting of proteinopathic neurodegenerative diseases. *Neurochem Int*, 145:105011.
- [194] Dash R, Ali MC, Jahan I, Munni YA, Mitra S, Hannan MA, et al. (2021). Emerging potential of cannabidiol in reversing proteinopathies. *Ageing Res Rev*, 65:101209.
- [195] Mizushima N, Komatsu M (2011). Autophagy: renovation of cells and tissues. *Cell*, 147(4):728-41.
- [196] Levine B, Mizushima N, Virgin HW (2011). Autophagy in immunity and inflammation. *Nature*, 469(7330):323-35.
- [197] Vazza G, Picelli S, Bozzato A, Mostacciolo ML (2003). Identification and characterization of C3orf6, a new conserved human gene mapping to chromosome 3q28. *Gene*, 314:113-20.
- [198] Lin Y, Li Z, Wang Y, Tian T, Jia P, Ye Y, et al. (2022). CCDC50 suppresses NLRP3 inflammasome activity by mediating autophagic degradation of NLRP3. *EMBO Rep*, 23(5):e54453.
- [199] Farfing A, Engel F, Seiffert M, Hartmann E, Ott G, Rosenwald A, et al. (2009). Gene knockdown studies revealed CCDC50 as a candidate gene in mantle cell lymphoma and chronic lymphocytic leukemia. *Leukemia*, 23(11):2018-26.
- [200] Hou P, Yang K, Jia P, Liu L, Lin Y, Li Z, et al. (2021). A novel selective autophagy receptor, CCDC50, delivers K63 polyubiquitination-activated RIG-I/MDA5 for degradation during viral infection. *Cell Res*, 31(1):62-79.
- [201] Cruchaga C, Kauwe JS, Harari O, Jin SC, Cai Y, Karch CM, et al. (2013). GWAS of cerebrospinal fluid tau levels identifies risk variants for Alzheimer's disease. *Neuron*, 78(2):256-68.
- [202] Skaria AP (2022). The economic and societal burden of Alzheimer disease: managed care considerations. *Am J Manag Care*, 28(10 Suppl):S188-S196.
- [203] Wong W (2020). Economic burden of Alzheimer disease and managed care considerations. *Am J Manag Care*, 26(8 Suppl):S177-S183.
- [204] Holbrook JA, Jarosz-Griffiths HH, Caseley E, Lara-Reyna S, Poulter JA, Williams-Gray CH, et al. (2021). Neurodegenerative Disease and the NLRP3 Inflammasome. *Front Pharmacol*, 12:643254.
- [205] Anderson FL, Biggs KE, Rankin BE, Havrda MC (2023). NLRP3 inflammasome in neurodegenerative disease. *Transl Res*, 252:21-33.
- [206] Chakraborty S, Kaushik DK, Gupta M, Basu A (2010). Inflammasome signaling at the heart of central nervous system pathology. *J Neurosci Res*, 88(8):1615-31.
- [207] Hung SC, Choi CH, Said-Sadier N, Johnson L, Atanasova KR, Sellami H, et al. (2013). P2X4 assembles with P2X7 and pannexin-1 in gingival epithelial cells and modulates ATP-induced reactive oxygen species production and inflammasome activation. *PLoS One*, 8(7):e70210.
- [208] Kanellopoulos JM, Almeida-da-Silva CLC, Rüttel Boudinot S, Ojcius DM (2021). Structural and Functional Features of the P2X4 Receptor: An Immunological Perspective. *Front Immunol*, 12:645834.
- [209] Li Q, Fu W, Yao J, Ji Z, Wang Y, Zhou Z, et al. (2014). Heme induces IL-1 β secretion through activating NLRP3 in kidney inflammation. *Cell Biochem Biophys*, 69(3):495-502.
- [210] Ducza L, Gajtkó A, Hegedűs K, Bakk E, Kis G, Gaál B, et al. (2023). Neuronal P2X4 receptor may contribute to peripheral inflammatory pain in rat spinal dorsal horn. *Front Mol Neurosci*, 16:1115685.
- [211] Du RH, Wu FF, Lu M, Shu XD, Ding JH, Wu G, et al. (2016). Uncoupling protein 2 modulation of the NLRP3 inflammasome in astrocytes and its implications in depression. *Redox Biol*, 9:178-187.
- [212] Zhou X, Xiao Q, Xie L, Yang F, Wang L, Tu J (2019). Astrocyte, a Promising Target for Mood Disorder Interventions. *Front Mol Neurosci*, 12:136.
- [213] Zang X, Zheng X, Hou Y, Hu M, Wang H, Bao X, et al. (2018). Regulation of proinflammatory monocyte activation by the kynurenine-AhR axis underlies immunometabolic control of depressive behavior in mice. *FASEB J*, 32(4):1944-1956.

- [214] Rolstad S, Jakobsson J, Sellgren C, Isgren A, Ekman CJ, Bjerke M, et al. (2015). CSF neuroinflammatory biomarkers in bipolar disorder are associated with cognitive impairment. *Eur Neuropsychopharmacol*, 25(8):1091-8.
- [215] Tsai PJ, Lai YH, Manne RK, Tsai YS, Sarbassov D, Lin HK (2023) Akt: a key transducer in cancer. *J Biomed Sci*, 29(1):76. Erratum in: *J Biomed Sci*, 2023 30(1):7.
- [216] Peng S, Gu JH, Dai CL, Iqbal K, Liu F, Gong CX (2022). AKT/GSK-3 β signaling is altered through downregulation of mTOR during cerebral Ischemia/Reperfusion injury. *Mol Biol Rep*, 49(5):3955-3964.
- [217] Stefaniak J, O'Brien J (2016). Imaging of neuroinflammation in dementia: a review. *J Neurol Neurosurg Psychiatry*, 87(1):21-8.
- [218] Passamonti L, Tsvetanov KA, Jones PS, Bevan-Jones WR, Arnold R, Borchert RJ (2019). Neuroinflammation and Functional Connectivity in Alzheimer's Disease: Interactive Influences on Cognitive Performance. *J Neurosci*, 39(36):7218-7226.
- [219] Ng A, Tam WW, Zhang MW, Ho CS, Husain SF, McIntyre RS, et al. (2018). IL-1 β , IL-6, TNF- α and CRP in Elderly Patients with Depression or Alzheimer's disease: Systematic Review and Meta-Analysis. *Sci Rep*, 8(1):12050.
- [220] Cuello AC (2017). Early and Late CNS Inflammation in Alzheimer's Disease: Two Extremes of a Continuum? *Trends Pharmacol Sci*, 38(11):956-966.
- [221] Aziz N (2015). Measurement of Circulating Cytokines and Immune-Activation Markers by Multiplex Technology in the Clinical Setting: What Are We Really Measuring? *For Immunopathol Dis Therap*, 6(1-2):19-22.
- [222] Koelman L, Pivovarova-Ramich O, Pfeiffer AFH, Grune T, Aleksandrova K (2019). Cytokines for evaluation of chronic inflammatory status in ageing research: reliability and phenotypic characterisation. *Immun Ageing*, 16:11.
- [223] Zuliani G, Ranzini M, Guerra G, Rossi L, Munari MR, Zurlo A, et al. (2007). Plasma cytokines profile in older subjects with late onset Alzheimer's disease or vascular dementia. *J Psychiatr Res*, 41(8):686-93.
- [224] Belkhef M, Beder N, Mouhoub D, Amri M, Hayet R, Tighilt N, et al. (2018). The involvement of neuroinflammation and necroptosis in the hippocampus during vascular dementia. *J Neuroimmunol*, 320:48-57.
- [225] Engelhart MJ, Geerlings MI, Meijer J, Kiliaan A, Ruitenbergh A, van Swieten JC, et al. (2004). Inflammatory proteins in plasma and the risk of dementia: the rotterdam study. *Arch Neurol*, 61(5):668-72.
- [226] Heneka MT, Kummer MP, Stutz A, Delekate A, Schwartz S, Vieira-Saecker A, et al. (2013). NLRP3 is activated in Alzheimer's disease and contributes to pathology in APP/PS1 mice. *Nature*, 493(7434):674-8.
- [227] Baldwin AG, Brough D, Freeman S (2016). Inhibiting the Inflammasome: A Chemical Perspective. *J Med Chem*, 59(5):1691-710.
- [228] Bertinaria M (2021). Inflammasome Inhibitors. *Molecules*, 26(22):6912.
- [229] Coll RC, Robertson AA, Chae JJ, Higgins SC, Muñoz-Planillo R, Inserra MC, et al. (2015). A small-molecule inhibitor of the NLRP3 inflammasome for the treatment of inflammatory diseases. *Nat Med*, 21(3):248-55.
- [230] Dempsey C, Rubio Araiz A, Bryson KJ, Finucane O, Larkin C, Mills EL, et al. (2017). Inhibiting the NLRP3 inflammasome with MCC950 promotes non-phlogistic clearance of amyloid- β and cognitive function in APP/PS1 mice. *Brain Behav Immun*, 61:306-316.
- [231] Gordon R, Albornoz EA, Christie DC, Langley MR, Kumar V, Mantovani S, et al. (2018). Inflammasome inhibition prevents α -synuclein pathology and dopaminergic neurodegeneration in mice. *Sci Transl Med*, 10(465):eaah4066.
- [232] Piancone F, La Rosa F, Marventano I, Saresella M, Clerici M (2021). The Role of the Inflammasome in Neurodegenerative Diseases. *Molecules*, 26(4):953.
- [233] La Rosa F, Saresella M, Marventano I, Piancone F, Ripamonti E, Al-Daghri N, et al. (2019). Stavudine Reduces NLRP3 Inflammasome Activation and Modulates Amyloid- β Autophagy. *J Alzheimers Dis*, 72(2):401-412.
- [234] Gastaldi S, Boscaro V, Gianquinto E, Sandall CF, Giorgis M, Marini E, et al. (2021). Chemical Modulation of the 1-(Piperidin-4-yl)-1,3-dihydro-2H-benzo[d]imidazole-2-one Scaffold as a Novel NLRP3 Inhibitor. *Molecules*, 26(13): 3975.
- [235] Poli G, Fabi C, Bellet MM, Costantini C, Nunziangeli L, Romani L, et al. (2020). Epigenetic Mechanisms of Inflammasome Regulation. *Int J Mol Sci*, 21(16):5758.
- [236] Li D, Yang H, Ma J, Luo S, Chen S, Gu Q (2018). MicroRNA-30e regulates neuroinflammation in MPTP model of Parkinson's disease by targeting Nlrp3. *Hum Cell*, 31(2):106-115.
- [237] Zhou Y, Lu M, Du RH, Qiao C, Jiang CY, Zhang KZ, et al. (2016). MicroRNA-7 targets Nod-like receptor protein 3 inflammasome to modulate neuroinflammation in the pathogenesis of Parkinson's disease. *Mol Neurodegener*, 11:28.
- [238] Kaushal V, Dye R, Pakavathkumar P, Foveau B, Flores J, Hyman B, et al. (2015). Neuronal NLRP1 inflammasome activation of Caspase-1 coordinately regulates inflammatory interleukin-1-beta production and axonal degeneration-associated Caspase-6 activation. *Cell death and differentiation*, 22(10):1676-1686.
- [239] de Souza N (2013). Model organisms: Mouse models challenged. *Nat Methods*, 10(4):288.
- [240] Pound P, Ritskes-Hoitinga M (2018). Is it possible to overcome issues of external validity in preclinical animal research? Why most animal models are bound to fail. *J Transl Med*, 16(1):304.
- [241] Chiarini A, Armato U, Hu P, Dal Prà I (2020). Danger-Sensing/Patten Recognition Receptors and Neuroinflammation in Alzheimer's Disease. *Int J Mol Sci*, 21(23):9036.

- [242] van Dyck CH, Swanson CJ, Aisen P, Bateman RJ, Chen C, Gee M, et al. (2023). Lecanemab in Early Alzheimer's Disease. *N Engl J Med*, 388(1):9-21.
- [243] Jiang W, Li M, He F, Bian Z, Liu J, He Q, et al. (2016). Dopamine D1 receptor agonist A-68930 inhibits NLRP3 inflammasome activation and protects rats from spinal cord injury-induced acute lung injury. *Spinal Cord*, 54(11):951-956.
- [244] Wang S, Yao Q, Wan Y, Wang J, Huang C, Li D, et al. (2020). Adiponectin reduces brain injury after intracerebral hemorrhage by reducing NLRP3 inflammasome expression. *Int J Neurosci*, 130(3):301-308.
- [245] Aryanpour R, Zibara K, Pasbakhsh P, Jame'ei SB, Namjoo Z, Ghanbari A, et al. (2021). 17 β -Estradiol Reduces Demyelination in Cuprizone-fed Mice by Promoting M2 Microglia Polarity and Regulating NLRP3 Inflammasome. *Neuroscience*, 463:116-127.
- [246] Yang X, Sun J, Kim TJ, Kim YJ, Ko SB, Kim CK, et al. (2018). Pretreatment with low-dose fimasartan ameliorates NLRP3 inflammasome-mediated neuroinflammation and brain injury after intracerebral hemorrhage. *Exp Neurol*, 310:22-32
- [247] Soriano-Teruel PM, García-Laínez G, Marco-Salvador M, Pardo J, Arias M, DeFord C, et al. (2021). Identification of an ASC oligomerization inhibitor for the treatment of inflammatory diseases. *Cell Death Dis*, 12(12):1155.