

Facile Deoxygenation of Hydroxylated Flavonoids by Palladium-Catalysed Reduction of its Triflate Derivatives

József Kövér and Sándor Antus

Department of Organic Chemistry, University of Debrecen, H-4010 Debrecen, P. O. Box 20, Hungary

Reprint requests to Sándor Antus. Fax: +36-52-453-836. E-mail: antuss@tigris.klte.hu

Z. Naturforsch. **60b**, 792 – 796 (2005); received November 12, 2004

Dedicated to Professor András Lipták on the occasion of his 70th birthday


An efficient procedure to deoxygenate hydroxy substituted flavonoids, isoflavonoids and related compounds *via* their trifluoromethanesulfonates is presented. Their reduction with formic acid in the presence of a catalytic amount of palladium acetate, triethylamine and 1,3-bis(diphenylphosphanyl)propane (dppp) in DMF results in their *des*-hydroxy derivatives without affecting other functional groups.

Key words: Flavonoids, Reduction, Palladium(II) Acetate

Flavonoids and isoflavonoids possessing a 2- and 3-aryl substituted 1-benzopyran skeleton with various levels of saturation and oxidation, respectively, belong to the class of naturally occurring *O*-heterocyclic compounds [1] with wide range of biological importance due to their significant antioxidant [2], hepatoprotective [3], antifungal [4], antibacterial [5] and antiviral [6] activities.

Although, there are numerous independent methods available for the preparation of these natural products and the procedures of synthetic importance have been reviewed extensively [1, 7, 8], these conditions are mostly suitable for direct synthesis of their 7-hydroxy- or 5,7-dihydroxy derivatives and hence synthetic accessibility of their *des*-7-hydroxy derivatives is strongly limited. To overcome this problem several methods such as catalytic hydrogenation of *O*-aryl-N,N-dialkylisoureas by Pd/C [9], desulfurization of *S*-aryl-N,N-dialkyl thiocarbamates (prepared by Newman-Kwart rearrangement of *O*-aryl-N,N-dialkyl thiocarbamates) by Raney nickel [10] and reduction of *O*-aryl (or methyl)-*p*-toluenesulfonates by NaBH₄-NiCl₂ [11] or Raney nickel [12] have been recently introduced by Patonay *et al.* [13] in this field using 2,2-dimethyl-7-hydroxychromanone (**1a**) as the substrate.

Efficient procedures for the deoxygenation of simple phenols have been reported *via* their trifluoromethanesulfonates (triflates) by hydrogenolysis under heterogeneous conditions (Pd/C) [14] and Ni(0)- [15] or Pd(0)-catalyzed [16, 17] reduction under homoge-




Scheme 1.

neous conditions. As a continuation of our research program aiming at the application of Pd(0)-catalyzed transformation for the synthesis of *O*-heterocyclic compounds [18–20], it seemed promising to study also the deoxygenation of hydroxybenzopyranoid triflates by Pd(0)-catalyzed reduction.

In order to compare our results with those published by Patonay *et al.* [13], 2,2-dimethyl-7-hydroxychromanone (**1a**) and its derivatives **1b–h** were chosen as model compounds (Scheme 1) whose corresponding triflates **2a–h** were prepared in high yields (85–94%) by treatment with trifluoromethanesulfonic anhydride in the presence of triethylamine in dichloromethane at –15 °C. When the reaction of 2,2-dimethyl-chromanon-7-yl triflate (**2a**) with 1 equiv-

Table 1. Compounds **3a–h** prepared.

Substrate	Product	Time [min]	Yield ^a [%]	M. p. [°C]	Lit. m. p. [°C]	Lit. yield ^b [%]
2a	3a	50	93	88.5–89	89–90 [13]	80
2b	3b	25	90	68–69	96.5 [21]	82
2c	3c	30	91	oil	oil [22]	80
2d	3d	70	95	124–125	118–119 [23]	83
2e	3e	30	92	75–76	71–72 [23]	80
2f	3f	20	91	132–133	126–127 [23]	78
2g	3g	15	70.4	oil	oil [24]	41
2h	3h	30	75	88–89	89–90 [13]	70

^a Isolated yields; ^b ref. [13].

Scheme 2.

alent of formic acid as a reducing agent was conducted in *N,N*-dimethyl formamide (DMF) at 60 °C in the presence of palladium(II) acetate (2.2 mol-%) and triphenylphosphane (PPh₃) (4.2 mol-%) according to the method described by Cacchi *et al.* [17], quantitative formation of 2,2-dimethylchromanone (**1c**) was observed within 15 minutes. As shown in Table 1, the other related derivatives **2b–g** reacted readily to afford the corresponding reduction products **3b–g**, respectively, in excellent yields (70–95%). Ditriflate **2h** required the use of twice the amounts of the reagents to convert efficiently into **3a**. It is noteworthy that chloro and carbonyl groups are completely unaffected under these conditions and the yields have been found to be comparable with those published by Patonay *et al.* [13]

Table 2. Compounds **4c–11c** prepared.

Substrate	Product	Time [min]	Yield ^a [%]	M. p. [°C]	Lit. m. p. [°C]
4b	4c	60	83	57–58	60–62 [25]
5b	5c	90	83	56–57	60–62 [25]
6b	6c	60	85	96–97	97 [26]
7b	7c	30	90	134–135	133–134 [27]
8b	8c	60	93	108–109	86–87 [28]
9b	9c	60	90	150–151	154–156 [29]
10b	10c	30	94	75–76	75–78 [30]
11b	11c	90	79	66–67	64–65 [31]

^a Isolated yields.

but our procedure is significantly milder and simpler. Moreover, it has been also recognized that triflates **2a** and **2g** were less reactive towards the catalytic system using PPh₃ instead of dppp. In both cases, hydrolysis of triflates **2a, g** to **1a, g**, respectively, could also be observed besides their deoxygenation (**2a** → **3a**, **2g** → **3g**). In order to determine the scope and limitation of this deoxygenation procedure the transformation of the chalcones **4b** and **5b**, the flavone **6b**, the isoflavones **7b–9b**, the coumarin **10b** and chromone derivatives **11b** was also studied (Scheme 2). The reduction of these compounds took place very smoothly to give the desired deoxygenated derivatives **4c–11c**, respectively, in good yields (79–94%) and no byproducts could be detected (TLC) in any cases (Table 2).

In conclusion, these results have clearly indicated that the palladium(II) acetate/1,3-bis(diphenylphosphanyl)propane (dppp)/formic acid/triethylamine system in DMF is an efficient agent for deoxygenation of hydroxysubstituted flavonoids, isoflavonoids and related compounds *via* their trifluoromethanesulfonates.

Experimental Section

Melting points were determined on a Büchi 535 apparatus and are not corrected. ¹H NMR spectra were obtained on a Varian Gemini 200 NMR spectrometer in CDCl₃ with TMS as internal standard. EI-MS (70 eV) spectra were obtained with a VG TRIO-2 instrument. TLC was performed on Merck Kieselgel 60 F₂₅₄ pre-coated aluminium plates. Elemental analyses (C, H) were conducted using Carlo Erba 1106 EA instrument. 7-Hydroxyflavone (**6a**) was purchased from Sigma-Aldrich. The preparation of the following derivatives was described earlier: **1a** [22], **1b** [32], **1c** [22], **1d** [13], **1e** [33], **1f** [13], **1g** [34], **1h** [35], **4a**, **5a** [36], **7a**, **8a**, **9a** [37], **10a** [38], **11a** [39].

Aryl triflates. General procedure: 2,2-Dimethyl-7-(trifluoromethylsulfonyloxy)chromanone (**2a**)

To a solution of 7-hydroxy-2,2-dimethylchromanone [22] (2.40 g, 12.50 mmol) in 20 ml of dichloromethane and

3 ml of triethylamine at -15°C was slowly added trifluoromethanesulfonic anhydride (2.15 ml, 3.70 g, 13.10 mmol). The mixture was stirred at this temperature for 5 minutes and then was allowed to warm to 23°C and stirred until no starting material could be detected by TLC (30 minutes). The mixture was washed with conc. NaHCO_3 solution (2×30 ml) and conc. NaCl solution, dried (MgSO_4), and concentrated. The residue was purified by chromatography [silica gel, elution with *n*-hexane : ethyl acetate (3:1)] to afford **2a** as white crystals (3.80 g, 94%); m. p. $29-30^{\circ}\text{C}$. – ^1H NMR: $\delta = 1.45$ (s, 6H, 2-H), 2.75 (s, 2H, 3-H), 6.90 (m, 2H, 6-H, 8-H), 7.95 (d, $J = 9.28$ Hz, 1H, 5-H). – MS: m/e (%) = 324 (M^+ , 8), 309 (22), 269 (12), 176 (19), 149 (19), 107 (39), 79 (32), 69 (100). – $\text{C}_{12}\text{H}_{11}\text{F}_3\text{O}_5\text{S}$ (324.29): calcd. C 44.45, H 3.42; found C 44.40, H 3.49.

The compounds **2b–h** and **4b–11b** were prepared in an analogous manner.

7-Trifluoromethylsulfonyloxy-2,2,5-trimethylchromanone (2b)

(20 min, yield: 93%, oil). – ^1H NMR: $\delta = 1.45$ (s, 6H, 2,2- CH_3), 2.65 (s, 3H, 5- CH_3), 2.75 (s, 2H, 3-H), 6.65 (d, $J = 2.80$ Hz, 1H, 6-H or 8-H), 6.75 (d, $J = 2.80$ Hz, 1H, 6-H or 8-H). – MS: m/e (%) = 338 (M^+ , 26), 323 (10), 283 (16), 121 (25), 93 (11), 69 (100). – $\text{C}_{13}\text{H}_{13}\text{F}_3\text{O}_5\text{S}$ (338.30): calcd. C 46.15, H 3.87; found C 46.91, H 3.81.

7-Trifluoromethylsulfonyloxy-2,2,8-trimethylchromanone (2c)

(10 min, yield: 90%, white crystals, m. p. $53-54^{\circ}\text{C}$). – ^1H NMR: $\delta = 1.50$ (s, 6H, 2,2- CH_3), 2.25 (s, 3H, 8- CH_3), 2.75 (s, 2H, 3-H), 6.90 (d, $J = 9.13$ Hz, 1H, 5-H or 6-H), 7.80 (d, $J = 9.10$ Hz, 1H, 5-H or 6-H). – MS: m/e (%) = 338 (M^+ , 20), 323 (51), 283 (22), 190 (25), 149 (29), 93 (11), 69 (100). – $\text{C}_{13}\text{H}_{13}\text{F}_3\text{O}_5\text{S}$ (338.30): calcd. C 46.15, H 3.87; found C 46.31, H 3.60.

2,2-Dimethyl-5-methoxy-7-(trifluoromethylsulfonyloxy)-chromanone (2d)

(10 min, yield: 93%, oil). – ^1H NMR: $\delta = 1.45$ (s, 6H, 2,2- CH_3), 2.70 (s, 2H, 3-H), 3.90 (s, 3H, 5-OMe), 6.35 (d, $J = 2.17$ Hz, 1H, 6-H or 8-H), 6.50 (d, $J = 2.15$ Hz, 1H, 6-H or 8-H). – MS: m/e (%) = 354 (M^+ , 11), 339 (10), 299 (39), 206 (7), 138 (10), 83 (8), 69 (100). – $\text{C}_{13}\text{H}_{13}\text{F}_3\text{O}_6\text{S}$ (354.30): calcd. C 44.07, H 3.69; found C 44.25, H 3.62.

2,2-Dimethyl-6-methoxy-7-(trifluoromethylsulfonyloxy)-chromanone (2e)

(10 min, yield: 89%, white crystals, m. p. $87-88^{\circ}\text{C}$). – ^1H NMR: $\delta = 1.45$ (s, 6H, 2,2- CH_3), 2.75 (s, 2H, 3-H), 3.90 (s, 3H, 6-OMe), 6.90 (s, 1H, 8-H), 7.45 (s, 1H, 5-H). – MS:

m/e (%) = 354 (M^+ , 8), 339 (15), 299 (8), 206 (5), 165 (6), 138 (32), 69 (100). – $\text{C}_{13}\text{H}_{13}\text{F}_3\text{O}_6\text{S}$ (354.30): calcd. C 44.07, H 3.69; found C 44.50, H 3.59.

2,2-Dimethyl-8-methoxy-7-(trifluoromethylsulfonyloxy)-chromanone (2f)

(15 min, yield: 88%, oil). – ^1H NMR: $\delta = 1.55$ (s, 6H, 2,2- CH_3), 2.80 (s, 2H, 3-H), 4.05 (s, 3H, 8-OMe), 6.85 (d, $J = 8.80$ Hz, 1H, 6-H), 7.65 (d, $J = 8.80$ Hz, 1H, 5-H). – MS: m/e (%) = 354 (M^+ , 30), 339 (41), 299 (31), 206 (25), 107 (100), 109 (28), 69 (100). – $\text{C}_{13}\text{H}_{13}\text{F}_3\text{O}_6\text{S}$ (354.30): calcd. C 44.07, H 3.69; found C 44.32, H 3.70.

6-Chloro-2,2-dimethyl-7-(trifluoromethylsulfonyloxy)chromanone (2g)

(30 min, yield: 92%, white crystals, m. p. $58-58.5^{\circ}\text{C}$). – ^1H NMR: $\delta = 1.45$ (s, 6H, 2,2- CH_3), 2.75 (s, 2H, 3-H), 7.00 (s, 1H, 8-H), 8.00 (s, 1H, 5-H). – MS: m/e (%) = 358 (M^+ , 6), 343 (12), 303 (6), 210 (10), 141 (12), 113 (9), 69 (100). – $\text{C}_{12}\text{H}_{10}\text{ClF}_3\text{O}_5\text{S}$ (358.72): calcd. C 40.17, H 2.80; found C 40.30, H 2.95.

5,7-Bis(trifluoromethylsulfonyloxy)-2,2-dimethylchromanone (2h)

(30 min, yield: 85%, white crystals, m. p. $123-124^{\circ}\text{C}$). – ^1H NMR: $\delta = 1.51$ (s, 6H, 2,2- CH_3), 2.80 (s, 2H, 3-H), 6.73–6.74 (d, $J = 2.20$ Hz, 1H, 6-H or 8-H), 6.97–6.99 (d, $J = 2.29$ Hz, 1H, 6-H or 8-H). – $\text{C}_{13}\text{H}_{10}\text{F}_6\text{O}_8\text{S}_2$ (472.26): calcd. C 33.06, H 2.11; found C 33.21, H 2.16.

4'-(Trifluoromethylsulfonyloxy)chalcone (4b)

(20 min, yield: 94%, white crystals, m. p. $96-97^{\circ}\text{C}$). – ^1H NMR: $\delta = 7.14-7.55$ (m, 6H, aromatic-H, α -CH), 7.60–7.75 (m, 2H, aromatic-H), 7.80–8.15 (m, 3H, aromatic-H, β -CH). – MS: m/e (%) = 356 (M^+ , 42), 253 (76), 131 (100), 69 (72). – $\text{C}_{16}\text{H}_{11}\text{F}_3\text{O}_4\text{S}$ (356.32): calcd. C 53.93, H 3.11; found C 53.90, H 3.18.

4-(Trifluoromethylsulfonyloxy)chalcone (5b)

(15 min, yield: 95%, white crystals, m. p. $60-61^{\circ}\text{C}$). – ^1H NMR: $\delta = 7.35$ (d, $J = 8.72$ Hz, 2H, aromatic-H), 7.50 (d, $J = 5.99$ Hz, 2H, aromatic-H), 7.56–7.66 (m, 5H, aromatic-H, α - and β -CH), 8.03 (dd, $J_1 = 1.37$ Hz, $J_2 = 8.71$ Hz, 2H, aromatic-H). – MS: m/e (%) = 356 (M^+ , 25), 223 (66), 131 (77), 105 (100), 69 (82). – $\text{C}_{16}\text{H}_{11}\text{F}_3\text{O}_4\text{S}$ (356.32): calcd. C 53.93, H 3.11; found C 53.19, H 3.08.

7-(Trifluoromethylsulfonyloxy)flavone (6b)

(30 min, yield: 78%, white crystals, m. p. $137-138^{\circ}\text{C}$). – ^1H NMR: $\delta = 6.87$ (s, 1H, 3-H), 7.35 (dd, $J_1 = 2.25$ Hz, $J_2 =$

8.82 Hz, 1H, 6-H), 7.51–7.59 (m, 4H, aromatic-H), 7.91–7.96 (m, 2H, aromatic-H), 8.35 (d, $J = 8.80$ Hz, 1H, 5-H). – MS: m/e (%) = 370 (M^+ , 80), 273 (57), 268 (54), 237 (100), 135 (55), 69 (75). – $C_{16}H_9F_3O_5S$ (370.30): calcd. C 51.89, H 2.74; found C 51.01, H 2.79.

7-(Trifluoromethylsulfonyloxy)isoflavone (7b)

(45 min, yield: 89%, white crystals, m. p. 146–147 °C). – 1H NMR: $\delta = 7.26–7.57$ (m, 7H, aromatic-H), 8.06 (s, 1H, 2-H), 8.43 (d, $J = 8.84$ Hz, 1H, 5-H). – MS: m/e (%) = 370 (M^+ , 52), 273 (72), 268 (88), 237 (72), 135 (82), 69 (100). – $C_{16}H_9F_3O_5S$ (370.30): calcd. C 51.89, H 2.74; found C 52.01, H 2.86.

5-Hydroxy-2-methyl-7-(trifluoromethylsulfonyloxy)-isoflavone (8b)

(30 min, yield: 68%, white crystals, m. p. 128–129 °C). – 1H NMR: $\delta = 2.30$ (s, 3H, CH_3), 7.02 (s, 1H, 8-H), 7.35–7.44 (m, 6H, aromatic-H), 13.14 (s, 1H, 5-OH). – MS: m/e (%) = 400 (M^+ , 32), 331 (44), 303 (63), 284 (58), 267 (88), 151 (77), 69 (100). – $C_{17}H_{11}F_3O_6S$ (400.33): calcd. C 51.00, H 2.76; found C 51.17, H 2.85.

3',4'-Methylenedioxy-7-(trifluoromethylsulfonyloxy)-isoflavone (9b)

(30 min, yield: 73%, white crystals, m. p. 174–175 °C). – 1H NMR: $\delta = 6.01$ (s, 2H, $-OCH_2O-$), 6.89 (d, $J = 8.01$ Hz, 1H, 5'-H), 6.98 (dd, $J_1 = 1.66$ Hz, $J_2 = 8.01$ Hz, 1H, 6'-H), 7.08 (d, $J = 1.66$ Hz, 1H, 2'-H), 7.35 (dd, $J_1 =$

2.32 Hz, $J_2 = 8.83$ Hz, 1H, 6-H), 7.46 (d, $J = 2.32$ Hz, 1H, 8-H), 8.02 (s, 1H, 2-H), 8.42 (d, $J = 8.83$ Hz, 1H, 5-H). – MS: m/e (%) = 414 (M^+ , 38), 281 (88), 268 (75), 252 (72), 146 (51), 69 (100). – $C_{17}H_9F_3O_7S$ (414.32): calcd. C 49.28, H 2.18; found C 49.57, H 2.23.

4-Methyl-7-(trifluoromethylsulfonyloxy)coumarin (10b)

(30 min, yield: 91%, white crystals, m. p. 83–84 °C). – 1H NMR: $\delta = 2.47$ (d, $J = 1.18$ Hz, 3H, 4- CH_3), 6.36–6.38 (q, $J = 1.18$ Hz, 1H, 3-H), 7.22 (d, $J = 2.45$ Hz, 1H, 8-H), 7.29 (dd, $J_1 = 2.45$ Hz, $J_2 = 8.60$ Hz, 1H, 6-H), 7.70 (d, $J = 8.60$ Hz, 1H, 5-H). – MS: m/e (%) = 308 (M^+ , 46), 239 (40), 175 (74), 69 (100). – $C_{11}H_7F_3O_5S$ (308.23): calcd. C 42.86, H 2.29; found C 42.57, H 2.25.

Ethyl 7-(trifluoromethylsulfonyloxy)chromone-2-carboxylate (11b)

(30 min, yield: 66%, white crystals, m. p. 127–128 °C). – 1H NMR: $\delta = 1.45$ (t, $J = 7.14$ Hz, 3H, ethyl- CH_3), 4.49 (q, $J = 7.14$ Hz, 2H, ethyl- CH_2), 7.16 (s, 1H, 3-H), 7.37 (dd, $J_1 = 2.34$ Hz, $J_2 = 8.87$ Hz, 1H, 6-H), 7.65 (d, $J = 2.34$ Hz, 1H, 8-H), 8.31 (d, $J = 8.87$ Hz, 1H, 5-H). – MS: m/e (%) = 336 (M^+ , 20), 297 (62), 268 (82), 233 (54), 69 (100). – $C_{13}H_9F_3O_7S$ (366.27): calcd. C 42.63, H 2.47; found C 42.17, H 2.50.

Acknowledgement

We wish to thank the National Science Foundation (OTKA T049436) for financial support.

- [1] J. B. Harborn, T. J. Mabry, H. Mabry (eds): *The Flavonoids*, Chapman and Hall Ltd., London (1975).
- [2] P. G. Pietta, *J. Nat. Prod.* **63**, 1035 (2000).
- [3] H. Wagner, in J. L. Bear, E. Reinhardt (eds): *Plant Constituents with Antihepatotoxic Activity in Natural Products as Medicinal Agents*, p. 217, Hippokrates Verlag, Stuttgart (1980).
- [4] J. A. M. Cruickshank, *Aust. J. Biol. Chem.* **15**, 147 (1962).
- [5] M. Hattori, S. Hada, A. Watahiki, H. Ihara, Y. Z. Shu, N. Kakiuchi, T. Mizuno, T. Namba, *Chem. Pharm. Bull.* **34**, 3885 (1986).
- [6] C. Q. Hu, K. Ke, Q. Shi, R. E. Kilkuskie, K. H. Lee, *J. Nat. Prod.* **57**, 42 (1994).
- [7] I. M. Lockhart, in G. P. Ellis (ed.): *Chromenes, Chromanones and Chromones*, p. 207, John Wiley and Sons Ltd., New York (1977).
- [8] J. B. Harborn (ed.): *The Flavonoids, Advances in Research since 1986*, Chapman and Hall Ltd., London (1994).
- [9] E. Vowinkel, H. J. Baese, *Chem. Ber.* **107**, 1213 (1974).
- [10] H. Kwart, E. R. Evans, *J. Org. Chem.* **31**, 410 (1966).
- [11] F. Wang, K. Chiba, M. Tada, *J. Chem. Soc. Perkin Trans. 1*, 1897 (1992).
- [12] G. W. Kenner, M. A. Murray, *J. Chem. Soc.* 178 (1949).
- [13] P. Sebők, T. Tímár, T. Eszenyi, T. Patonay, *J. Org. Chem.* **59**, 6318 (1994).
- [14] A. G. Martinez, R. M. Alvarez, J. A. Aguirre, L. R. Subramanian, *J. Chem. Soc. Perkin Trans. 1*, 1595 (1986).
- [15] K. Sasaki, M. Sakai, Y. Sakakibara, K. Takagi, *Chem. Lett.* 2017 (1991).
- [16] S. Cacchi, E. Morera, G. Ortar, *Tetrahedron Lett.* **25**, 4821 (1986).
- [17] S. Cacchi, E. Morera, G. Ortar, *Org. Synth.* **68**, 138 (1990) and references therein.
- [18] A. L. Tőkés, Gy. Litkei, K. Gulácsi, S. Antus, E. Baitz-Gács, Cs. Szántay, L. L. Darkó, *Tetrahedron* **55**, 9283 (1999).
- [19] L. Kiss, S. Antus, *Heterocycl. Commun.* **6**, 309 (2000).
- [20] L. Kiss, G. Papp, F. Joó, S. Antus, *Heterocycl. Commun.* **7**, 417 (2001).

- [21] J. Cologne, R. Chambard, *Bull. Soc. Chim. Fr.* **20**, 573 (1953).
- [22] P. Sebők, J. Jekő, T. Tímár, J. Cs. Jászberényi, *Heterocycles* **38**, 2099 (1994).
- [23] F. Camps, J. Coll, O. Colomina, *J. Heterocyclic Chem.* **22**, 363 (1985).
- [24] H. J. Kabbe, A. Widdig, *Angew. Chem.* **21**, 254 (1982).
- [25] R. D. Offenbauer, S. F. Nelsen, *J. Org. Chem.* **33**, 775 (1968).
- [26] A. Nishinaga, H. Ando, K. Maruyama, T. Mashino, *Synthesis* 839 (1992).
- [27] A. Lévai, L. Balogh, *Pharmazie* **30**, 747 (1975).
- [28] A. Lévai, P. Sebők, *Synth. Commun.* **22**, 1735 (1992).
- [29] Gy. Litkei, R. Bognár, Z. Dinya, *Chim. Acad. Sci. Hung.* **71**, 403 (1972).
- [30] A. N. Brubaker, J. DeRuiter, W. L. Whitmer, *J. Med. Chem.* **29**, 1094 (1986).
- [31] G. P. Ellis, D. Shaw, *J. Med. Chem.* **15**, 865 (1972).
- [32] T. Tímár, J. Cs. Jászberényi, S. Hosztafi, *Acta. Chim. Hung.* **125**, 457 (1988).
- [33] F. Camps, J. Coll, A. Messguer, M. A. Pericas, S. Ricart, W. S. Bowers, D. M. Soderlund, *Synthesis* 725 (1980).
- [34] P. Sebők, T. Tímár, J. Cs. Jászberényi, J. Jekő, *Acta Chim. Hung.* **126**, 471 (1989).
- [35] T. Tímár, J. Cs. Jászberényi, *J. Heterocyclic Chem.* **25**, 871 (1988).
- [36] H. Bablich, St. Kostanecki, *Chem. Ber.* **29**, 233 (1897).
- [37] A. Lévai, P. Sebők, *Synth. Commun.* **22**, 1735 (1992).
- [38] R. N. Lacey, *J. Chem. Soc.* 854 (1954).
- [39] P. S. Manchand, R. A. Micheli, S. J. Saposnik, *Tetrahedron* **48**, 9391 (1992).