

Optimization and Control in Rubella Transmission Dynamics: A Boundedness-Preserving Numerical Model with Vaccination

Samiullah Salim¹, Fazal Dayan¹, Muhammad Azizur Rehman¹, Husam A. Neamah^{2*}

¹Department of Mathematics, School of Science, University of Management and Technology, Lahore, 54000, Pakistan.

²Department of Electrical Engineering and Mechatronics, Faculty of Engineering, University of Debrecen, Óttemető Utca 2-4, 4028 Debrecen, Hungary

*Corresponding Authors: Email: husam@eng.unideb.hu

Abstract: Rubella is an infectious disease that can spread globally. It spreads in the subtropics as well as the tropics. Although it is commonly thought to be a non-fatal condition, there are several circumstances in which it can be fatal. Pregnant women infected with the Rubella virus face a high risk of fetal development. The main goal of this study is to look into a model for the spread of Rubella while considering a vaccination campaign as a control measure. The positivity and boundedness of the re-infection rubella transmission and vaccination model are investigated. The local and global stability of the equilibrium points is examined. The sensitivity of parameters is also investigated. Three different techniques forward Euler, RK-4, and non-standard finite difference (NSFD) method are developed for the numerical solution, and their simulation results are examined. Among these three, the NSFD method is superior due to its convergence and positive behavior for all step size values. For some values of the step size, the other two techniques failed to produce positivity and convergent solutions. Analytically, the proposed model's convergence, positivity, boundedness, and consistency are investigated. Finally, the impact of the Rubella vaccine on infected populations has been examined, revealing that vaccination is one of the most effective ways to prevent rubella transmission.

Keywords: Rubella; sensitivity; stability; convergence; consistency; positivity.

1 Introduction

German measles, also known as rubella, is an infectious viral infection caused by the rubella virus. When an infected person sneezes or coughs, the virus is typically spread through airborne droplets. The disease can also be passed from a pregnant woman to her unborn child, which can result in serious birth defects and complications. A mild fever, rash, and swollen lymph nodes are common Rubella symptoms. The symptoms are often minor and resolve within a week or two. The virus poses a considerable risk to pregnant women because it can cause congenital rubella syndrome (CRS) in their unborn children. CRS can cause various issues, including hearing and vision difficulties, heart defects, and developmental delays. Immunization is critical in preventing rubella because it protects not just the individual but also fosters herd immunity, which protects vulnerable people such as infants and those with impaired immune systems. Rubella symptoms include a low fever, sore throat, headache, and a characteristic rash that spreads from the face to the body, usually lasting a few days. Additional symptoms include swollen lymph nodes, joint aches, and red eyes. Rubella normally cures spontaneously and without consequences, but it has a significant impact on pregnant women because the virus can cross the placenta and impair fetal development, leading to CRS. Rubella has no specific therapy and usually goes away on its own within a short period. Treatment is generally aimed at reducing symptoms like temperature and discomfort. Rubella control is primarily based on prevention. Rubella is highly preventable with the MMR vaccine, which all children should receive as part of their usual vaccination plan. Adults who have not been vaccinated or have never had the disease should consider getting vaccinated, especially if they intend to visit locations where Rubella is common [1-4]. The virus that causes the sickness flourishes in both tropical and subtropical climates. Rubella has the potential to be severely damaging to a developing fetus when pregnant. The primary modes of transmission for postnatal rubella are direct contact with contaminated nasopharyngeal secretions and inhalation of virus-carrying airborne droplets. In susceptible pregnant women, the virus can penetrate the placenta and propagate throughout the fetus's vascular system. Nonimmune pregnant women who have measles risk harming their fetuses and neonates. Prenatal immunization can help prevent the

sickness. Vaccination is not advised during pregnancy due to the presence of live, weakened viruses, which could potentially pose risks to the fetus [5].

Gao et al. [6] examined the influence of rubella vaccination on the disease's epidemiology in Australia. Thompson conducted a systematic review of the literature, identifying quantitative models of measles and rubella transmission dynamics and highlighting crucial findings pertinent to future modeling endeavors [7]. Thompson and Badizadegan developed a dynamic disease model for the propagation of measles and rubella. They subsequently applied this model to 180 WHO member states and three additional regions, collectively representing over 99.5% of the global population in 2013 [8]. The model took into account seasonality, age-specific mixing patterns, and the potential for preferential mixing among under-vaccinated subgroups, all factors that could contribute to disparities in vaccination coverage affecting disease transmission. Saito et al. applied this model to rubella notification data from 2012 to 2013, presenting a discrete-time and spatial epidemic model [9]. In a separate study, Alleman et al. conducted a rubella antibody seroprevalence assessment using serum samples collected in 2008-2009 from 1605 pregnant women aged 15-46 attending seven prenatal care facilities across three regions in the Democratic Republic of the Congo [10].

The rubella virus was widespread across all three regions, with most cases of viral exposure and seroconversion occurring before the age of 15. However, between 10% and 20% of the mothers were at risk of contracting the rubella virus and having a child with CRS. Wu et al. used an age-structured transmission model to compare the effectiveness of seven different vaccine approaches in lowering the CRS illness burden in East Java [11]. The model predictions were tested to see how resistant they were to changes in vaccine coverage and other important epidemiological parameters. Chen et al. [12] performed a phylogenetic study on rubella virus isolates and constructed a phylogenetic tree based on the World Health Organization standard sequencing window. Gerberry and Milner extend the SIR model to include individuals who have the disease but are unable to transmit it to others to analyze the dynamics of the latent class. For the model, both a homoclinic bifurcation and the existence of Hopf bifurcations are demonstrated [13]. Baleanu et al. used the Caputo-Fabrizio fractional derivative to investigate the rubella disease model and developed a three-step Adams-Bashforth scheme for the mathematical solution of the liver model. The existence and uniqueness of the solution were argued using fixed point theory [14]. Buonomo has explored an SEIR epidemic model with vertical transmission where the optimal control theory has been applied to assess the vaccination strategy on the model dynamics. The optimality system was derived and solved numerically [15]. Ahmad et al. came up with a deterministic SEITR model of Rubella by applying the Non-Standard Finite Difference (NSFD) method to achieve the numerical solution of the disease [16]. Xu et al. studied a Rubella model with a fractal-fractional exponential decay kernel [17]. Al Qurashi has researched the Rubella virus by fractal-fractional differential and integral operators [18]. Tilahun proposed both a deterministic and stochastic model to discuss Rubella dynamics by considering vertical transmission and environmental factors [19]. Rasit and Tuncer applied a measles model that tends to forecast measles in Turkey from 1970 to 2021 [20]. Bhavithra and Devi analyzed the dynamics of a measles pandemic with the help of a fuzzy SIR model [21]. Khan et al. studied the discrete-time phytoplankton-zooplankton model with some impacts of external toxicity in the phytoplankton population [22]. Abbas et al. have proposed a model representing the dynamics of methamphetamine and performed local and global stability and sensitivity analysis [23]. Alqahtani et al. presented deterministic and stochastic models for the study of the dynamics of MERS-CoV transmission and its epidemic potential [24]. Sheergojri et al. offered a fuzzy Gompertz growth model to reduce uncertainty in modeling tumor population dynamics [25]. Li et al. investigated the Sasa-Satsuma equation using ideas from planar dynamical theory and the beta differential operator. The bifurcation, equilibrium, and sensitivity of the system are also analyzed [26]. Xu et al. studied the plankton population dynamical system accompanying delay [27]. Xu et al. investigated fractional neural networks involving delays [28, 29]. Baber et al. investigated the stochastic Chen-Lee-Liu equation numerically and analytically [30]. Xu et al. studied the dynamics and chaotic behavior of a fish farm model containing populations of nutrients and mussels with different kernels insight of fractal-fractional operator [31]. Li and Wu examined SCIR models for COVID-19 dynamics with and without immigration to derive conditions for stability and properties at the Hopf bifurcation [32]. Tong et al. discussed deterministic and stochastic models for rumor propagation that also incorporate media coverage and age-dependent education [33]. The dynamics for a reaction-diffusion SEIR epidemic model driven by a mass action

infection mechanism in a heterogeneous environment are analyzed [34]. Researchers also investigated the local and global stabilities of the disease dynamical models, [35-41], just to mention a few.

The NSFD theory was first proposed by Mickens [42]. It plays a crucial role in the mathematical and numerical modeling of diseases. It has proven to be one of the most valuable strategies in recent years [43]. One of the most important aspects is the complete absence of the fundamental numerical instabilities that plague traditional finite difference schemes. Numerous differential equations have been solved using the EFD and NSFD techniques [44-46] for example. Verma and Kayenat investigated the generalized Burgers-Huxley (GBH) equation using the NSFD approach. The maximum error of NSFD solutions is compared to the maximum error of several other methods to demonstrate the superiority of the proposed method. The CPU time for all computations is also compared, with the results demonstrating that the NSFD scheme produces accurate results in a matter of seconds, saving time [47]. Raza et al. explored a stochastic HIV/AIDS model within a bisexual community, employing NSFD theory, which incorporates counseling and antiviral medication. The findings suggest that the stochastic model proves to be more practical compared to the deterministic model [48]. The NSFD stochastic scheme retains all of the disease dynamical model's important attributes. Iqbal et al. investigated NSFD theory-based fractional order HIV/AIDS transmission [49]. Ahmed et al. [50] developed a new HIV CD4+T cell reaction-diffusion model. Allehiany et al. [51] applied the NSFD scheme to study a Covid-19 model incorporating fuzziness for its numerical resolution. Similarly, Alhebshi et al. [52] explored a computer virus model utilizing fuzzy criteria. The investigated models undergo equilibrium and reproduction analysis. Both the forward Euler method and NSFD schemes were utilized for numerical solutions. It was noted that while the NSFD scheme maintained stability, convergence, and positivity, the forward Euler method did not uphold these fundamental characteristics of epidemic models.

The primary objective of this study is to investigate a dependable implicit numerical integration method for modeling rubella disease. We contend that the tool for the proposed implicit numerical integration technique is trustworthy, efficient, and encompasses all the dynamic properties necessary for modeling long-term illness behavior. The current work is interesting since it is the first to study the dynamics of the rubella virus disease using NSFD conditions. This work additionally investigates the constructed scheme's convergence, positivity, boundedness, and consistency. Vaccination plays a crucial role in preventing rubella. Rubella vaccination, typically given as part of the MMR vaccine, is highly effective in preventing rubella infection. The vaccine contains a weakened live virus that stimulates the body's immune system to produce antibodies to fight the virus. By providing immunity to the virus, the rubella vaccine can prevent the spread of the disease, particularly to vulnerable populations, such as pregnant women, who can experience serious complications if they contract rubella during pregnancy. The effects of vaccination on infected populations are also studied in this work. The novelty of the paper consists of constructing, implementing, and analyzing a first-order numerical scheme under the novel conditions of the NSFD senses. This approach is carefully framed to a specific model of the transmission dynamics of Rubella. To the best of our knowledge, this model has never been examined using NSFD conditions; hence, it constitutes a novel contribution to the literature.

2 Rubella Virus Transmission and Vaccination Model

We considered the rubella model [53].

$$\frac{dS}{dt} = \alpha N - pqS - (1 - pq)\beta SI - \mu S, \quad (1)$$

$$\frac{dE}{dt} = (1 - pq)\beta SI + \beta RI - \delta E - \mu E, \quad (2)$$

$$\frac{dI}{dt} = \delta E - \gamma I - \mu I, \quad (3)$$

$$\frac{dR}{dt} = \gamma I - \beta RI - \mu R, \quad (4)$$

$$\frac{dV}{dt} = pqS - \mu V. \quad (5)$$

Here, the parameters S , E , I , R , V and N are used to denote susceptible, exposed, infectious, recovered, vaccinated and total populations respectively. The variables used above are described in Table 1 and Fig. 1 shows the flowchart of the above model.

Table 1: The description of the Parameter

Parameter	Description
α	Birth rate
μ	Rate of natural death
p	percentage of those at risk who have had vaccinations
q	The effectiveness level of vaccination
β	Rate of contact
δ	Rate of transformation from latent to infectious
γ	Recovery rate




Figure 1. Flowchart of the model

2.1 Positivity and Boundedness of the Model

For the positivity, consider the system (1-5) as follows.

$$\frac{dS}{dt} \Big|_{S=0} = \alpha N \geq 0.$$

$$\frac{dE}{dt} \Big|_{E=0} = (1 - pq)\beta SI + \beta RI \geq 0.$$

$$\frac{dI}{dt} \Big|_{I=0} = \delta E \geq 0,$$

$$\frac{dR}{dt} \Big|_{R=0} = \gamma I \geq 0.$$

$$\frac{dV}{dt} \Big|_{V=0} = pqS \geq 0.$$

as desired.

For boundedness, consider the population function as follows at any time t .

$$\frac{dN}{dt} = \frac{dS}{dt} + \frac{dE}{dt} + \frac{dI}{dt} + \frac{dR}{dt} + \frac{dV}{dt},$$

$$\frac{dN}{dt} = \alpha N - \mu N,$$

$$\frac{dN}{dt} = N(\alpha - \mu),$$

$$\frac{dN}{dt} \leq N(\alpha - \mu).$$

Since, $S(t) + E(t) + I(t) + R(t) + V(t) \leq N$,

$$N(t) \leq N(0)e^{(\alpha - \mu)t}, \quad t \geq 0,$$

as desired.

2.2 Stability Analysis

The model has a DFE point $p_0 = (\frac{\alpha}{pq+\mu}, 0, 0, 0, \frac{pq\alpha}{(pq+\mu)\mu})$ and an EE point $p(s^*, e^*, i^*, r^*, v^*)$ as an inherent balance. A second equilibrium has no simple form, so it is only stated by $(s^*, e^*, i^*, r^*, v^*)$. As a result, the analysis was limited to an equilibrium that was devoid of sickness. Jacobian of the system (1-5) is

$$J = \begin{bmatrix} -pq - (1-pq)\beta I - \mu & 0 & -(1-pq)\beta S & 0 & 0 \\ (1-pq)\beta I & -\delta - \mu & (1-pq)\beta S + \beta R & \beta I & 0 \\ 0 & \delta & -\gamma - \mu & 0 & 0 \\ 0 & 0 & \gamma - \beta R & -\mu & 0 \\ pq & 0 & 0 & 0 & -\mu \end{bmatrix}$$

and

$$J(p_0) = \begin{bmatrix} -pq - \mu & 0 & \frac{-\alpha\beta(1-pq)}{pq+\mu} & 0 & 0 \\ 0 & -\delta - \mu & \frac{\alpha\beta(1-pq)}{pq+\mu} & 0 & 0 \\ 0 & \delta & -\gamma - \mu & 0 & 0 \\ 0 & 0 & \gamma & -\mu & 0 \\ pq & 0 & 0 & 0 & -\mu \end{bmatrix}$$

The eigenvalues of the above matrix are $\lambda_1 = \lambda_2 = -\mu$, $\lambda_3 = -\mu - pq$, $\lambda_{4,5} = -\frac{1}{2(pq+\mu)}(A \pm \sqrt{B})$, where $A = (pq + \mu)(\gamma + \delta + 2\mu)$ and $B = (pq + \mu)^2(\delta - \gamma)^2 + (pq + \mu)(1 - pq)4\alpha\beta\delta$ [53]. Since all eigenvalues are less than one, it indicates that the system (1-5) is locally stable at the DEF point p_0 .

Theorem 1: The system at DFE $(S^1, E^1, I^1, R^1, V^1) = (\frac{\alpha}{pq+\mu}, 0, 0, 0, \frac{pq\alpha}{(pq+\mu)\mu})$ is globally asymptotically stable if $R_0 < 1$.

Proof: Consider the Lyapunov function, $U: \Omega \rightarrow R$ as follows:

$$U = \left(S - S^1 - S^1 \ln \frac{S}{S^1} \right) + \left(E - E^1 - E^1 \ln \frac{E}{E^1} \right) + \left(I - I^1 - I^1 \ln \frac{I}{I^1} \right) + \left(R - R^1 - R^1 \ln \frac{R}{R^1} \right) + \left(V - V^1 - V^1 \ln \frac{V}{V^1} \right),$$

$$\frac{dU}{dt} = \left(1 - \frac{S^1}{S} \right) \frac{dS}{dt} + \frac{dE}{dt} + \frac{dI}{dt} + \frac{dR}{dt} + \left(1 - \frac{V^1}{V} \right) \frac{dV}{dt},$$

$$\frac{dU}{dt} = \left(1 - \frac{S^1}{S} \right) (\alpha - pqS - (1-pq)\beta SI - \mu S) + (1-pq)\beta SI + \beta RI - \delta E - \mu E + \delta E - \gamma I - \mu I + \gamma I - \beta RI - \mu R + \left(1 - \frac{V^1}{V} \right) (pqS - \mu V),$$

$$\frac{dU}{dt} = -\frac{\alpha(S-S^1)^2}{SS^1} - \frac{pqS(V-V^1)^2}{VV^1},$$

$$\frac{dU}{dt} = -\left(\frac{\alpha(S-S^1)^2}{SS^1} + \frac{pqS(V-V^1)^2}{VV^1} \right) < 1.$$

From the above calculation, it can be observed that $\frac{dU}{dt} < 0$, if $R_0 < 1$ and $\frac{dU}{dt} = 0$ if $S = S^1, E = E^1, I = I^1, R = R^1, V = V^1$. As a result, the system is globally asymptotically stable at C_1 .

2.3 Sensitivity Analysis

The basic reproduction number of the studied model [53] is

$$R_0 = \frac{(1-pq)\alpha\beta\delta}{(pq+\mu)(\mu+\delta)(\mu+\gamma)}.$$

The sensitivity of a parameter χ can be calculated as [54]

$$\xi(\chi) = \frac{\chi}{R_0} \frac{\partial R_0}{\partial \chi}.$$

To find the sensitivity of the parameter β we have,

$$\xi(\beta) = \frac{\beta}{R_0} \frac{\partial R_0}{\partial \beta} = \frac{\beta}{\frac{(1-pq)\alpha\beta\delta}{(pq+\mu)(\mu+\delta)(\mu+\gamma)}} \frac{\partial \frac{(1-pq)\alpha\beta\delta}{(pq+\mu)(\mu+\delta)(\mu+\gamma)}}{\partial \beta} = 1.$$

$$\xi(\alpha) = \frac{\alpha}{\frac{(1-pq)\alpha\beta\delta}{(pq+\mu)(\mu+\delta)(\mu+\gamma)}} \frac{\partial \frac{(1-pq)\alpha\beta\delta}{(pq+\mu)(\mu+\delta)(\mu+\gamma)}}{\partial \alpha} = 1.$$

$$\xi(\delta) = \frac{\delta}{\frac{(1-pq)\alpha\beta\delta}{(pq+\mu)(\mu+\delta)(\mu+\gamma)}} \frac{\partial \frac{(1-pq)\alpha\beta\delta}{(pq+\mu)(\mu+\delta)(\mu+\gamma)}}{\partial \delta} = 1.$$

$$\begin{aligned} \xi(p) &= \frac{p}{\frac{\alpha\beta\delta - pq\alpha\beta\delta}{pq\mu^2 + pq\mu\gamma + \mu^3 + \mu^2\gamma + pq\mu\delta + pq\gamma\delta + \mu^2\delta + \mu\delta\gamma}} \frac{\partial \left(\frac{\alpha\beta\delta - pq\alpha\beta\delta}{pq\mu^2 + pq\mu\gamma + \mu^3 + \mu^2\gamma + pq\mu\delta + pq\gamma\delta + \mu^2\delta + \mu\delta\gamma} \right)}{\partial p}, \\ &= \frac{p(-\alpha\beta\delta q)((pq+\mu)(\mu+\delta)(\mu+\gamma) - (q\mu^2 + q\mu\gamma + q\mu\delta + q\gamma\delta)(\alpha\beta\delta - \alpha\beta\delta pq))}{(pq+\mu)(\mu+\delta)(\mu+\gamma)}. \end{aligned}$$

$$\begin{aligned} \xi(q) &= \frac{q}{\frac{\alpha\beta\delta - pq\alpha\beta\delta}{pq\mu^2 + pq\mu\gamma + \mu^3 + \mu^2\gamma + pq\mu\delta + pq\gamma\delta + \mu^2\delta + \mu\delta\gamma}} \frac{\partial \left(\frac{\alpha\beta\delta - pq\alpha\beta\delta}{pq\mu^2 + pq\mu\gamma + \mu^3 + \mu^2\gamma + pq\mu\delta + pq\gamma\delta + \mu^2\delta + \mu\delta\gamma} \right)}{\partial q}, \\ &= \frac{p(-\alpha\beta\delta p)((pq+\mu)(\mu+\delta)(\mu+\gamma) - (p\mu^2 + p\mu\gamma + p\mu\delta + p\gamma\delta)(\alpha\beta\delta - \alpha\beta\delta pq))}{(pq+\mu)(\mu+\delta)(\mu+\gamma)(1-pq)}. \end{aligned}$$

$$\xi(\mu) = \frac{\mu}{\frac{(1-pq)\alpha\beta\delta}{(pq+\mu)(\mu+\delta)(\mu+\gamma)}} \frac{\partial \frac{(1-pq)\alpha\beta\delta}{(pq+\mu)(\mu+\delta)(\mu+\gamma)}}{\partial \mu} = \frac{-\mu((2pq\mu + pq\gamma + 3\mu^2 + 2\mu\gamma + pq\delta + 2\mu\delta + \gamma\delta)(\alpha\beta\delta - \alpha\beta\delta pq))}{(pq+\mu)(\mu+\delta)(\mu+\gamma)(1-pq)(\alpha\beta\delta)}.$$

$$\xi(\gamma) = \frac{\gamma}{\frac{(1-pq)\alpha\beta\delta}{(pq+\mu)(\mu+\delta)(\mu+\gamma)}} \frac{\partial \frac{(1-pq)\alpha\beta\delta}{(pq+\mu)(\mu+\delta)(\mu+\gamma)}}{\partial \gamma} = \frac{-\gamma((pq\mu + \mu^2 + pq\delta + \mu\delta)(\alpha\beta\delta - \alpha\beta\delta pq))}{(pq+\mu)(\mu+\delta)(\mu+\gamma)(1-pq)(\alpha\beta\delta)}.$$

The calculated findings indicate that β , α and δ are sensitive parameters. Increasing sensitive parameters increases R_0 , and vice versa. On the other side, the parameters p , μ and γ are insensitive and any increase or decrease in these variables do not affect the value of R_0 .

3 Numerical Modelling

In this section, we will develop three different numerical schemes for the solution of the studied model.

3.1 Forward Euler Method

Considering the system of equations (1-5), we develop a forward Euler scheme. Constructing a numerical scheme for the forward Euler involves making the following supposition.

$$S(t) \approx S_n, E(t) \approx E_n, I(t) \approx I_n, V(t) \approx V_n \text{ and } R(t) \approx I_n.$$

$$S^{n+1} = S^n + h(\alpha - pqS^n - (1-pq)\beta S^n I^n - \mu S^n), \quad (6)$$

$$E^{n+1} = E^n + h((1-pq)\beta S^n I^n + \beta R^n I^n - \delta E^n - \mu E^n), \quad (7)$$

$$I^{n+1} = I^n + h(\delta E^n - \gamma I^n - \mu I^n), \quad (8)$$

$$R^{n+1} = R^n + h(\gamma I^n - \beta R^n I^n - \mu R^n), \quad (9)$$

$$V^{n+1} = V^n + h(pqS^n - \mu V^n). \quad (10)$$

3.2 Runge- Kutta Scheme of Fourth Order (R-K4 Method)

Step 1

$$\begin{aligned} k_1 &= h(\alpha - pqS^n - (1 - pq)\beta S^n I^n - \mu S^n), \\ l_1 &= h((1 - pq)\beta S^n I^n + \beta R^n I^n - \delta E^n - \mu E^n), \\ m_1 &= h(\delta E^n - \gamma I^n - \mu I^n), \\ n_1 &= h(\gamma I^n - \beta R^n I^n - \mu R^n), \\ o_1 &= h(pqS^n - \mu V^n). \end{aligned}$$

Step 2

$$\begin{aligned} k_2 &= h(\alpha - pq(S^n + \frac{k_1}{2}) - (1 - pq)\beta(S^n + \frac{k_1}{2})(I^n + \frac{m_1}{2}) - \mu(S^n + \frac{k_1}{2})), \\ l_2 &= h((1 - pq)\beta(S^n + \frac{k_1}{2})(I^n + \frac{m_1}{2}) + \beta(R^n + \frac{n_1}{2})(I^n + \frac{m_1}{2}) - \delta(E^n + \frac{l_1}{2}) - \mu(E^n + \frac{l_1}{2})), \\ m_2 &= h(\delta(E^n + \frac{l_1}{2}) - \gamma(I^n + \frac{m_1}{2}) - \mu(I^n + \frac{m_1}{2})), \\ n_2 &= h(\gamma(I^n + \frac{m_1}{2}) - \beta(R^n + \frac{n_1}{2})(I^n + \frac{m_1}{2}) - \mu(R^n + \frac{n_1}{2})), \\ o_2 &= h(pq(S^n + \frac{k_1}{2}) - \mu(V^n + \frac{o_1}{2})). \end{aligned}$$

Step 3

$$\begin{aligned} k_3 &= h(\alpha - pq(S^n + \frac{k_2}{2}) - (1 - pq)\beta(S^n + \frac{k_2}{2})(I^n + \frac{m_2}{2}) - \mu(S^n + \frac{k_2}{2})), \\ l_3 &= h((1 - pq)\beta(S^n + \frac{k_2}{2})(I^n + \frac{m_2}{2}) + \beta(R^n + \frac{n_2}{2})(I^n + \frac{m_2}{2}) - \delta(E^n + \frac{l_2}{2}) - \mu(E^n + \frac{l_2}{2})), \\ m_3 &= h(\delta(E^n + \frac{l_2}{2}) - \gamma(I^n + \frac{m_2}{2}) - \mu(I^n + \frac{m_2}{2})), \\ n_3 &= h(\gamma(I^n + \frac{m_2}{2}) - \beta(R^n + \frac{n_2}{2})(I^n + \frac{m_2}{2}) - \mu(R^n + \frac{n_2}{2})), \\ o_3 &= h(pq(S^n + \frac{k_2}{2}) - \mu(V^n + \frac{o_2}{2})). \end{aligned}$$

Step 4

$$\begin{aligned} k_4 &= h(\alpha - pq(S^n + k_3) - (1 - pq)\beta(S^n + k_3)(I^n + m_3) - \mu(S^n + k_3)), \\ l_4 &= h((1 - pq)\beta(S^n + k_3)(I^n + m_3) + \beta(R^n + n_3)(I^n + m_3) - \delta(E^n + l_3) - \mu(E^n + l_3)), \\ m_4 &= h(\delta(E^n + m_3) - \gamma(I^n + m_3) - \mu(I^n + m_3)), \\ n_4 &= h(\gamma(I^n + m_3) - \beta(R^n + n_3)(I^n + m_3) - \mu(R^n + n_3)), \\ o_4 &= h(pq(S^n + k_3) - \mu(V^n + o_3)). \end{aligned}$$

Final Step

$$S^{n+1} = S^n + \frac{1}{6}(k_1 + 2k_2 + 2k_3 + k_4), \quad (11)$$

$$E^{n+1} = E^n + \frac{1}{6}(l_1 + 2l_2 + 2l_3 + l_4), \quad (12)$$

$$I^{n+1} = I^n + \frac{1}{6}(m_1 + 2m_2 + 2m_3 + m_4), \quad (13)$$

$$R^{n+1} = R^n + \frac{1}{6}(n_1 + 2n_2 + 2n_3 + n_4), \quad (14)$$

$$V^{n+1} = V^n + \frac{1}{6}(o_1 + 2o_2 + 2o_3 + o_4). \quad (15)$$

3.3 NSFD Scheme

$$S^{n+1} = \frac{S^n + h\alpha}{1 + h(pq + (1 - pq)\beta I^n + \mu)}, \quad (16)$$

$$E^{n+1} = \frac{S^n + h((1-pq)\beta S^n I^n + \beta R^n I^n)}{1+h(\delta+\mu)}, \quad (17)$$

$$I^{n+1} = \frac{I^n + h\delta E^n}{1+h(\gamma+\mu)}, \quad (18)$$

$$R^{n+1} = \frac{R^n + h\gamma I^n}{1+h(\beta I^n + \mu)}, \quad (19)$$

$$V^{n+1} = \frac{V^n + hhpqS^n}{(1+h\mu)}. \quad (20)$$

3.4 Convergence Analysis

In this section, we will conduct the convergence analysis of the NSFD scheme applied to the SEIRV model, focusing on the DFE point. Jacobian of the NSFD scheme is

$$J = \begin{bmatrix} \frac{1}{1+h(pq+(1-pq)\beta I+\mu)} & 0 & -\frac{h(1-pq)(S+h\alpha)}{(1+h(pq+(1-pq)\beta I+\mu))^2} & 0 & 0 \\ \frac{h(1-pq)\beta I}{1+h(\delta+\mu)} & \frac{1}{1+h(\delta+\mu)} & \frac{h((1-pq)\beta S+\beta R)}{1+h(\delta+\mu)} & \frac{h\beta I}{1+h(\delta+\mu)} & 0 \\ 0 & \frac{h\delta}{1+h(\gamma+\mu)} & \frac{1}{1+h(\gamma+\mu)} & 0 & 0 \\ 0 & 0 & \frac{h\gamma(1+h(\beta I+\mu)-h\beta(R+h\gamma I))}{(1+h(\beta I+\mu))^2} & \frac{1}{1+h(\beta I+\mu)} & 0 \\ \frac{hpq}{1+h\mu} & 0 & 0 & 0 & \frac{1}{1+h\mu} \end{bmatrix}.$$

$$J(p_0) = \begin{bmatrix} \frac{1}{1+h(pq+\mu)} & 0 & -\frac{h(1-pq)(\frac{\alpha}{pq+\mu}+h\alpha)}{(1+h(pq+\mu))^2} & 0 & 0 \\ 0 & \frac{1}{1+h(\delta+\mu)} & \frac{h((1-pq)\beta(\frac{\alpha}{pq+\mu}))}{1+h(\delta+\mu)} & 0 & 0 \\ 0 & \frac{h\delta}{1-h(\gamma+\mu)} & \frac{1}{1-h(\gamma+\mu)} & 0 & 0 \\ 0 & 0 & \frac{h\gamma(1+h\mu)}{(1+h\mu)^2} & \frac{1}{1+h\mu} & 0 \\ \frac{hpq}{1+h\mu} & 0 & 0 & 0 & \frac{1}{1+h\mu} \end{bmatrix}.$$

Three eigenvalues of the above matrix are $\lambda_1 = \lambda_2 = \frac{1}{1+\mu}$, $\lambda_3 = \frac{1}{1+h(pq+\mu)}$. The following matrix can be used to get the remaining eigenvalues.

$$A = \begin{bmatrix} \frac{1}{1+h(\delta+\mu)} & \frac{h((1-pq)\beta(\frac{\alpha}{pq+\mu}))}{1+h(\delta+\mu)} \\ \frac{h\delta}{1-h(\gamma+\mu)} & \frac{1}{1-h(\gamma+\mu)} \end{bmatrix}.$$

Now, by using the Bauer F lemma [55], we have

$$\lambda^2 - c\lambda + d = 0,$$

where c is the trace and d is the determinant of the above matrix. Here, $d < 0$, $1 + c + d > 0$ and $1 - c + d > 0$ as desired.

3.5 Positivity Analysis

Theorem: Assume that S, E, I, R and V , the state variables of the NSFD scheme are all positive at $t = 0$; furthermore, if all of the parameters are positive, then $S^{n+1} \geq 0, E^{n+1} \geq 0, I^{n+1} \geq 0, R^{n+1} \geq 0, V^{n+1} \geq 0$.

Proof: Taking into account the state variables S, E, I, R and V of the NSFD scheme (16–20) as,

$$\begin{aligned}
S^{n+1} &= \frac{S^n + h\alpha}{1 + h(pq + (1-pq)\beta I^n + \mu)}, \\
E^{n+1} &= \frac{S^n + h((1-pq)\beta S^n I^n + \beta R^n I^n)}{1 + h(\delta + \mu)}, \\
I^{n+1} &= \frac{I^n + h\delta E^n}{1 + h(\gamma + \mu)}, \\
R^{n+1} &= \frac{R^n + h\gamma I^n}{1 + h(\beta I^n + \mu)}, \\
V^{n+1} &= \frac{V^n + hhpqS^n}{(1 + h\mu)}.
\end{aligned}$$

By combining all the equations in the aforementioned system with $n = 0$, we arrive at the following expression.

$$\begin{aligned}
S^1 &= \frac{S^0 + h\alpha}{1 + h(pq + (1-pq)\beta I^0 + \mu)} \geq 0, \\
E^1 &= \frac{S^0 + h((1-pq)\beta S^0 I^0 + \beta R^0 I^0)}{1 + h(\delta + \mu)} \geq 0, \\
I^1 &= \frac{I^0 + h\delta E^0}{1 + h(\gamma + \mu)} \geq 0, \\
R^1 &= \frac{R^0 + h\gamma I^0}{1 + h(\beta I^0 + \mu)} \geq 0, \\
V^1 &= \frac{V^0 + hhpqS^0}{(1 + h\mu)} \geq 0.
\end{aligned}$$

Now, by substituting $n = 1$ in, we can proceed to the next step.

$$\begin{aligned}
S^2 &= \frac{S^1 + h\alpha}{1 + h(pq + (1-pq)\beta I^1 + \mu)} \geq 0, \\
E^2 &= \frac{S^1 + h((1-pq)\beta S^1 I^1 + \beta R^1 I^1)}{1 + h(\delta + \mu)} \geq 0, \\
I^2 &= \frac{I^1 + h\delta E^1}{1 + h(\gamma + \mu)} \geq 0, \\
R^2 &= \frac{R^1 + h\gamma I^1}{1 + h(\beta I^1 + \mu)} \geq 0, \\
V^2 &= \frac{V^1 + hhpqS^1}{(1 + h\mu)} \geq 0.
\end{aligned}$$

Assume that the given equations maintain positivity for $n = 2, 3, 4, \dots, n-1$, i.e. $S^{n+1} \geq 0, E^{n+1} \geq 0, I^{n+1} \geq 0, R^{n+1} \geq 0, V^{n+1} \geq 0$; for $n = 2, 3, 4, \dots, n-1$.

We will now analyze the positivity for a random positive integer n in Z . We find that

$$\begin{aligned}
S^{n+1} &= \frac{S^n + h\alpha}{1 + h(pq + (1-pq)\beta I^n + \mu)} \geq 0, \\
E^{n+1} &= \frac{S^n + h((1-pq)\beta S^n I^n + \beta R^n I^n)}{1 + h(\delta + \mu)} \geq 0, \\
I^{n+1} &= \frac{I^n + h\delta E^n}{1 + h(\gamma + \mu)} \geq 0, \\
R^{n+1} &= \frac{R^n + h\gamma I^n}{1 + h(\beta I^n + \mu)} \geq 0, \\
V^{n+1} &= \frac{V^n + hhpqS^n}{(1 + h\mu)} \geq 0.
\end{aligned}$$

Thus, for any positive integer values of n , the suggested approach ensures positivity for the state variables S, E, I, R and V .

3.6 Boundedness of the NSFD Scheme

Theorem: Let S^0, E^0, I^0, R^0 and V^0 are finite, such that $S^0 + E^0 + I^0 + R^0 + V^0 \leq 1$. Moreover, in the model μ, β, V, α are all positive. Then, $S^{n+1}, E^{n+1}, I^{n+1}$, and R^{n+1} are bounded by recrossing defined real constant b_{n+1} such that $S^{n+1}, E^{n+1}, I^{n+1}$ and $R^{n+1} < b_{n+1} \forall n \in Z^+$ where $b_{n+1} = 5b_n + h\mu + h\beta b_n^2 + hVb_n + hab_n$ and $b_1 = 5 + h\mu + h\beta S^0 I^0 + hVE^0 + h\alpha I^0$.

Proof: Taking into account the state variables S, E, I, R and V of the NSFD scheme (16–20) as,

$$S^{n+1}(1 + h(pq + (1 - pq)\beta I^n + \mu)) = S^n + h\alpha,$$

$$E^{n+1}(1 + h(\delta + \mu)) = E^n + h((1 - pq)\beta S^n I^n + \beta R^n I^n),$$

$$I^{n+1}(1 + h(\gamma + \mu)) = I^n + h\delta E^n,$$

$$R^{n+1}(1 + h(\beta I^n + \mu)) = R^n + h\gamma I^n,$$

$$V^{n+1}(1 + h\mu) = V^n + hpqS^n.$$

Adding all of the equations in the preceding system yields the following expression.

$$S^{n+1} + E^{n+1} + I^{n+1} + R^{n+1} + V^{n+1} + S^{n+1}(h(pq + (1 - pq)\beta I^n + \mu)) + E^{n+1}(h(\delta + \mu)) + I^{n+1}(h(\gamma + \mu)) + R^{n+1}(h(\beta I^n + \mu)) + V^{n+1}(h\mu) = S^n + h\alpha + E^n + h((1 - pq)\beta S^n I^n + \beta R^n I^n) + I^n + h\delta E^n + R^n + h\gamma I^n + V^n + hpqS^n,$$

$$S^{n+1} + E^{n+1} + I^{n+1} + R^{n+1} + V^{n+1} + S^{n+1}(h(pq + (1 - pq)\beta I^n + \mu)) + E^{n+1}(h(\delta + \mu)) + I^{n+1}(h(\gamma + \mu)) + R^{n+1}(h(\beta I^n + \mu)) + V^{n+1}(h\mu) = S^n + E^n + I^n + R^n + V^n + h\alpha + h((1 - pq)\beta S^n I^n + \beta R^n I^n) + h\delta E^n + h\gamma I^n + hpqS^n. \quad (21)$$

Using mathematical induction theory and restrictions placed on the parameters and state variables, we show that the nonstandard numerical scheme is bounded. We get the following when we substitute $n = 0$ in equation (21).

$$(S^1 + E^1 + I^1 + R^1 + V^1) + S^1(1 + h(pq + (1 - pq)\beta I^0 + \mu)) + E^1(1 + h(\delta + \mu)) + I^1(1 + h(\gamma + \mu)) + R^1(1 + h(\beta I^0 + \mu)) + V^1(1 + h\mu) = S^0 + E^0 + I^0 + R^0 + V^0 + h\alpha + h((1 - pq)\beta S^0 I^0 + \beta R^0 I^0) + h\delta E^0 + h\gamma I^0 + hpqS^0,$$

$$(S^1 + E^1 + I^1 + R^1 + V^1) + S^1(1 + h(pq + (1 - pq)\beta I^0 + \mu)) + E^1(1 + h(\delta + \mu)) + I^1(1 + h(\gamma + \mu)) + R^1(1 + h(\beta I^0 + \mu)) + V^1(1 + h\mu) < b_1,$$

$$\Rightarrow (S^1 + E^1 + I^1 + R^1 + V^1) + S^1(1 + h(pq + (1 - pq)\beta I^0 + \mu)) + E^1(1 + h(\delta + \mu)) + I^1(1 + h(\gamma + \mu)) + R^1(1 + h(\beta I^0 + \mu)) + V^1(1 + h\mu) < 5 + h\mu + h\beta S^0 I^0 + hVE^0 + h\alpha I^0,$$

$$\Rightarrow S^1 < b_1,$$

$$E^1 < b_1,$$

$$I^1 < b_1,$$

$$R^1 < b_1,$$

$$V^1 < b_1,$$

Now, by substituting $n = 1$ in equation (21) to reach the following step.

$$(S^2 + E^2 + I^2 + R^2 + V^2) + S^2(1 + h(pq + (1 - pq)\beta I^1 + \mu)) + E^2(1 + h(\delta + \mu)) + I^2(1 + h(\gamma + \mu)) + R^2(1 + h(\beta I^1 + \mu)) + V^2(1 + h\mu) < b_2,$$

$$\Rightarrow (S^2 + E^2 + I^2 + R^2 + V^2) + S^2(1 + h(pq + (1 - pq)\beta i^1 + \mu)) + E^2(1 + h(\delta + \mu)) + I^2(1 + h(\gamma + \mu)) + R^2(1 + h(\beta i^2 + \mu)) + V^2(1 + h\mu) < b_{n+1} = 5b_n + h\mu + h\beta b_n^2 + hVb_n + h\alpha b_n ,$$

$$\Rightarrow S^2 < b_2,$$

$$E^2 < b_2,$$

$$I^2 < b_2,$$

$$R^2 < b_2,$$

$$V^2 < b_2.$$

Now, let us assume that, for positive integers 2, 3, 4, ..., n - 1, the expression (21) is bounded i.e. $S^n < b_n, E^n < b_n, I^n < b_n, R^n < b_n, V^n < b_n, \forall n \in \{2, 3, 4, \dots, n - 1\}$. Now, we look into the boundedness of an integer n.

$$(S^{n+1} + E^{n+1} + I^{n+1} + R^{n+1} + V^{n+1}) + S^{n+1}(1 + h(pq + (1 - pq)\beta I^n + \mu)) + E^{n+1}(1 + h(\delta + \mu)) + I^{n+1}(1 + h(\gamma + \mu)) + R^{n+1}(1 + h(\beta I^n + \mu)) + V^{n+1}(1 + h\mu) = S^n + E^n + I^n + R^n + V^n + h\alpha + h((1 - pq)\beta S^n I^n + \beta r^n I^n) + h\delta E^n + h\gamma I^n + hpqS^n < b_n,$$

$$S^{n+1} < b_{n+1},$$

$$E^{n+1} < b_{n+1},$$

$$I^{n+1} < b_{n+1},$$

$$R^{n+1} < b_{n+1},$$

$$V^{n+1} < b_{n+1}.$$

Therefore, for all positive values of , $S^{n+1}, E^{n+1}, I^{n+1}, R^{n+1}$, and V^{n+1} are constrained within a real number denoted by b_{n+1} .

3.7 Consistency Analysis

This section employs Taylor's series expansion to carry out the consistency analysis of the NSFD scheme. First, we use the Taylor's series expansion (TSE) of S^{n+1} to the first equation of the implicit numerical integration technique system (16-20).

$$S^{n+1} = S^n + h \frac{dS}{dt} + \frac{h^2}{2!} \frac{d^2S}{dt^2} + \frac{h^3}{3!} \frac{d^3S}{dt^3} + \dots,$$

in the following expression:

$$(S^n + h \frac{dS}{dt} + \frac{h^2}{2!} \frac{d^2S}{dt^2} + \frac{h^3}{3!} \frac{d^3S}{dt^3} + \dots)(1 + h(pq + (1 - pq)\beta I^n + \mu)) = S^n + h\alpha,$$

$$S^n + S^n h(pq + (1 - pq)\beta I^n + \mu) + h \frac{dS}{dt}(1 + h(pq + (1 - pq)\beta I^n + \mu)) + \frac{h^2}{2!} \frac{d^2S}{dt^2}(1 + h(pq + (1 - pq)\beta I^n + \mu)) + \frac{h^3}{3!} \frac{d^3S}{dt^3}(1 + h(pq + (1 - pq)\beta I^n + \mu)) + \dots = S^n + h\alpha,$$

We get the following by applying $h \rightarrow 0$,

$$S^n(pq + (1 - pq)\beta I^n + \mu) + \frac{dS}{dt} = \alpha,$$

$$\frac{dS}{dt} = \alpha - S^n(pq + (1 - pq)\beta I^n + \mu),$$

$$\frac{dS}{dt} = \alpha - (pq + (1 - pq)\beta I^n + \mu)S^n.$$

This result shows that our discretized equation is in agreement with the initial equation of the system (1-5). Similarly, we take the following equation and apply the TSE of E^{n+1} :

$$E^{n+1} = E^n + h \frac{dE}{dt} + \frac{h^2}{2!} \frac{d^2E}{dt^2} + \frac{h^3}{3!} \frac{d^3E}{dt^3} + \dots,$$

In the following expression:

$$E^{n+1}(1 + h(\delta + \mu)) = E^n + h((1 - pq)\beta S^n I^n + \beta R^n I^n),$$

$$(E^n + h \frac{dE}{dt} + \frac{h^2}{2!} \frac{d^2E}{dt^2} + \frac{h^3}{3!} \frac{d^3E}{dt^3} + \dots)(1 + h(\delta + \mu)) = E^n + h((1 - pq)\beta S^n I^n + \beta R^n I^n),$$

We get the following by applying $h \rightarrow 0$,

$$E^n + \frac{dE}{dt} = (1 - pq)\beta S^n I^n + \beta R^n I^n,$$

$$\frac{dE}{dt} = (1 - pq)\beta S^n I^n + \beta R^n I^n - E^n.$$

$$\frac{dE}{dt} = (1 - pq)\beta SI + \beta RI - E.$$

Similarly, applying TSE on the third equation of the developed NSFD scheme, we obtain as follows

$$I^{n+1} = I^n + h \frac{dI}{dt} + \frac{h^2}{2!} \frac{d^2I}{dt^2} + \frac{h^3}{3!} \frac{d^3I}{dt^3} + \dots,$$

In the following expression:

$$I^{n+1}(1 + h(\gamma + \mu)) = I^n + h\delta E^n,$$

$$(I^n + h \frac{dI}{dt} + \frac{h^2}{2!} \frac{d^2I}{dt^2} + \frac{h^3}{3!} \frac{d^3I}{dt^3} + \dots)((1 + h(\gamma + \mu))) = I^n + h\delta E^n,$$

We get the following by applying $h \rightarrow 0$,

$$I^n + \frac{dI}{dt} = \delta E^n,$$

$$\Rightarrow \frac{dI}{dt} = \delta E - I.$$

Similarly, TSE of the next variable is

$$R^{n+1} = R^n + h \frac{dR}{dt} + \frac{h^2}{2!} \frac{d^2R}{dt^2} + \frac{h^3}{3!} \frac{d^3R}{dt^3} + \dots,$$

Equation four of the NSFD scheme gives us

$$R^{n+1}(1 + h(\beta I^n + \mu)) = R^n + h\gamma I^n,$$

$$(R^n + h \frac{dR}{dt} + \frac{h^2}{2!} \frac{d^2R}{dt^2} + \frac{h^3}{3!} \frac{d^3R}{dt^3} + \dots)(1 + h(\beta I^n + \mu)) = R^n + h\gamma I^n,$$

We get the following by applying $h \rightarrow 0$,

$$R^n + \frac{dR}{dt} = \delta E^n,$$

$$\Rightarrow \frac{dR}{dt} = \delta E - R.$$

The TSE for the last equation of the studied model is

$$V^{n+1} = V^n + h \frac{dV}{dt} + \frac{h^2}{2!} \frac{d^2V}{dt^2} + \frac{h^3}{3!} \frac{d^3V}{dt^3} + \dots,$$

From equation (20) of the NSFD scheme, we have

$$V^{n+1}(1 + h\mu) = V^n + hpqS^n,$$

$$(V^n + h \frac{dV}{dt} + \frac{h^2}{2!} \frac{d^2V}{dt^2} + \frac{h^3}{3!} \frac{d^3V}{dt^3} + \dots)(1 + h\mu) = V^n + hpqS^n,$$

Applying $h \rightarrow 0$ results in the following,

$$V^n \mu + \frac{dV}{dt} = pqS^n,$$

$$\Rightarrow \frac{dV}{dt} = pqS - v\mu.$$

As a result, the ODES above system and our discretized implicit numerical integration technique are consistent.

4 Numerical Simulations

Tables 2 and 3 show the values of the variables and parameters utilized in the numerical simulations.

Table 2: Values of variable

Variable	Value
S	0.3
E	0.05
I	0.12
R	0.33
V	0.2

Table 3: Values of Parameter

Parameter	Value
α	0.015
μ	0.015
γ	0.01
δ	0.85
p	0.3
q	0.85




Figure 2. Subpopulations using Euler scheme

Figure 2 depicts the graphical behaviors of the subpopulations using the Euler technique. It can be observed that the Euler scheme converges at small step sizes and begins nonphysical oscillations, producing negative values when the step size increases somewhat. Negative values in these models are meaningless as the population cannot be negative. This behavior of the graphs shows that the approach is not suitable for representing rubella illness dynamics.




Figure 3. Subpopulations using the RK-4 scheme

Figure 3 represents the population dynamics using the RK-4 method. The method is stable, and gives positive and convergent solutions at small values of the time step sizes. However, higher step size yield negative outcomes, which are not feasible for population dynamics and this behavior shows that the RK-4 method is also not suitable for representing rubella illness dynamics.




Figure 4. Subpopulations using the NSFD scheme

The subpopulations using the NSFD method are shown in Figure 4. The NSFD scheme remains positive and convergent despite changing step size values. We can conclude that the NSFD scheme is a reliable tool for reflecting rubella virus disease dynamics. These results demonstrate that the NSFD method is a reliable tool while analyzing the dynamics of rubella transmission.




Figure 5: Effects of vaccination on the infected population

Fig. 5 depicts the effects of vaccination. It can be seen that vaccination is inversely proportional to infection, with an increase in vaccination decreasing infection and vice versa. It can be concluded that vaccination is the most effective method of preventing rubella transmission. The MMR vaccine is extremely effective against rubella infection. The vaccine contains live, weakened viruses that stimulate the immune system to produce anti-rubella antibodies. When a large proportion of the population is immunized against rubella, herd immunity develops, which aids in the prevention of the virus's spread.

5 Conclusion

Rubella is typically a mild infectious disease presenting with a rash and swollen lymph nodes for about three days. The illness spreads in the air and through direct contact. It is found among children and adults, though it can be quite severe and risky for pregnant women and their fetuses when developing in the uterus. Vaccination is an important tool in preventing the spread of Rubella and protecting vulnerable populations. This study presents an accurate and trustworthy numerical solution for a Rubella epidemic model using an NSFD approach. Two equilibrium points that can be used to symbolize DFE and endemic, respectively, have been established by the model from a mathematical perspective. Local and global stability for the DFE point has been determined. The sensitivity of parameters is investigated. The model is solved numerically with three different techniques. The graphical results of the developed three methods are compared. According to the simulation, neither presented the conventional, well-known methods Euler and RK-4 are capable of producing reliable results even at very small step sizes. On the other hand, the suggested method gives accurate and convergent solutions for all choices of the values of the step sizes. Furthermore, the NSFD scheme's positivity, convergence, boundedness, and consistency, are investigated. The scheme preserves positivity which is the main feature of these types of models as the compartments in the model consist of the population which cannot be negative. So negative values are meaningless in these models. The Euler and RK-4 methods give negative values at some choices of the values of the step size which makes these methods not suitable for the study of disease dynamics of the studied model. The effects of the Rubella vaccination are also being investigated. Rubella vaccination is typically administered as part of routine childhood immunizations.

Furthermore, vaccination is critical in preventing rubella transmission. It not only protects those who receive the vaccine, but it also contributes to herd immunity, lowering the likelihood of outbreaks and protecting vulnerable populations who are unable to receive the vaccine. The benefits of rubella vaccination extend beyond the prevention of transmission and congenital rubella syndrome. Vaccination also provides economic benefits, such as lowering the cost of disease treatment and complications, as well as reducing lost productivity due to illness. Vaccination is substantially less expensive than treating rubella and its consequences, such as hospitalization for pregnant women with rubella or infants born with CRS. Maintaining high vaccination coverage rates is important for achieving and maintaining herd immunity and protecting vulnerable populations. The current work mainly focuses on the construction of an NSFD scheme for the solution of the Rubella virus disease dynamics model. The work can be extended to fractional, fuzzy, delayed and stochastic senses and many more directions.

Credit authorship contribution statement:

Samiullah: Conceptualization, Methodology, Software, Writing – original draft. **Fazal Dayan:** Data curation, Project administration, Supervision, Formal analysis, Visualization, Investigation. **Muhammad Azizur Rehman:** Validation, Writing – original draft; **Husam A. Neamah:** Supervision, Investigation, Formal analysis, Resources, Writing – review and editing

Declaration of competing interest: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Availability of Data and Materials: The datasets used and/or analyzed during the current study are available from the corresponding author upon reasonable request.

References

1. Wesselhoft C. Rubella (German measles) and congenital deformities. *New England journal of medicine* 1949; 240(7):258-61. <https://doi.org/10.1056/NEJM194902172400706>
2. Edmunds WJ, Van de Heijden OG, Eerola M, Gay NJ. Modelling rubella in Europe. *Epidemiology & Infection* 2000;125(3):617-34. <https://doi.org/10.1017/S0950268800004660>
3. Edlich R, Winters KL, Long III WB. Rubella and congenital rubella (German measles). *Journal of long-term effects of medical implants* 2005;15(3). <https://doi.org/10.1615/JLongTermEffMedImplants.v15.i3.80>
4. Leung AK, Hon KL, Leong KF. Rubella (German measles) revisited. *Hong Kong Medical Journal* 2019; 25(2):134. <https://doi.org/10.12809/hkmj187785>
5. White SJ, Boldt KL, Holditch SJ, Poland GA, Jacobson RM. Measles, mumps, and rubella. *Clinical obstetrics and gynecology* 2012; 55(2):550-9. <https://doi.org/10.1097/GRF.0b013e31824df256>
6. Gao Z, Wood JG, Burgess MA, Menzies RI, McIntyre PB, MacIntyre CR. Models of strategies for control of rubella and congenital rubella syndrome—A 40 year experience from Australia. *Vaccine*. 2013 ;31(4):691-7. <https://doi.org/10.1016/j.vaccine.2012.11.043>

7. Thompson KM. Evolution and use of dynamic transmission models for measles and rubella risk and policy analysis. *Risk Analysis* 2016;36(7):1383-403. <https://doi.org/10.1111/risa.12637>
8. Thompson KM, Badizadegan ND. Modeling the transmission of measles and rubella to support global management policy analyses and eradication investment cases. *Risk Analysis* 2017;37(6):1109-31. <https://doi.org/10.1111/risa.12831>
9. Saito MM, Nishiura H, Higuchi T. Reconstructing the transmission dynamics of rubella in Japan, 2012-2013. *PloS one*. 2018;13(10):e0205889. <https://doi.org/10.1371/journal.pone.0205889>
10. Alleman MM, Wannemuehler KA, Hao L, Perelygina L, Icenogle JP, Vynnycky E et al. Estimating the burden of rubella virus infection and congenital rubella syndrome through a rubella immunity assessment among pregnant women in the Democratic Republic of the Congo: Potential impact on vaccination policy. *Vaccine*. 2016; 34(51):6502-11. <https://doi.org/10.1016/j.vaccine.2016.10.059>
11. Wu Y, Wood J, Khandaker G, Waddington C, Snelling T. Informing rubella vaccination strategies in East Java, Indonesia through transmission modelling. *Vaccine*. 2016; 34(46):5636-5642. <https://doi.org/10.1016/j.vaccine.2016.09.010>
12. Chen M, Zhu Z, Liu D, Huang G, Huang F, Wu J, Zhang T, Xu W, Pang X. Rubella epidemic caused by genotype 1E rubella viruses in Beijing, China, in 2007–2011. *Virology journal* 2013; 10:1-10. <https://doi.org/10.1186/1743-422X-10-122>
13. Gerberry DJ, Milner FA. An SEIQR model for childhood diseases. *Journal of Mathematical Biology* 2009; 59:535-61. <https://doi.org/10.1007/s00285-008-0239-2>
14. Baleanu D, Mohammadi H, Rezapour S. A mathematical theoretical study of a particular system of Caputo–Fabrizio fractional differential equations for the Rubella disease model. *Advances in Difference Equations* 2020; 2020:1-9. <https://doi.org/10.1186/s13662-020-02614-z>
15. Buonomo B. A simple analysis of vaccination strategies for rubella. *Mathematical Biosciences & Engineering* 2011;8(3):677-687. [10.3934/mbe.2011.8.677](https://doi.org/10.3934/mbe.2011.8.677)
16. Ahmad W, Butt AI, Akhtar N, Rafiq M, Gohar M, Idrees Z, Ahmad N. Developing computationally efficient optimal control strategies to eradicate Rubella disease. *Physica Scripta* 2024; 99(3):035202. [10.1088/1402-4896/ad1fc0](https://doi.org/10.1088/1402-4896/ad1fc0)
17. Xu C, Saifullah S, Ali A. Theoretical and numerical aspects of Rubella disease model involving fractal fractional exponential decay kernel. *Results in Physics* 2022; 34:105287. <https://doi.org/10.1016/j.rinp.2022.105287>
18. Al Qurashi MM. Role of fractal-fractional operators in modeling of rubella epidemic with optimized orders. *Open Physics* 2020; 18(1):1111-20. <https://doi.org/10.1515/phys-2020-0217>
19. Tilahun GT, Tolasa TM, Asfaw MD, Wole GA. Stochastic and deterministic models for Rubella dynamics with two doses of vaccination and vertical transmission. *Discrete Dynamics in Nature and Society*. 2024; 2024(1):9697951. <https://doi.org/10.1155/2024/9697951>
20. Rasit OI, Tuncer N, Martcheva M. Mathematical model of measles in Turkey. *Journal of Biological Systems* 2024:1-30. <https://doi.org/10.1142/S0218339024500323>

-
21. Bhavithra HA, Devi SS. Feasibility and stability analysis for basic measles model using fuzzy parameter. *Contemporary mathematics* 2024;897-912. <https://doi.org/10.37256/cm.5120242428>
 22. Khan MS, Samreen M, Gómez-Aguilar JF, Pérez-Careta E. On the qualitative study of a discrete-time phytoplankton-zooplankton model under the effects of external toxicity in phytoplankton population. *Heliyon*. 2022;8(12). <https://doi.org/10.1016/j.heliyon.2022.e12415>
 23. Abbas S, Gómez-Aguilar JF, ur Rahman G, Asif M. Modeling the mechanisms of methamphetamine abuse transmission with immigration impact and relapse. *The European Physical Journal Plus*. 2023; 138(6):484. <https://doi.org/10.1140/epjp/s13360-023-04122-x>
 24. Alqahtani H, Badshah Q, Sakhi S, ur Rahman G, Gómez-Aguilar JF. Qualitative aspects and sensitivity analysis of MERS-Corona epidemic model with and without noise. *Physica Scripta* 2023; 98(12):125018. [10.1088/1402-4896/ad0bb6](https://doi.org/10.1088/1402-4896/ad0bb6)
 25. Sheergojri AR, Iqbal P, Bhat SA, Khaliq R, Gómez-Aguilar JF. Uncertainty analysis of the tumour population logistic model in a fuzzy context. *Annals of Mathematical Sciences and Applications*. 2024; 9(2):309-39. [10.4310/AMSA.2024.v9.n2.a2](https://doi.org/10.4310/AMSA.2024.v9.n2.a2)
 26. Li P, Shi S, Xu C, Rahman MU. Bifurcations, chaotic behavior, sensitivity analysis and new optical solitons solutions of Sasa-Satsuma equation. *Nonlinear Dynamics* 2024;112(9):7405-15. <https://doi.org/10.1007/s11071-024-09438-6>
 27. Xu C, Ou W, Cui Q, Pang Y, Liao M, Shen J, Baber MZ, Maharajan C, Ghosh U. Theoretical exploration and controller design of bifurcation in a plankton population dynamical system accompanying delay. *Discrete and Continuous Dynamical Systems-S*. 2024, doi: 10.3934/dcdss.2024036.
 28. Xu C, Lin J, Zhao Y, Cui Q, Ou W, Pang Y, Liu Z, Liao M, Li P. New results on bifurcation for fractional-order octonion-valued neural networks involving delays. *Network: Computation in Neural Systems* 2024;1-53. <https://doi.org/10.1080/0954898X.2024.2332662>
 29. Xu C, Zhao Y, Lin J, Pang Y, Liu Z, Shen J, Liao M, Li P, Qin Y. Bifurcation investigation and control scheme of fractional neural networks owning multiple delays. *Computational and Applied Mathematics* 2024;43(4):1-33. <https://doi.org/10.1007/s40314-024-02718-2>
 30. Baber MZ, Ahmed N, Xu C, Iqbal MS, Sulaiman TA. A computational scheme and its comparison with optical soliton solutions for the stochastic Chen–Lee–Liu equation with sensitivity analysis. *Modern Physics Letters B* 2024; 2450376. <https://doi.org/10.1142/S0217984924503767>
 31. Xu C, Farman M, Shehzad A. Analysis and chaotic behavior of a fish farming model with singular and non-singular kernel. *International Journal of Biomathematics* 2023;2350105, DOI:10.1142/S179352452350105X
 32. Li H, Wu Y. Dynamics of SCIR Modeling for COVID-19 with Immigration. *Complexity*. 2022;2022(1):9182830. <https://doi.org/10.1155/2022/9182830>
 33. Tong X, Jiang H, Chen X, Li J, Cao Z. Deterministic and stochastic evolution of rumor propagation model with media coverage and class-age-dependent education. *Mathematical Methods in the Applied Sciences* 2023;46(6):7125-39. <https://doi.org/10.1002/mma.8959>

34. Lei C, Li H, Zhao Y. Dynamical behavior of a reaction-diffusion SEIR epidemic model with mass action infection mechanism in a heterogeneous environment. *Discrete Contin. Dyn. Syst. - B.* 2024;29(7):3163-98.
35. Zhang D, Chen F. Global bifurcations and single-pulse homoclinic orbits of a plate subjected to the transverse and in-plane excitations. *Math. Methods Appl. Sci.* 2017;40(12):4338-49.
36. He X, Qiu J, Li X, Cao J. A brief survey on stability and stabilization of impulsive systems with delayed impulses. *Discrete and Continuous Dynamical Systems - S* 2022;15(7):1797-821. <https://doi.org/10.3934/dcdss.2022080>
37. Xing R, Xiao M, Zhang Y, Qiu J. Stability and Hopf bifurcation analysis of an (n+ m)-neuron double-ring neural network model with multiple time delays. *Journal of Systems Science and Complexity.* 2022;35(1):159-78. <https://doi.org/10.1007/s11424-021-0108-2>
38. Zou K, Li X, Wang N, Lou J, Lu J. Stability and stabilization of delayed neural networks with hybrid impulses. *Complexity.* 2020;2020(1):8712027. <https://doi.org/10.1155/2020/8712027>
39. Shao P, Shateyi S. Stability analysis of SEIRS epidemic model with nonlinear incidence rate function. *Mathematics.* 2021;9(21):2644. <https://doi.org/10.3390/math9212644>
40. Wang H, Qi W, Zhang L, Cheng J, Kao Y. Stability and stabilization for positive systems with semi-Markov switching. *Applied Mathematics and Computation* 2020; 379:125252. <https://doi.org/10.1016/j.amc.2020.125252>
41. Jiang X, Chen X, Chi M, Chen J. On Hopf bifurcation and control for a delay systems. *Applied Mathematics and Computation* 2020;370:124906. <https://doi.org/10.1016/j.amc.2019.124906>
42. R.E. Mickens, *Nonstandard Finite Difference Models of Differential Equations*, World Scientific, Singapore (1993), <https://doi.org/10.1142/2081>.
43. Mickens RE. *Applications of nonstandard finite difference schemes*. World Scientific, Singapore ; 2000.
44. Verma AK, Kayenat S. On the convergence of Mickens' type nonstandard finite difference schemes on Lane-Emden type equations. *Journal of Mathematical Chemistry* 2018; 56:1667-706. <https://doi.org/10.1007/s10910-018-0880-y>
45. Kumar Verma A, Kayenat S. On the stability of Micken's type NSFD schemes for generalized Burgers Fisher equation. *Journal of Difference Equations and Applications* 2019;25(12):1706-37. <https://doi.org/10.1080/10236198.2019.1689236>
46. Verma AK, Pandit B, Verma L, Agarwal RP. A review on a class of second order nonlinear singular BVPs. *Mathematics.* 2020;8(7):1045. <https://doi.org/10.3390/math8071045>
47. Verma AK, Kayenat S. An efficient Mickens' type NSFD scheme for the generalized Burgers Huxley equation. *Journal of Difference Equations and Applications* 2020;26(9-10):1213-46. <https://doi.org/10.1080/10236198.2020.1812594>
48. Raza A, Rafiq M, Baleanu D, Shoaib Arif M, Naveed M, Ashraf K. Competitive numerical analysis for stochastic HIV/AIDS epidemic model in a two-sex population. *IET Systems Biology* 2019;13(6):305-15. <https://doi.org/10.1049/iet-syb.2019.0051>

-
49. Iqbal Z, Ahmed N, Baleanu D, Adel W, Rafiq M, Rehman MA, Alshomrani AS. Positivity and boundedness preserving numerical algorithm for the solution of fractional nonlinear epidemic model of HIV/AIDS transmission. *Chaos, Solitons & Fractals* 2020;134:109706. <https://doi.org/10.1016/j.chaos.2020.109706>
 50. Ahmed N, Rafiq M, Adel W, Rezazadeh H, Khan I, Nisar KS. Structure preserving numerical analysis of HIV and CD4+ T-cells reaction diffusion model in two space dimensions. *Chaos, Solitons & Fractals* 2020;139:110307. <https://doi.org/10.1016/j.chaos.2020.110307>
 51. Allehiyany FM, Dayan F, Al-Harbi FF, Althobaiti N, Ahmed N, Rafiq M, Raza A, Elamin M. Bio-inspired numerical analysis of COVID-19 with fuzzy parameters. *Computers, Materials & Continua* 2022;72(2):3213-29. <https://doi.org/10.32604/cmc.2022.025811>
 52. Alhebshi RM, Ahmed N, Baleanu D, Fatima U, Dayan F, Rafiq M, Raza A, Ahmad MO, Mahmoud EE. Modeling of computer virus propagation with fuzzy parameters. *Computers, Materials & Continua* 2023;74(3): 5663-5678. <https://doi.org/10.32604/cmc.2023.033319>
 53. Prawoto BP, Abadi A, Artiono R. Dynamic of re-infection Rubella transmission model with vaccination. In *AIP Conference Proceedings* 2020; (Vol. 2264, No. 1). AIP Publishing. <https://doi.org/10.1063/5.0025674>
 54. Chitnis N, Hyman JM, Cushing JM. Determining important parameters in the spread of malaria through the sensitivity analysis of a mathematical model. *Bulletin of mathematical biology* 2008; 70:1272-96. <https://doi.org/10.1007/s11538-008-9299-0>
 55. F. Brauer and C. Castillo-Chavez, *Mathematical Models in Population Biology and Epidemiology*, vol. 40. in *Texts in Applied Mathematics*, vol. 40. New York, NY: Springer, 2012. doi: 10.1007/978-1-4614-1686-9.