



Once you have Acrobat Reader open on your computer, click on the [Comment](#) tab at the right of the toolbar:



This will open up a panel down the right side of the document. The majority of tools you will use for annotating your proof will be in the [Annotations](#) section, pictured opposite. We've picked out some of these tools below:



### 1. [Replace \(Ins\)](#) Tool – for replacing text.




Strikes a line through text and opens up a text box where replacement text can be entered.

#### How to use it

- Highlight a word or sentence.
- Click on the [Replace \(Ins\)](#) icon in the Annotations section.
- Type the replacement text into the blue box that appears.

standard framework for the analysis of microeconomic activity. Nevertheless, it also led to the development of a number of strategic approaches. The number of competitors in an industry is that the structure of the industry is a key component. The main components of the industry are the number of firms and the level, are exogenous variables. The important words on entry by firms (M henceforth) we open the 'black b



### 2. [Strikethrough \(Del\)](#) Tool – for deleting text.



Strikes a red line through text that is to be deleted.

#### How to use it

- Highlight a word or sentence.
- Click on the [Strikethrough \(Del\)](#) icon in the Annotations section.

there is no room for extra profits as mark-ups are zero and the number of firms (net) values are not determined by market structure. Blanchard ~~and Kiyotaki~~ (1987), perfect competition in general equilibrium. The effects of aggregate demand and supply shocks in the classical framework assuming monopoly. The number of firms is an exogenous number of firms

### 3. [Add note to text](#) Tool – for highlighting a section to be changed to bold or italic.



Highlights text in yellow and opens up a text box where comments can be entered.

#### How to use it

- Highlight the relevant section of text.
- Click on the [Add note to text](#) icon in the Annotations section.
- Type instruction on what should be changed regarding the text into the yellow box that appears.

dynamic responses of mark-ups are consistent with the VAR evidence

sation of the industry. The number of firms in the industry is a key component. The main components of the industry are the number of firms and the level, are exogenous variables. The important words on entry by firms (M henceforth) we open the 'black b



### 4. [Add sticky note](#) Tool – for making notes at specific points in the text.




Marks a point in the proof where a comment needs to be highlighted.

#### How to use it

- Click on the [Add sticky note](#) icon in the Annotations section.
- Click at the point in the proof where the comment should be inserted.
- Type the comment into the yellow box that appears.

and supply shocks. Most of the firms in the industry are small. The number of firms in the industry is a key component. The main components of the industry are the number of firms and the level, are exogenous variables. The important words on entry by firms (M henceforth) we open the 'black b




5. **Attach File** Tool – for inserting large amounts of text or replacement figures.




Inserts an icon linking to the attached file in the appropriate place in the text.

How to use it

- Click on the **Attach File** icon in the Annotations section.
- Click on the proof to where you'd like the attached file to be linked.
- Select the file to be attached from your computer or network.
- Select the colour and type of icon that will appear in the proof. Click OK.




6. **Drawing Markups** Tools – for drawing shapes, lines and freeform annotations on proofs and commenting on these marks. Allows shapes, lines and freeform annotations to be drawn on proofs and for comment to be made on these marks.



How to use it

- Click on one of the shapes in the Drawing Markups section.
- Click on the proof at the relevant point and draw the selected shape with the cursor.
- To add a comment to the drawn shape, move the cursor over the shape until an arrowhead appears.
- Double click on the shape and type any text in the red box that appears.



# Platelet-derived growth factor receptor $\beta$ (PDGFR $\beta$ ) immunohistochemistry highlights activated bone marrow stroma and is potentially predictive for fibrosis progression in prefibrotic myeloproliferative neoplasia

Gábor Méhes, Alexandar Tzankov,<sup>1</sup> Konnie Hebeda,<sup>2</sup> Ioannis Anagnostopoulos,<sup>3</sup>

<sup>3</sup>László Krenács<sup>4</sup> & Judit Bedekovics

Department of Pathology, University of Debrecen, Debrecen, Hungary, <sup>1</sup>Institute of Pathology, University Hospital, Basel, Switzerland, <sup>2</sup>Department of Pathology, Radboud University Nijmegen Medical Centre, Nijmegen, the Netherlands, <sup>3</sup>Institute of Pathology, Charité-University Medicine Berlin, Campus Charité Mitte, Berlin, Germany, and

<sup>4</sup>Laboratory of Tumor Pathology and Molecular Diagnostics, Szeged, Hungary

<sup>2</sup> Date of submission xxxx

Accepted for publication xxxxx

<sup>1</sup> Published online Article Accepted xxxxx

Méhes G, Tzankov A, Hebeda K, Anagnostopoulos I, Krenács L, Bedekovics J

(2015) *Histopathology* DOI: 10.1111/his.12704

## Platelet-derived growth factor receptor $\beta$ (PDGFR $\beta$ ) immunohistochemistry highlights activated bone marrow stroma and is potentially predictive for fibrosis progression in prefibrotic myeloproliferative neoplasia

**Aims:** Myelofibrosis is the result of aberrant stromal activity which is determined routinely by reticulin staining in bone marrow biopsies. As matrix fibres are the product of activated fibroblasts, we analysed fibre accumulation compared to stromal cell activity during myelofibrosis progression using the fibroblast activation marker platelet-derived growth factor receptor  $\beta$  (PDGFR $\beta$ ) by immunohistochemistry.

**Methods and results:** Initial and follow-up bone marrow biopsies from 84 patients with myeloproliferative neoplasia, including 55 cases with primary myelofibrosis, were evaluated from five haematopathology centres. The stromal mass was measured by conventional reticulin staining [myelofibrosis (MF) grade, 0–3] and PDGFR $\beta$ -positive cells using a novel

PDGFR $\beta$  scoring system (0–3). Results were correlated for prediction of progression. The MF grade and the PDGFR $\beta$  score showed excellent correlation (Spearman's  $r = 0.83$ ,  $P < 0.0001$ ). Elevated PDGFR $\beta$  scores (higher than MF-grade) predicted myelofibrosis progression in total with 43% sensitivity and 57% specificity, and short-term (within 1 year) progression with 82% sensitivity and 53% specificity. Progression of prefibrotic disease to manifest myelofibrosis could be forecast with 90% sensitivity and 75% specificity. **Conclusion:** PDGFR $\beta$  highlights stromal cell activation in marrow fibrosis, which is closely related to matrix accumulation, indicating a direct clinical impact especially in prefibrotic myeloproliferative disorders.

**Keywords:** bone marrow, immunohistochemistry, platelet-derived growth factor receptor, primary myelofibrosis

### Introduction

Myelofibrosis (MF) is the result of bone marrow stromal cell activation occurring frequently in different

pathological conditions of the bone marrow.<sup>1–3</sup> Accumulation of the non-haematopoietic stroma consists of both mesenchymal cellular components and interstitial matrix, the latter composed predominantly of reticulin and collagen fibres.<sup>4,5</sup> While the matrix fibre deposition is determined routinely using reticulin silver staining, much less attention is paid to the

Address for correspondence: G Méhes, Department of Pathology, University of Debrecen, H-4032 Debrecen, Nagyerdei krt. 98, Hungary. e-mail: gabor.mehes@med.unideb.hu

Dispatch: 4.5.15	CE: Rama Prabha
No. of pages: 8	PE: Alex
WILEY	
12704	Manuscript No.
H I S	Journal Code
	

1 activation and proliferation of stromal cells. Fibro-  
 2 blasts responsible for the production of reticulin and  
 3 collagen fibres are not prominent in the normal bone  
 4 marrow<sup>4</sup> and become numerous only upon stimula-  
 5 tion from the microenvironment.<sup>6,7</sup> Several growth  
 6 factors, including megakaryocyte-derived basic fibro-  
 7 blastic growth factor (bFGF), transforming growth  
 8 factor (TGF)- $\beta$  and platelet-derived growth factor  
 9 (PDGF), have been proposed to be the most important  
 10 inducers of fibroblast activation and fibre production  
 11 during the process of MF.<sup>8–11</sup> Growth factors act by  
 12 specific triggering of the appropriate surface mem-  
 13 brane receptor pathways in the target cells.<sup>12,13</sup> Pre-  
 14 vious studies have demonstrated that platelet-derived  
 15 growth factor receptors (PDGFR) show a differential  
 16 expression in the mesenchymal cell types of the bone  
 17 marrow and that the PDGFR $\beta$  subunit is expressed  
 18 predominantly by activated fibroblasts in a cell type-  
 19 specific manner.<sup>6,12</sup> Moreover, the extent of the  
 20 PDGFR $\beta$  subunit demonstrated by immunohistochem-  
 21 istry (IHC) in bone marrow biopsy samples correlated  
 22 closely with the degree of matrix fibre accumulation  
 23 in MF.<sup>6</sup> Notably, despite the strong correlation  
 24 between stromal cell activation and fibre accumula-  
 25 tion, fibroblast-related PDGFR $\beta$  immunostaining  
 26 proved to be more prominent than the amount of  
 27 fibre accumulation in selected cases, suggesting a pro-  
 28 gressive potential of the stromal changes.

29 As PDGFR $\beta$  appeared to be useful for the character-  
 30 ization of MF, the question was raised as to whether  
 31 the histological demonstration of activated fibroblasts  
 32 might principally indicate ongoing fibrosis, and thus  
 33 predict the course of MF progression. For this pur-  
 34 pose, a statistically significant number of sequential  
 35 bone marrow samples from the same patients need to  
 36 be evaluated, which are rarely provided. To evaluate  
 37 the utility of PDGFR $\beta$  assessment by IHC and to  
 38 address its predictive potential in the long- and short-  
 39 term follow-up of primary myelofibrosis (PMF), a re-  
 40 spective multicentre study was initiated with the  
 41 participation of the members of the European Bone  
 42 Marrow Working Group. Initial and follow-up bone  
 43 marrow biopsy samples featuring different degrees of  
 44 MF were collected and evaluated for the fibroblast  
 45 activation marker PDGFR $\beta$  using IHC and for the  
 46 fibre products highlighted by conventional reticulin  
 47 silver staining.

## 51 Materials and methods

52 Archival cases of 84 patients having had multiple  
 53 bone marrow examinations were collected by the

following participants: the Institute of Pathology at  
 the University Hospital Basel, Switzerland, the Rad-  
 4 boud University Nijmegen Medical Center, the Neth-  
 5 erlands, the Charité – University Medicine, Berlin,  
 6 Germany, the Laboratory of Tumor Pathology and  
 7 Molecular Diagnostics, Szeged, Hungary and the  
 8 Department of Pathology, University of Debrecen,  
 9 Hungary. Clinical and histological diagnoses and  
 10 basic patient data, including time of sampling, were  
 11 provided in an anonymized fashion. Detailed evalua-  
 12 tion of clinical and laboratory parameters was not  
 13 the aim of this study. Histopathological evaluation of  
 14 the slides was performed in accordance with the gen-  
 15 eral ethical regulations and with the approval of the  
 16 local ethical committees.

17 A total of 193 biopsy samples were obtained from  
 18 84 patients, including 55 PMF, eight essential  
 19 thrombocythaemia (ET), five polycythaemia vera  
 20 (PV), three chronic myeloid leukaemia (CML) and a  
 21 further three unclassifiable myeloproliferative neoplas-  
 22 ias (MPN-U) and four overlapping MPN/myelodys-  
 23 plastic neoplasias (MDS). Myeloproliferative neoplasia  
 24 was associated with other bone marrow disorders or  
 25 conditions in six additional cases [lymphoproliferative  
 26 disorders, granulocyte–macrophage colony-stimulat-  
 27 ing factor (GM-CSF) therapy]. Usually two serial sam-  
 28 ples per case represented the course of the disease,  
 29 although for some cases three or more biopsies were  
 30 provided. The mean follow-up time was 34.4 months  
 31 (range = 2–151 months).

32 Within the study collection, PMF represented the  
 33 largest homogeneous disease group which could be  
 34 evaluated separately. Altogether, 126 initial and fol-  
 35 low-up biopsy samples from 55 PMF cases were anal-  
 36 ysed. The mean follow-up time was 30.1 months  
 37 (range = 2–143 months).

38 Detailed evaluation of stromal fibrosis using reticu-  
 39 lin silver staining and cellular PDGFR $\beta$  expression fol-  
 40 lowing immunohistochemistry was performed at the  
 41 Department of Pathology, University of Debrecen,  
 42 Hungary. To specifically demonstrate PDGFR $\beta$ -posi-  
 43 tive bone marrow cells, samples were incubated with  
 44 the anti-PDGFR $\beta$  (ab-32570; Abcam, Cambridge, UK)  
 45 primary monoclonal antibody at a dilution of 1:100,  
 46 as described previously.<sup>6</sup> Antibody binding was visu-  
 47 alized by the Dako EnVision FLEX/HRP and FLEX  
 48 DAB3 Chromogen detection system (Dako, Glostrup,  
 49 Denmark). The PDGFR $\beta$  score was determined by  
 50 microscopic analysis according to a previously pro-  
 51 posed scoring system.<sup>6</sup> The reticulin grade staining  
 52 was determined following Gömöri's reticulin staining  
 53 according to the European Consensus on grading  
 bone marrow fibrosis.<sup>5</sup> To enhance reproducibility,

both stainings were evaluated in a blinded fashion using the same criteria by two histologists independently, and the results were compared. In case of disagreement a consensus decision was achieved.

Statistical evaluation and graphs were made using the GraphPad Prism software. Mean and standard deviations were calculated for each group and analysed with Student's *t*-test. *P*-values <0.05 were considered statistically significant. Correlations between data sets were obtained using linear regression analysis.

## Results

### RETICULIN GRADE AND PDGFR $\beta$ EXPRESSION CORRELATE STRONGLY IN MF IRRESPECTIVE OF THE TYPE OF MPN


IHC preparations were generally of high quality and optimal for microscopic analysis, irrespective of the admission site, and semiquantitative microscopic evaluations were performed in all 193 samples from 84 patients using the previously established scoring system (Figure 1). The relation of MF grade and PDGFR $\beta$  score for each sample was assessed and a strong correlation was found between the amount of accumulated fibres (MF grade) and the activated stromal cell component expressed by the PDGFR $\beta$  score (Spearman's  $r = 0.83$ ,  $P < 0.0001$ ) (Figure 2A). Evaluation of the 126 samples from 55 cases with PMF diagnosis gave similar results (Spearman's  $r = 0.86$ ,

$P < 0.0001$ ) (Figure 2B). This correlation was statistically constant through all grades of PMF when the initial (Spearman's  $r = 0.84$ ,  $P < 0.0001$ ) and the follow-up samples (Spearman's  $r = 0.82$ ,  $P < 0.0001$ ) were evaluated separately.


### PREDICTION OF PROGRESSION IN ALL MF SAMPLES USING THE PDGFR $\beta$ SCORING SYSTEM

At the time of diagnosis, 14 of 84 cases showed MF-0, 30 cases MF-1, 21 cases MF-2 and 19 cases MF-3, while the PDGFR $\beta$  score was PDGFR $\beta$ -0 in four cases, PDGFR $\beta$ -1 in 31 cases, PDGFR $\beta$ -2 in 24 cases and PDGFR $\beta$ -3 in 25 cases. The PDGFR $\beta$  score was higher than the MF grade in 29 of 84 cases (34.5%), equal to the MF grade in 51 of 84 cases (60.7%) and lower than the MF grade in only four of 84 cases (4.7%). Thirty-four of the total 84 cases showed progression in MF grade during the whole follow-up period, while 40 cases did not show any change. The remaining 10 cases showed regression/reduction of fibrosis. Of these 10 cases, in three cases anti-MF medication was applied, while in a further three cases stem cell transplantation was performed; in the four remaining cases no clinical data were available. Representative morphological changes related to fibrosis progression are presented in Figure 3. We evaluated the relation of initial PDGFR $\beta$  scores with MF progression defined by the increase of MF grade during the follow-up period. Cases with no change or a decrease in fibre content during follow-up were


COLOR



**Figure 1.** Platelet-derived growth factor receptor  $\beta$  (PDGFR $\beta$ ) immunohistochemistry in bone marrow trephine biopsies highlights myelofibrosis-related stromal cell activation. Positive staining represents stromal fibroblasts in different amounts in parallel with reticulin fibre accumulation. A four-grade (0–3) scoring system was uniformly applicable for all initial and follow-up samples collected in our collaborative study (A = score 0, B = score 1, C = score 2, D = score 3).



**Figure 2.** Graphical presentation of the correlation between myelofibrosis (MF) grades and platelet-derived growth factor receptor  $\beta$  (PDGFR $\beta$ ) scores of the same bone marrow samples. The reticulin grade and the PDGFR $\beta$  immunohistochemistry (IHC) score were determined and matched in the 193 samples from all the 84 cases collected within this study (A) and in the 126 samples from the 55 primary myelofibrosis (PMF) cases (B).




**Figure 3.** Examples of the microscopic changes in the successive bone marrow samples from a case with progressive primary myelofibrosis (PMF). Hypercellularity [haematoxylin and eosin (H&E) staining], no fibre accumulation (MF-0, reticulin staining) and mild stromal cell activation [platelet-derived growth factor receptor  $\beta$  (PDGFR $\beta$ ) score-1, immunohistochemistry] was visible in the initial biopsy sample (upper row). The same cellularity accompanied with mild increase in fibrosis (MF-1) and with extended stromal cell activation (PDGFR $\beta$  score-2) indicated progression in the follow-up sample obtained 7 months later (lower row).

considered as non-progressive. Nineteen of the 84 cases (22.6%) had already presented with fully developed myelofibrosis (MF-3) at the time of the initial diagnosis, and thus further progression was no longer

possible. The statistical analysis was performed after their exclusion. Figure 4A shows the outcome of MF in cases with higher ( $n = 28$ ), equal ( $n = 34$ ) and lower ( $n = 3$ ) PDGFR $\beta$  scores compared to MF grade

COLOR



**Figure 4.** Probability of progression respective to the platelet-derived growth factor receptor  $\beta$  (PDGFR $\beta$ ) score. The initially higher score did not predict fibrosis progression in the total sample collection after the removal of MF-3 cases (A). The same result was seen when short-term follow-up (12 months or less) was considered (B). Increase in myelofibrosis (MF) grade was defined as progression, while cases with decrease or no change in the MF grade were considered as non-progressive.

during the whole follow-up period. Using the log-rank test there was no significant difference in the probability of fibrosis progression in the different PDGFR $\beta$  subgroups ( $P = 0.4547$ ). To evaluate the short-term effect of stromal cell PDGFR $\beta$  expression, the same analysis was performed for cases having follow-up samples within the initial 12 months after diagnosis (26 cases). Eleven of the 26 (42.31%) matched short-term sample pairs showed progression, while in 15 cases (57.69%) there was no progression. The log-rank test did not show any significant difference in the probability of fibrosis progression in these short-term follow-up series ( $P = 0.7398$ ) (Figure 4B). PDGFR $\beta$  scores higher than the MF grade did not predict the progression of MF in general ( $n = 65$ ), with appropriate sensitivity (43%) and specificity (57%) after the exclusion of MF-3 cases on a long-term basis (Table 1). In the short-term analysis, considering only the initial 12 months follow-up time ( $n = 26$ ), the elevated PDGFR $\beta$  score proved to be predictive of progression, with 82% sensitivity and 53% specificity (Table 1).

Most interestingly, prefibrotic cases (initial MF-0) showing an elevated PDGFR $\beta$  score (ranging 1–2+) turned frequently to fibrotic cases, as measured by reticulin staining in the repeated biopsy. The progression from the prefibrotic to the fibrotic status could be predicted with 90% sensitivity and 75% specificity based on the initially higher PDGFR staining (Table 1). The prefibrotic samples evaluated here were of heterogeneous origin, including ET (two cases), PV (two cases), PMF (six cases), CML (two cases) and MPN-U (two cases). No other objective clinical (gender, age, medication) or histological (bone marrow involvement, cellularity, inflammation) parameter potentially influencing the extent and dynamics of the bone marrow stromal reaction could be identified.

**Table 1.** Predictive value of the platelet-derived growth factor receptor  $\beta$  (PDGFR $\beta$ ) scoring for myelofibrosis (MF) progression


	All cases	All cases within 1 year follow-up	All prefibrotic cases
Number of cases	65	26	14
TP	15	9	9
TN	17	8	3
FP	13	7	1
FN	20	2	1
Sensitivity	43% (15/35)	82% (9/11)	90% (9/10)
Specificity	57% (17/30)	53% (8/15)	75% (3/4)

TP = true positive; TN = true negative; FP = false positive; FN = false negative.

Specificity and sensitivity was calculated considering all evaluated cases for the full length of follow-up, for the short-term (1 year) follow-up and separately for prefibrotic cases only. End-stage (MF-3) fibrosis was excluded, as these cases were not subjects of further progression.

PREDICTIVE VALUE OF PDGFR $\beta$  IN SELECTED CASES OF PRIMARY MYELOFIBROSIS

The group of 55 cases with a diagnosis of PMF was the largest and most homogeneous disease group within our multicentre study. At the time of diagnosis, six PMF cases proved to be prefibrotic (MF-0) and 20 cases showed MF-1, 11 cases MF-2 and 18 cases MF-3 reticulin grade. At the same time, the PDGFR $\beta$  score was PDGFR $\beta$ -0 in one case, PDGFR $\beta$ -1 in 18



**Figure 5.** Probability of progression in primary myelofibrosis (PMF) reflected by the platelet-derived growth factor receptor  $\beta$  (PDGFR $\beta$ ) score. The initially higher score did not predict progression of fibrosis during the full-length (A) or short-term (B) follow-ups. Initial MF-3 cases were excluded as these were not subjects of further progression.

cases, PDGFR $\beta$ -2 in 15 cases and PDGFR $\beta$ -3 in 21 cases. During disease follow-up, 24 of 55 cases (43.6%) with PMF remained unchanged, while 24 cases (43.6%) showed MF progression and seven cases (12.8%) regression/reduction of the MF degree. The stromal cell reaction measured by the PDGFR $\beta$  score remained unchanged in 33 of 55 cases (60.0%), 16 showed an increase (29.1%) and six cases (10.1%) a decrease during the total follow-up period.

The PDGFR $\beta$  score was higher than the MF grade in 14 of 55 cases (25.5%), equal to the MF grade in 39 of 55 cases (70.9%) and lower than the MF grade in only two of 55 cases (3.6%). Cases with a fully developed myelofibrosis (MF-3) at the time of diagnosis (18 of the 55) were excluded from further evaluation regarding progression. Figure 5A shows the progression of MF in cases with higher ( $n = 16$ ), equal ( $n = 20$ ) and lower ( $n = 1$ ) PDGFR $\beta$  scores compared to the MF grade. There was no significant difference in the probability of fibrosis progression in the different subgroups ( $P = 0.3905$ ). The dynamics of PMF on a short-term basis (within 12 months' follow-up history, a total of 15 cases) was also not associated statistically with the PDGFR $\beta$  scores ( $P = 0.7915$ ), despite a prominent increase in selected cases with progression (Figure 5B).

In PMF, a higher PDGFR $\beta$  score than MF grade predicted the progression of MF with 40% sensitivity and 53% specificity after the exclusion of MF-3 cases ( $n = 37$ ) (Table 2). The short-term ( $n = 15$ ) progression prediction was effective with 83% sensitivity and 44% specificity (Table 2). The six prefibrotic PMF cases enrolled into this study did not allow a conclusion regarding the predictive utility of PDGFR $\beta$  scoring.

The potential effect of the initial PDGFR $\beta$  finding on the follow-up parameters of MF was also evaluated (Table 3). While the correlation between MF

grade and PDGFR $\beta$  score was reproducibly strong within the same bone marrow samples at any time during our investigations, the initial PDGFR $\beta$  score correlated only moderately with either the MF grade or PDGFR $\beta$  score of the endpoint follow-up sample.

## Discussion

Although a previous report also demonstrated PDGFR expression in thrombopoietic cells, our earlier studies and current observations revealed that PDGFR $\beta$  expression is limited to the endosteal and the perivascular stromal niche and reticular fibroblasts in the bone marrow.<sup>14</sup> These differences may be explained by the specificity of applied antibodies. PDGFR $\beta$  is a dynamically expressed and potentially targetable tyrosine-kinase receptor of this set of mesenchymal cell types. The role of PDGFR signalling has been studied extensively experimentally<sup>15</sup> and its role could be demonstrated in fibrotic disorders, such as in local and systemic sclerosis and in neoplasias of mesenchymal origin.<sup>16–19</sup> The increased amounts of PDGFR $\beta$ -expressing fibroblasts also proved to be a characteristic feature of the different degrees of myelofibrosis. Using PDGFR $\beta$  immunohistochemistry, the topography and dynamics of the newly generated fibroblasts can be followed in the diseased bone marrow. Although the link between receptor expression and matrix fibre synthesis is not completely clear, the total amount of the PDGFR $\beta$  immunopositivity was associated strongly with reticulin fibre deposition, which could also be reproduced by the present study. According to current opinion, the degree of MF is an important clinical parameter reflecting disease activity. The amount and distribution of reticulin and collagen fibres, however, is a result of the activity and proliferation of stromal fibroblasts. For this reason, we speculated that the cellular activation marker

**Table 2.** Specificity and sensitivity of the elevated platelet-derived growth factor receptor  $\beta$  (PDGFR $\beta$ ) score to predict progression in primary myelofibrosis (PMF)

	PMF cases	PMF cases within 1 year follow-up	Prefibrotic PMF cases
Number of cases	37	15	6
TP	8	5	4
TN	9	4	0
FP	8	5	1
FN	12	1	1
Sensitivity	40% (8/20)	83% (5/6)	(4/5)
Specificity	53% (9/17)	44% (4/9)	(0/1)

TP = true positive; TN = true negative; FP = false positive; FN = false negative.

Initially, MF-3 cases were excluded, as these were not subjects of further progression. Analysis was performed for all PMF cases within the full follow-up range and also for cases having short-term follow-up. Prefibrotic PMF could not be evaluated separately due to low case number ( $n = 6$ ).

PDGFR $\beta$  might be useful to highlight ongoing cellular processes occurring in association with or even in advance of stromal fibre deposition.

Using a microscopic scoring system, detailed evaluation of 193 bone marrow samples from five European centres clearly validated the close relationship between the conventionally determined MF grade and the fibroblast activation marker PDGFR $\beta$ . Moreover, elevated PDGFR $\beta$  scores indicated stromal activity in a significant portion of the samples, also disclosing early lesions lacking prominent reticulin fibrosis. According to our evolving concept, fibroblast activation may also reflect early phases of stromal reaction, the result of which can only be identified in subsequent bone marrow samples as clinically relevant fibrosis. In our comparison, however, no strong predictive role could be stated for the actual PDGFR $\beta$  status in myeloproliferative disease – even if the follow-up period was limited to only 1 year. Conversely, patients at risk for progressive bone marrow disease could be identified with a sensitivity of 90% and specificity of 75% among cases presenting initially with MF-0 (prefibrotic) if the PDGFR $\beta$  score was elevated (Table 1). Unfortunately, the low number of cases did not allow us to reproduce the same predictive effect

**Table 3.** Correlation of the initial platelet-derived growth factor receptor  $\beta$  (PDGFR $\beta$ ) score with the follow-up status in primary myelofibrosis (PMF); PDGFR $\beta$  score correlated well with the MF grade of the same sample, but regression coefficients did not support a strong relation with the endpoint bone marrow status

	Correlations with initial beta score ( $n = 55$ )		
	MF initial	MF endpoint	Beta endpoint
Spearman's $r$	0.8463	0.4002	0.4931
$P$ -value (two-tailed)	<0.0001	0.0025	0.0001

when PMF was evaluated separately (six prefibrotic/55 PMF cases).

According to current opinion, MF is itself an accompanying condition appearing with variable dynamics in myeloproliferative disease. The course of myelofibrosis is influenced by many factors, and the appreciation of MF progression is difficult and somewhat limited. Early fibrosis may occur in different degrees throughout the bone marrow, frequently with only focal presentation in biopsy samples. In particular, the MF-2 grade may be associated with significant heterogeneity and a broad spectrum in parenchymal reticulin accumulation, making a single-parameter histological evaluation and statistical consideration problematic. The dynamics of fibrosis in bone marrow disease are also reliant upon the local cytokine/growth factor stimuli depending on the composition of the neoplastic microenvironment. Conversely, the fibre production may be affected significantly by the treatment applied. Due to this complexity, fibroblast activation and extracellular fibre accumulation represent two different biological aspects of the same pathological process. As a major finding of our study, we validated the generally strong linear association between bone marrow fibrosis and PDGFR $\beta$ -expressing mesenchymal cells. Further, the elevated PDGFR $\beta$  expression in initial prefibrotic bone marrow lesions indicated progression in the absence of fibre deposition. According to our concept, up-regulation of the PDGF receptor for optimal ligand binding is required very early in advance of significant stromal proliferation and matrix production, a feature that can be demonstrated uniformly in parallel with the neoplastic transformation of the bone marrow. Thus, increased PDGFR $\beta$  expression can be expected in all phases and even before manifest myelofibrosis (MF-0). Finally, the fibrotic process

may be gradually tuned down due to the loss of paracrine stimuli in end-stage fibrotic bone marrow disease (MF-3). For these reasons, the use of stromal activation markers, together with the classical disease characteristics, offers a completely new insight into the pathological processes against a background of ongoing stromal fibrosis. In clinical practice, PDGFR $\beta$  IHC provides valuable additional information when the reticulin staining is difficult to interpret or when immunostainings are preferred to classical histochemistry. Interpretation of the staining pattern requires only minimum effort and the proposed scoring is easily comparable with the reticulin-based grading.

In the current study we focused on the pathological progression of MF as an endpoint of our investigations, which covered mainly histological parameters. Clinical correlations regarding the PDGFR expression similar to the reticulin MF grade are highly difficult to measure due to the heterogeneous nature and the different, individually adapted clinical management of the myeloproliferative neoplasias. Probably for the same reasons, despite a consistent correlation within the same sample, an association between the initial and endpoint MF grade or PDGFR score could not be demonstrated (Table 3).

In summary, despite the above-mentioned biological-histological variables, PDGFR $\beta$  immunohistochemistry proved to be a robust indicator of marrow stromal reaction that could be measured reproducibly in bone marrow biopsies. PDGFR $\beta$  highlights the cellular aspects of fibrosis that are related closely to the reticulin and collagen accumulation. Its potential value to predict progression in early, pre-fibrotic myeloproliferative neoplasias could also be demonstrated based on a small number of retrospective follow-up cases which, however, calls for validation in a larger sample collection.

## Acknowledgements

This work was supported by the TÁMOP-4.2.2.A-11/1/KOV-2012-0045 'Research network on vascular biology/medicine' project grant. Judit Bedekovics was a research fellow of the 'National Excellence Program – Elaborating and operating an inland student and researcher personal support system' (TÁMOP 4.2.4. A/2-11-1-2012-0001) subsidized by the European Union and co-financed by the European Social Fund.

## Conflicts of interest

The authors report no potential conflicts of interest.

## References

1. Kuter DJ, Bain B, Mufti G *et al.* Bone marrow fibrosis: pathophysiology and clinical significance of increased bone marrow stromal fibres. *Br. J. Haematol.* 2007; **139**: 351–362.
2. Koopmans SM, van Marion AM, Schouten HC. Myeloproliferative neoplasia: a review of clinical criteria and treatment. *Neth. J. Med.* 2012; **70**: 159–167.
3. Thiele J, Kvasnicka HM. Myelofibrosis in chronic myeloproliferative disorders – dynamics and clinical impact. *Histol. Histopathol.* 2006; **21**: 1367–1378.
4. Koury MJ, Lichtman MA. Structure of the marrow and the hematopoietic microenvironment. In Lichtman MA ed. *Williams manual of hematology*. 8th ed. New York: McGraw Hill Professional, 2010; 39–74.
5. Thiele J, Kvasnicka HM, Facchetti F *et al.* European consensus on grading bone marrow fibrosis and assessment of cellularity. *Haematologica* 2005; **90**: 1128–1132.
6. Bedekovics J, Kiss A, Beke L *et al.* Platelet derived growth factor receptor-beta (PDGFRbeta) expression is limited to activated stromal cells in the bone marrow and shows a strong correlation with the grade of myelofibrosis. *Virchows Arch.* 2013; **463**: 57–65.
7. Tripodo C, Sangaletti S, Piccaluga PP *et al.* The bone marrow stroma in hematological neoplasms – a guilty bystander. *Nat. Rev. Clin. Oncol.* 2011; **8**: 456–466.
8. Reilly JT. Pathogenesis of idiopathic myelofibrosis: role of growth factors. *J. Clin. Pathol.* 1992; **45**: 461–464.
9. Bonner JC. Regulation of PDGF and its receptors in fibrotic diseases. *Cytokine Growth Factor Rev.* 2004; **15**: 255–273.
10. Bock O, Loch G, Busche G *et al.* Aberrant expression of platelet-derived growth factor (PDGF) and PDGF receptor-alpha is associated with advanced bone marrow fibrosis in idiopathic myelofibrosis. *Haematologica* 2005; **90**: 133–134.
11. Border WA, Noble NA. Transforming growth factor beta in tissue fibrosis. *N. Engl. J. Med.* 1994; **331**: 1286–1292.
12. Heldin CH, Westermark B. Mechanism of action and *in vivo* role of platelet-derived growth factor. *Physiol. Rev.* 1999; **79**: 1283–1316.
13. Funa K, Uramoto H. Regulatory mechanisms for the expression and activity of platelet-derived growth factor receptor. *Acta Biochim. Pol.* 2003; **50**: 647–658.
14. Yoon SY, Tefferi A, Li CY. Cellular distribution of platelet-derived growth factor, transforming growth factor-beta, basic fibroblast growth factor, and their receptors in normal bone marrow. *Acta Haematol.* 2000; **104**: 151–157.
15. Rajkumar VS, Shiwen X, Bostrom M *et al.* Platelet-derived growth factor-beta receptor activation is essential for fibroblast and pericyte recruitment during cutaneous wound healing. *Am. J. Pathol.* 2006; **169**: 2254–2265.
16. Pinzani M, Milani S, Herbst H *et al.* Expression of platelet-derived growth factor and its receptors in normal human liver and during active hepatic fibrogenesis. *Am. J. Pathol.* 1996; **148**: 785–800.
17. Diang XC, Ma LN, Li YF *et al.* Association between serum platelet-derived growth factor BB and degree of liver damage, fibrosis and hepatitis B e antigen (HBeAg) status in CHB patients. *Hepatogastroenterology* 2012; **59**: 2357–2360.
18. Nishioka Y, Azuma M, Kishi M *et al.* Targeting platelet-derived growth factor as a therapeutic approach in pulmonary fibrosis. *J. Med. Invest.* 2013; **60**: 175–183.
19. Klareskog L, Gustafsson R, Scheynius A *et al.* Increased expression of platelet-derived growth factor type B receptors in the skin of patients with systemic sclerosis. *Arthritis Rheum.* 1990; **33**: 1534–1541.

## Author Query Form




Journal: HIS

Article: 12704

Dear Author,

During the copy-editing of your paper, the following queries arose. Please respond to these by marking up your proofs with the necessary changes/additions. Please write your answers on the query sheet if there is insufficient space on the page proofs. Please write clearly and follow the conventions shown on the attached corrections sheet. If returning the proof by fax do not write too close to the paper's edge. Please remember that illegible mark-ups may delay publication.

Many thanks for your assistance.

Query reference	Query	Remarks
1	WILEY: Please insert the published online date for online accepted publication.	
2	WILEY: Please supply date of submission/acceptance.	
3	AUTHOR: Please confirm that given names (red) and surnames/family names (green) have been identified correctly.	
4	AUTHOR: Please check that authors and their affiliations are correct.	
5	AUTHOR: Please advise if you already submitted the accomplished Colour Work Agreement form of your article. If not, please complete the form at the end of this proof and send it to OPI via post or courier. Thank you	

# Colour Work Agreement Form

Form No: SN\_Sub2000\_X\_CoW

**If you have any problems or queries please contact the Production Editor for guidance (details can be found in the Instructions for Authors)**

If you have submitted a manuscript that contains colour figures, it is the policy of the journal for authors to meet the full cost of colour reproduction. John Wiley & Sons Group of companies require you to complete this form. Please fill in your contact details and then complete the table as instructed.

**Note:** this form must be filled in for **all** manuscripts that have been submitted with colour work, even if you subsequently choose not to publish either in print or online in colour.

**We are unable to process your manuscript until we receive your instructions. Please return this form as quickly as possible.**

Journal name: \_\_\_\_\_

Corresponding Author: \_\_\_\_\_

Manuscript No/Title: \_\_\_\_\_

Who should be billed for this charge? Please supply below the name and the address of the payee.

Payee Name: \_\_\_\_\_

Payee Address: \_\_\_\_\_

\_\_\_\_\_

Fax: \_\_\_\_\_ Tel: \_\_\_\_\_

e-mail: \_\_\_\_\_

**Please now complete this table (Remember: only one 'X' per figure).**

Figure Number	Column 1 Colour in print and online*	Column 2 Colour online ONLY	Column 3 Black & White in print and online
Costs	1st and 2nd figures = Free Subsequent figures = £60 each	Free	Free
1			
2			
3			
4			
5			
6			
7			
8			
9			
10			
Total number of figures of each type			

*Quick cost lookup table for colour figures in print and online:			
No. of colour figs	Cost	No. of colour figs	Cost
1	Free	6	£240
2	Free	7	£300
3	£60	8	£360
4	£120	9	£420
5	£180	10	£480

Column 1 total £ \_\_\_\_\_

VAT (if applicable) £ \_\_\_\_\_

**Total: £** \_\_\_\_\_

PAGE 1

<<<For internal use ONLY>>>				
JPCMS No	Journal acronym	Volume	Issue	Print run
Production Editor			Author Correspondence Attached Y <input type="checkbox"/> / N <input type="checkbox"/>	

## WHAT YOU NEED TO DO

Please complete the table on page 1, placing an 'X' in the relevant column for each figure. The costs are shown at the top of each column. Please calculate the total cost for each category at the foot of the table, and then add VAT if applicable (see below).

### All customers MUST quote their VAT numbers

**Customers in the UK:** add VAT at the applicable rate. For the current UK VAT Rate, please refer to <http://www.hmrc.gov.uk/vat/forms-rates/rates/rates.htm#1>.

### Customers elsewhere in the EU:

- *Customers registered for VAT.* Please enter your VAT details below (please note VAT will not be charged). If no VAT registration number is supplied, VAT will be charged.

VAT registration number: \_\_\_\_\_

### If Exempt From VAT – Please attach a copy of the certificate to this form.

- *Customers NOT registered for VAT:* Add VAT at the applicable rate.

**Customers outside the EU:** No VAT applicable.

---

### Payment by Credit Card (MasterCard/VISA/American Express)

John Wiley & Sons accepts payment in three major currencies:

Pounds Sterling, United States Dollars and the Euro. Please **indicate** which currency you would like to pay in .

Please note this section **must** be completed. If no preference is given, the payment will be taken in Pounds Sterling (£).

Pounds Sterling (£)  / United States Dollars\* (US\$)  / Euros\* (€)  (tick one)

\*Conversions from US\$ or € will be made at our monthly rate.

Name on card \_\_\_\_\_ Expiry Date \_\_\_\_\_ / \_\_\_\_\_

Cardholder address \_\_\_\_\_

Please debit my credit card number: 

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

With the sum of \_\_\_\_\_

Signature of cardholder \_\_\_\_\_ Date DD / MM / YYYY

**Note:** payment by credit card can only be accepted if **all** sections are completed. Please use BLOCK CAPITALS where possible.

### Payment by Purchase Order

If you are asking your university or institution to pay we require purchase order details

I attach order number \_\_\_\_\_ dated \_\_\_\_\_

**Please post or courier all pages of your completed form to:**

#### Customer Services (OPI)

**John Wiley & Sons Ltd, European Distribution Centre**

**New Era Estate**

**Oldlands Way**

**Bognor Regis**

**West Sussex**

**PO22 9NQ**

**Please note that electronic or faxed copies cannot be accepted. For queries, please contact the production editor for your journal. Contact details can be found in the instructions for authors.**

**Please note that funds will be collected after the article is published in print.**

---

#### Contact Details of the person completing this form in case of any queries:

Print Name: \_\_\_\_\_ Tel No: \_\_\_\_\_

Signature: \_\_\_\_\_ Date: DD / MM / YYYY

E-mail Address: \_\_\_\_\_ Manuscript No: \_\_\_\_\_