

P318

Stroke and thrombolysis: an old disease with a new approach

N Catorze, L Pessoa, M Abu Hazima, F Carrilho
Centro Hospitalar Médio Tejo, Abrantes, Portugal
Critical Care 2011, 15(Suppl 1):P318 (doi: 10.1186/cc9738)

Introduction The incidence of stroke doubles for every decade after 45 years of age, and 70% of these events occur in the over 65s. A rational approach with a thrombolysis protocol can diminish this clinical and social burden.

Methods In the past 20 months all patients with acute stroke were referred to ICU staff for evaluation and compliance to effective thrombolysis until 4.30 hours from the onset of symptoms. All clinicians were advised and triage in the ED was adapted using NIHSS.

Results In this period 152 patients were evaluated and 34 (22.4%) were eligible for reperfusion treatment. Men were more prevalent than women (70.6 vs. 29.4%) and age was distributed between 29 and 82 years. Risk factors were equally distributed (Table 1). Twenty-nine patients (88%) received thrombolysis within 3 hours of symptoms onset and 19 (63%) got better NIHSS after treatment. Eleven patients (37%) never recovered. Five out of 34 patients (12%) were treated in the 3 to 4.30 hours window and three received benefit. All deaths were related to ischemia progression. Table 2 presents complications during the ICU stay.

Table 1 (abstract P318). Stroke risk factors

Factor	n
High blood pressure	25
AF	4
>Lipids	6
Diabetes	4
>BMI	6
Smoke	10

Table 2 (abstract P318). Complications during the ICU stay

Complication	n
Bradycardia	8
Pneumonia	5
Hemorrhage	4
Death	4

Conclusions The clinicians' compliance and patients' reference to dedicated teams (stroke teams) resulted in the treatment of 22.4% of observed patients (1 to 11% in the literature). Some complications could be avoided with simple measures. This protocol should continue and should be emphasized.

Reference

1. Alteplase for the Treatment of Acute Ischaemic Stroke [http://www.nice.org.uk/nicemedia/live/11618/33974/33974.pdf]

P319

Cerebral vasoreactivity is not impaired in patients with severe sepsis

S Szatmári, Z Fülep, P Sárkány, C Antek, P Siró, C Molnár, B Fülesdi
University of Debrecen, Hungary
Critical Care 2011, 15(Suppl 1):P319 (doi: 10.1186/cc9739)

Introduction In a previous report it was observed that acetazolamide-induced cerebrovascular reactivity is impaired in patients with sepsis-associated encephalopathy without organ dysfunction [1]. The aim of the present work was to assess whether patients suffering from severe sepsis also have these impaired cerebrovascular responses.

Methods Patients fulfilling the criteria of clinical sepsis and showing at least two organ dysfunctions other than the brain were included (n = 14). Nonseptic persons without previous diseases affecting cerebral vasoreactivity served as controls (n = 20). Transcranial Doppler blood flow velocities were measured at rest and at 5, 10, 15 and 20 minutes after intravenous administration of 15 mg/kg BW acetazolamide. The time course of the acetazolamide effect on cerebral blood flow velocity (cerebrovascular reactivity) and the maximal vasodilatory effect of acetazolamide (cerebrovascular reserve capacity (CRC)) were compared among the groups.

Results Mean blood flow velocity in the middle cerebral artery was lower (41.7 ± 13.3 cm/second) in septic patients at rest than in controls (58.2 ± 12.0 cm/second, P < 0.01). Pulsatility indices were higher among septic patients at rest (1.56 ± 0.79) than in controls (0.85 ± 0.20, P < 0.01). Assessment of the time course of the vasomotor reaction showed that patients with sepsis reacted in similar fashion and extent to the vasodilatory stimulus than did control persons. When assessing the maximal vasodilatory ability of the cerebral arterioles to acetazolamide during vasomotor testing, we found that patients with sepsis reacted to a similar extent to the drug than did control subjects (CRC controls: 46.2 ± 15.9%, CRC SAE: 63.2 ± 28.4%).

Conclusions Cerebral vasoreactivity to acetazolamide is not impaired in patients with severe sepsis. Our data suggest that the reaction of the cerebral arterioles to vasoactive stimuli changes along with the severity of the septic process.

Reference

1. Szatmari et al.: Crit Care 2010, 14:R50.

P320

Significance of admission temperature and impact on mortality in critically ill neurological patients

F Rincon¹, C Schorr², C Hunter², B Milcareck², R Dellinger², J Parrillo², S Zanotti²

¹Thomas Jefferson University, Philadelphia, PA, USA; ²Cooper University Hospital, Camden, NJ, USA

Critical Care 2011, 15(Suppl 1):P320 (doi: 10.1186/cc9740)

Introduction The purpose of this study is to test the hypothesis that hyperthermia is associated with increased mortality after neurological injury using a robust multicenter ICU database.

Methods A multicenter cohort study using the Project IMPACT critical care database of ICUs at 120 US hospitals between 2003 and 2008. Patient inclusion criteria were age older than 17 years, acute neurological injury within 24 hours of admission (acute ischemic stroke (AIS), subarachnoid hemorrhage (SAH), intracerebral hemorrhage (ICH), subdural hematoma (SDH), and traumatic brain injury (TBI)), and admission to the ICU. Patients were divided into three main groups based on definitions of hyperthermia and hypothermia in the ICU. Hyperthermia was defined as temperature greater than 37.5°C, hypothermia as a temperature lower than 36.5°C, and normothermia, not classified as hyperthermia or hypothermia. The outcome measure was in-hospital mortality.

Results Over the 8-year period, the Project IMPACT database contained data on more than 700,000 ICU admissions. We found 16,889 patients that met the inclusion criteria. The mean age was 61 ± 19 years, 9,339 (56%) were male, and 12,634 (76%) were white. Of these, 3,081 (18%) had AIS, 2,413 (14%) had SAH, 4,315 (26%) had ICH, 2,748 (16%) had SDH, and 4,317 (26%) had TBI. The mean admission temperature was 37.5 ± 3°C and the overall mortality was 3,628/16,676 (22%). Of the total cohort, 7,878 (47%) had hyperthermia, 689 (4%) had hypothermia, and 8,167 (49%) were normothermic. The hyperthermia group had a high in-hospital mortality (2,180/7,822 (28%)) compared with normothermia (1,169/8,167 (14%)) but the hypothermia group had significantly higher in-hospital mortality (279/687 (41%)). In a preliminary multivariate model controlling for potential confounders (age and gender), hyperthermia (OR, 1.2; 95% CI, 1.1 to 1.23) and hypothermia (OR, 1.9; 95% CI, 1.7 to 2.1) increased the odds of hospital mortality.

Conclusions Among critically ill neurological patients admitted to the ICU, hyperthermia and hypothermia are associated with increased