



Research Article

Flower-visiting assemblages of arable lands and orchards in the eastern Pannonian Lowland, studied with volatile traps

Aletta Ősz[‡], Kálmán Szanyi[§], Dóra Arnóczkyné Jakab[§], Szabolcs Szanyi[§], Sándor Koczor[‡],
Miklós Tóth[‡], István Szarukán[§], Antal Nagy[§]

[‡] Plant Protection Institute, Centre for Agricultural Research, Hungarian Research Network, Herman Otto u. 15., H-1022, Budapest, Hungary

[§] Faculty of Agriculture and Food Sciences and Environmental Management, Institute of Plant Protection, University of Debrecen, H-4032, Debrecen, Hungary

Corresponding author: Aletta Ősz (osz.aletta@atk.hun-ren.hu)

Academic editor: Grégoire Noël

Received: 09 Oct 2025 | Accepted: 31 Dec 2025 | Published: 04 Feb 2026

Citation: Ősz A, Szanyi K, Arnóczkyné Jakab D, Szanyi S, Koczor S, Tóth M, Szarukán I, Nagy A (2026) Flower-visiting assemblages of arable lands and orchards in the eastern Pannonian Lowland, studied with volatile traps. Biodiversity Data Journal 14: e174461. <https://doi.org/10.3897/BDJ.14.e174461>

Abstract

Pollinator assemblages of different habitats are widely studied, but sampling methodologies and covering of lesser-known pollinators and other flower-visitor taxa (e.g., floriphagous species) need improvement to obtain reliable data on the structure and functioning of these communities. We assessed flower-visiting insect assemblages in arable lands and apple orchards of the eastern Pannonian Lowland using phenylacetaldehyde-based volatile traps originally developed for pest monitoring. In seven sites, 27,539 individuals belonging to 123 species and seven insect orders were recorded. On average, over 90% of the captured insects were potential pollinators, with a higher proportion in orchards (98%) than in arable lands (85%). In orchards, the dominant groups were Hymenoptera, Diptera, Lepidoptera and Neuroptera, while in arable lands, Lepidoptera were most abundant. Floriphagous and pest species were more numerous in arable lands. These results demonstrate that phenylacetaldehyde-baited traps are easy to use, standardized, and sufficiently efficient for assessing of flower-visiting assemblages. Beyond their application in pest monitoring, they can

provide reliable data on the flower-visiting and pollinator assemblages in different natural and agricultural habitats.

Keywords

phenylacetaldehyde, pollinator assemblages, alfalfa field, apple orchard, agroecosystem, pest monitoring, floral scent

Introduction

Insect pollination is an essential ecological service, that plays a key role in global food production and the maintenance of biodiversity (Kevan et al. 1990, Klein et al. 2007, Aizen et al. 2009). Consequently, numerous studies have investigated pollinators in both natural and agricultural habitats (Brunet and Stewart 2010, Ambaw and Workiyie 2020). However, most of these surveys primarily focus on *major* pollinator groups – particularly bees – while other flower-visiting insect taxa remained comparatively understudied (Cecen et al. 2008). Nevertheless, a wide range of less conspicuous or less well-known flower-visitor insect groups (e.g., lepidopterans, flies, beetles, etc.) also contribute considerably to plant pollination (Steffan-Dewenter et al. 2005, Garibaldi et al. 2013, Rauf et al. 2021, Rácz et al. 2023). In addition, taxa belong to other flower-visiting functional groups such as species feeding on nectar, pollen and/or different parts of the flowers and buds, and even their predators and parasites are also less known, and their interactions are also not revealed. Therefore, comprehensive surveys that encompass the wide spectrum of flower-visiting assemblages are essential to advance our understanding of insects-plant interactions (Biesmeijer et al. 2006, O'Connor et al. 2019).

Moreover, the methodological heterogeneity among studies often limits the comparability of results across different habitats (Thompson et al. 2021). Some sampling methods focus exclusively on particular flower-visiting taxa – such as pan traps or insect nets (Eeraerts and Meeus 2025) – while others rely on labour-intensive and frequently subjective sampling approaches, including visual observations, transect walks, or focal plant surveys (Sarma et al. 2025). These observer-based techniques require substantial field effort, depend heavily on the observer's expertise, and can underrepresent small, fast-moving, or nocturnal species (Roy et al. 2016). Their effectiveness is also strongly influenced by environmental conditions, leading to inconsistent detectability and limited comparability across studies (Sarma et al. 2025). This highlights the need for standardized methods that can be applied broadly in diverse ecological contexts (Kearns C. A. and Inouye D. W. 1993, O'Connor et al. 2019).

Volatile-baited traps, originally developed for monitoring agricultural pests, have undergone considerable development and methodological advancement in the last decades (Eby et al. 2013, Landolt et al. 2013, Szanyi et al. 2024). Due to their efficiency in the case of a broad spectrum of insect taxa, these traps are increasingly used in faunistic and ecological studies as standardized sampling tools (Henning et al. 1992,

Andrews et al. 2007, Nagy et al. 2023a, Nagy et al. 2023b, Szanyi et al. 2024). Their advantages include continuous sampling independent of observer bias, high efficiency in capturing both dominant and less conspicuous taxa, and applicability across various habitats.

The efficiency of volatile baited traps for attraction of lepidopteran and hemipteran pests was previously studied by our research group. During these experiments, the phenylacetaldehyde-based lures – imitating odors of flowers – proved effective not only for their target pest taxa (Szanyi et al. 2019, Szalárdi et al. 2021), but also for several non-target flower-visiting groups, including non-pest Lepidoptera (Szanyi et al. 2019, Szanyi et al. 2024), Diptera (Nagy et al. 2023b), Orthoptera (Nagy et al. 2023a), Hymenoptera (Segers et al. 2023) and Neuroptera (Szanyi et al. 2025).

In the present study, we provide data on the flower-visiting insect assemblages of agricultural habitats of the northeastern Pannonian Lowland based on the non-target catches of phenylacetaldehyde-baited traps. Additionally, we compare the composition and diversity of these assemblages living in two different agricultural habitat types – arable lands and orchards – during different flowering periods.

Material and methods

Study sites and sampling

We studied the efficiency of different phenylacetaldehyde-based lures between 2019 and 2022 at seven sites including four alfalfa fields and three commercial apple orchards in the Pannonian Lowland of Hungary (Fig. 1).




Figure 1. [doi](#)

Locations of the sampling sites. Red circles=arable lands, yellow triangles=orchards.

Experiments in alfalfa fields:

Püspökladány, Hajdú-Bihar County (GPS: 47.1814N, 21.0913E). Baits: phenylacetaldehyde, 1-phenylacetaldehyde, 4-methoxycinnamaldehyde, phenylacetaldehyde + 4-methoxycinnamaldehyde, phenylacetaldehyde + 1-phenylacetaldehyde + 4-methoxycinnamaldehyde + 1-nonanol, phenylacetaldehyde + 1,4-dimethoxybenzene, phenylacetaldehyde + benzyl alcohol + 6-methyl-5-hepten-2-one, phenylacetaldehyde + 1,4-dimethoxybenzene + benzyl alcohol + 6-methyl-5-hepten-2-one, and unbaited traps. Number of repetitions: 10. Date: from 03/07 to 31/08, 2019.

Gégény, Szabolcs-Szatmár Bereg County (GPS: 48.1608218N, 21.9350966E). Baits: phenylacetaldehyde, phenylacetaldehyde + anetol, phenylacetaldehyde + anisacet, phenylacetaldehyde + betion, phenylacetaldehyde + acetacid, unbaited control traps. Number of repetitions: 6. Date: from 09/07 to 31/08, 2020.

Nyírtelek, Szabolcs-Szatmár Bereg County (GPS: 48.000133N, 21.600674E). Baits: phenylacetaldehyde, phenylacetaldehyde + 4-methoxycinnamaldehyde, phenylacetaldehyde + 1,4-dimethoxybenzene, phenylacetaldehyde + 1,4-dimethoxybenzene, phenylacetaldehyde + 4-methoxycinnamaldehyde + 1,4-dimethoxybenzene, unbaited control traps. Number of repetitions: 7. Date: from 22/06 to 31/08, 2021.

Gyomaendrőd, Békés County (GPS: 46.54569N, 20.52989E). Baits: phenylacetaldehyde, phenylacetaldehyde + hexanal, phenylacetaldehyde + (E)-2-hexanal, phenylacetaldehyde + alpha-pyrene, unbaited control traps. Number of repetitions: 5. Date: from 14/07 to 31/08, 2022

Experiments in apple orchards:

Komoró, Szabolcs-Szatmár Bereg County (GPS: 48.292307N, 22.133233E),

Ajak, Szabolcs-Szatmár Bereg County (GPS: 48.163577N, 22.059229E), and

Nyírpazony, Szabolcs-Szatmár Bereg County (GPS: 47.978482N, 21.824846E). Investigated habitat: apple orchard. Baits: phenylacetaldehyde + eugenol + benzyl acetate, unbaited control traps. Number of repetitions: 3. Date: from 23/04 to 06/05, 2022.

The samplings were carried out in the flowering period of the chosen cultures. CSALOMON® VARL+ baited traps (Plant Protection Institute, CAR, Budapest, Hungary), available at www.csalomontraps.com (accessed on 11 May 2025), were used in each sampling sites (Fig. 2.), with phenylacetaldehyde-based synthetic lures that imitate the scent of strongly fragrant, mostly white flowers (Szanyi et al. 2019, Nagy et al. 2022). The tested lures were enclosed in 1.5 × 1.5 cm polyethylene bags with a 0.2 mm wall thickness, allowing the volatiles to diffuse through the bag's walls (Tóth M. et al. 2020). In apple orchards, the traps were placed on branches of trees, with a spacing of 20 m between them at heights of 1.8–2.0 m. In alfalfa fields, traps were placed at the field margins on 1 m height mounted on wooden stakes and spaced 15 m apart. Traps were checked and emptied weekly in the apple orchards, and twice a week in the alfalfa fields.

The baits were replaced every four weeks. To kill the trapped insects, we used Vaportape® II pesticide strips designed specifically for insect traps.



Figure 2. [doi](#)

CSALOMON® VARL+ baited trap used to collect the flower visiting insects.

Evaluation of samples and data analysis

Samples were stored at -20 °C in a fridge until identification in a laboratory. The identification of the insects caught was made at species level, when it was possible, based on the keys of Wagner (1952) and Schwartz M. D and Footitt R. G (1998) for heteropterans, Harz (1957), Harz (1960) and Harz (1975) for orthopterans, Aspoeck et al. (1980) and Thierry et al. (1998) for neuropterans, Móczár (1985), Goulet and Hubert (1993) and Bossert (2015) for hymenopterans, Papp and Darvas (2000) and Tóth (2017) for dipterans, and Varga Z. (2010) for lepidopterans.

Relative frequencies (RF%) of insect groups were calculated using all collected specimens, including those identified only at the genus or family level. To evaluate the composition and mean ratios (MR%) of different taxonomical and functional groups, we compared both the total number and mean number of individuals (individuals/sampling sites). Two functional group classifications were applied. The first classification was

based on the presumed reason attraction to floral scent: potential pollinators were considered species attracted by floral rewards for the pollination (e.g., nectar, pollen), while floriphagous species (e.g., Miridae, Cerambycidae) were those attracted due to their consumption of floral tissues. The second classification distinguished species by pest status: species capable of causing serious damage to the host plant at any phenological stage were categorized as pests, whereas those not causing such damage were considered non-pests. Data normality and variance homogeneity were assessed using Q-Q plots and Levene’s test, respectively. As assumptions were not met, the non-parametric Kruskal–Wallis test was used. Where significant differences were detected, pairwise comparisons were conducted using the Mann–Whitney U-test. All statistical analyses were performed using SPSS 21.0 (Ketskeméty et al. 2011).

All records collected for the present work were published through GBIF (Ősz et al. 2025).

Results

A total of 30,595 insects were caught using volatile traps baited with phenylacetaldehyde-based lures (Table 1, Suppl. material 1). Of these, 27,539 individuals of 123 flower-visiting species were identified to species level, belonging to eight insect orders. Lepidoptera was notably the most species-rich order, with 98 species, and was dominated by the family Noctuidae, with 52 species of 9 subfamilies. The further most diverse groups were Hymenoptera (S=9) and Hemiptera (S=8), while Diptera, Orthoptera, Coleoptera, and Mecoptera were represented by only a few species.

Table 1. Checklist by habitat types (A=arable land, O=orchard), including information on the occurrence (1=presence), pest status (PE=pest, causes damage to crops at any developmental stage; NPE=non-pest, does not cause serious damage) and pollinator role of the species (PO=known pollinator; PPO=potential pollinator; FLO=floriphagous, feeds on the flowers). Protected species in Hungary are marked with an asterisk (*).						
Order	Family	Species	A	O	PE	PO
Coleoptera	Cerambycidae	<i>Plagionotus floralis</i> (Pallas, 1773)	1		PE	FLO
Diptera	Bombyliidae	<i>Bombylius major</i> (Linnaeus, 1758)		1	NPE	PO
Diptera	Syrphidae	<i>Episyrphus balteatus</i> (De Geer, 1776)		1	NPE	PO
Diptera	Syrphidae	<i>Eristalis tenax</i> (Linnaeus, 1758)		1	NPE	PO
Diptera	Syrphidae	<i>Melanostoma scalare</i> (Fabricius, 1794)		1	NPE	PO
Hemiptera	Miridae	<i>Adelphocoris lineolatus</i> (Goeze, 1778)	1		PE	FLO
Hemiptera	Miridae	<i>Adelphocoris quadripunctatus</i> (Fabricius, 1794)	1		PE	FLO
Hemiptera	Miridae	<i>Adelphocoris reichellii</i> (Fieber, 1836)	1		PE	FLO
Hemiptera	Miridae	<i>Adelphocoris seticornis</i> (Fabricius, 1775)	1		PE	FLO

Order	Family	Species	A	O	PE	PO
Hemiptera	Miridae	<i>Lygus gemellatus</i> (Herrich-Schaeffer, 1835)	1		PE	FLO
Hemiptera	Miridae	<i>Lygus pratensis</i> (Linnaeus, 1758)	1		PE	FLO
Hemiptera	Miridae	<i>Lygus rugulipennis</i> (Poppius, 1911)	1		PE	FLO
Hemiptera	Miridae	<i>Polymerus vulneratus</i> (Panzer, 1805)	1		PE	FLO
Hymenoptera	Apidae	<i>Apis mellifera</i> (Linnaeus, 1758)	1	1	NPE	PO
Hymenoptera	Apidae	<i>Bombus argillaceus</i> (Scopoli, 1763)*		1	NPE	PO
Hymenoptera	Apidae	<i>Bombus hortorum</i> (Linnaeus, 1761)		1	NPE	PO
Hymenoptera	Apidae	<i>Bombus lapidarius</i> (Linnaeus, 1758)		1	NPE	PO
Hymenoptera	Apidae	<i>Bombus pascuorum</i> (Scopoli, 1763)		1	NPE	PO
Hymenoptera	Apidae	<i>Bombus ruderatus</i> (Fabricius, 1775)*		1	NPE	PO
Hymenoptera	Apidae	<i>Bombus sylvarum</i> (Linnaeus, 1761)		1	NPE	PO
Hymenoptera	Apidae	<i>Bombus terrestris</i> (Linnaeus, 1758)		1	NPE	PO
Hymenoptera	Vespidae	<i>Vespa crabro</i> (Linnaeus, 1758)		1	PE	FLO
Lepidoptera	Crambidae	<i>Cydalima perspectalis</i> (Walker, 1859)	1		PE	PPO
Lepidoptera	Crambidae	<i>Haritala ruralis</i> (Scopoli, 1763)	1		NPE	PPO
Lepidoptera	Crambidae	<i>Loxostege sticticalis</i> (Linnaeus, 1761)	1		PE	PPO
Lepidoptera	Crambidae	<i>Nomophila noctuella</i> (Denis & Schiffermüller, 1775)	1		NPE	PPO
Lepidoptera	Crambidae	<i>Nymphula nitidulata</i> (Hufnagel, 1767)	1		NPE	PPO
Lepidoptera	Crambidae	<i>Sitochroa verticalis</i> (Linnaeus, 1758)	1		NPE	PPO
Lepidoptera	Erebidae	<i>Amata phegea</i> (Linnaeus, 1758)	1		NPE	PO
Lepidoptera	Erebidae	<i>Dysauxes ancilla</i> (Linnaeus, 1767)	1		NPE	PPO
Lepidoptera	Erebidae	<i>Dysgonia algira</i> (Linné, 1767)	1		NPE	PPO
Lepidoptera	Erebidae	<i>Eilema lurideola</i> (Zincken, 1817)	1		NPE	PPO
Lepidoptera	Noctuidae	<i>Euclidia glyphica</i> (Linnaeus, 1758)	1		NPE	PO
Lepidoptera	Erebidae	<i>Herminia tarsipennalis</i> (Treitschke, 1835)	1		NPE	PPO
Lepidoptera	Erebidae	<i>Hypena proboscidalis</i> (Linnaeus, 1758)	1		NPE	PPO
Lepidoptera	Erebidae	<i>Hypena rostralis</i> (Linnaeus, 1758)	1		NPE	PPO
Lepidoptera	Erebidae	<i>Lygephila craccae</i> (Denis & Schiffermüller, 1775)	1		NPE	PPO
Lepidoptera	Erebidae	<i>Lymantria dispar</i> (Linnaeus, 1758)	1		PE	PPO
Lepidoptera	Erebidae	<i>Prodotis stolidia</i> (Fabricius, 1775)	1		NPE	PPO

Order	Family	Species	A	O	PE	PO
Lepidoptera	Erebidae	<i>Rivula sericealis</i> (Scopoli, 1763)	1		NPE	PPO
Lepidoptera	Erebidae	<i>Scoliopteryx libatrix</i> (Linnaeus, 1758)		1	NPE	PPO
Lepidoptera	Gelechiidae	<i>Sitotroga cerealella</i> (Olivier, 1789)	1		PE	PPO
Lepidoptera	Gelechiidae	<i>Stomopteryx radicalis</i> (Falkovitsh & Bidzilya, 2003)	1		NPE	PPO
Lepidoptera	Gelechiidae	<i>Camptogramma bilineata</i> (Linnaeus, 1758)	1		NPE	PPO
Lepidoptera	Geometridae	<i>Chiasmia clathrata</i> (Linnaeus, 1758)	1		PE	PO
Lepidoptera	Geometridae	<i>Cidaria fulvata</i> (Forster, 1771)	1		NPE	PPO
Lepidoptera	Geometridae	<i>Ematurga atomaria</i> (Linnaeus 1758)	1		PE	PPO
Lepidoptera	Geometridae	<i>Epirrhoe alternata</i> (Müller, 1764)	1		NPE	PPO
Lepidoptera	Geometridae	<i>Idaea aversata</i> (Linnaeus, 1758)	1		NPE	PPO
Lepidoptera	Geometridae	<i>Idaea rusticata</i> (Denis & Schiffermüller, 1775)	1		NPE	PPO
Lepidoptera	Geometridae	<i>Ligdia adustata</i> (Denis & Schiffermüller, 1775)	1		NPE	PPO
Lepidoptera	Geometridae	<i>Peribatodes rhomboidaria</i> (Denis & Schiffermüller, 1775)	1		PE	PPO
Lepidoptera	Geometridae	<i>Peribatodes secundaria</i> (Denis & Schiffermüller, 1775)	1		NPE	PPO
Lepidoptera	Geometridae	<i>Plemyria rubiginata</i> (Denis & Schiffermüller, 1775)	1		NPE	PPO
Lepidoptera	Geometridae	<i>Pseudopanthera macularia</i> (Linnaeus, 1758)	1		NPE	PPO
Lepidoptera	Geometridae	<i>Tephрина arencearia</i> (Denis & Schiffermüller, 1775)	1		PE	PPO
Lepidoptera	Lasiocampidae	<i>Lasiocampa trifolii</i> (Denis & Schiffermüller, 1775)	1		NPE	PPO
Lepidoptera	Lycaenidae	<i>Polyommatus icarus</i> (Rottemburg, 1775)	1		NPE	PO
Lepidoptera	Noctuidae	<i>Acontia lucida</i> (Hufnagel, 1766)	1		NPE	PPO
Lepidoptera	Noctuidae	<i>Acontia trabealis</i> (Scopoli, 1763)	1	1	NPE	PPO
Lepidoptera	Noctuidae	<i>Aedia leucomelas</i> (Linnaeus, 1758)	1		NPE	PPO
Lepidoptera	Noctuidae	<i>Abrostola tripartita</i> (Hufnagel, 1766)	1	1	NPE	PO
Lepidoptera	Noctuidae	<i>Abrostola triplasia</i> (Linnaeus, 1758)	1		NPE	PO
Lepidoptera	Noctuidae	<i>Agrotis bigramma</i> (Esper, 1790)	1		PE	PPO
Lepidoptera	Noctuidae	<i>Agrotis exclamationis</i> (Linnaeus, 1758)	1		PE	PPO
Lepidoptera	Noctuidae	<i>Agrotis segetum</i> (Denis & Schiffermüller, 1775)	1		PE	PPO
Lepidoptera	Noctuidae	<i>Apamea sordens</i> (Hufnagel, 1766)	1		NPE	PPO
Lepidoptera	Noctuidae	<i>Apterogeton ypsilon</i> (Denis & Schiffermüller, 1775)	1		NPE	PPO
Lepidoptera	Noctuidae	<i>Autographa gamma</i> (Linnaeus, 1758)	1	1	PE	PO

Order	Family	Species	A	O	PE	PO
Lepidoptera	Noctuidae	<i>Axylia putris</i> (Linnaeus, 1761)	1		NPE	PPO
Lepidoptera	Noctuidae	<i>Calophasia lunula</i> (Hufnagel, 1766)	1		NPE	PPO
Lepidoptera	Noctuidae	<i>Caradrina morpheus</i> (Hufnagel, 1766)	1		NPE	PPO
Lepidoptera	Noctuidae	<i>Chrysodeixis chalcites</i> (Esper, 1789)	1		PE	PO
Lepidoptera	Noctuidae	<i>Cirrhia icteritia</i> (Hufnagel, 1766)	1		NPE	PPO
Lepidoptera	Noctuidae	<i>Cornutiplusia circumflexa</i> (Linnaeus, 1767)	1		NPE	PO
Lepidoptera	Noctuidae	<i>Cucullia chamomillae</i> (Denis & Schiffermüller, 1775)		1	NPE	PPO
Lepidoptera	Noctuidae	<i>Cucullia umbratica</i> (Linnaeus, 1758)	1		NPE	PO
Lepidoptera	Noctuidae	<i>Diachrysia chrysis</i> (Linnaeus, 1758)	1	1	NPE	PO
Lepidoptera	Noctuidae	<i>Diachrysia stenochrysis</i> (Warren, 1913)	1		NPE	PO
Lepidoptera	Noctuidae	<i>Eupsilia transversa</i> (Hufnagel, 1766)	1		NPE	PPO
Lepidoptera	Noctuidae	<i>Euxoa temera</i> (Hübner, 1808)	1		PE	PPO
Lepidoptera	Noctuidae	<i>Hadena bicruris</i> (Hufnagel, 1766)	1		NPE	PPO
Lepidoptera	Noctuidae	<i>Hadula trifolii</i> (Hufnagel, 1766)	1		PE	PO
Lepidoptera	Noctuidae	<i>Helicoverpa armigera</i> (Hübner, 1808)	1	1	PE	PO
Lepidoptera	Noctuidae	<i>Heliopsis maritima</i> (Draudt, 1938)	1		PE	PO
Lepidoptera	Noctuidae	<i>Heliopsis viriplaca</i> (Hufnagel, 1766)	1		PE	PO
Lepidoptera	Noctuidae	<i>Hoplodrina ambigua</i> (Denis & Schiffermüller, 1775)	1		NPE	PPO
Lepidoptera	Noctuidae	<i>Hoplodrina octogeneria</i> (Goeze, 1781)	1		NPE	PPO
Lepidoptera	Noctuidae	<i>Hydraecia micacea</i> (Esper, 1789)	1		NPE	PPO
Lepidoptera	Noctuidae	<i>Lacanobia oleracea</i> (Linnaeus, 1758)	1		PE	PPO
Lepidoptera	Noctuidae	<i>Macdunnoughia confusa</i> (Stephens, 1850)	1	1	PE	PPO
Lepidoptera	Noctuidae	<i>Mamestra brassicae</i> (Linnaeus, 1758)	1		PE	PPO
Lepidoptera	Noctuidae	<i>Mesapamea secalis</i> (Linnaeus, 1758)	1		PE	PPO
Lepidoptera	Noctuidae	<i>Mesoligia furuncula</i> (Denis & Schiffermüller, 1775)	1		NPE	PPO
Lepidoptera	Noctuidae	<i>Mythimna albipuncta</i> (Denis & Schiffermüller, 1775)	1		NPE	PPO
Lepidoptera	Noctuidae	<i>Mythimna l-album</i> (Linnaeus, 1767)	1		NPE	PPO
Lepidoptera	Noctuidae	<i>Mythimna pallens</i> (Linnaeus, 1758)	1		PE	PPO
Lepidoptera	Noctuidae	<i>Mythimna vitellina</i> (Hübner, 1808)	1		PE	PPO
Lepidoptera	Noctuidae	<i>Noctua interposita</i> (Hübner, 1790)	1		PE	PPO

Order	Family	Species	A	O	PE	PO
Lepidoptera	Noctuidae	<i>Noctua janthina</i> (Borkhausen, 1792)	1		PE	PPO
Lepidoptera	Noctuidae	<i>Noctua pronuba</i> (Linnaeus, 1758)	1		PE	PPO
Lepidoptera	Noctuidae	<i>Orthosia cruda</i> (Denis & Schiffermüller, 1775)		1	PE	PPO
Lepidoptera	Noctuidae	<i>Pyrrhia umbra</i> (Hufnagel, 1766)	1		NPE	PPO
Lepidoptera	Noctuidae	<i>Rhyacia simulans</i> (Hufnagel, 1766)	1		NPE	PPO
Lepidoptera	Noctuidae	<i>Shargacucullia verbasci</i> (Linnaeus, 1758)	1		NPE	PPO
Lepidoptera	Noctuidae	<i>Thalophila matura</i> (Hufnagel, 1766)	1		NPE	PPO
Lepidoptera	Noctuidae	<i>Trachea atriplicis</i> (Linnaeus, 1758)	1		NPE	PPO
Lepidoptera	Noctuidae	<i>Tyta luctuosa</i> (Denis & Schiffermüller, 1775)	1		NPE	PPO
Lepidoptera	Noctuidae	<i>Xestia c-nigrum</i> (Linnaeus, 1758)	1		PE	PPO
Lepidoptera	Noctuidae	<i>Xestia xanthographa</i> (Denis & Schiffermüller, 1775)	1		NPE	PPO
Lepidoptera	Nymphalidae	<i>Araschnia levana</i> (Linnaeus, 1758)	1		NPE	PO
Lepidoptera	Nymphalidae	<i>Coenonympha pamphilus</i> (Linnaeus, 1758)	1		NPE	PO
Lepidoptera	Nymphalidae	<i>Nymphalis c-album</i> (Linnaeus, 1758)*	1		NPE	PO
Lepidoptera	Pieridae	<i>Colias croceus</i> (Geoffroy in Fourcroy, 1785)	1		NPE	PO
Lepidoptera	Pyalidae	<i>Etiella zinckenella</i> (Treitschke, 1832)	1		PE	PPO
Lepidoptera	Pyalidae	<i>Hypsopygia costalis</i> (Fabricius, 1775)	1		PE	PPO
Lepidoptera	Pyalidae	<i>Ostrinia nubilalis</i> (Hübner, 1796)	1		PE	PPO
Lepidoptera	Pyalidae	<i>Salebria semirubella</i> (Scopoli, 1763)	1		PE	PPO
Lepidoptera	Sesiidae	<i>Synanthedon vespiformis</i> (Linnaeus, 1761)	1		NPE	PO
Mecoptera	Panorpidae	<i>Panorpa communis</i> (Linnaeus, 1758)	1		NPE	PPO
Orthoptera	Acrididae	<i>Calliptamus italicus</i> (Linnaeus, 1758)	1		PE	FLO
Orthoptera	Gryllidae	<i>Oecanthus pellucens</i> (Scopoli, 1763)	1		NPE	PPO
Orthoptera	Tettigoniidae	<i>Tettigonia viridissima</i> (Linnaeus, 1758)	1		NPE	PPO

Most identified species were common and widespread in Hungary. The *major* part of the assemblage consisted of agricultural pest species (e.g., *Hemiptera* spp., *Ostrinia nubilalis* *Autographa gamma*, *Agrotis* spp., *Helicoverpa armigera*), primarily associated with alfalfa fields. In contrast, the majority of individuals and species caught in apple orchards were considered beneficial. Additionally, two protected species (*Bombus argillaceus*, *B. ruderatus*) were recorded in the apple orchards, and one (*Nymphalis c-album*) in the alfalfa fields.

Although Lepidoptera had the highest species-richness, they accounted for 47.25% of all specimens, with 21.75% belonging to the Noctuidae family. Within Noctuidae, the Plusiinae (RF%=53.79), and Heliiothinae (RF%=37.50) subfamilies were dominant. Lepidoptera was followed by Hymenoptera (RF%=15.76), *Hemiptera* (RF%=6.95), and Diptera (RF%=3.64).

Notable differences were found between the flower visiting assemblages of the two habitat types. Arable lands supported a more diverse assemblage, with 24,834 specimens representing 108 species, compared to only 2,705 individuals and 22 species in the orchards.

In arable lands, Lepidoptera dominated species-richness ($S=95$), with other groups contributing only a few species each. However, Lepidoptera accounted for only 67.83% of individuals. Hymenoptera (RF%=15.97), *Hemiptera* (RF%=10.10) and Neuroptera (RF%=4.93) were also relatively abundant given their species numbers.

In orchards, only 22 species were recorded. Almost half belonged to Hymenoptera ($S=9$), while Lepidoptera, Diptera, and Neuroptera were represented by only a few species. Relative frequencies followed a similar pattern: Hymenoptera were dominant (RF%=53.13), followed by Diptera (RF%=34.44), Neuroptera (RF%=6.12), and Lepidoptera (RF%=6.29).

Across all habitats, the mean ratio (MR%) of potential pollinators (MR%=90.462 $\pm$ 15.049) was significantly higher than that of floriphagous insects (MR%=9.537 $\pm$ 15.049) (Fig. 3). Similarly, the mean ratio of pest species (MR%=29.361 $\pm$ 28.860) was significantly lower than that of non-pest species (MR%=70.638 $\pm$ 28.860).




Figure 3. [doi](#)

Mean ratio (ratio/location [%]) of potential pollinator, floriphagous (A), pest, and non-pest species (B). Means with the same letter within a diagram are not significantly different at $P=0.05$ by Mann-Whitney U non-parametric test.

Comparing habitat types, both the mean ratio of potential pollinators ($MR\%=98.239\pm1.049$) and non-pest species ($MR\%=95.420\pm0.691$) were significantly higher in orchards than in arable lands (potential pollinators: $MR\%=84.629\pm18.610$; non-pests: $MR\%=52.052\pm24.305$) (Fig. 4). Accordingly, arable lands had a much higher ratio of pests ($MR\%=47.947\pm24.305$) compared to orchards ($MR\%=4.579\pm0.691$).




Figure 4. [doi](#)

Mean ratio of potential pollinator, floriphagous, pest, and non-pest species in the different habitat types (orchard and arable land). Means with the same letter within a diagram are not significantly different at $P=0.05$ by Mann-Whitney U non-parametric test.

The dominant pollinator groups in orchards were Hymenoptera ($MR\%=51.537\pm4.275$), Diptera ($MR\%=34.453\pm2.913$), and Lepidoptera ($MR\%=6.313\pm0.564$) and Neuroptera ($MR\%=5.934\pm2.913$). In arable lands, the most abundant pollinators were Lepidoptera ($MR\%=63.445\pm28.774$), Hymenoptera ($MR\%=15.876\pm27.294$), and Neuroptera ($MR\%=4.209\pm6.031$).

Floriphagous and pest species had significantly higher mean ratios in arable lands (floriphagous: $MR\%=15.371\pm18.610$; pests: $MR\%=47.947\pm24.305$) than in orchards (floriphagous: $MR\%=1.760\pm1.049$; pests: $MR\%=4.579\pm0.691$). The most abundant pest groups in arable lands were Lepidoptera ($MR\%=32.576\pm16.283$) and *Hemiptera* ($MR\%=15.222\pm18.697$), with the latter also comprising the only notable floriphagous group. In orchards, the most abundant pest groups were Lepidoptera

(MR%=2.819±0.424) and Hymenoptera (MR%=1.760±1.049), the latter also being the sole floriphagous group.

Discussion

A total of 123 flower-visiting insect species were recorded in arable lands and apple orchards of the eastern Pannonian Lowland. The insect assemblages included a wide range of taxa from multiple orders, reflecting the diverse communities occurring in these agroecosystems. Notably, three protected species – *Bombus argillaceus*, *B. ruderatus*, and *Nymphalis c-album* – were recorded, underscoring the ecological importance of these habitats.

The observed species richness, especially within Lepidoptera, highlights the attractiveness of phenylacetaldehyde as a floral scent. The dominance of the Noctuidae family within Lepidoptera is consistent with their known responsiveness to floral volatiles (Szanyi et al. 2019, Szanyi et al. 2024). Nonetheless, Hymenoptera and Diptera also contributed substantially, especially in orchards, suggesting that these groups play there the *major* role in pollination.

Our findings demonstrate that phenylacetaldehyde-based volatile traps, originally designed for pest monitoring, are highly effective at capturing a broad spectrum of flower-visiting insects belonging several orders. This supports earlier findings (Landolt et al. 2013, Hesler 2016, Nagy et al. 2023a, Nagy et al. 2023b, Szanyi et al. 2024) and highlights the potential of such traps for broader ecological and faunistic applications beyond pest surveillance.

The study revealed notable differences in the composition of flower-visiting insect communities in arable lands and orchards. Arable lands supported a more diverse assemblage, which may be partly due to the higher sampling effort and longer sampling duration in these habitats, as the flowering stage of the investigated arable crop – alfalfa – is much longer than that of the apple orchard. However, the relative frequency and ratio patterns suggest that these differences are not solely methodological. From a functional perspective, potential pollinator species made up the majority of the assemblages in both habitat types, although their proportion was significantly higher in orchards. Accordingly, pest and floriphagous species were more abundant in arable lands.

These observed differences underline the importance of multi-taxa monitoring in agroecosystems. Focusing solely on bees or model pollinators would overlook important contributors to pollination and ecosystem functioning. Moreover, the detection of protected species like *Bombus argillaceus*, *B. ruderatus* and *Nymphalis c-album* reinforces the conservation value of these habitats and the relevance of including non-target data collected with volatile traps in faunistic studies.

In conclusion, standardized volatile-baited traps can be a valuable tool for studying flower-visiting insects and their interactions with plants in various ecosystems. Their

ability to provide consistent and replicable data makes them especially suitable for long-term monitoring and comparative studies in both natural and agricultural habitats.

Acknowledgements

Supported by the University of Debrecen Program for Scientific Publication.

Hosting institution

Faculty of Agriculture and Food Sciences and Environmental Management, Institute of Plant Protection, University of Debrecen

Author contributions

AŐ and AN conducted the experiments. SZSZ, DAJ, SK, AŐ identified the collected insect material. KSZ analysed data and conducted statistical analyses. AŐ, KSZ, MT, ISZ and AN wrote the manuscript. All authors read and approved the manuscript.

Conflicts of interest

The authors have declared that no competing interests exist.

References

- Aizen M, Garibaldi LA, Cunningham SA, Klein AM (2009) How much does agriculture depend on pollinators? Lessons from long-term trends in crop production. *Annals of Botany* 103 (9): 1579-1588. <https://doi.org/10.1093/aob/mcp076>
- Ambaw M, Workiye M (2020) Evaluation and demonstration of the role of honey bees on seed yield of alfalfa (*Medicago sativa* FL77) in Kulumsa, Ethiopia. *Journal of Entomology and Zoology Studies* 2269-2272.
- Andrews ES, Theis N, Adler LS (2007) Pollinator and herbivore attraction to *Cucurbita* floral volatiles. *Journal of Chemical Ecology* 33 (9): 1682-1691. <https://doi.org/10.1007/s10886-007-9337-7>
- Aspöck H, Aspöck U, Hoelzel H, Rausch H (1980) Die Neuropteren Europas : eine zusammenfassende Darstellung der Systematik, Oekologie und Chorologie der Neuropteroidea (Megaloptera, Raphidioptera, Planipennia) Europas. Krefeld : Goecke & Evers
- Biesmeijer JC, Roberts SPM, Reemer M, Ohlemüller R, Edwards M, Peeters T, Schaffers AP, Potts SG, Kleukers R, Thomas CD, Settele J, Kunin WE (2006) Parallel declines in pollinators and insect-pollinated plants in Britain and the Netherlands. *Science* 313 (5785): 351-354. <https://doi.org/10.1126/science.1127863>

- Bossert S (2015) Recognition and identification of bumblebee species in the *Bombus lucorum*-complex (Hymenoptera, Apidae) – A review and outlook. Deutsche Entomologische Zeitschrift 62 (1): 19-28. <https://doi.org/10.3897/dez.62.9000>
- Brunet J, Stewart CM (2010) Impact of bee species and plant density on alfalfa pollination and potential for gene flow. Psyche: A Journal of Entomology 2010: 1-7. <https://doi.org/10.1155/2010/201858>
- Cecen S, Gurel F, Karaca A (2008) Impact of honeybee and bumblebee pollination on alfalfa seed yield. Acta Agriculturae Scandinavica, Section B - Plant Soil Science 58 (1): 77-81. <https://doi.org/10.1080/09064710701214627>
- Eby C, Gardiner MGT, Gries R, Judd GJR, Khaskin G, Gries G (2013) Phenylacetaldehyde attracts male and female apple clearwing moths, *Synanthedon myopaeformis*, to inflorescences of showy milkweed, *Asclepias speciosa*. Entomologia Experimentalis et Applicata 147 (1): 82-92. <https://doi.org/10.1111/eea.12045>
- Eraerts M, Meeus I (2025) Different pollinator sampling methods measure distinct pollinator communities in a mass-flowering crop, which respond differently to the composition of the surrounding landscape. Web Ecology 25: 47-57. [In English]. <https://doi.org/10.5194/we-25-47-2025>
- Garibaldi LA, Steffan-Dewenter I, Winfree R, Aizen MA, Bommarco R, Cunningham SA, Kremen C, Carvalheiro LG, Harder LD, Afik O, Bartomeus I, Benjamin F, Boreux V, Cariveau D, Chacoff NP, Dudenhöffer JH, Freitas BM, Ghazoul J, Greenleaf S, Hipólito J, Holzschuh A, Howlett B, Isaacs R, Javorek SK, Kennedy CM, Krewenka KM, Krishnan S, Mandelik Y, Mayfield MM, Motzke I, Munyuli T, Nault BA, Otieno M, Petersen J, Pisanty G, Potts SG, Rader R, Ricketts TH, Rundlöf M, Seymour CL, Schüepp C, Szentgyörgyi H, Taki H, Tschamtké T, Vergara CH, Viana BF, Wanger TC, Westphal C, Williams N, Klein AM (2013) Wild pollinators enhance fruit set of crops regardless of honey bee abundance. Science 339 (6127): 1608-1611. <https://doi.org/10.1126/science.1230200>
- Goulet H, Hubert JF (1993) Hymenoptera of the world. An identification guide to families. Research Branch, Agricultural Canada Publication. Canada Communication Group-Publishing, Ottawa
- Harz K (1957) Die geradflügler mitteleuropas. VEB Gustav Fischer Verlag
- Harz K (1960) Geradflügler oder Orthopteren (Blattodea, Mantodea, Saltatoria, Dermapter). InF. Dahl (Ed.), Die Tierwelt Deutschland und der angrenzenden Meeresküste nach ihren Merkmalen und nach ihrer Lebensweise. VEB Gustav Fischer Verlag, 213-214 pp.
- Harz K (1975) Die Orthoptera Europas. Dr. W. Junk N. V. Publishers.
- Henning JA, Peng Y-, Montague MA, Teuber LR (1992) Honey Bee (Hymenoptera: Apidae) Behavioral response to primary alfalfa (Rosales: Fabaceae) floral volatiles. Journal of Economic Entomology 85 (1): 233-239. <https://doi.org/10.1093/jee/85.1.233>
- Hesler LS (2016) Volatile semiochemicals increase trap catch of green lacewings (Neuroptera: Chrysopidae) and flower flies (Diptera: Syrphidae) in corn and soybean plots. Journal of Insect Science 16 (1). <https://doi.org/10.1093/jisesa/iew057>
- Kearns C. A., Inouye D. W. (1993) Techniques for pollination biologists. University Press of Colorado, Boulder, CO.
- Ketskeméty L, Izsó L, Tóth EK (2011) Bevezetés az IBM SPSS Statistics Programrendszerbe. Artéria Stúdió Kft: Budapest, Hungary

- Kevan PG, Clark EA, Thomas VG (1990) Insect pollinators and sustainable agriculture. *American Journal of Alternative Agriculture* 5 (1): 13-22. <https://doi.org/10.1017/S0889189300003179>
- Klein AM, Vaissière BE, Cane JH, Steffan-Dewenter I, Cunningham SA, Kremen C, Tscharntke T (2007) Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society B: Biological Sciences* 274 (1608): 303-313. <https://doi.org/10.1098/rspb.2006.3721>
- Landolt PJ, Tóth M, Meagher RL, Szarukán I (2013) Interaction of acetic acid and phenylacetaldehyde as attractants for trapping pest species of moths (Lepidoptera: Noctuidae). *Pest Management Science* 69 (2): 245-249. <https://doi.org/10.1002/ps.3381>
- Móczár L (1985) Hártáyszárnyúak – Hymenoptera. In: Móczár L (Ed.) *Állathatározó*. Vol. 2. Tankönyvkiadó, Budapest, 263-500 pp.
- Nagy A, Szarukán I, Szalárdi T, Szanyi S, Jósvai JK, Tóth M (2022) Addition of 4-oxoisophorone improves performance of bisexual lure for *Autographa gamma* (L.) (Lepidoptera: Noctuidae). *Journal of Applied Entomology* 146 (3): 328-334. <https://doi.org/10.1111/jen.12958>
- Nagy A, Ősz A, Tóth M, Rácz IA, Kovács S, Szanyi S (2023a) Nontarget catches of traps with chemical lures may refer to the flower-visitation, probable pollination, and feeding of bush crickets (Ensifera: Tettigoniidae). *Ecology and Evolution* 13 (7). <https://doi.org/10.1002/ece3.10249>
- Nagy A, Katona P, Molnár A, Rádai Z, Tóth M, Szanyi K, Szanyi S (2023b) Wide range of brachyceran fly taxa attracted to synthetic and semi-synthetic generic noctuid lures and the description of new attractants for Sciomyzidae and Heleomyzidae families. *Insects* 14 (8). <https://doi.org/10.3390/insects14080705>
- O'Connor RS, Kunin WE, Garratt MPD, Potts SG, Roy HE, Andrews C, Jones CM, Peyton JM, Savage J, Harvey MC, Morris RKA, Roberts SPM, Wright I, Vanbergen A, Carvell C (2019) Monitoring insect pollinators and flower visitation: The effectiveness and feasibility of different survey methods. *Methods in Ecology and Evolution* 10 (12): 2129-2140. <https://doi.org/10.1111/2041-210X.13292>
- Ősz A, Szanyi K, Jakab DA, Szanyi S, Koczor S, Tóth M, Szarukán I, Nagy A (2025) A dataset of flower-visiting assemblages in the eastern Pannonian Lowland. Version 1.1. *Biodiversity Data Journal*. Occurrence dataset. URL: https://ipt.pensoft.net/resource?r=a_dataset_of_flower-visiting_assemblages_in_the_eastern_annonian_lowland&v=1.1
- Papp L, Darvas B (2000) Contributions to a manual of Palaearctic Diptera, Vol. 1. General and Applied Dipterology. Science Herald, Budapest.
- Rácz IA, Szanyi S, Nagy A (2023) Review on flower-visiting behaviour of orthopterans and setting priorities for further studies. *Biologia Futura* 74 (4): 393-400. <https://doi.org/10.1007/s42977-024-00203-9>
- Rauf A, Saeed S, Ali M, T. N, Muhammad H (2021) Comparative efficiency of native insect pollinators in reproductive performance of *Medicago sativa* L. in Pakistan. *Insects* 12 (11). <https://doi.org/10.3390/insects12111029>
- Roy H, Baxter E, Saunders A, Pocock MO (2016) Focal Plant Observations as a Standardised Method for Pollinator Monitoring: Opportunities and Limitations for Mass Participation Citizen Science. *PLoS ONE* 11 (3): 1-14. [In english]. <https://doi.org/10.1371/journal.pone.0150794>
- Sarma AK, Neog B, Deka MK, Carabet A, Stef R (2025) Variable Transect Method Outperformed in Sampling Hymenopteran Flower Visitors in *Brassica campestris* L. var.

- toria Ecosystem. *Agronomy* 15 (1281): 1-20. [In English]. <https://doi.org/10.3390/agronomy15061281>
- Schwartz M. D, Footitt R. G (1998) Revision of the Nearctic species of the genus *Lygus* Hahn, with a review of the Palearctic species (Heteroptera: Miridae). Associated Publishers, Gainesville, FL.
 - Segers A, Noël G, Delanglez L, Caparros Megido R, Francis F (2023) Impacts of Semiochemical Traps Designed for *Bruchus rufimanus* Boheman 1833 (Coleoptera: Chrysomelidae) on Nontarget Beneficial Entomofauna in Field Bean Crops. *Insects* 14 (2) (153). [In English]. <https://doi.org/10.3390/insects14020153>
 - Steffan-Dewenter I, Potts SG, Packer L (2005) Pollinator diversity and crop pollination services are at risk. *Trends in Ecology & Evolution* 20 (12): 651-652. <https://doi.org/10.1016/j.tree.2005.09.004>
 - Szalárdi T, Nagy A, Oláh F, Tóth M, Koczor S (2021) Preliminary data on attractiveness of phenylacetaldehyde-based lures on economically important plant bug pests (Hemiptera: Miridae). *International Journal of Horticultural Science* 27: 87-94. <https://doi.org/10.31421/ijhs/27/2021/8259>
 - Szanyi K, Nagy A, Ősz A, Ábrahám L, Molnár A, Tóth M, Szanyi S (2025) Nontarget catches of green and brown lacewings (Insecta: Neuroptera: Chrysopidae, Hemerobiidae) collected by light- and volatile-baited traps in the transcarpathian lowland (W Ukraine). *Insects* 16 (1). <https://doi.org/10.3390/insects16010074>
 - Szanyi S, Molnár A, Kozák L, Szalárdi T, Varga Z, Tóth M, Nagy A (2019) Nyírségi Macroheterocera együttesek vizsgálata illatanyagcsapdák alkalmazásával. *Erdészettudományi Közlemények* 9 (1): 51-68. <https://doi.org/10.17164/EK.2019.004>
 - Szanyi S, Molnár A, Szanyi K, Tóth M, Jósvai JK, Varga Z, Nagy A (2024) Semiochemical-baited traps as a new method supplementing light traps for faunistic and ecological studies of Macroheterocera (Lepidoptera). *Scientific Reports* 14 (1). <https://doi.org/10.1038/s41598-024-71109-8>
 - Thierry D, Cloupeau R, Jarry M, Canard M (1998) Discrimination of the West-Palearctic *Chrysoperla* Steinmann species of the carnea Stephens group by means of claw morphology (Neuroptera, Chrysopidae). *Acta Zool. Fenn.* 209: 255-262.
 - Thompson A, Frenzel M, Schweiger O, Musche M, Groth T, Roberts SM, Kuhlmann M, Knight T (2021) Pollinator sampling methods influence community patterns assessments by capturing species with different traits and at different abundances. *Ecological Indicators* 132 (108284): 1-10. [In English]. <https://doi.org/10.1016/j.ecolind.2021.108284>
 - Tóth M., Szarukán I., Nagy A., Gém F., Nyitrai R., Kecskés Z., Krakkó L., Jósvai J. K., Béla I. (2020) Félpszintetikus "biszex" csalétek kártevő rovarok nőtényeinek és hímjeinek fogására. *Növényvédelem* 51: 197-205.
 - Tóth S (2017) Képes zengőléghatározó (Diptera: Syrphidae). *e-Acta Naturalia Pannonica* 15: 1-228. <https://doi.org/10.24369/EANP.2017.15.1>
 - Varga Z. (2010) Magyarország nagylepkéi. Heterocera Press, Budapest.
 - Wagner E (1952) Blindwanzen oder Miriden. *Die Tierwelt Deutschlands*. Verlag von Gustav Fischer

Supplementary material

Suppl. material 1: Relative frequencies

Authors: Ősz A., Szanyi K., Arnóczkyné Jakab D., Szanyi Sz., Koczor S., Tóth M., Szarukán I., Nagy A.

Data type: Occurences

Brief description: Relative frequencies (RF%) of insect orders, families (only in the case of Hymenoptera and Lepidoptera orders) and subfamilies (only in the case of Noctuidae family) calculated based on all caught individuals and individual numbers of insects identified to species level.

[Download file](#) (21.42 kb)