

# Path properties of dilatively stable processes and singularity of their distributions

ENDRE IGLÓI<sup>\*, $\diamond$</sup>  and MÁTYÁS BARCZY<sup>\*</sup>

<sup>\*</sup> University of Debrecen, Faculty of Informatics, Pf. 12, H-4010 Debrecen, Hungary,  
Tel.: 06-52-512900, Fax: 06-52-512996

e-mails: igloi@tigris.unideb.hu (E. Iglói), barczy.matyas@inf.unideb.hu (M. Barczy),

$\diamond$  Corresponding author.

*2010 Mathematics Subject Classifications:* 60G18, 60G17, 60G30, 60G22, 60J60

*Key words and phrases:* dilatively stable processes, self-similar processes, sample path properties, Hölder continuity, singularity

## Abstract

First, we present some results about the Hölder continuity of the sample paths of so called dilatively stable processes which are certain infinitely divisible processes having a more general scaling property than self-similarity. As a corollary, we obtain that the most important  $(H, \delta)$ -dilatively stable limit processes (e.g., the LISOU and the LISCBI processes, see Iglói [4]) almost surely have a local Hölder exponent  $H$ . Next we prove that, under some slight regularity assumptions, any two dilatively stable processes with stationary increments are singular (in the sense that their distributions have disjoint supports) if their parameters  $H$  are different. We also study the more general case of not having stationary increments. Throughout the paper we specialize our results to some basic dilatively stable processes such as the above-mentioned limit processes and the fractional Lévy process.

## 1 Introduction

Path properties are important features of a stochastic process, this fact does not call for an explanation. Given a set of stochastic processes, the same is true for the singularity of their distributions on the space of sample paths (by which we understand that under certain conditions any two such distributions have disjoint supports).

A well-known feature enabling a process to have nice path properties is self-similarity. Systematic investigations of the sample path properties of self-similar processes had been

begun by Vervaat's classical paper [22], and continued by Vervaat [23], Takashima [20] and Watanabe and Yamamuro [24] (the latter one gives laws of the iterated logarithm for multidimensional self-similar processes with independent increments). Maejima [11], Kono and Maejima [8] and Samorodnisky [15] (among others) treated the sample path properties of some special self-similar processes. However, there are very few results in the literature about the singularity properties of distributions of self-similar processes. To the authors' knowledge the only paper concerning this subject is that of Prakasa Rao [13], which states that the distributions of two fractional Brownian motions (FBMs) with different Hurst parameters are singular with respect to each other. The question of singularity for other types of processes, for instance, for diffusion processes deserved more attention, see, e.g., Ben-Ari and Pinsky [1], Jacod and Shiryaev [5] and Zhang [25].

In this paper we show that the so-called dilatively stable processes, introduced by Iglói [4], can have nice sample path and singularity properties. Dilative stability is a generalization of self-similarity. For the comparison of these two notions, we give both definitions.

**1.1 Definition.** Let  $\alpha > 0$ . A process  $\{X(t), t \geq 0\}$  starting from zero (i.e.,  $X(0) = 0$ ) is called  $\alpha$ -self-similar if it is not identically zero and fulfills the scaling relation

$$\forall T > 0 : X(T \cdot) \stackrel{fd}{\sim} T^\alpha X(\cdot), \quad (1.1)$$

where  $\stackrel{fd}{\sim}$  denotes that the finite-dimensional distributions are the same.

Dilative stability, see Iglói [4], is an analogous property of certain infinitely divisible processes (all finite-dimensional distributions are infinitely divisible) involving a scaling also in the convolution exponent. The exact definition is as follows.

**1.2 Definition.** Let  $\alpha > 0$  and  $\delta \leq 2\alpha$ . A process  $\{X(t), t \geq 0\}$  starting from zero is said to be  $(\alpha, \delta)$ -dilatively stable if its finite-dimensional distributions are infinitely divisible,  $X(1)$  is non-Gaussian,  $X(t)$  has finite moments of all orders for all  $t \geq 0$ , and fulfills the scaling relation

$$\forall T > 0 : X(T \cdot) \stackrel{fd}{\sim} T^{\alpha-\delta/2} X^{\otimes T^\delta}(\cdot). \quad (1.2)$$

Here for all  $c > 0$ , we denote by  $X^{\otimes c} = \{X(t), t \geq 0\}^{\otimes c}$  the  $c$ -th convolution power of  $\{X(t), t \geq 0\}$ , that is,  $\{X(t), t \geq 0\}^{\otimes c}$  is a process the finite dimensional distributions of which are the  $c$ -th convolution powers of the corresponding ones of  $\{X(t), t \geq 0\}$ .

In Appendix A we give more insight to the properties of a dilatively stable process that are supposed in Definition 1.2. We also note that in Kaj [7, Section 3.6] one can find a somewhat similar, but not so general concept called aggregate-similarity. One of the main differences between the concept of dilative stability and aggregate-similarity

is that the later one defined only in the case of  $n$ -th convolution powers with  $n \in \mathbb{N}$  and when the relation  $2\alpha - \delta = 2$  holds for the parameters (which is in fact the most important case, see the parametrization of the LIS processes in Example 1.3).

Let us observe that a not identically zero process  $\{X(t), t \geq 0\}$  starting from zero and fulfilling the scaling relation (1.2) with  $\alpha > 0$  and  $\delta = 0$  is an  $\alpha$ -self-similar process, roughly speaking  $(\alpha, 0)$ -dilative stability is just  $\alpha$ -self-similarity. Hence dilative stability is a generalization of self-similarity. Many results, such as Lamperti's theorems, can be transferred from the self-similar to the dilatively stable case, see Iglói [4]. Self-similar processes are fixed points of their renormalization operators (see Taqqu [21]), and the same is true for dilatively stable processes (see, Iglói [4, Theorem 2.8.4 (DS)]). Accordingly, the possible limit processes in linear rescaling (i.e., in self-similar renormalization limit theorems) are the self-similar processes, see Lamperti [10], and if there is a rescaling also in the convolution exponent (i.e., in dilatively stable renormalization limit theorems) the possible limit processes are the dilatively stable ones, see, Iglói [4, Theorem 2.2.7]. This is why dilative stability is important.

Important examples for dilatively stable processes are non-Gaussian fractional Lévy processes (FLPs) (i.e., FLPs, where the underlying two-sided Lévy process is non-Gaussian, but possibly with a Gaussian component) having zero mean and finite moments of all orders. FLPs were originally introduced by Benassi et al. [2] and Marquardt [12]. For historical fidelity we note that in [2] a FLP is called a moving-average fractional Lévy motion. We also recall that Marquardt [12, Theorem 4.4] proved that a FLP with an underlying two-sided Lévy process having zero mean, finite second moment and not having a Brownian component cannot be self-similar. However, by Iglói [4, Example 2.1.7], a non-Gaussian FLP (considered only on  $[0, \infty)$ ) having zero mean and finite moments of all orders is  $(H, 1)$ -dilatively stable with stationary increments, where  $H \in (1/2, 1)$  is the so-called Hurst parameter or long memory parameter (see also Kaj [7, page 212]). Roughly speaking, non-Gaussian FLPs are not self-similar but they belong to a wider class of processes, to the class of dilatively stable processes, which also underlines the importance of dilative stability.

The main contributions of this paper relate to dilatively stable processes with stationary increments, the general properties of which are treated by Iglói [4, Section 2.7]. Here we point out that in this case the range of the parameter  $H \doteq \alpha$  is  $(0, 1]$ , see, Iglói [4, Theorem 2.7.1 (DS) 1)]. In case  $H = 1$ , i.e., in case of a  $(1, \delta)$ -dilatively stable process we have  $\delta = 0$  (i.e., the process is self-similar) and it takes the form  $X(t) = tX(1)$ ,  $t \geq 0$ , almost surely (a.s.), i.e., it is degenerate. Indeed, by Iglói [4, Theorem 2.7.1 (DS) 2)], the process is degenerate, and if we suppose on the contrary that  $\delta \neq 0$ , then, by (1.2) (with  $T \doteq c^{1/\delta}$ ,  $c > 0$ ),

$$X(c^{1/\delta}) \sim c^{\frac{1}{\delta}(1-\delta/2)} X^{\otimes c}(1), \quad c > 0,$$

which yields that

$$c^{\frac{1}{\delta}}X(1) \sim c^{\frac{1}{\delta}-\frac{1}{2}}X^{\otimes c}(1), \quad \text{i.e.,} \quad X(1) \sim \frac{1}{\sqrt{c}}X^{\otimes c}(1), \quad c > 0.$$

Hence  $X(1)$  is Gaussian with mean zero (which follows by considering characteristic functions), in contradiction with Definition 1.2.

Moreover, an  $(H, \delta)$ -dilatively stable process  $\{X(t), t \geq 0\}$  with stationary increments has the same covariance function as a FBM with parameter  $H$  (apart from a constant factor):

$$\text{Cov}(X(t_1), X(t_2)) = \frac{1}{2}D^2X(1) (t_1^{2H} + t_2^{2H} - |t_1 - t_2|^{2H}), \quad t_1, t_2 \geq 0,$$

see Iglói [4, Theorem 2.7.2] (where the parameter  $H$  can not be one, however the proof given there works also in case  $H = 1$ ).

Moments and cumulants provide a very handy tool throughout when dealing with dilative stability. Using this tool in Section 2 we obtain some results for the almost sure limit behavior of dilatively stable processes at zero, at infinity and, in case of processes with stationary increments, at any point, see Lemmas 2.3, 2.5 and 2.6. Then, by the help of the Kolmogorov–Chentsov theorem, we characterize dilatively stable processes with stationary increments from the point of view of Hölder continuity of their sample paths, see Theorem 2.7. This characterization has a well-known self-similar analogue presented in Remark 2.11. Section 3 contains some results about the singularity of the distributions of dilatively stable processes, induced on the space of continuous functions. Though a dilatively stable process does not admit automatically continuous sample paths almost surely, in the last section we will restrict ourselves to such processes. We note that an  $(H, \delta)$ -dilatively stable process with stationary increments and zero mean such that either  $\delta \neq 2H$  or  $H > 1/2$  has a continuous modification, see Corollary 2.10. Using the path properties proved in Section 2, we will find that, under some slight regularity conditions, the distributions of dilatively stable processes with different parameters  $\alpha$  are pairwise singular, see Theorem 3.1 and Theorem 3.2 (this latter one is for the stationary increments case). Theorem 3.2 can be carried out to self-similar processes with finite absolute moment and stationary increments, implying, particularly, a new (and simple) proof of the pairwise singularity of FBMs with different Hurst parameters, see Theorem 3.5.

Throughout the paper we will specialize our results to some particular dilatively stable processes with stationary increments presented in the next example.

**1.3 Example.** *The following processes are dilatively stable with stationary increments, see Iglói [4]:*

- *LISOU process (limit of integrated superposition of Ornstein–Uhlenbeck type processes),*

- *LISCBI process* (limit of **i**ntegrated superposition of **c**ontinuous state **b**ranched processes with **i**mmigration),
- *LISDLG process* (limit of **i**ntegrated superposition of **d**iffusion processes with **l**inear generator; this is a particular LISCBI process),
- *non-Gaussian FLP* (**f**ractional **L**évy **p**rocess), i.e., FLP defined on  $[0, \infty)$ , where the underlying two-sided Lévy process is non-Gaussian (but possibly with a Gaussian component) having zero mean and finite moments of all orders.

These processes have a parameter  $H$ , which can now be only in the interval  $(1/2, 1)$ , hence it can be called the Hurst parameter or long memory parameter. The LISOU, LISCBI and LISDLG processes with parameter  $H$  are  $(H, 2H - 2)$ -dilatively stable, while the non-Gaussian FLP with parameter  $H$  is  $(H, 1)$ -dilatively stable, and all of these processes have the same covariance function as a FBM with parameter  $H$ .

We will suppose throughout that the processes have zero mean. This is a natural assumption when dealing with path properties, since the mean function, as a deterministic function, can be handled separately. One can also check that the subtraction of the mean function preserves dilative stability. In addition, if an  $(H, \delta)$ -dilatively stable process has stationary increments, then its mean function is automatically zero, unless  $H + \delta/2 = 1$ , see Iglói [4, Theorem 2.7.1 (DS) 3)].

## 2 Path properties

Throughout in this paper  $I \subseteq [0, \infty)$  denotes an interval. Let  $\gamma \in (0, 1]$ . A function  $f : I \rightarrow \mathbb{R}$  is called *locally  $\gamma$ -Hölder continuous* if for every bounded subinterval  $J \subseteq I$ ,

$$\sup_{\substack{t, s \in J, \\ t \neq s}} \frac{|f(t) - f(s)|}{|t - s|^\gamma} < \infty,$$

see, e.g., Revuz and Yor [14, page 26]. Clearly, local  $\gamma_1$ -Hölder continuity implies local  $\gamma_2$ -Hölder continuity if  $1 \geq \gamma_1 \geq \gamma_2 > 0$ . This relation gives rise to the following notion of local Hölder exponent. The value

$$\Gamma_f \doteq \sup_{\gamma \in (0, 1]} \{ \gamma : f \text{ is locally } \gamma\text{-Hölder continuous} \}$$

is called the (*optimal*) *local Hölder exponent* of a function  $f$  and set to 0 if  $f$  is not locally Hölder continuous. This notion is similar to the notion of the optimal Hölder index at a point, see, e.g., Jaffard [6] or Fleischmann, Mytnik and Wachtel [3].

In what follows by the expression that ‘an infinitely divisible distribution has a Gaussian component’ we mean that in its Lévy-Khintchine representation the Gaussian part has positive variance.

In some cases we will refer to the following assumptions.

**2.1 Assumption.** If  $\{X(t), t \geq 0\}$  is an  $(\alpha, \delta)$ -dilatively stable process with  $\delta \geq 0$ , then the distribution of  $X(1)$  (equivalently, the distribution of  $X(t)$  for any  $t > 0$ ) has a Gaussian component.

**2.2 Assumption.** If  $\{X(t), t \geq 0\}$  is an  $(\alpha, \delta)$ -dilatively stable process with  $\delta \leq 0$ , then the distribution of  $X(1)$  (equivalently, the distribution of  $X(t)$  for any  $t > 0$ ) has a Gaussian component.

Note that the above two assumptions are the dual of each other with respect to  $\delta$ , and we will always indicate explicitly which of them is used. Furthermore, when Assumption 2.1 is supposed and the parameter  $\delta$  of an  $(\alpha, \delta)$ -dilatively stable process is negative, then this assumption does not come into play, and a similar statement holds for Assumption 2.2 and an  $(\alpha, \delta)$ -dilatively stable process with positive  $\delta$ . The intrinsic reason for considering only the non-negative values  $\delta$  in Assumption 2.1 is that the LISOU, LIS-CBI and LISDLG processes (the dilatively stable processes with stationary increments presented in Example 1.3) may not have a Gaussian component and their parameter  $\delta$  is negative. However, this also means that Assumption 2.2 may not hold for these particular dilatively stable processes, and from this specific point of view Assumptions 2.1 and 2.2 are not exactly the dual of each other. On the other side, we call the attention that these assumptions are not too restrictive, since, as it is easy to see, the independent sum of a Gaussian  $\alpha$ -self-similar process with zero mean and an  $(\alpha, \delta)$ -dilatively stable process remains  $(\alpha, \delta)$ -dilatively stable. Particularly, the independent sum of an  $(H, \delta)$ -dilatively stable process and a FBM with parameter  $H$  remains  $(H, \delta)$ -dilatively stable, see Iglói [4, Proposition 2.7.5]. Thus, we can make a Gaussian component preserving the dilative stability. Finally, we remark that under the non-trivial cases of Assumption 2.1 or Assumption 2.2, i.e., when there exists a Gaussian component, the distribution of  $X(1)$  is absolutely continuous (see, e.g., Sato [16, Lemma 27.1]), and hence  $P(X(1) = 0) = 0$  in this case.

The following lemma treats the sample path behaviour at zero of a dilatively stable process. In what follows by a zero-sequence we mean a sequence of real numbers converging to 0.

**2.3 Lemma.** *There exists a zero-sequence  $(t_n)_{n \in \mathbb{N}}$  with positive terms such that for any  $(\alpha, \delta)$ -dilatively stable process  $\{X(t), t \geq 0\}$  with zero mean, the following assertions hold.*

(i) *If  $\kappa < \alpha$ , then*

$$\limsup_{n \rightarrow \infty} \frac{|X(t_n)|}{t_n^\kappa} = 0 \quad \text{a.s.} \quad (2.1)$$

(ii) *If  $\kappa > \alpha$  and Assumption 2.1 holds, then*

$$\limsup_{n \rightarrow \infty} \frac{|X(t_n)|}{t_n^\kappa} = \infty \quad \text{a.s.} \quad (2.2)$$

Further, for any sequence  $(t_n)_{n \in \mathbb{N}}$  with positive terms such that

$$\limsup_{n \rightarrow \infty} \sqrt[n]{t_n} < 1, \quad (2.3)$$

the assertions of parts (i) and (ii) hold.

**Proof.** Let  $(t_n)_{n \in \mathbb{N}}$  be a sequence with positive terms such that (2.3) holds. Then the root test yields that the series  $\sum_{n=1}^{\infty} t_n$  is convergent, and hence  $(t_n)_{n \in \mathbb{N}}$  is a zero-sequence.

(i) By (1.2), we have

$$D^2 X(t_n) = D^2(t_n^{\alpha-\delta/2} X^{\otimes t_n^\delta}(1)) = t_n^{2\alpha-\delta+\delta} D^2 X(1) = t_n^{2\alpha} D^2 X(1), \quad n \in \mathbb{N}.$$

Hence the Markov inequality yields that

$$\sum_{n=1}^{\infty} \mathbb{P} \left( \frac{|X(t_n)|}{t_n^\kappa} > \varepsilon \right) \leq \frac{D^2 X(1)}{\varepsilon^2} \sum_{n=1}^{\infty} t_n^{2(\alpha-\kappa)} < \infty, \quad \varepsilon > 0,$$

where the convergence of the series is a consequence of (2.3). Applying the Borel–Cantelli lemma we obtain that  $\lim_{n \rightarrow \infty} X(t_n)/t_n^\kappa = 0$  a.s., which implies (2.1). We remark that in this case any sequence  $(t_n)_{n \in \mathbb{N}}$  with positive terms satisfying (2.3) is obviously universal.

(ii) Using (1.2), we have

$$\frac{|X(t_n)|}{t_n^\kappa} \sim t_n^{\alpha-\kappa} \frac{|X^{\otimes t_n^\delta}(1)|}{t_n^{\delta/2}}, \quad n \in \mathbb{N}. \quad (2.4)$$

First we show that the right-hand side (and hence the left-hand side) of (2.4) converges in probability to infinity along some appropriate subsequence. We consider the three cases corresponding to the sign of  $\delta$  separately.

• If  $\delta < 0$ , then  $X^{\otimes t_n^\delta}(1)/t_n^{\delta/2}$  converges in distribution to a normal distribution with variance  $\mathbb{E}(X(1))^2$ . Indeed,  $\lim_{n \rightarrow \infty} t_n^\delta = \infty$ , and

$$X^{\otimes t_n^\delta}(1) \sim \xi_1 + \cdots + \xi_{\lfloor t_n^\delta \rfloor} + \eta_n,$$

where  $\lfloor \cdot \rfloor$  denotes the integer part,  $\xi_1, \dots, \xi_{\lfloor t_n^\delta \rfloor}$  and  $\eta_n$  are independent random variables such that  $\xi_1, \dots, \xi_n$  has a common distribution as  $X(1)$  has and  $\eta_n \sim X^{\otimes(t_n^\delta - \lfloor t_n^\delta \rfloor)}(1)$ . By the central limit theorem we get that

$$\frac{\xi_1 + \cdots + \xi_{\lfloor t_n^\delta \rfloor}}{t_n^{\delta/2}}$$

converges in distribution to a normal distribution with variance  $\mathbb{E}(X(1))^2$ , while the remainder term  $\eta_n/t_n^{\delta/2}$  converges to 0 in  $L^2$  as  $n \rightarrow \infty$ . Therefore the right-hand



side of (2.4) converges in probability to infinity, and hence so does the left-hand side. (Here we used the fact that if  $\zeta_n, n \in \mathbb{N}$ , and  $\zeta$  are non-negative random variables such that  $\zeta_n$  converges to  $\zeta$  in distribution,  $\zeta$  has a continuous distribution function [hence  $\mathbf{P}(\zeta > 0) = 1$ ] and  $(c_n)_{n \in \mathbb{N}}$  is a sequence of positive real numbers converging to infinity, then  $c_n \zeta_n$  converges in probability to infinity as  $n \rightarrow \infty$ , i.e.,  $\forall M > 0 : \lim_{n \rightarrow \infty} \mathbf{P}(c_n \zeta_n < M) = 0$ .)

- If  $\delta = 0$ , then the right-hand side of (2.4) is  $t_n^{\alpha-\kappa} |X(1)|$ . Using Assumption 2.1 and the fact that if at least one term of an independent (finite) sum of random variables has a continuous distribution, then also the sum itself has a continuous distribution (see, e.g., Sato [16, Lemma 27.1]), we have  $\mathbf{P}(X(1) = 0) = 0$ , i.e.,  $X(1)$  has no atom at zero. Hence  $t_n^{\alpha-\kappa} |X(1)|$  converges almost surely to infinity, which yields that the left-hand side of (2.4) converges in probability to  $\infty$ .

- If  $\delta > 0$ , then, since

$$\mathbf{D}^2 \left( \frac{X^{\otimes t_n^\delta}(1)}{t_n^{\delta/2}} \right) = \mathbf{D}^2 X(1) = \mathbf{E}(X(1))^2, \quad n \in \mathbb{N}, \quad (2.5)$$

the tightness of the sequence of distributions of  $X^{\otimes t_n^\delta}(1)/t_n^{\delta/2}$ ,  $n \in \mathbb{N}$ , follows by the Markov inequality. Indeed, with the notation

$$K_\varepsilon \doteq \left[ -\sqrt{\mathbf{E}(X(1))^2/\varepsilon}, \sqrt{\mathbf{E}(X(1))^2/\varepsilon} \right], \quad \varepsilon > 0,$$

for all  $n \in \mathbb{N}$ , we have

$$\mathbf{P} \left( \frac{X^{\otimes t_n^\delta}(1)}{t_n^{\delta/2}} \in \mathbb{R} \setminus K_\varepsilon \right) = \mathbf{P} \left( \left| \frac{X^{\otimes t_n^\delta}(1)}{t_n^{\delta/2}} \right| > \sqrt{\frac{\mathbf{E}(X(1))^2}{\varepsilon}} \right) \leq \varepsilon.$$

Therefore, by Prohorov's theorem, there exists a subsequence  $X^{\otimes t_{n_k}^\delta}(1)/t_{n_k}^{\delta/2}$ ,  $k \in \mathbb{N}$ , which converges in distribution. Using Assumption 2.1 and the Lévy–Khintchine formula we get that for all  $k \in \mathbb{N}$ ,  $X^{\otimes t_{n_k}^\delta}(1)/t_{n_k}^{\delta/2}$  has a Gaussian component, the distribution of which is the same as the distribution of the Gaussian component of  $X(1)$ . Indeed, if  $G$  denotes the Gaussian component of  $X(1)$ , then  $G^{\otimes t_{n_k}^\delta}/t_{n_k}^{\delta/2}$  is the Gaussian component of  $X^{\otimes t_{n_k}^\delta}(1)/t_{n_k}^{\delta/2}$  and it has the same distribution as  $G$  (here we also use that  $G$  has zero mean since  $X(1)$  has zero mean). Thus the limiting distribution of this subsequence also has a Gaussian component. Therefore the limiting distribution is continuous and hence it has no atom at zero, and, as it was explained earlier in the proof of the case  $\delta < 0$ , this fact ensures that the right-hand side of (2.4) converges along the subsequence  $(t_{n_k})_{k \in \mathbb{N}}$  in probability to infinity, hence so does the left-hand side (along the subsequence  $(t_{n_k})_{k \in \mathbb{N}}$ ).

We call the attention that in the proof of the case  $\delta > 0$  (i.e., in the last case) we do not use that  $\delta$  is positive, but Assumption 2.1 comes into play in this case. This also



shows that the proof of the case  $\delta > 0$  remains still valid for the case  $\delta < 0$  under the additional assumption that  $X(1)$  has a Gaussian component. (However, the proof of the case  $\delta < 0$  presented above does not work for the case  $\delta > 0$ .)

In each of the above three cases (denoted by bullets) we obtained that a subsequence of the left-hand side of (2.4) (in case of  $\delta \leq 0$  the whole sequence) converges in probability to infinity. Thus, by the Riesz lemma, there is some subsequence of the left-hand side of (2.4), which converges to infinity almost surely. Indeed, Riesz's lemma can be proved for a sequence of random variables  $(\zeta_n)_{n \in \mathbb{N}}$  converging in probability to  $\infty$ , as follows. Since for all  $k \in \mathbb{N}$  there exists some  $n_k \in \mathbb{N}$  such that  $P(\zeta_{n_k} < k) < 1/2^k$ , the Borel–Cantelli lemma yields that

$$P(\zeta_{n_k} < k \text{ for infinitely many } k \geq 1) = 0.$$

Hence  $P(\liminf_{k \rightarrow \infty} \zeta_{n_k} = \infty) = 1$  which yields that  $P(\lim_{k \rightarrow \infty} \zeta_{n_k} = \infty) = 1$ . This implies (2.2) and we also have the universality of any sequence  $(t_n)_{n \in \mathbb{N}}$  with positive terms satisfying (2.3).  $\square$

**2.4 Remark.** We note that in part (i) of Lemma 2.3 one can also write  $\lim$  instead of  $\limsup$ . The reason for writing  $\limsup$  is that we will use only this later on. We also remark that the case  $\kappa = \alpha$  is not covered by Lemma 2.3, since we do not need it and we can not address any result in this case.

The following lemma shows that the behaviour of the sample paths at infinity can be characterized similarly as their behaviour at zero.

**2.5 Lemma.** *There exists a sequence  $(t_n)_{n \in \mathbb{N}}$  converging to infinity such that for any  $(\alpha, \delta)$ -dilatively stable process  $\{X(t), t \geq 0\}$  with zero mean the following assertions hold.*

(i) *If  $\kappa < \alpha$  and Assumption 2.2 holds, then*

$$\limsup_{n \rightarrow \infty} \frac{|X(t_n)|}{t_n^\kappa} = \infty \quad \text{a.s.} \quad (2.6)$$

(ii) *If  $\kappa > \alpha$ , then*

$$\limsup_{n \rightarrow \infty} \frac{|X(t_n)|}{t_n^\kappa} = 0 \quad \text{a.s.} \quad (2.7)$$

*Further, for any sequence  $(t_n)_{n \in \mathbb{N}}$  with positive terms such that  $\limsup_{n \rightarrow \infty} \sqrt[n]{1/t_n} < 1$ , the assertions of parts (i) and (ii) hold.*

**Proof.** One can use the same arguments as in the proof of Lemma 2.3, but with some changes. Namely, the sequence  $(t_n)_{n \in \mathbb{N}}$  can be the reciprocal of the sequence (with positive terms satisfying (2.3)) used in the proof of Lemma 2.3. Then the proof of statement

(ii) corresponds to that of statement (i) in Lemma 2.3, while part (i) goes the same way as part (ii) in Lemma 2.3, just the cases  $\delta < 0$  and  $\delta > 0$  have to be interchanged.  $\square$

In the stationary increments case Lemma 2.3 can be formulated not only for zero-sequences but also for sequences converging to a non-negative real number, see as follows.

**2.6 Lemma.** *For any  $t_0 \geq 0$  there exists a sequence  $(t_n)_{n \in \mathbb{N}}$  converging to  $t_0$  such that  $t_n \neq t_0$ ,  $n \in \mathbb{N}$ , and for any  $(H, \delta)$ -dilatively stable process  $\{X(t), t \geq 0\}$  with stationary increments and zero mean the following assertions hold.*

(i) *If  $\kappa < H$ , then*

$$\limsup_{n \rightarrow \infty} \frac{|X(t_n) - X(t_0)|}{|t_n - t_0|^\kappa} = 0 \quad \text{a.s.} \quad (2.8)$$

(ii) *If  $\kappa > H$  and Assumption 2.1 holds, then*

$$\limsup_{n \rightarrow \infty} \frac{|X(t_n) - X(t_0)|}{|t_n - t_0|^\kappa} = \infty \quad \text{a.s.} \quad (2.9)$$

Further, for any sequence  $(t_n)_{n \in \mathbb{N}} \doteq (t_0 + \tilde{t}_n)_{n \in \mathbb{N}}$ , where  $(\tilde{t}_n)_{n \in \mathbb{N}}$  is a sequence with positive terms such that  $\limsup_{n \rightarrow \infty} \sqrt[n]{\tilde{t}_n} < 1$ , the assertions of parts (i) and (ii) hold.

**Proof.** The process  $Y(t) \doteq X(t + t_0) - X(t_0)$ ,  $t \geq 0$ , is  $(H, \delta)$ -dilatively stable with zero mean such that the distribution of  $Y(1)$  has a Gaussian component if  $X(1)$  has. Hence Lemma 2.3 yields that there exists a zero-sequence  $(\tilde{t}_n)_{n \in \mathbb{N}}$  with positive terms such that if  $\kappa < H$ , then

$$\limsup_{n \rightarrow \infty} \frac{|X(\tilde{t}_n + t_0) - X(t_0)|}{\tilde{t}_n^\kappa} = 0 \quad \text{a.s.},$$

and if  $\kappa > H$  and Assumption 2.1 holds, then

$$\limsup_{n \rightarrow \infty} \frac{|X(\tilde{t}_n + t_0) - X(t_0)|}{\tilde{t}_n^\kappa} = \infty \quad \text{a.s.}$$

With the definition  $t_n \doteq \tilde{t}_n + t_0$ ,  $n \in \mathbb{N}$ , we have the assertions of the lemma. (This also shows that in (2.8) and (2.9) one can write  $(t_n - t_0)^\kappa$  instead of  $|t_n - t_0|^\kappa$ .)  $\square$

The above lemmas, interesting in their own rights, will be used mainly in the next section, but Lemma 2.6 will appear also in the proof of the following theorem. The rest of the section deals with Hölder continuity of the sample paths of dilatively stable processes with stationary increments.

**2.7 Theorem.** *Let  $\{X(t), t \geq 0\}$  be an  $(H, \delta)$ -dilatively stable process with stationary increments and zero mean such that either  $\delta \neq 2H$  or  $H > 1/2$ . Then  $\{X(t), t \geq 0\}$  has a continuous modification, the sample paths of which are locally  $\gamma$ -Hölder continuous*

- (i) for every  $\gamma \in (0, H)$  if  $\delta < 0$ ,
- (ii) for every  $\gamma \in (0, H - \delta/2)$  if  $0 \leq \delta < 2H$ ,
- (iii) for every  $\gamma \in (0, H - 1/2)$  if  $\delta = 2H$  and  $H > 1/2$ .

Moreover, under Assumption 2.1, for the local Hölder exponent  $\Gamma_X$  of the sample paths of the above continuous modification of the process  $\{X(t), t \geq 0\}$  we have

- (i)  $\Gamma_X = H$  if  $\delta < 0$ ,
- (ii)  $\Gamma_X \in [H - \delta/2, H]$  if  $0 \leq \delta < 2H$ ,
- (iii)  $\Gamma_X \in [H - 1/2, H]$  if  $\delta = 2H$  and  $H > 1/2$ .

**Proof.** We are going to apply the Kolmogorov–Chentsov theorem see, e.g., Revuz and Yor [14, Chapter I, Theorem 2.1]. Therefore we have to prove that  $\{X(t), t \geq 0\}$  satisfies Kolmogorov’s condition: there exist constants  $c, p, q > 0$  such that

$$\mathbb{E}|X(t) - X(s)|^p \leq c|t - s|^{1+q}, \quad s, t \geq 0. \quad (2.10)$$

Then, by the Kolmogorov–Chentsov theorem,  $\{X(t), t \geq 0\}$  has a continuous modification which is locally  $\gamma$ -Hölder continuous for every  $\gamma \in (0, q/p)$ . Clearly, it is sufficient for (2.10) to hold for every  $s, t \geq 0$  for which  $0 \leq t - s < 1$ , so in what follows we suppose that  $t$  and  $s$  are of these kinds. Indeed, in this case for all  $n \in \mathbb{N} \cup \{0\}$ ,

$$\mathbb{E}|X^{(n)}(t) - X^{(n)}(s)|^p \leq c|t - s|^{1+q}, \quad s, t \geq 0,$$

where

$$X^{(n)}(t) \doteq \begin{cases} X(n/2) & \text{if } t \leq n/2, \\ X(t) & \text{if } n/2 < t < n/2 + 1, \\ X(n/2 + 1) & \text{if } t \geq n/2 + 1, \end{cases} \quad n \in \mathbb{N} \cup \{0\},$$

and hence for all  $n \in \mathbb{N} \cup \{0\}$ , the process  $\{X^{(n)}(t) : t \geq 0\}$  has a continuous modification which is locally  $\gamma$ -Hölder continuous for every  $\gamma \in (0, q/p)$ . The desired property of  $\{X(t) : t \geq 0\}$  follows by that  $[0, \infty) = \bigcup_{n \in \mathbb{N} \cup \{0\}} [n/2, n/2 + 1]$ .

Now, let  $p$  be a positive even number. Using the stationary increments property and the relation between moments and cumulants we obtain

$$\mathbb{E}|X(t) - X(s)|^p = \mathbb{E}(X(t - s))^p = \sum_{\Pi} \prod_{B \in \Pi} \text{Cum}_{n_B}(X(t - s)), \quad (2.11)$$

where  $\Pi$  runs through the list of all partitions of a set of size  $p$ ,  $B \in \Pi$  means that  $B$  is one of the blocks into which the set is partitioned given the partition  $\Pi$ ,  $n_B$  is the size of the set  $B$  (in notation:  $n_B = |B|$ ) and  $\text{Cum}_k(X(t - s))$  denotes the cumulant

of order  $k$  of  $X(t-s)$ , see, e.g., Shiryaev [17, page 292, formula (46)]. Observe that  $\mathbb{E}X(t-s) = 0$  implies that for any  $\Pi$  and any one element block  $B \in \Pi$ , we have  $\text{Cum}_{n_B}(X(t-s)) = \text{Cum}_1(X(t-s)) = \mathbb{E}X(t-s) = 0$ , which yields that for any partition  $\Pi$  the sum on the right hand side of (2.11) has at most  $p/2$  terms different from 0. Using the dilative stability relation (1.2) we can continue (2.11) as follows:

$$\begin{aligned} \mathbb{E}|X(t) - X(s)|^p &= \sum_{\Pi} \prod_{B \in \Pi} \text{Cum}_{n_B}(X(t-s)) \\ &= \sum_{\Pi} \prod_{B \in \Pi} (t-s)^{(H-\delta/2)n_B+\delta} \text{Cum}_{n_B}(X(1)) \\ &= (t-s)^{(H-\delta/2)p} \sum_{\Pi} \left( (t-s)^{\delta|\Pi|} \prod_{B \in \Pi} \text{Cum}_{n_B}(X(1)) \right). \end{aligned} \quad (2.12)$$

It is important to observe that for each  $\Pi$ , the product  $\prod_{B \in \Pi} \text{Cum}_{n_B}(X(1))$  is non-negative. Indeed, if  $n_B$  is even, then  $\text{Cum}_{n_B}(X(1)) \geq 0$ , since the cumulant of order greater than or equal to 2 of  $X(1)$  is the moment of the same order of the Lévy measure in the Lévy–Khintchine representation of the distribution of  $X(1)$  (plus the variance of the Gaussian component if  $n_B = 2$ ), see Steutel and Van Harn [18, Chapter IV, Theorem 7.4]. A cumulant  $\text{Cum}_{n_B}(X(1))$  can be negative for an odd  $n_B$ , however, the number of blocks  $B \in \Pi$ , with an odd size  $n_B$ , must be even, since  $p$  is even. This yields the nonnegativity of  $\prod_{B \in \Pi} \text{Cum}_{n_B}(X(1))$ . At this point the proof separates into three cases corresponding to the three parts of the statement of the theorem.

- If  $\delta < 0$ , then (2.12) can be continued in the following way:

$$\begin{aligned} \mathbb{E}|X(t) - X(s)|^p &= (t-s)^{(H-\delta/2)p} \sum_{\Pi} \left( (t-s)^{\delta|\Pi|} \prod_{B \in \Pi} \text{Cum}_{n_B}(X(1)) \right) \\ &\leq (t-s)^{(H-\delta/2)p} \sum_{\Pi} \left( (t-s)^{\delta p/2} \prod_{B \in \Pi} \text{Cum}_{n_B}(X(1)) \right) \\ &= \mathbb{E}(X(1))^p (t-s)^{Hp}, \end{aligned}$$

where at the inequality we used the facts that  $0 \leq t-s < 1$  and  $|\Pi| \leq p/2$  for all  $\Pi$  not having one-element blocks; and the last equality follows by (2.11). Choosing  $p > 1/H$ , we conclude, by the Kolmogorov–Chentsov theorem, that  $\{X(t), t \geq 0\}$  has a continuous modification which is locally  $\gamma$ -Hölder continuous for every

$$0 < \gamma < \frac{Hp-1}{p} = H - \frac{1}{p}.$$

By letting  $p \rightarrow \infty$  we have finished the proof of the case (i).

- If  $0 \leq \delta < 2H$ , the proof proceeds similarly. Using (2.12) we obtain

$$\begin{aligned} \mathbb{E}|X(t) - X(s)|^p &= (t-s)^{(H-\delta/2)p} \sum_{\Pi} \left( (t-s)^{\delta|\Pi|} \prod_{B \in \Pi} \text{Cum}_{n_B}(X(1)) \right) \\ &\leq (t-s)^{(H-\delta/2)p} \sum_{\Pi} \left( (t-s)^0 \prod_{B \in \Pi} \text{Cum}_{n_B}(X(1)) \right) \\ &= \mathbb{E}(X(1))^p (t-s)^{(H-\delta/2)p}. \end{aligned}$$

For  $p > 1/(H - \delta/2)$ , the Kolmogorov–Chentsov theorem ensures that  $\{X(t), t \geq 0\}$  has a continuous modification which is locally  $\gamma$ -Hölder continuous for every

$$0 < \gamma < \frac{(H - \delta/2)p - 1}{p} = H - \frac{\delta}{2} - \frac{1}{p}.$$

Letting  $p \rightarrow \infty$  as above, we obtain the statement (ii).

- If  $\delta = 2H$  and  $H > 1/2$ , it is enough to use the second moment:

$$\mathbb{E}|X(t) - X(s)|^2 = \mathbb{E}(X(t-s))^2 = \mathbb{E}(X^{\otimes(t-s)^{2H}}(1))^2 = \mathbb{E}(X(1))^2 (t-s)^{2H},$$

to conclude that (by the Kolmogorov–Chentsov theorem)  $\{X(t), t \geq 0\}$  has a continuous modification which is locally  $\gamma$ -Hölder continuous for every

$$0 < \gamma < \frac{2H - 1}{2} = H - \frac{1}{2},$$

which is the statement (iii).

Finally, the statements  $(\widetilde{i}) - (\widetilde{iii})$  follow from part (ii) of Lemma 2.6, using also that for a  $(H, \delta)$ -dilatively stable process with stationary increments, the range of the parameter  $H$  is  $(0, 1]$  (see the Introduction).  $\square$

**2.8 Remark.** The key in the proof of Theorem 2.7 was the Kolmogorov condition, which is known to be not a necessary condition for having a continuous modification of a stochastic process, see Stoyanov [19, p. 219]. It is not a necessary condition for having a locally Hölder continuous modification either, as one can see using a modification of the counterexample of Stoyanov [19, p. 220]. Namely, if  $\{W(t), t \geq 0\}$  is a FBM with parameter  $H$ , then  $X(t) \doteq \exp(W^3(t))$ ,  $t \geq 0$ , has infinite moments, hence the Kolmogorov condition does not make sense. However, as  $\{W(t), t \geq 0\}$  has a continuous modification which is locally  $\gamma$ -Hölder continuous for every  $\gamma \in (0, H)$  (see Remark 2.11), we have for every bounded subinterval  $J \subseteq [0, \infty)$ ,

$$|X(t) - X(s)| \leq c_1 |W^3(t) - W^3(s)| \leq c_2 |W(t) - W(s)| \leq c_3 |t - s|^\gamma, \quad s, t \in J,$$

with some (random) constants  $c_1, c_2, c_3$ , hence  $\{X(t), t \geq 0\}$  is a.s. locally  $\gamma$ -Hölder continuous for every  $\gamma \in (0, H)$ . (Here one can choose universal constants  $c_i$ ,  $i = 1, 2, 3$ , i.e., which do not depend on the specific choices of  $s, t \in J$ , since  $W$  is almost surely bounded on the bounded interval  $J$ .) Therefore it is reasonable to ask how strong the assertions (ii) and (iii) of Theorem 2.7 are. By Benassi et al. [2, Proposition 3.2], if  $\gamma > H - 1/2$ , then on any interval the sample paths of a non-Gaussian FLP with parameter  $H$  are not  $\gamma$ -Hölder continuous with positive probability  $p > 0$ . Furthermore, if the Lévy measure (control measure) of the Lévy process in the defining integral of the non-Gaussian FLP is infinite, then  $p = 1$ . This example shows that part (ii) of Theorem 2.7 cannot be strengthened in general, in the sense that, as the above shows, there exists an  $(H, \delta)$ -dilatively stable process with stationary increments and zero mean for which  $0 \leq \delta < 2H$  and  $\Gamma_X = H - \delta/2$  (namely, a non-Gaussian FLP with parameter  $H$  for which  $\delta = 1$ ), i.e., the left endpoint of the interval  $[H - \delta/2, H]$  can be reached. The authors do not know whether the right endpoint of the interval  $[H - \delta/2, H]$  can be reached. We have only a partial result in case of  $\delta = 0$ . Namely, by part (ii) of Theorem 2.7 with  $\delta = 0$ , we see that if  $X$  is a  $H$ -self-similar process with stationary increments, having non-Gaussian, infinitely divisible finite-dimensional distributions, finite moments of all orders, and a Gaussian component, then it a.s. admits a local Hölder exponent  $H$ .

Next we formulate two corollaries of Theorem 2.7.

**2.9 Corollary.** For the LISOU, LISCBI and LISDLG processes with parameter  $H$  (see Example 1.3) one can apply parts (i) and (i) of Theorem 2.7 (remember,  $1/2 < H < 1$ , hence  $\delta = 2H - 2 < 0$ ), hence the local Hölder exponent  $\Gamma_X$  of these processes is a.s.  $H$ . On the other hand, the non-Gaussian FLP with parameter  $H$  (but possibly with a Gaussian component, see again Example 1.3) is a.s. locally  $\gamma$ -Hölder continuous for every  $0 < \gamma < H - 1/2$  by part (ii) of Theorem 2.7 (since we have  $1/2 < H < 1$ , hence  $\delta = 1 < 2H$ ), which is known by Benassi et al. [2, Proposition 3.2] or Marquardt [12, Theorem 4.3]. Particularly, all four processes have continuous modifications.

**2.10 Corollary.** Let  $\{X(t), t \geq 0\}$  be an  $(H, \delta)$ -dilatively stable process with stationary increments and zero mean such that either  $\delta \neq 2H$  or  $H > 1/2$ . Then  $\{X(t), t \geq 0\}$  has a continuous modification.

**2.11 Remark.** The self-similar analogue of Theorem 2.7 is a well-known and easy consequence of the Kolmogorov–Chentsov theorem. Namely, for a  $H$ -self-similar process  $\{X(t), t \geq 0\}$  with stationary increments and finite moments of all orders, we have  $E|X(t) - X(s)|^p = |t - s|^{pH} E|X(1)|^p$ ,  $p > 0$ , from which we obtain, by the same way as in the proof of Theorem 2.7, that  $\{X(t), t \geq 0\}$  has a continuous modification, the sample paths of which are locally  $\gamma$ -Hölder continuous for every  $\gamma \in (0, H)$ . For completeness, we note that the zero mean condition in Theorem 2.7 is automatically satisfied

by a  $H$ -self-similar process with  $H \neq 1$ , stationary increments and finite absolute moment, see, e.g., Vervaat [22, Auxiliary Theorem 3.1]. In the special case  $H = 1$  (and  $\delta = 0$ ) it is not sure that we have a zero mean process, but it is a degenerate case, i.e.,  $X(t) = tX(1)$ ,  $t \geq 0$ , a.s., see, e.g., Iglói [4, Theorem 2.7.1 (SS) 2)], and hence the corresponding assertions of parts (ii) and (ii) of Theorem 2.7 hold readily.

Further, assuming that  $P(X(1) = 0) = 0$ , i.e., the distribution of  $X(1)$  has no atom at zero, the proof of Lemma 2.3 in the case of  $\delta = 0$  shows that the local Hölder exponent  $\Gamma_X$  equals a.s.  $H$ .

### 3 Singularity of the distributions

In what follows  $C(I)$  will denote the set of continuous functions on a closed interval  $I \subseteq [0, \infty)$  with the local uniform topology and Borel  $\sigma$ -algebra  $\mathcal{B}(C(I))$ . If the processes  $\{X_1(t), t \in I\}$  and  $\{X_2(t), t \in I\}$  have sample paths in  $C(I)$  a.s., then we say that they are singular on  $I$  (in notation:  $X_1 \perp X_2$ ), if their distributions  $P_{X_1}$  and  $P_{X_2}$  on  $(C(I), \mathcal{B}(C(I)))$  are singular (in notation:  $P_{X_1} \perp P_{X_2}$ ), i.e., there exists a set  $A \in \mathcal{B}(C(I))$  such that  $P_{X_1}(A) = 1$  and  $P_{X_2}(A) = 0$ .

**3.1 Theorem.** *Let  $I \subseteq [0, \infty)$  be a closed interval and  $\{X_1(t), t \geq 0\}$ ,  $\{X_2(t), t \geq 0\}$  be dilatively stable processes with parameters  $(\alpha_1, \delta_1)$  and  $(\alpha_2, \delta_2)$ , respectively, both processes with zero mean, and having sample paths in  $C(I)$  a.s. Assume that one of the following two conditions holds:*

- $\inf\{t : t \in I\} = 0$  (i.e., the left endpoint of  $I$  is zero) and the above two processes satisfy Assumption 2.1.
- $\sup\{t : t \in I\} = \infty$  (i.e.,  $I$  is unbounded) and the above two processes satisfy Assumption 2.2.

*Then  $\alpha_1 \neq \alpha_2$  implies  $X_1 \perp X_2$ .*

**Proof.** Assume that the left endpoint of  $I$  is zero, and Assumption 2.1 holds for the processes  $X_1$  and  $X_2$ . Let  $(t_n)_{n \in \mathbb{N}}$  be a sequence for which the assertions of Lemma 2.3 hold, and define the following two subsets of  $C(I)$ :

$$A_i \doteq \left\{ f \in C(I) : \limsup_{n \rightarrow \infty} \frac{|f(t_n)|}{t_n^\kappa} = 0 \text{ for all } \kappa \in \mathbb{Q} \cap (0, \alpha_i) \right. \\ \left. \text{and } \limsup_{n \rightarrow \infty} \frac{|f(t_n)|}{t_n^\kappa} = \infty \text{ for all } \kappa \in \mathbb{Q} \cap (\alpha_i, \infty) \right\}, \quad i = 1, 2,$$



where  $\mathbb{Q}$  denotes the set of rational numbers. Considering the decompositions

$$A_i = \bigcap_{\kappa \in \mathbb{Q} \cap (0, \alpha_i)} \left( \bigcap_{m=1}^{\infty} \bigcup_{p=1}^{\infty} \bigcap_{r=p}^{\infty} \left\{ f \in C(I) : \frac{|f(t_r)|}{t_r^\kappa} < \frac{1}{m} \right\} \right) \\ \bigcap_{\kappa \in \mathbb{Q} \cap (\alpha_i, \infty)} \left( \bigcap_{m=1}^{\infty} \bigcap_{p=1}^{\infty} \bigcup_{r=p}^{\infty} \left\{ f \in C(I) : \frac{|f(t_r)|}{t_r^\kappa} > m \right\} \right), \quad i = 1, 2,$$

and using that for each  $\kappa \in \mathbb{Q}$  and  $n \in \mathbb{N}$ , the mapping  $C(I) \ni f \mapsto f(t_n)/t_n^\kappa$  is continuous, we get  $A_i \in \mathcal{B}(C(I))$ ,  $i = 1, 2$ . One can argue in another way, namely, by Lemma 2.3, for all  $\kappa \in \mathbb{Q} \cap (0, \alpha_i)$  [resp.  $\kappa \in \mathbb{Q} \cap (\alpha_i, \infty)$ ], we have

$$\left\{ f \in C(I) : \limsup_{n \rightarrow \infty} \frac{|f(t_n)|}{t_n^\kappa} = 0 \right\} \quad \left[ \text{resp.} \quad \left\{ f \in C(I) : \limsup_{n \rightarrow \infty} \frac{|f(t_n)|}{t_n^\kappa} = \infty \right\} \right]$$

is in  $\mathcal{B}(C(I))$ . Hence, by Lemma 2.3,  $P_{X_i}(A_i) = 1$ ,  $i = 1, 2$ . Since  $\alpha_1 \neq \alpha_2$  we have  $A_1 \cap A_2 = \emptyset$ , hence the assertion follows.

In the other case the proof is analogous, but we have to refer to Lemma 2.5 instead of Lemma 2.3.  $\square$

The next theorem is the counterpart of Theorem 3.1 for processes with stationary increments, in which case the closed interval  $I$  can be arbitrary.

**3.2 Theorem.** *Let  $I \subseteq [0, \infty)$  be a closed interval, and  $\{X_1(t), t \geq 0\}$ ,  $\{X_2(t), t \geq 0\}$  be dilatively stable processes with stationary increments and parameters  $(H_1, \delta_1)$  and  $(H_2, \delta_2)$ , respectively, both processes with zero mean and having sample paths in  $C(I)$  a.s. such that they satisfy Assumption 2.1. Then  $H_1 \neq H_2$  implies  $X_1 \perp X_2$ .*

**Proof.** One can argue similarly to the proof of Theorem 3.1. Namely, let  $t_0 \in I$ ,  $(t_n)_{n \in \mathbb{N}}$  be a sequence for which the assertions of Lemma 2.6 hold, and for  $i = 1, 2$ ,

$$A_i \doteq \left\{ f \in C(I) : \limsup_{n \rightarrow \infty} \frac{|f(t_n) - f(t_0)|}{|t_n - t_0|^\kappa} = 0 \text{ for all } \kappa \in \mathbb{Q} \cap (0, \alpha_i) \right. \\ \left. \text{and } \limsup_{n \rightarrow \infty} \frac{|f(t_n) - f(t_0)|}{|t_n - t_0|^\kappa} = \infty \text{ for all } \kappa \in \mathbb{Q} \cap (\alpha_i, \infty) \right\}.$$

Hence the assertion follows by Lemma 2.6.  $\square$

**3.3 Remark.** *If either  $\delta \neq 2H$  or  $H > 1/2$ , then in Theorem 3.2 the assumption of having a.s. continuous sample paths is automatically fulfilled. Indeed, by Corollary 2.10, apart from the exceptional case  $\delta = 2H$  and  $0 < H \leq 1/2$ , every dilatively stable process with stationary increments and zero mean has a continuous modification.*

**3.4 Example.** By Theorem 3.2, any two LISOU processes with different parameters  $H$  are singular on any closed interval  $I \subseteq [0, \infty)$ , because Assumption 2.1 is trivially fulfilled (since  $\delta = 2H - 2 < 0$ ). The same is true for the LISCBI, and particularly, for the LISDLG processes, since they have the same parameter of dilative stability as the LISOU process. It also follows that any two of these processes of three types are singular if their parameters  $H$  are different. However, for the non-Gaussian FLP we have  $\delta = 1 \geq 0$ , hence Assumption 2.1 is non-trivial, and Theorem 3.2 states that two non-Gaussian FLPs, both having Gaussian components, are singular if their parameters  $H$  are different.

Theorem 3.2 applies to the case  $\delta = 0$ , i.e., for  $(\alpha, 0)$ -dilatively stable processes too. These processes are exactly those self-similar, infinitely divisible processes, which have finite moments of all orders and non-Gaussian one-dimensional distributions (except for that at zero). But of course, there exist self-similar processes without these properties, e.g. the FBM. For this reason, Theorem 3.2 does not automatically apply to self-similar processes in general. However, let us observe that not all of the three properties above are utilized in the proof: neither the non-Gaussianity, nor the moments (or cumulants) of order higher than two are in use (recall the proofs of Lemma 2.3 and Theorem 3.2 in the case of  $\delta = 0$ ), and in case of  $\delta = 0$  we had nothing to do with infinite divisibility (since the convolution exponent  $T^\delta$  in (1.2) is 1 in this case). In fact, not even finite second moments are needed for Theorem 3.2 to apply to the self-similar case. Indeed, the only place where moments appear in case of  $\delta = 0$ , is the Markov inequality in the proof of part (i) of Lemma 2.3, where one can use, e.g., the absolute moment instead of the second order one. Let us also observe that when  $\delta = 0$ , Assumption 2.1 (i.e., the existence of a Gaussian component) is utilized only in a way that it follows that the dilatively stable process in question does not have an atom at zero. So, if we replace dilatively stable processes by self-similar ones possessing the following properties, then Theorem 3.2 remains true, and reads as follows.

**3.5 Theorem.** Let  $I \subseteq [0, \infty)$  be a closed interval, and  $\{X_1(t), t \geq 0\}$ ,  $\{X_2(t), t \geq 0\}$  be self-similar processes with stationary increments and parameters  $H_1, H_2 \in (0, 1)$ , both processes with finite absolute moment and having sample paths in  $C(I)$  a.s. such that neither the distribution of  $X_1(1)$  nor that of  $X_2(1)$  has an atom at zero. Then  $H_1 \neq H_2$  implies  $X_1 \perp X_2$ .

Note that under the conditions Theorem 3.5, the processes  $X_1$  and  $X_2$  have zero mean, see, Vervaat [22, Auxiliary Theorem 3.1].

The most important particular case of Theorem 3.5 sounds as follows.

**3.6 Corollary.** Two FBMs with different parameters  $H$  are singular.

In fact, this latter result is known, see Prakasa Rao [13], where the proof is based on a Baxter type theorem of Kurchenko [9].

**3.7 Remark.** A consequence of the above theorems on singularity (which are in fact based on Lemmas 2.3 [or 2.5] and 2.6) is that the parameter  $\alpha$ , or  $H$  in the stationary increments case, can be estimated without error, as long as we have a continuous-time sample path available (or at least its values at the time points of a sequence  $(t_n)_{n \in \mathbb{N}}$  appearing in Lemmas 2.3 [2.5] and 2.6). Note also that in Lemmas 2.3 [2.5] and 2.6 not just the existence of an appropriate sequence  $(t_n)_{n \in \mathbb{N}}$  is guaranteed, but we also give an example for such a sequence, which is important from the point of view of practical applications.

Finally, we recall a known result about variation of sample paths of self-similar processes, the generalization of which may serve as a future task.

**3.8 Remark.** If  $\{X(t), t \geq 0\}$  is a  $H$ -self-similar process with stationary increments, finite absolute moment and  $H < 1$  then the sample paths of  $X$  have no bounded variation on any (bounded) interval a.s., see Vervaat [22, Theorem 3.3]. As a possible future task one can study variation of sample paths of dilatively stable processes.

## A On the definition of dilatively stability

First we note that, at the first view, the Definition 1.2 of dilative stability is a little bit different from Definition 2.1.3 in Iglói [4], since the right continuity of the  $n$ -th order ( $n \geq 2$ ,  $n \in \mathbb{N}$ ) cumulant function  $c_n : [0, \infty) \rightarrow \mathbb{R}$ ,  $c_n(t) := \text{Cum}_n(X(t))$ ,  $t \geq 0$ , is not supposed. However, it follows since  $c_n(t) = t^{(\alpha-\delta/2)n+\delta} c_n(1)$ ,  $t \geq 0$ . Even so, this does not mean that Definition 2.1.3 in Iglói [4] contains a superfluous condition (since it defines dilative stability in a more general setting) and it turns out to be equivalent to our definition, see Iglói [4, Theorem 2.2.1].

Next we note that there is a slight redundancy in Definition 1.2 in the sense that the property that a dilatively stable process starts from 0 follows from the other properties. More precisely, if a process  $\{X(t), t \geq 0\}$  satisfies the properties listed in Definition 1.2 except that it starts from 0, then  $P(X(0) = 0) = 1$ . Indeed, by (1.2),  $X(0) \sim T^{\alpha-\delta/2} X^{\otimes T^\delta}(0)$  (where  $\sim$  denotes equality in distribution), which yields that

$$EX(0) = T^{\alpha-\delta/2} EX^{\otimes T^\delta}(0) = T^{\alpha-\delta/2} T^\delta EX(0) = T^{\alpha+\delta/2} EX(0), \quad T > 0,$$

and

$$D^2 X(0) = T^{2\alpha-\delta} D^2 X^{\otimes T^\delta}(0) = T^{2\alpha-\delta} T^\delta D^2 X(0) = T^{2\alpha} D^2 X(0), \quad T > 0.$$

If  $\delta \neq -2\alpha$ , then this implies that  $EX(0) = 0$  and  $D^2 X(0) = 0$ , yielding that  $P(X(0) = 0) = 1$ . If  $\delta = -2\alpha$ , then, by (1.2),  $X(0) \sim T^{2\alpha} X^{\otimes T^{-2\alpha}}(0)$  for all  $T > 0$ , or equivalently  $cX(0) \sim X^{\otimes c}(0)$  for all  $c > 0$ , which yields that the distribution of  $X(0)$  is strictly 1-stable. Using that a non-degenerate (strictly) 1-stable distribution

does not have a finite first moment, our assumption that  $X(0)$  has (finite) moments of all orders implies that  $\mathbb{P}(X(0) = C) = 1$  with some  $C \in \mathbb{R}$ . For completeness, we also note that there is a slight redundancy in Definition 1.1 too, in the sense that if a process  $\{X(t), t \geq 0\}$  satisfies the scaling property (1.1), then  $\mathbb{P}(X(0) = 0) = 1$ .

Finally, we call the attention that the non-Gaussianity condition in Definition 1.2, namely, that  $X(1)$  (or, equivalently,  $X(t)$ , for some  $t > 0$ ) is non-Gaussian is crucial in the sense that it is extensively used in the proofs both in the present paper and in Iglói [4]. Note also that this condition ensures that all the higher-dimensional distributions are also non-Gaussian.

## Acknowledgements

M. Barczy has been supported by the NKTH-OTKA-EU FP7 (Marie Curie action) co-funded 'MOBILITY' Grant No. OMFB-00610/2010, and by the Hungarian Scientific Research Fund under Grant No. OTKA T-079128.

## References

- [1] I. BEN-ARI, R. G. PINSKY, Absolute continuity/singularity and relative entropy properties for probability measures induced by diffusions on infinite time intervals, *Stochastic Process. Appl.* **115**(2) (2005), 179–206.
- [2] A. BENASSI, S. COHEN, J. ISTAS, On roughness indices for fractional fields, *Bernoulli* **10**(2) (2004), 357–373.
- [3] K. FLEISCHMANN, L. MYTNIK, V. WACHTEL, Optimal local Hölder index for density states of superprocesses with  $(1 + \beta)$ -branching mechanism, *Ann. Probab.* **38**(3) (2010), 1180–1220.
- [4] E. IGLÓI, *Dilative Stability*, Ph.D. Thesis, University of Debrecen, Faculty of Informatics, 2008.  
URL: <http://www.inf.unideb.hu/valseg/dolgozok/igloi/dissertation.pdf>
- [5] J. JACOD, A. N. SHIRYAEV, *Limit theorems for stochastic processes*, 2nd edition, Springer-Verlag, Berlin, 2003.
- [6] S. JAFFARD, The multifractal nature of Lévy processes, *Probab. Theory Relat. Fields* **114**(2) (1999), 207–227.
- [7] I. KAJ, Limiting fractal random processes in heavy-tailed systems. *In: Fractals in Engineering, New Trends in Theory and Applications*, 199–218. Eds. J. Levy-Lehel, E. Lutton, Springer-Verlag, London, 2005.

- [8] N. KONO, M. MAEJIMA, Hölder Continuity of Sample Paths of Some Self-Similar Stable Processes, *Tokyo J. Math.* **14**(1) (1991), 93–100.
- [9] O. O. KURCHENKO, A strongly consistent estimate of the Hurst parameter for a fractional Brownian motion, *Theor. Probab. Math. Statist.* **67** (2003), 97–106.
- [10] J. W. LAMPERTI, Semi-stable stochastic processes, *Trans. Amer. Math. Soc.* **104** (1962), 62–78.
- [11] M. MAEJIMA, A self-similar process with nowhere bounded sample paths, *Z. Wahrsch. Verw. Gebiete* **65** (1983), 115–119.
- [12] T. MARQUARDT, Fractional Lévy processes with an application to long memory moving average processes, *Bernoulli* **12** (2006), 1099–1126.
- [13] B. L. S. PRAKASA RAO, Singularity of fractional Brownian motions with different Hurst indices, *Stoch. Anal. Appl.* **26**(2) (2008), 334–337.
- [14] D. REVUZ, M. YOR, *Continuous Martingales and Brownian Motion*, 3rd edition, corrected 2nd printing, Springer-Verlag, Berlin, 2001.
- [15] G. SAMORODNITSKY, Possible sample paths of self-similar  $\alpha$ -stable processes, *Statist. Probab. Lett.* **19**(3) (1994), 233–237.
- [16] K.-I. SATO, *Lévy Processes and Infinitely Divisible Distribution*, Cambridge University Press, The Edinburgh Building, Cambridge, 1999.
- [17] A. N. SHIRYAEV, *Probability*, 2nd ed., Springer, New York, 1996.
- [18] F. W. STEUTEL, K. VAN HARN, *Infinite Divisibility of Probability Distributions on the Real Line*, Pure and Applied Mathematics, Marcel Dekker 259, New York, 2004.
- [19] J. STOYANOV, *Counterexamples in Probability*, Wiley, 1997.
- [20] K. TAKASHIMA, Sample path properties of ergodic self-similar processes, *Osaka J. Math.* **26** (1989), 159–189.
- [21] M. S. TAQQU, Fractional Brownian motion and long-range dependence. In: *Theory and applications of long-range dependence*, pp. 5–38, Birkhäuser Boston, Boston, MA, 2003.
- [22] W. VERVAAT, Sample path properties of self-similar processes with stationary increments, *Ann. Probab.* **13**(1) (1985), 1–27.

- [23] W. VERVAAT, Properties of general self-similar processes, *Bull. Int. Statist. Inst.* **52**(4) (1987), 199–216.
- [24] T. WATANABE, M. YAMAMURO, Limsup behaviors of multi-dimensional selfsimilar processes with independent increments, *Alea* **7** (2010), 79–116.
- [25] S. ZHANG, Singularity of two diffusions on  $\mathcal{C}_\infty$ , *Statist. Probab. Lett.* **19**(2) (1994), 143–145.