

Journal Pre-proof

Effect of Chitosan Based Edible Coating in Management of Post Harvest Losses in Papaya: A Comprehensive Review

Harmanjot Singh , Jasleen Kaur Bhasin , Kshirod Kumar Dash , Rafeeya Shams , Ayaz Mukarram shaikh , Kovács Béla

PII: S2772-5022(24)00068-4
DOI: <https://doi.org/10.1016/j.afres.2024.100456>
Reference: AFRES 100456



To appear in: *Applied Food Research*

Received date: 12 April 2024
Revised date: 20 July 2024
Accepted date: 22 July 2024

Please cite this article as: Harmanjot Singh , Jasleen Kaur Bhasin , Kshirod Kumar Dash , Rafeeya Shams , Ayaz Mukarram shaikh , Kovács Béla , Effect of Chitosan Based Edible Coating in Management of Post Harvest Losses in Papaya: A Comprehensive Review, *Applied Food Research* (2024), doi: <https://doi.org/10.1016/j.afres.2024.100456>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2024 Published by Elsevier B.V.
This is an open access article under the CC BY-NC-ND license
(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Highlights

- Chitosan is a natural biopolymer formed from chitin found in shrimp and crab shells.
- Chitosan coating is biodegradable and eco-friendly.
- Chitosan is a natural preservative as well as a clarifying agent.
- Chitosan coating increases shelf life, and maintains freshness of papaya.
- Edible coating decrease oxidation and degeneration of papaya by restricting oxygen.

Journal Pre-proof

Effect of Chitosan Based Edible Coating in Management of Post Harvest Losses in Papaya: A Comprehensive Review

Harmanjot Singh^{1#}, Jasleen Kaur Bhasin^{1#}, Kshirod Kumar Dash^{2*}, Rafeeya Shams¹, Ayaz Mukarram shaikh³, Kovács Béla^{3*}

¹Department of Food Technology and Nutrition, Lovely Professional University, Phagwara, Punjab, India.

²Department of Food Processing Technology, Ghani Khan Choudhury Institute of Engineering and Technology, Malda, West Bengal, India

³Faculty of Agriculture, Food Science and Environmental Management Institute of Food Science, University of Debrecen, Debrecen 4032, Hungary

#First author and second author have equal contributions.

Corresponding Author:

Kshirod Kumar Dash email: kshirod@tezu.ernet.in

Kovacs Béla email: kovacsb@agr.unideb.hu

Abstract

Chitosan is a biopolymer abundantly present in the exoskeletons of crustaceans like shrimp and crabs. Chitosan is derived from chitin by a process called deacetylation, which includes demineralization, deproteinization, and deacetylation steps. Its many attributes, including biodegradability, biocompatibility, and antibacterial action, have made it extensively useful in food processing sector. Chitosan, due to its different reactive groups (primary -OH, secondary -OH, and -NH₂), has been discovered to impede the growth of wide range of fungi and both gram-positive and gram-negative bacteria. This is achieved through electrostatic interactions with the cell wall, cell membrane, and cytoplasmic components of the microorganisms. Papaya is a significant tropical fruit belonging to the genus *Carica*. Post-harvest losses in papaya are high, owing to factors such as physical damage, microbial proliferation, and moisture depletion. Utilizing chitosan coating, either alone or in combination with other preservative components, has been discovered to effectively mitigate these variables, resulting in decreased post-harvest losses and enhanced papaya shelflife. The review analyzes the impact of chitosan coating on the antibacterial characteristics, quality preservation, and

decrease of post-harvest losses in papaya. Applying a chitosan coating on papaya has great potential in enhancing the sustainability of papaya production and supply.

Keywords: Chitosan; edible coating; papaya; antimicrobial properties; post harvest loss

1. Introduction

Papaya is the fruit belonging to the sole species in the genus *Carica* of the family *Caricaceae*, which is short-lived and perennial in nature and is typically propagated from seeds (Devaki *et al.*, 2015). It is a climacteric and tropical fruit which is known for its rich nutritional value. It may also be described as a berry-like fruit or as the “fruit of the common man” (Prajapati *et al.*, 2017). The shape of the fruit is almost round to oval in shape, and somewhat club-shaped, which is dimensionally 15–50 cm in length, 10–20 cm in width, and weighs around 9–10 kg (Devaki *et al.*, 2015). Today, it is widely dispersed throughout the tropical, spreading to the subtropical zone of 32° north and south latitude, which includes more than 60 countries worldwide, including India, Indonesia, Brazil, Mexico, Nigeria, Dominican Republic, United States, and Australia (Rodrigues *et al.*, 2021; Anitha *et al.*, 2018). Furthermore, tropical and subtropical regions of the world now harvest and consume the most fruits from this plant (Santos *et al.*, 2014; Vij & Prashar, 2015).

The flesh of papaya is delicious and comes in a variety of colours and textures, including yellow, orange, and red. Ideally, papaya flesh contains around 0.6% protein, 10-13% sugar, and 85–90% water (Prasad & Paul, 2021). It contains glucose, fructose, and sucrose, among other sugars and has comparatively low calories (32 Kcal/100g ripe fruits), making it an optimal and favoured fruit for weight reduction (Krishna *et al.*, 2008). Additionally, it contains abundant amount of vitamins such as vitamins A and C, riboflavin, folate, thiamine, pantothenic acid, and niacin and minerals such as calcium, iron, potassium, copper and fibres as well (Prajapati *et al.*, 2017). Papaya is a natural source of papain, a digestive enzyme that is a common industrial ingredient in brewing, meat tenderisation, medicines, cosmetics, and beauty products (Devaki *et al.*, 2015). It is well known for its high pectin content and can be consumed either raw or in processed forms such as jelly, candy, jam, and pickles (Prasad & Paul, 2021; Tan *et al.*, 2020). Numerous components of the papaya plant, including the roots, leaves, peels, latex, flowers, fruits, and seeds, have nutritional and therapeutic value (Ali *et al.*, 2012). Papaya leaf and seed extract contains a variety of phytochemicals such as flavonoids, phytosterols, carotenoids, alkaloids,

phenolic compounds, and cyanogenic compounds (benzyl glucosinolate), which have been reported to provide ameliorative effects for medical conditions such as diabetes mellitus, hepatic and renal problems, fertility, hyperglycemia, amoebic dysentery, and antitumor activities (Tan *et al.*, 2020; Archampong *et al.*, 2019; Olcum *et al.*, 2020; Doan *et al.*, 2020; Singh *et al.*, 2019; Odhong *et al.*, 2014).

The mature papaya fruit softens quickly, gets readily infected by diseases, and is vulnerable to additional post-harvest losses, such as injury from chilling due to exposure at low temperatures (Mahmud *et al.*, 2008). Papaya is susceptible to numerous ailments brought on by fungi, bacteria, nematodes, and viruses, which significantly reduces its productivity. Papaya anthracnose, which is caused by *Colletotrichum gloeosporioides*, is known to be the most serious of all diseases, causing significant losses to papaya fruits during transportation and storage (Prajapati *et al.*, 2017). Due to this reason, it has a very short post-harvest shelf-life of about 7–14 days under greenhouse conditions (Marpudi *et al.*, 2011). The percentage of wastage is greatly increased in developing countries, where the correct technologies for the storage of fruits and vegetables are lacking (FAO, 2011). The susceptibility of fresh produce to post-harvest diseases and deterioration of quality attributes increases after harvest and during prolonged storage as a result of physiological and biochemical changes in the commodities. These changes can favour the development of post-harvest pathogens and the incidence of post-harvest diseases, which are the major causes of losses through the supply chain. Therefore, the food sector places its primary focus on appropriate preservation techniques. Inadequate and costly solutions for food preservation have led scientists to create natural preservatives which are safe, effective, and acceptable (Huq *et al.*, 2015). Keeping in mind the relatively long time storage and transportation, the use of biologically derived preservatives with compliance with health and safety regulations can bring a great solution for the preservation of the fruits (Romanazzi *et al.*, 2018).

One such strategy that has been of recent interest is the use of edible coatings in the fresh fruit industry to reduce the deleterious effects that could take place on intact fruit tissues, which are usually subject to minimal processing. It is a novel technique for food preservation that creates physical barriers on the surface of fruits and vegetables, preventing moisture and solute movement and slowing down respiration, gas exchange, and oxidative reaction rates in order to increase the shelf-life (Arnon *et al.*, 2014). Biopolymer coating materials are designed to transport active substances such as antibrowning agents, colourants, flavourings, nutrients, spices, and antimicrobial chemicals in order to increase product shelf life and lower the danger of

pathogen growth on food surfaces (Pranoto *et al.*, 2005). As a technology, the application of edible coatings with antifungal qualities has become popular for preserving fruits and vegetables against fungi that cause post-harvest deterioration. Studies from a variety of fields have shown that edible coatings have an antibacterial effect against *Botrytis cinerea*, *Colletotrichum* spp., *Penicillium* spp., and *Alternaria* spp (Wardana *et al.*, 2021; Landi *et al.*, 2021; Rajestary *et al.*, 2021). Coatings can change the composition of the atmosphere surrounding the fruit, which results in creating a barrier to gas exchange, such as oxygen, carbon dioxide, and ethylene, which are involved in the respiration process. Different edible coatings have been reported to preserve the nutritional value of fruits similar to or even better than conventional packaging.

Application of chitosan treatment at the pre-harvest or post-harvest stages has been suggested to be an appropriate replacement for the usage of synthetic fungicides. By maintaining the general quality of the various fresh commodities can aid in preventing post-harvest fruit diseases and extending storage life (Bautista-Baos *et al.*, 2006). Chitosan (poly b-(1-4) N acetyl-d-glucosamine) has been identified as providing an ideal coating with antimicrobial properties that can induce plant defence responses when applied to vegetal tissues (Devlieghere *et al.*, 2004). On the other hand, chitosan coating also offers a substrate for the incorporation of additional useful natural food additives, which may enhance its antibacterial capabilities and stop the quality of the fruit from degrading (Vargas *et al.*, 2008). The United States Food and Drug Administration (USFDA) has certified chitosan as a “Generally Recognised as Safe” (GRAS) food additive; therefore, chitosan treatment in the fresh produce industry is safe for consumers and the environment (USFDA, 2013). The review analyzes the characteristics and preservation impact of chitosan-based edible coating in reducing post-harvest losses in papaya. The effects of applying chitosan coating on the antibacterial properties, and quality maintenance in papaya. The use of a chitosan coating on papaya has significant potential to improve the sustainability of papaya production and supply.

2. Production Rate of Papaya and Factors Influencing its Production

Papaya is currently grown in more than 37 nations throughout the world, with India, Brazil, Mexico, Nigeria, Indonesia, Ethiopia, Thailand, Peru, Columbia, Guatemala, and the Philippines being the top producers (Tridge.com, 2021). These countries together accounted for approximately 68% of papaya production worldwide (FAOSTAT, 2021). India is the world’s top papaya-growing nation, having an annual production of 5989,000 metric tons (MT) from an area covered by 138,000 hectares and a productivity of 43.40 MT/h (Anonymous, 2018). It is grown

in practically all tropical and subtropical areas of India, and the top producing states are Andhra Pradesh, Gujarat, Karnataka, Madhya Pradesh, and Tamil Nadu (APEDA, 2021; Prasad & Paul, 2021). Papaya is exported in the form of dried or fresh produce to various countries such as the Netherlands, Nepal, Germany, UAE, Saudi Arabia, Qatar, Kuwait, and other nations, with a total annual export profit of around 4150 lakh Indian rupees from 9996 MT of papaya export (Anonymous, 2018). The top exporting countries of papaya include Mexico, which has been reported to be the leading exporter of papaya (33.4%) followed by Brazil (15.4%), Guatemala (7.6%), United States (7.6%), and the Netherlands (7.2%) while the United States continues to be the top importer of papaya with 37.2% of the total import value, followed by Germany (10.1%), Portugal (7.1%), Canada (6.9%), and other nations (Tridge.com, 2021).

Papaya, being a tropical plant, is highly vulnerable to frost, and its cultivation is confined to regions between 32° N and 32° S of the Equator, where the optimum temperature range for growing papayas lies between 25°C and 30°C (Saran *et al.*, 2019). Exposing papaya plants to flooding for 48 hours proves fatal, while even brief exposure to 0°C can cause damage and prolonged cold spells without overhead sprinkling can result in the death of the plants (DAIS, 2009). Although mature papaya trees can endure temperatures as low as -2°C, they are usually grown in areas where the average daily minimum temperature during mid-winter never falls below 5°.

Strong, cool, hot, and dry winds are not preferred and can damage papaya plants in open and high-lying areas (Saran *et al.*, 2016). To ensure successful plantation establishment, a shelterbelt is utilized with full sunshine, along with employing techniques like staking, earthing up for trunk support, and windbreaks to mitigate wind damage (Saran *et al.*, 2015). Papayas thrive across a diverse range of soils, often developing a robust taproot shortly after planting. Optimal growth occurs in well-drained sandy-loam soil enriched with sufficient organic matter, which is crucial for successful cultivation (Saran *et al.*, 2019). In regions with high rainfall and inadequate drainage, prolonged saturation can lead to plant demise, necessitating the use of raised beds and drainage ditches (Saran *et al.*, 2015). Mulching and incorporating organic material, such as planting a cover crop like legumes, serve as valuable sources of organic matter and is implemented approximately six months before transplanting (DAIS, 2009).

Maintaining soil pH between 6.0 and 6.5 is ideal for papaya growth, and addressing deficiencies in acidic soils below pH 5.5 is achieved with calcium carbonate or lime, while sulphur is effective for correcting deficiencies in alkaline soils above pH 8.0 (DaMatta *et al.*, 2007; Medina *et al.*, 2003). To optimize papaya orchard growth, a weed-free environment is essentially

maintained which is achieved through various techniques like herbicide application, hand weeding, mulching, and deep hoeing, either individually or in combination (DaMatta *et al.*, 2007). Inorganic materials such as plastic mulches are utilized for weed control, with plastic mulch combined with mounding identified as the most effective treatment, also aiding in preventing root rot disease (Saran *et al.*, 2015).

Earthing up at the 4th, 6th, and 8th months post-transplanting before the monsoon season helps prevent overcrowding and waterlogging and this practice is conducted within a 30 cm radius around the plants, which promotes their upright growth in sandy-loam soils (Saran *et al.*, 2014). Staking, using bamboo or other sticks, is implemented during the fruit development stage to protect the plant from wind damage and manage the fruit load, especially following heavy rainfall (Saran *et al.*, 2014).

Thinning involves the removal of excess fruits, and it is done to prevent damage caused by overcrowding and reduces the risk of pressure injury and competition for space and nutrients, particularly in dwarf cultivars like Pusa Dwarf and Pusa Nanha (Saran *et al.*, 2014; Saran *et al.*, 2015). Upon the onset of flowering, retaining about 5–10% of male plants in the orchards ensures adequate pollination, considering that hermaphrodite plants bear complete flowers (Saran *et al.*, 2014). Employing cheesecloth, glassine paper, or wax paper bags for bagging can effectively control fruit fly infestations in small plantings (Saran *et al.*, 2019). Cold weather conditions may disrupt pollination and lead to the shedding of unfertilized female flowers. To ensure a healthy harvest, side shoots are removed from the stem during the initial crop and prune old, dry, diseased leaves, and petioles, as these are standard practices (Saran *et al.*, 2014). Pollarding, a technique involving the reduction of plant height, is done for one-year-old ratoon crops to mitigate vulnerability to wind damage and improve yield and productivity (Prakash *et al.*, 2014).

3. Factors causing post harvest losses in papaya

Even though papaya has an immense production rate with the highest productivity among all the fruits grown in India, around 40-60% of the total production at various papaya growing regions is wasted due to the perishable nature of papaya, leading to deterioration in quality and quantity of the product which affects the marketing as well as the income of the farmers (Prasad & Paul, 2021). As shown in Fig. (3), Post-harvest loss in papaya can be attributed to various factors, including pre-harvest factors, environmental conditions, inadequate marketing channels,

improper storage and transportation, post-harvest pests and diseases, chilling injury, physiological disorders, and senescence (Prasad & Paul, 2021; Prajapati *et al.*, 2017; Paull *et al.*, 1997). These factors can lead to significant changes in the fruit's quality and nutritional value, as well as its appearance, texture, firmness, and flavour, which can further negatively impact the fruit's marketability and reduce its economic value (Mashau *et al.*, 2012).

In a study conducted by Gajanana *et al.* (2010), papaya variety Taiwan 786, which is grown in the major papaya producing states of Andhra Pradesh and marketed in Bangalore, was studied to estimate the rate of post harvest losses, causes of these losses at different stages of handling and to study its impact on marketing efficiency. The results showed that the total post-harvest losses were calculated to be roughly 25.49%, with field losses of 1.66%, transit losses of 4.12%, ripening losses of 8.22% at the market level, and retail losses of 11.49%. It was observed that losses in the fields were primarily brought on by immature and small-sized fruits, malformations, and harvesting damage, while bruises and pressing injuries led to transit loss at the market level. The main reasons for loss during ripening were found to be anthracnose and fruit rot caused by *alternaria* and *phytophthora*, which are two of the main post-harvest diseases which can lead to loss of quality, deeming them unfit for sale and consumption (Esguerra *et al.*, 2020).

A similar study was conducted by Dadi *et al.*, (2020) to assess the post-harvest losses of papaya at the wholesaler and retailer levels in Jimma Town, South Western Ethiopia. The results of the study revealed that at wholesale level, the overall loss was 21.75%, which was further classified as 12.5% during transportation and 9.25% during storage, while retailers experienced a loss of approximately 15.6% and all of these losses were attributed to factors such as inappropriate transportation, inadequate storage conditions, and a lack of suitable selling places, which often leads to fruit softening, rotting, wounding, and compactness. The study concluded that a total of 37.35% of papaya fruits were lost across the two marketing channels in Jimma town. These results were in accordance with another study conducted by Paull *et al.* (1997) for examining the marketability and shelf life of papaya fruit delivered at New York terminal markets by Hawaii's packer-shippers, and they observed that the papayas were subjected to a range of disorders, including mechanical injury, over-ripeness, and parasitic diseases. Additionally, the most common causes of loss included anthracnose rot, affecting 62% of shipments; bruise damage, affecting 22%, and overripe fruit, affecting 48%. Other causes of loss included soft fruit, chilling injury, and other diseases, affecting a total of 35% of shipments inspected. According to Souza *et al.* (2014), papaya fruit exhibits high ethylene production when harvested at a physiologically ripe stage, owing to its increased respiratory activity, and the duration of this process may vary

depending on the genotype and harvest stage. The consequent increase in ethylene production triggers alterations in the fruit's skin and pulp colour, flavour, and pulp firmness, as well as the production of volatile aromatic compounds during ripening (Galo *et al.*, 2014). Due to their high sugar content, soft pulp, high water activity, low acidity, and rapid ripening process, papaya fruits are conducive to the establishment of pathogen-caused infections (Gonzalez-Aguilar *et al.*, 2009; Tabassum and Khan, 2020).

4. Physiochemical composition of chitosan

As indicated in Table 1, the moisture content of chitosan, which is extracted from different sources, ranges from 0.0004 to 9.23%. Commercial chitosan products consist of moisture, which is less than 10% (Li *et al.*, 1992). The variance in moisture content may be due to different animal species and methods used. Due to the hygroscopic nature of chitosan, the moisture content can be affected by absorption during the storage period as it has a great capacity to form hydrogen bonds with water through both its hydroxyl and amino groups (Ssekatawa *et al.*, 2021; Khan *et al.*, 2002). The ash content of chitosan is found to be between 0.03 and 11.77%, and this mainly depends on the starting material and composition. According to Kumari *et al.*, 2017, it was reported that chitosan extracted from crabs had more ash content and protein as compared to chitosan extracted from fish and shrimp sources. The ash content of a high quality grade chitosan should be less than 1%, therefore, it acts as an indicator for checking the effectiveness of the demineralisation process of chitosan for the removal of calcium carbonate (Hossain & Iqbal, 2014). Chitosan containing high amounts of ash content may also affect its solubility, consequently contributing to lower viscosity and average molecular weight, and it can also affect other more important characteristics of the final product (Kumari *et al.*, 2017). The protein content ranges from 1.99 to 48.55%, and this percentage in chitosan can differ based on the source of chitin, the precursor material of chitosan and may also indicate the effectiveness of the deproteinisation process during the chitosan production (Kumari *et al.*, 2017; Hossain & Iqbal, 2014). The fibre content in chitosan ranges from 8.74 to 55.8%, and this difference may be due to variation in the degree of deacetylation process applied for the production of chitosan from chitin where the acetyl groups in chitin are removed using an alkaline solution resulting in a reduction in the fibre content of the material, as well as changes in other physicochemical and functional properties (Miraftab *et al.*, 2011).

The C content, N content, and C/N ratio are important parameters in the characterisation of chitosan, as they provide information about the chemical composition and structure of the

material. The C content in chitosan indicates the level of carbon in the material, which is a key component of the glucosamine units that make up chitosan (Islam *et al.*, 2022). According to Ibitoye *et al.* (2018), the C content in chitosan was found to be around 40.97% in commercial shrimp and 38.98% in house crickets. Similarly, Kumari *et al.* (2017) reported the C content in chitosan synthesised from fish scales, crab, and shrimp shells to be in the range of 34.89 - 48.60%. The N content, on the other hand, refers to the level of nitrogen present in the material, which indicates the level of purity in chitosan and also points out the effectiveness of the deproteinisation process (Ibitoye *et al.*, 2018). A lower amount of nitrogen signifies the minimum residual protein remaining in chitosan (Majtán *et al.* 2007, Ivshina *et al.* 2009). Ssekatawa *et al.* (2021) reported the N content in chitosan extracted from Nile perch scales to be 7.0%, while the N content in chitosan extracted from banana weevils was found to be 6.9%. Ibitoye *et al.* (2018) reported the N content in chitosan extracted from house crickets to be 5.932%, while the N content in chitosan extracted from commercial shrimps was found to be 6.182%. The C/N ratio in chitosan provides information about the degree of deacetylation, which is an important parameter that affects various properties of chitosan (Isa *et al.*, 2012). A low C/N ratio indicates the presence of impurities in the derived chitosan (Yen *et al.*, 2009). According to Kumari *et al.* (2017), the C/N ratio of chitosan derived from fish scales, shrimp and crab was found to be 7.62%, 7.79% and 6.20%, respectively. Similarly, Ibitoye *et al.* (2018) reported that the C/N ratio in chitosan extracted from house cricket was found to be 6.571%, while the C/N ratio in chitosan extracted from commercial shrimps was found to be 6.63%. Ssekatawa *et al.* (2021) also reported the C/N ratio in chitosan extracted from Ugandan edible mushrooms to be in the range of 5.92-6.80%.

One of the most significant physicochemical features of chitosan is its molecular weight, which influences a variety of physicochemical and biological behaviours, including mucoadhesion, hydrophilicity, viscosity, moisture absorption, biodegradability, and antibacterial activity (Aranaz *et al.*, 2009). According to Struszczyk *et al.* (2002), the average molecular weight of commercial chitosan ranges from 100 kDa to 1200 kDa. This is in accordance with the results obtained by Renuka *et al.* (2019), who reported that the average molecular weight of chitosan determined by using intrinsic viscosity was 110.64 kDa. Similarly, Ssekatawa *et al.* (2021) found that the average molecular weight ranged from 291 kDa for Nile perch scales chitosan to 348 kDa for mushroom chitosan. The process of deacetylation may lead to the lower molecular weight of chitosan (Szymańskav & Winnicka, 2015), which has a greater ability to penetrate bacterial cell walls compared to high molecular weight chitosan. As a result, it can effectively

infiltrate the cell and interact with the DNA, leading to the inhibition of the transcription process and subsequently disrupting protein synthesis. Thus, low molecular weight chitosan possesses potent antimicrobial activity. (Yilmaz, 2020; Sudarshan *et al.*, 1992). In contrast, high molecular weight chitosan can exhibit antibacterial activity at higher concentrations by binding to negatively charged bacterial cell wall components through electrostatic interactions, thereby creating an impermeable coating around the cell, which restricts the movement of materials into and out of the cell. On the other hand, chitosan with a moderate molecular weight has been demonstrated to possess superior anti-cholesterol activity when compared to chitosan with a higher molecular weight (Kara & Stevens, 2002).

The solubility of chitosan ranges from 48.30 to 98.40% and is a crucial parameter for assessing its quality. The degree of deacetylation, which directly impacts the solubility of chitosan, is a key determinant of its overall quality (Samar *et al.*, 2015). Kumari *et al.* (2017) reported that the solubility of fish chitosan was higher (78%) than crab (60%) and shrimp chitosan (70%). On the other hand, Hossain & Iqbal (2014) discovered that the solubility of chitosan ranged from 48.3 to 97.65%, and this variation was due to the influence of various critical factors such as temperature and deacetylation reaction time, alkali concentration, the ratio of chitin to alkali solution, as well as particle size. The solubility of chitosan is dependent upon the removal of the acetyl group during the deacetylation process, and a lower degree of deacetylation may have an unfavourable impact on the outcomes (No *et al.*, 2003).

The fat binding capacity (FBC) and water binding capacity (WBC) of chitosan are important functional properties that can affect its potential applications. Renuka *et al.* (2019) reported that chitosan extracted from *Parapeneopsis stylifera* shrimp shells had a FBC of 331.28% and a WBC of 637.33%. The authors attributed these properties to the degree of deacetylation of chitosan and its molecular weight. Similarly, Kumari *et al.* (2017) synthesised chitosan from fish scales, crab, and shrimp shells and found that the FBC and WBC were influenced by the source of chitosan. Shrimp shell chitosan had the highest FBC and WBC values among the three sources tested. Hossain and Iqbal (2014) observed that the FBC and WBC of chitosan were found to be 427.98% and 537.29%. The fat binding capacity (FBC) of chitosan is affected by factors such as the crystallinity and presence of salt-forming groups and residual protein. Additionally, the FBC may be influenced by the viscosity of chitosan, with lower viscosity leading to lower FBC (Rout, 2001; Knorr, 1982). These findings have been reported in previous studies, with Rout (2001) highlighting that FBC and WBC were influenced by reversing the sequence of steps such as demineralisation and deproteinisation.

5. Method of Extraction of Chitosan

A large number of edible coatings are available, and among them, chitosan is the most common. Chitosan (β - (1,4)-2-amino-2-deoxy-D-glucose) is a natural biopolymer obtained by deacetylation of chitin, which is the second most important polysaccharide in nature after cellulose and is present in the exoskeleton structure of marine invertebrates, insects, as well as fungi, algae, and yeast (Neto *et al.*, 2022; Gao *et al.*, 2022).

The traditional process of extracting chitosan involves three main stages, namely demineralisation, deproteinisation, and deacetylation (Knidri *et al.*, 2018). The extraction of Chitosan from Crustaceans is presented in Fig. (1). Demineralisation is carried out in dilute hydrochloric acid to remove calcium carbonate and calcium chloride, which are the primary inorganic components of crustacean exoskeletons. The resulting material is then washed, filtered, and dried overnight. The release of carbon dioxide (CO₂) during the demineralisation process serves as a reliable indicator of the amount of mineral content and the level of interaction between the acid and mineral matter, which is influenced by the type of species being utilised (Kumari *et al.*, 2015). Deproteinisation involves treating the material with dilute sodium hydroxide to remove proteins, and complete deproteinisation can be confirmed by the absence of any colour in the solution when left undisturbed overnight (Kumari *et al.*, 2015). This is followed by washing with distilled water to neutralise chitin and drying it in the oven at around 110°C. The resulting purified chitin can be used as an excellent source of animal feed (Benhabiles *et al.*, 2013). Deacetylation involves the removal of the acetyl group to convert chitin to chitosan, which is achieved by treating the material with concentrated sodium or potassium hydroxide at high temperatures. From a chemical perspective, both acids and alkalis are viable options for deacetylating chitin, however, due to the high susceptibility of glycosidic bonds to acid, alkali deacetylation is more commonly employed (Hajji *et al.*, 2014). Chitosan can be extracted from chitin using either heterogeneous or homogeneous approach. In the heterogeneous approach, chitin is typically subjected to treatment with a concentrated hot solution of NaOH for several hours. As a result, chitosan is produced as an insoluble residue with deacetylation levels ranging from approximately 85% to 99%. On the other hand, the homogeneous method involves preparing alkali chitin by dispersing chitin in concentrated NaOH solution at room temperature for at least three hours, followed by dissolution in crushed ice at around 0°C. This method yields soluble chitosan with an average degree of acetylation ranging from 48% to 55% (Hajji *et al.*, 2014). According to Kurita, (2006), the deacetylation process of chitin can be significantly

expedited by immersing it in a concentrated solution of sodium hydroxide at room temperature before applying heat. Following the reaction, the resulting material is subjected to multiple washes with distilled water until it reaches a neutral pH level, and then it is dried overnight in an oven. Several factors involved in the deacetylation process can influence the characteristics of the resulting chitosan (Li *et al.*, 1992). Rege & Block, (1999) investigated the impact of temperature, processing duration, and mechanical shear on chitosan properties, highlighting that temperature and processing duration notably affect the degree of deacetylation (DA) and molecular weight (MW). Another study conducted by Tolaimate *et al.*, (2000) emphasized the significant influence of temperature and the repetition of alkaline steps on chitosan DA. These studies, conducted using a traditional one-variable-at-a-time approach, collectively suggest that chitosan MW and DA are primarily influenced by NaOH concentration, reaction time, temperature, and the repetition of alkaline steps.

Even though the chemical extraction method of chitosan is the most widely used method available due to its low cost and suitability to mass production, this extraction method has several drawbacks, including, contamination of wastewater with chemical residues, and increased purification costs (Younes & Rinaudo, 2015). The growing focus on environmentally friendly extraction techniques, guided by principles of green chemistry, has led to increased interest in using enzymes and microorganisms for chitosan extraction. Enzymatic extraction methods offer advantages such as high reproducibility, shorter processing times, simpler handling, reduced solvent usage, and lower energy requirements. Chitin deacetylase is an enzyme extracted from various fungi (Araki & Ito, 1975; Martinou *et al.*, 1993; Gao *et al.*, 1995; Alfonso *et al.*, 1995) and insect species (Sundara *et al.*, 1982), and this enzyme facilitates the breakdown of N-acetamido bonds within chitin, resulting in the formation of chitosan. However, in a study conducted by Martinou *et al.*, (1995), the efficacy of chitin deacetylase isolated from the fungus *M. rouxii*, in chitosan production was evaluated by using chitin as a substrate in both its crystalline and amorphous states, and it was reported that the deacetylation levels remained notably low (<10%), suggesting that the enzyme exhibits limited effectiveness on insoluble chitins. Therefore, it's worth noting that enzymatic extraction methods are currently limited to laboratory-scale studies.

6. Properties and importance of chitosan as an edible coating

The major property of chitosan is dictated by the presence of three different functional groups (primary —OH, secondary —OH and —NH₂) and its water solubility in acidic pH. Due to the

presence of reactive groups, it inhibits the growth of a wide variety of bacteria and fungi (Hosseinnejad & Jafari, 2016). Chitosan has a wide range of uses and can be used to create different formulations (Fatehi *et al.*, 2010). To increase the security, functionality, and quality of fruits and vegetables, chitosan-based edible coatings can also be utilised as transporters of food ingredients such as antimicrobials, texture enhancers, and nutraceuticals. It is one of the most used edible coatings due to its biocompatibility, biodegradability, and bioactivity since it is a powerful material that can be applied in human medicine, cosmetics, and agriculture. Chitosan can induce host defences, exhibit antibacterial activity towards fungi that cause decay, and create a semi-permeable layer on a treated surface when applied to fruits and vegetables (Romanazzi *et al.*, 2018). This edible coating has been used extensively to preserve fresh fruits and vegetables after harvest.

In recent years, there has been a rise in the scientific literature discussing chitosan-based edible coatings. This can be explained by the significance of chitosan in plant protection as a natural fungicide and plant-defence enhancer, as well as by the fact that it is used all over the world to extend the shelf life of a variety of fruits and vegetables. By changing the permeability of O₂ intake and CO₂ production, this biopolymer can create a semipermeable layer on the surfaces of fruits and vegetables, which lowers respiration rates and boosts antioxidant activity (Shah & Hashmi, 2020). Chitosan has demonstrated inhibitory effects on a variety of post-harvest fungal infections and exhibits broad-spectrum antibacterial activities (Rajestary *et al.*, 2021). In a study by Novita *et al.* (2012), it was shown that chitosan has excellent fungicidal qualities and can be used to extend the shelf life of tomatoes. It can operate as an exogenous elicitor inducing activity of many defence-related enzymes in papaya fruit, as demonstrated by Landi *et al.* (2021). Additionally, Hamdayanty *et al.*, (2012) discovered that chitosan coating with a concentration of 0.75% was efficient in reducing disease infestation and the severity of damage, as well as inhibiting fruit ripening in papayas over a period of six days.

7. Challenges associated with the use of chitosan in food preservation

Even though chitosan can be extensively utilized for preserving various food products due to its distinctive properties, its application in the food sector is significantly restricted by its poor solubility and inadequate mechanical strength (Yadav *et al.*, 2023). Nonetheless, solubility can be enhanced through various modifications, resulting in new derivatives with better physicochemical properties and a broader range of applications.

Another challenge associated with chitosan application in food preservation was studied by (Hu & Gänzle, 2018), where the authors observed that the effectiveness of chitosan in reducing microbial cell counts is limited to a 2.5 log (CFU per g) decrease, regardless of the food matrix or the method of application. This statement is supported by the findings of Leleu et al. (2011), who demonstrated that coating eggs with a 2% chitosan solution did not prove lethal to *Salmonella Enteritidis*, however, when the chitosan solution was applied to eggshells and dried before bacterial inoculation, it provided a protective barrier that reduced *Salmonella* penetration. Similar results were obtained by Moradi *et al.*, (2011) where the authors reported that chitosan films were not bactericidal but inhibited the growth of *L. monocytogenes* on ready-to-eat sausage slices. However, in contrast to these studies, a study conducted by Anacarso *et al.*, (2011) reported a reduction of over 5 log (CFU per g) of *Listeria monocytogenes* on apples and grapes treated with a 2% (w/v) chitosan solution. This significant antilisterial effect is likely due to the smooth surfaces of the fruits, which create a high local concentration of chitosan and promote strong interactions between the bacterial cells and chitosan (Hu & Gänzle, 2018). According to Hu & Gänzle, (2018), food components such as NaCl, proteins, and starch can negatively impact chitosan's effectiveness by neutralizing its positive charge, therefore, chitosan's ability to inactivate pathogens in food is generally restricted to a reduction of 1–2 log (CFU per g), posing a significant challenge for its use as a broad-spectrum food preservative.

8. Mechanism of Preservative Action of Chitosan against Bacteria

Chitosan is a cationic biopolymer with antibacterial characteristics that are influenced by various factors such as pH, concentration, molecular weight, degree of polymerisation, and cross-linking (Elsabee & Abdou, 2013). Chitosan solution is highly stable over a long period of time; however, its stability in neutral pH is highly important for exhibiting antimicrobial activity against a wide variety of foodborne pathogens (Alishahi & Aïder, 2012). Electrostatic interactions with the bacterial cell wall, cell membrane, and cytoplasmic components are the main modes of action for antibacterial activity.

Chitosan has been found to be effective against both gram-positive and gram-negative bacteria. Gram-negative bacteria like *Escherichia coli* have an asymmetric lipid-protein bilayer (lipopolysaccharide, LPS) that makes up their outer membrane (OM). The divalent cations (Ca^{2+} and Mg^{2+}) in the outer membrane have a crucial role in maintaining the stability of the LPS molecules' core anionic charges (Khan *et al.*, 2015). It can be hypothesised that chitosan replaces the divalent cations from their binding sites and reduces the interaction between the LPS

molecules, causing membrane disruption and cell lysis due to penetration (through electrostatic interaction) of positively charged chitosan through the cell membrane of gram-negative bacteria. Unlike gram-negative bacteria, the gram-positive bacteria do not have an outer membrane. Hence, chitosan, as a polycationic long-chain molecule, can adhere better with gram-positive bacterial members such as *Staphylococcus aureus*. Due to this reason, gram-positive bacteria are more effectively inhibited by chitosan than gram-negative bacteria. According to the literature, gram-positive bacteria contain poly-anionic surface polymers such as teichoic acid and lipoteichoic acid, which interact with internal molecules to impair critical bacterial functions and activities (Aranda-Martinez *et al.*, 2016). Raafat *et al.*, (2008) reported simultaneous permeation of the cell membrane to small cellular components, coupled to a significant membrane depolarisation. No concomitant cell wall biosynthesis was observed. Later, they analysed multiple changes in the expression profile of *S. aureus* SG11 genes, which are involved with the regulation of stress and autolysis and the genes involved with energy metabolism and postulated a possible mechanism for chitosan's activity. According to Chung & Chen (2008), the removal ratios of chitosan for *S. aureus* protoplasts and *E. coli* spheroplasts were significantly higher than those for intact cells during the first 3–4 h of contact time, indicating chitosan-cell wall interaction is more intense than other cell membranes.

In another study, chitosan in gel form was used in the antibacterial tests carried out by turbidity and well inhibition zone, showing chitosan consistently more active against the gram-positive *S. aureus* than gram-negative *E. coli* (Goy *et al.*, 2016). Morimoto *et al.*, (2001) reported that chitosan derivatives have better specific binding activity on the cell wall of *Pseudomonas aeruginosa*. Lal *et al.*, (2013) reported the interaction of chitosan with cell surface polymers such as teichoic acid of gram-positive bacteria, which is consistent with the fact that binding of chitosan with the lipopolysaccharide layer of a gram-negative bacteria cell wall would not significantly affect the susceptibility. However, adherence due to electrostatic interaction may cause secondary effects on the cytoplasmic membrane, such as disruption of the cell membrane, which finally results in leakage of small cellular components. The similarity between the antibacterial profiles and patterns of chitosan and those of two other control substances, polymyxin and ethylene diamine tetraacetic acid (EDTA), verified the amino group assisted mechanism of chitosan. Helander *et al.*, (2001) detailed the specific binding mechanism of chitosan on gram-negative bacteria that relates weakening of the barrier function of the outer membrane. Chemical and electrophoretic analyses of free cell supernatants of chitosan-treated cell suspensions showed that the interaction of chitosan with *E. coli* and *S. typhimurium*

involved no release of the LPS or other membrane lipids. This was further evidenced by using highly cationic mutants of *S. typhimurium*, which was more resistant to chitosan than parent strains. In the same study, they found chitosan caused extensive cell surface alterations and covered the outer membrane with vesicular structures, as shown in their electron microscopy study (Helander *et al.*, 2001). The activity of chitosan on gram-positive and gram-negative bacteria was again evidenced and established that chitosan in acid pH is extensively protonated and bound with carboxyl and phosphate groups of the bacterial surface, which offers potential sites for electrostatic binding of chitosan (Li *et al.*, 2016). It should be noted that chitosan shows broad spectrum activity on microorganisms except the fungi, which contain chitosan as a wall constituent. Chitosan's antimicrobial activity (as a preservative) is often limited to food, such as fruit and vegetable products with low protein and NaCl content (Roller and Covill, 1999). The antibacterial activity of chitosan against gram-negative (left) and gram positive (right) bacteria is presented in Fig. (2).

9. Preservative effect of chitosan edible coating on shelf life of papaya

The utilisation of chitosan in the preservation of perishable fruits such as papaya is extensively explored in the form of edible coating, which refers to the application of a thin film directly onto the surface of the targeted product to provide protection (Duan *et al.*, 2019). Edible coatings or films serve as a protective barrier surrounding fruits and vegetables, which are also safe for consumption along with the coated product (Kerch, 2015). Table 2 shows various studies which have investigated the effect of chitosan based edible coating alone or in combination with other added values in managing the post harvest losses in papaya and increasing its shelf life while retaining its sensory attributes for a longer time.

A study conducted by Dotto *et al.*, 2015 investigated the effect of chitosan solution on the microbiological shelf life extension of papaya. The researchers coated papaya fruit samples using chitosan solutions having different molecular weights (150 kDa and 300 kDa) and stored them for 20 days under ambient conditions. It was observed that the application of chitosan solutions (150 kDa and 300 kDa) on papaya fruits resulted in decreased levels of mesophilic bacteria and yeasts/moulds compared to untreated fruits, especially after 5 and 7 days of storage, respectively. Similar observations were reported by Escamilla-García *et al.* (2018), where researchers coated the papaya sample with chitosan and oxidised starch and observed that the logarithmic values of the total coliforms, yeasts, mesophilic bacteria, and fungi were significantly higher in the

uncoated papaya samples compared to the coated ones. Additionally, the growth rate of these microorganisms was observed to be slower in the coated papaya samples compared to the uncoated ones. This indicates that chitosan solution possesses antibacterial and antifungal properties when used on papaya fruits, which are believed to be due to the cationic properties of chitosan, which can affect the composition and permeability of microorganisms' cell membranes (Bautista-Baños *et al.*, 2013). Chitosan's antimicrobial effects can also be attributed to its ability to reduce the respiratory activity and activity of certain enzymes in microorganisms (Aquino *et al.*, 2015).

Another study by Vilaplana *et al.* (2019) evaluated the impact of chitosan post-harvest dip treatment (different concentrations ranging from 0.5-3%) on papaya fruits during storage at room temperature ($28\pm 1^{\circ}\text{C}$) and cold storage ($12\pm 1^{\circ}\text{C}$). The results indicated that during a storage period of 7 and 18 days at ambient temperature and cold storage, respectively, untreated papaya fruits experienced a maximum weight loss of 9.90% and 8.97%, while papaya fruits coated with 3% chitosan exhibited a significantly lower weight loss of 5.1% and 4.5% respectively. These observations are in accordance with the results obtained by Bhanushree *et al.* (2018), which indicated that chitosan treated fruits (3% w/v) had the lowest weight loss and lowest moisture loss as compared to the untreated samples. The formation of a barrier against moisture and gases can result in reductions in weight loss and respiration rate, therefore delaying the onset of spoilage and extending the shelf-life of the product (Barrera *et al.*, 2015; Chien *et al.*, 2007). It was also observed that the softening of papaya fruits during storage was delayed by the application of chitosan, which resulted in lower firmness loss compared to untreated fruits. This observation may be attributed to the film-forming properties of chitosan, which can slow down the ripening process and thus minimise fruit softening. Similar results were obtained in other studies where papaya fruits were treated with chitosan in combination with ascorbic acid (Zhou *et al.*, 2022), beeswax emulsion (Mukdisari *et al.*, 2016), Tuna skin gelatin and black tea extract (Sekarina *et al.*, 2023), pullulan coating (Zhang *et al.*, 2019) and oxidised starch (Escamilla-García *et al.*, 2018) respectively.

Papaya fruits treated with chitosan alone or along with some added value (ascorbic acid, oxidised starch, pullulan and propolis extract respectively) recorded lesser change in colour than untreated fruits, and this is evident from the colour (L, a^* and b^*) values at the end of storage period (Vilaplana *et al.*, 2019; Zhou *et al.*, 2022; Zhang *et al.*, 2019; Bhanushree *et al.*, 2018; Barrera *et al.*, 2015). The colour loss occurs in untreated fruits due to the degradation of chlorophyll by ethylene gas and respiratory enzymes of papaya. The application of chitosan coating to papaya

creates a reduced oxygen environment on the fruit surface, which leads to the inhibition of respiratory enzymes and ethylene synthetases, and this is achieved more effectively when chitosan is combined with multilayered coatings (Niu *et al.*, 2019). These metabolic changes resulted in a delayed maturation process and increased stability of the fruit's cell wall, providing protection to chloroplasts and chlorophylls (Allanigue *et al.*, 2017).

Various studies have shown that chitosan combined with other preservative components can more effectively manage post harvest losses in papaya as compared to sole chitosan treatment. Zhang *et al.* (2022) studied the combined effect of chitosan and ascorbic acid in maintaining the post-harvest quality of papaya by coating the papaya samples with different concentrations, including 1.5% (w/v) ascorbic acid, 1.0% (w/v) chitosan, and 1.5% (w/v) ascorbic acid + 1.0% (w/v) chitosan respectively. It was reported that chitosan combined ascorbic acid treatment significantly reduced H₂O₂ content (26.23% – 66.26%) and MDA levels (28.06 %–43.36%) at all storage dates as compared to control and also resulted in inhibition of softening enzyme activity and gene expression level of cell wall degrading enzymes such as CX, PG, PME, and β -GAL due to the presence of ascorbic acid, which activates or preserves the antioxidant enzyme activity, leading to scavenging of the reactive oxygen species which are responsible for oxidative damage (Soares *et al.*, 2021). Similar results were obtained by Sekarina *et al.* (2023), where the researchers coated minimally processed papaya with edible coatings of chitosan - fish skin gelatine containing black tea extract in different concentrations (0, 5, 10 and 15%, respectively) and reported that the antioxidant activity was found to be slightly higher in chitosan-tuna fish gelatin extract combined with 15% black tea extract (59.26-72.62%) as compared to control samples (66.97-72%) and this may due to presence of tea polyphenols and chitosan, as both have strong antioxidant properties especially in terms of the ability to scavenge hydroxyl radicals and reactive oxygen species (Anggraini *et al.*, 2016). It was also observed that microbial growth in the control papaya was faster as compared to the treatment sample with the addition of black tea extract, and this may be due to the antimicrobial effect of tea polyphenols and chitosan. Chitosan can interact with negatively charged residues on the surface of microbial cells by an electrostatic attraction so that it flocculates and absorbs on the surface of microorganisms, then inhibits the physiological metabolism of microorganisms (Badawy *et al.*, 2017). On the other hand, tea polyphenols have the ability to cause coagulation of structural proteins, as well as binding to deoxyribonucleic acid molecules. Additionally, they can induce damage to the cell membranes and walls of microorganisms, ultimately leading to the inhibition of their growth (Bansal *et al.*, 2013). Another study conducted by Barrera *et al.* (2015) investigated the effect of chitosan

combined with propolis extract for post harvest treatment of papaya where papaya samples were treated with two different concentrations: 1% chitosan (control) and 1% chitosan combined with 5% propolis extract. The results indicated that the deterioration index was lower in fruits treated with chitosan combined with propolis extract as compared to control, and this may be due to a synergistic antimicrobial effect between propolis and chitosan due to which the treatment fruits presented moderate damage, indicating a higher resistance to deterioration compared to the control papaya. Chitosan-coated films demonstrate a wide-ranging ability to inhibit bacterial growth, and when combined with propolis extract, the antibacterial effectiveness is further improved against all the pathogens tested, demonstrating the fungicidal and bactericidal effect of propolis along with chitosan (Torlak & Sert, 2013). Also, the fruits treated with chitosan combined with propolis extract exhibited a significant increase in chlorophyll concentration towards the end of their storage period, as compared to the control fruits. This outcome provides evidence that the fruits coated with chitosan combined with propolis experienced a delayed pigmentation process and a probable inhibition of ripening.

10. Conclusion and future scope

In conclusion, chitosan coating has been shown to be a promising approach to managing post-harvest losses in papaya. It has been found to have antimicrobial properties, preserve the quality of the fruit, and reduce water loss, all of which contribute to an extended shelf life and a decrease in overall losses. Therefore, the use of chitosan coating can contribute to the sustainability of papaya production and supply, as well as improve the economic viability of farmers. Future research could focus on optimising the use of chitosan coating by investigating the effect of different concentrations and application methods on papaya and combining it with other preservative components to boost its effect. Moreover, exploring the potential of combining chitosan coating with other preservation techniques, such as modified atmosphere packaging or cold storage, could provide a more effective and integrated solution to post-harvest losses. Additionally, the potential health benefits of chitosan-coated papaya, such as increased antioxidant activity, could be further investigated. Overall, continued research on the use of chitosan coating in the management of post-harvest losses in papaya could have significant implications for the food industry and global food security.

Funding

Funding provided by University of Debrecen. Funding Project No. TKP2021-NKTA-32 has been implemented with support from the National Research, Development and Innovation Fund of Hungary, financed under the TKP2021-NKTA funding scheme.

Availability of data and materials

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Declarations Competing interests

The authors declare no competing interests

References

Alfonso, C., Nuero, O. M., Santamaría, F., & Reyes, F. (1995). Purification of a heat stable chitin deacetylase from *Aspergillus nidulans* and its role in cell wall degradation. *Current Microbiology*, 30, 49–54. <https://doi.org/10.1007/BF00294524>

Allanigue, D.K.A., Sabulase, V.C., Hernandez, H.P., & Serrano, E.P. (2017). The effect of chitosan-based nanocomposite coating on the post-harvest life of papaya (*Carica papaya* L.) fruits. *Philippine Agricultural Scientist*, 100(3), 233-242. <https://doi.org/10.1016/j.scienta.2016.11.017>

Ali, A., Devarajan, S., Waly, M. I., Essa, M. M., & Rahman, M. S. (2012). Nutritional and medicinal values of papaya (*Carica papaya* L.). In *Natural Products and Their Active Compounds on Disease Prevention* (pp. 307-324). Nova Science Publishers, Inc.

Alishahi, A., & Aider, M. (2011). Applications of Chitosan in the Seafood Industry and Aquaculture: A Review. *Food and Bioprocess Technology*, 5(3), 817–830. doi:10.1007/s11947-011-0664-x

Anacarso, I., De Niederhäusern, S., Iseppi, R., Sabia, C., Bondi, M., & Messi, P. (2011). Antilisterial activity of chitosan and Enterocin 416K1 in artificially contaminated RTE products. *Food Control*, 22(12), 2076–2080. <https://doi.org/10.1016/j.foodcont.2011.06.001>

Anitha, B, Raghu, N, Gopenath, T.S., Karthikeyan, M, Gnanasekaran, A, Chandrashekrappa, G.K. and Basalingappa, K. M. (2018). Medicinal Uses of Carica Papaya. *Journal of Natural & Ayurvedic Medicine*. <https://doi.org/10.23880/jonam-16000144>

Anonymous, (2018). 1st Advance Estimate, Indian Horticulture Database. National Horticulture Board, Ministry of Agriculture, Government of India, Gurgaon, India.

Anggraini, T., Febrianti, F., Aisman, & Ismanto, S. D. (2016). Black tea with averrhoa bilimbi l extract: A healthy beverage. *Agriculture and Agricultural Science Procedia*, 9, 241-252. <https://doi.org/10.1016/j.aaspro.2016.02.141>

APEDA Agriexchange, (2021). India production of PAPAYA. https://agriexchange.apeda.gov.in/India%20Production/India_Productions.aspx?cat=fruit&hscod e=1051

Aquino, A. B., Blank, A. F., & Santana, L. C. L. A. (2015). Impact of edible chitosan cassava starch coatings enriched with *Lippia gracilis* Schauer genotype mixtures on the shelf life of guavas (*Psidium guajava* L.) during storage at room temperature. *Food Chemistry*, 171, 108-116. <https://doi.org/10.1016/j.foodchem.2014.08.112>

Araki, Y., & Ito, E. (1975). A pathway of chitosan formation in *Mucor rouxii*: Enzymatic deacetylation of chitin. *European Journal of Biochemistry*, 189, 249–253.

Aranaz, I., Mengibar, M., Harris, R. B. S., Panos, I., Miralles, B., Acosta, N., Galed, G., & Heras, A. (2009). Functional Characterisation of Chitin and Chitosan. *Current Chemical Biology*, 3(2), 203–230. <https://doi.org/10.2174/187231309788166415>

Aranda-Martinez, A., Lopez-Moya, F., & Lopez-Llorca, L. V. (2016). Cell wall composition plays a key role on sensitivity of filamentous fungi to chitosan. *Journal of Basic Microbiology*, 56(10), 1059–1070. <https://doi.org/10.1002/jobm.201500775>

Archampong, T. N., Asmah, R. H., Richards, C., Martin, V. J., Bayliss, C. D., Botão, E., David, L., Beleza, S., & Carrilho, C. (2019). Gastro-duodenal disease in Africa: Literature review and clinical data from Accra, Ghana. *World Journal of Gastroenterology*, 25(26), 3344–3358. <https://doi.org/10.3748/wjg.v25.i26.3344>

Arnon, H., Zaitsev, Y., Porat, R., & Poverenov, E. (2014). Effects of carboxymethyl cellulose and chitosan bilayer edible coating on post-harvest quality of citrus fruit. *Post-harvest Biology and Technology*, 87, 21–26. doi:10.1016/j.postharvbio.2013.08.007

Barrera, E., Gil, J., Restrepo, A., Mosquera, K., & Durango, D. (2015). A coating of chitosan and propolis extract for the post-harvest treatment of papaya (*Carica papaya* L. cv. Hawaiiiana). *Revista Facultad Nacional de Agronomía*, 68(2), 7667–7678. doi:10.15446/rfnam.v68n2.50982.

Bautista-Baños, S., Sivakumar, D., Bello-Pérez, A., Arce, R. V., & Hernández-López, M. (2013). A review of the management alternatives for controlling fungi on papaya fruit during the post-harvest supply chain. *Crop Protection*, 49, 8-20. <https://doi.org/10.1016/j.cropro.2013.03.002>

Bautista-Baños, S., Hernández-López, M., Trejo-Tapia, J. L., Hernández-Lauzardo, A. N., Bautista-Cerón, M. K., & Melo-Giorgana, G. E. (2005). Effect of chitosan on in-vitro development and morphology of two isolates of *Colletotrichum gloeosporioides* Penz. *Mexican Journal of Phytopathology*, 23, 62-67. doi: 10.1007/s10658-009-9439-6.

Bansal, S., Choudhary, S., Sharma, M., Kumar, S. S., Lohan, S., Bhardwaj, V., Syan, N., & Jyoti, S. (2013). Tea: A native source of antimicrobial agents. *Food Research International*, 53(2), 568-584. <https://doi.org/10.1016/j.foodres.2013.01.032>

Badawy, M. E. I., Rabea, E. I. A. M., El-Nouby, M., Ismail, R. I. A., & Taktak, N. E. M. (2017). Strawberry shelf life, composition, and enzymes activity in response to edible chitosan coatings. *International Journal of Fruit Science*, 17(2), 117-136. <https://doi.org/10.1080/15538362.2016.1219290>

Benhabiles, M. S., Abdi, N., Drouiche, N., Lounici, H., Pauss, A., Goosen, M. F. A., & Mameri, N. (2013). Protein recovery by ultrafiltration during isolation of chitin from shrimp shells *Parapenaeus longirostris*. *Food Hydrocolloids*, 32(1), 28–34. doi:10.1016/j.foodhyd.2012.11.035.

Bhanushree, L. S., Vasudeva, K. R., Suresha, G. J., Sadananda, G. K., Mohamad Tayeebulla, H., & Halesh, G. K. (2018). Influence of chitosan on post-harvest behavior of papaya (*Carica papaya* L.) fruits under different storage conditions. *Journal of Pharmacognosy and Phytochemistry*, 7(2), 2010-2014. doi: 10.22271/phyto.2018.v7.isp2c.8787.

Chung, Y., & Chen, C. (2008). Antibacterial characteristics and activity of acid-soluble chitosan. *Bioresource Technology*, 99(8), 2806–2814. <https://doi.org/10.1016/j.biortech.2007.06.044>

Chien, P. J., Sheu, F., & Yang, F. H. (2007). Effects of edible chitosan coating on quality and shelf life of sliced mango fruit. *Journal of Food Engineering*, 78(1), 225-229. <https://doi.org/10.1016/j.jfoodeng.2005.09.022>.

Dadi, T. L., Habtamu, G. M., & Dejene, T. B. (2020). Assessment of Papaya Post-harvest Loss at Wholesaler and Retailer levels in Jimma Town, South Western Ethiopia. *International Journal of Plant Breeding and Crop Science*, 7(3), 900-908. doi: 10.15744/ijpbcs.2020.08.011

DAIS, (2009). Cultivating papayas. Department of Agriculture, Forestry and Fisheries, Directorate Agricultural Information Services, Pretoria, South Africa. Retrieved from www.nda.agric.za/docs/Infopaks/papayas.pdf

DaMatta, F. M., Ronchi, C. P., Maestri, M., & Barros, R. S. (2007). Ecophysiology of coffee growth and production. *Brazilian Journal of Plant Physiology*, 19(4), 485–510. <https://doi.org/10.1590/s1677-04202007000400014>

Devaki, C. S., Samreen, F., & Prakash, J. (2015). A Review on Composition, Processed Products and Medicinal Uses of Papaya (*Carica Papaya* L.). *International Journal of Food, Nutrition and Dietetics*. <https://doi.org/10.21088/ijfnd.2322.0775.3315.3>

Devlieghere, F., Vermeulen, A., & Debevere, J. (2004). Chitosan: antimicrobial activity, interactions with food components and applicability as a coating on fruit and vegetables. *Food Microbiology*, 21(6), 703–714. <https://doi.org/10.1016/j.fm.2004.02.008>

Divya, K., Rebello, S., & Jisha, M. S. (2014). A Simple and Effective Method for Extraction of High Purity Chitosan from Shrimp Shell Waste. *International Conference on Advances in Applied Science & Environmental Engineering*, December, 141-145. doi: 10.15242/IICBE.C1214059

Doan, M., Huynh, M., Pham, A. T., Chau, N. D. Q., & Le, P. T. K. (2020). Extracting Seed Oil and Phenolic Compounds from Papaya Seeds by Ultrasound-assisted Extraction Method and Their Properties. *Chemical Engineering Transactions*, 78, 493–498. <https://doi.org/10.3303/cet2078083>

Dotto, G. L., Vieira, M. L. G., & Pinto, L. A. A. (2015). Use of chitosan solutions for the microbiological shelf life extension of papaya fruits during storage at room temperature. *LWT - Food Science and Technology*, 64(1), 126–130. doi:10.1016/j.lwt.2015.05.042

Duan, C., Meng, X., Meng, J., Khan, I. A., Wang, L., Khan, A., An, X., Zhang, J., Huq, T., & Ni, Y. (2019). Chitosan as A Preservative for Fruits and Vegetables: A Review on Chemistry and Antimicrobial Properties. *Journal of Bioresources and Bioproducts*, 4(1), 11–21. <https://doi.org/10.21967/jbb.v4i1.189>

Elsabee, M. Z., & Abdou, E. S. (2013). Chitosan based edible films and coatings: A review. *Materials Science and Engineering: C*, 33(4), 1819–1841. doi:10.1016/j.msec.2013.01.010

Escamilla-García, M., Rodríguez-Hernández, M. A., Hernández-Hernández, H., Delgado-Sánchez, L. F., García-Almendárez, B. E., Amaro-Reyes, A., & Regalado, C., (2018). Effect of an Edible Coating Based on Chitosan and Oxidized Starch on Shelf Life of *Carica papaya* L., and Its Physicochemical and Antimicrobial Properties. *Coatings*, 8(9), 318. <https://doi.org/10.3390/coatings8090318>.

Esguerra, E., Rapusas, R., & Rolle, R. S. (2020). Post-Harvest Management of Papaya to Ensure Quality and Safety - Guidance for Stakeholders in the Horticultural Supply Chain. Rome: FAO. doi: 10.4060/ca9452en.

FAO, (2011). Global food losses and food waste. In: International Congress —“SAVE FOOD!” 16-17 May, Düsseldorf, Germany. Retrieved from <http://www.fao.org/3/i2697e/i2697e.pdf>.

FAOSTAT, (2021). Food and Agriculture Organization of the United Nations: Production/Crops - Papayas. Retrieved from <http://www.fao.org/faostat/en/#data/QC/visualize>

Fatehi, P., Kititerakun, R., Ni, Y., & Xiao, H. (2010). Synergy of CMC and modified chitosan on strength properties of cellulosic fiber network. *Carbohydrate Polymers*, 80(1), 208–214. doi:10.1016/j.carbpol.2009.11.012.

Gajanana, T., Sudha, M., Saxena, A. K., & Dakshinamoorthy, V. (2010). Post harvest handling, marketing and assessment of losses in papaya. *Acta Horticulturae*, 851, 519–526. <https://doi.org/10.17660/actahortic.2010.851.79>

Galo, J.D.Q.B., De Souza, M.L., Kusdra, J.F., Mattiuz, C.F.M., (2014). Post-harvest storage of papaya ‘Sunrise Solo’ with the use of chitosan. *Revista Brasileira De Fruticultura*. 36, 305–312. <https://doi.org/10.1590/0100-2945-194/13>

Gao, X. D., Katsumoto, T., & Onodera, K. (1995). Purification and characterization of chitin deacetylase from *Absidia coerulea*. *Journal of Biochemistry*, 117, 257–263. <https://doi.org/10.1093/jb/117.2.257>.

Gao, N., Du, W., Zhang, M., Ling, G., & Zhang, P. (2022). Chitosan-modified biochar: Preparation, modifications, mechanisms and applications. *International Journal of Biological Macromolecules*, 209, 31–49. <https://doi.org/10.1016/j.ijbiomac.2022.04.006>

Goy, R. C., Morais, S. T. B., & Assis, O. B. G., (2016). Evaluation of the antimicrobial activity of chitosan and its quaternized derivative on *E. coli* and *S. aureus* growth. *Revista Brasileira de Farmacognosia*, 26(1), 122-127. <https://doi.org/10.1016/j.bjp.2015.09.010>

Gonzalez-Aguilar, G. A., Valenzuela-Soto, E., Lizardi-Mendoza, J., Goycoolea, F., Martínez-Téllez, M. A., Villegas-Ochoa, M. A., Monroy-García, I. N., & Ayala-Zavala, J. F. (2009). Effect of chitosan coating in preventing deterioration and preserving the quality of fresh-cut papaya “Maradol”. *Journal of the Science of Food and Agriculture*, 89(1), 15-23. doi:10.1002/jsfa.3405

Hajji, S., Younes, I., Ghorbel-Bellaaj, O., Hajji, R., Rinaudo, M., Nasri, M., & Jellouli, K. (2014). Structural differences between chitin and chitosan extracted from three different marine sources. *International Journal of Biological Macromolecules*, 65, 298–306. <https://doi.org/10.1016/j.ijbiomac.2014.01.045>

Hamdayanty, Yunita, R., Amin, N. N., & Damayanti, T. A. (2012). Pemanfaatan chitosan untuk mengendalikan antraknosa pada papaya (*Colletotricum gloeosporioides*) dan meningkatkan daya simpan buah. *Jurnal Fitopatologi Indonesia*, 8, 97-102. <https://doi.org/10.14692/jfi.8.2.97>

Helander, I. M., Nurmiaho-Lassila, E. L., Ahvenainen, R., Rhoades, J., & Roller, S. (2001). Chitosan disrupts the barrier properties of the outer membrane of Gram-negative bacteria. *International Journal of Food Microbiology*, 71(2-3), 235-244. [https://doi.org/10.1016/s0168-1605\(01\)00609-2](https://doi.org/10.1016/s0168-1605(01)00609-2)

Hossain, M., & Iqbal, A. (2014). Production and characterisation of chitosan from shrimp waste. *Journal of Bangladesh Agricultural University*, 12(1), 153-160.

Hosseinnejad, M. T., & Jafari, S. M. (2016). Evaluation of different factors affecting antimicrobial properties of chitosan. *International Journal of Biological Macromolecules*, 85, 467–475. <https://doi.org/10.1016/j.ijbiomac.2016.01.022>

Hu, Z., & Gänzle, M. G. (2018). Challenges and opportunities related to the use of chitosan as a food preservative. *Journal of Applied Microbiology*, 126(5), 1318–1331. <https://doi.org/10.1111/jam.14131>

Huq, T., Vu, K. D., Riedl, B., Bouchard, J., & Lacroix, M. (2015). Synergistic effect of gamma (γ)-irradiation and microencapsulated antimicrobials against *Listeria monocytogenes* on ready-to-eat (RTE) meat. *Food Microbiology*, 46, 507–514. doi:10.1016/j.fm.2014.09.013

Ibitoye, E. B., Lokman, I. H., Hezme, M. N. M., Goh, Y. M., Zuki, A. B. Z., & Jimoh, A. A. (2018). Extraction and physicochemical characterisation of chitin and chitosan isolated from house cricket. *Biomedical Materials*, 13(2), 025009. doi:10.1088/1748-605x/aa9dde.

Isa, M. T., Ameh, A., Tijjani, M., & Adama, K. K. (2012). Extraction and characterisation of chitin and chitosan from Nigerian shrimps. *International Journal of Biological and Chemical Sciences*. <https://doi.org/10.4314/ijbcs.v6i1.40>

Islam, M. B., Khalekuzzaman, M., Kabir, S. B., & Hossain, M. R. (2022). Shrimp waste-derived chitosan harvested microalgae for the production of high-quality biocrude through hydrothermal liquefaction. *Fuel*, 320, 123906. <https://doi.org/10.1016/j.fuel.2022.123906>

Ivshina, T. N., Artamonova, S. D., Ivshin, V. P., & Sharnina, F. F. (2009). Isolation of the chitin-glucan complex from the fruiting bodies of *mycothallus*. *Applied Biochemistry and Microbiology*, 45, 31-38. <https://doi.org/10.1134/S0003683809010042>

Kara, A., & Stevens, R. (2002). Characterisation of biscuit fired bone china body microstructure. Part I: XRD and SEM of crystalline phases. *Journal of the European Ceramic Society*, 22(5), 731-736. [https://doi.org/10.1016/S0955-2219\(01\)00382-5](https://doi.org/10.1016/S0955-2219(01)00382-5)

Kerch, G. (2015). Chitosan films and coatings prevent losses of fresh fruit nutritional quality: a review. *Trends in Food Science & Technology*, 46(2): 159–166. doi: 10.1016/j.tifs.2015.10.010.

Khan, A., Vu, K. D., Riedl, B., & Lacroix, M. (2015). Optimisation of the antimicrobial activity of nisin, Na-EDTA and pH against gram-negative and gram-positive bacteria. *LWT - Food Science and Technology*, 61(1), 124-129. <https://doi.org/10.1016/j.lwt.2014.11.035>.

Khan, T. A., Peh, K. K., & Ch'ng, H. S. (2002). Reporting degree of deacetylation values of chitosan: The influence of analytical methods. *Journal of Pharmacy and Pharmaceutical Sciences*, 5(3), 205-212. <https://doi.org/10.18433/jpps.v5i3.225>

Knidri, H. E., Belaabed, R., Addaou, A., Laajeb, A., & Lahsini, A. (2018). Extraction, chemical modification and characterisation of chitin and chitosan. *International Journal of Biological Macromolecules*, 120, 1181–1189. <https://doi.org/10.1016/j.ijbiomac.2018.08.139>

Knorr, D. 1982. Functional properties of chitin and chitosan. *Journal of Food Science*. 47(2): 593-595.

Krishna, K., Paridhavi, M., & Patel, J. A. (2008). Review on nutritional, medicinal and pharmacological properties of Papaya (*Carica papaya* Linn.). *NPR Vol. 7(4)* [July-August 2008].

Kumari, S., Kumar Annamareddy, S. H., Abanti, S., & Kumar Rath, P. (2017). Physicochemical properties and characterisation of chitosan synthesised from fish scales, crab and shrimp shells. *International Journal of Biological Macromolecules*, 104, 1697–1705. [doi:10.1016/j.ijbiomac.2017.04.119](https://doi.org/10.1016/j.ijbiomac.2017.04.119)

Kumari, S., Rath, P., Sri Hari Kumar, A., & Tiwari, T. N. (2015). Extraction and characterisation of chitin and chitosan from fishery waste by chemical method. *Environmental Technology & Innovation*, 3, 77–85. [doi:10.1016/j.eti.2015.01.002](https://doi.org/10.1016/j.eti.2015.01.002)

Kurita, K. (2006). Chitin and Chitosan: Functional Biopolymers from Marine Crustaceans. *Marine Biotechnology*, 8(3), 203–226. <https://doi.org/10.1007/s10126-005-0097-5>

Lal, J., Gupta, S. K., Thavaselvam, D., & Agarwal, D. D. (2013). Biological activity, design, synthesis and structure activity relationship of some novel derivatives of curcumin containing sulfonamides. *European Journal of Medicinal Chemistry*, 64, 579-588. <https://doi.org/10.1016/j.ejmech.2013.03.012>

Landi, L., Peralta-Ruiz, Y. Y., Chaves-López, C., & Romanazzi, G. (2021). Chitosan Coating Enriched with *Ruta graveolens* L. Essential Oil Reduces Postharvest Anthracnose of Papaya

(*Carica papaya* L.) and Modulates Defense-Related Gene Expression. *Frontiers in Plant Science*, 12. <https://doi.org/10.3389/fpls.2021.765806>

Leleu, S., Herman, L., Heyndrickx, M., De Reu, K., Michiels, C., De Baerdemaeker, J., & Messens, W. (2011). Effects on *Salmonella* shell contamination and trans-shell penetration of coating hens' eggs with chitosan. *International Journal of Food Microbiology*, 145(1), 43–48. <https://doi.org/10.1016/j.ijfoodmicro.2010.11.023>

Li, J. H., Wu, Y. G., & Zhao, L. Q. (2016). Antibacterial activity and mechanism of chitosan with ultra-high molecular weight. *Carbohydrate Polymers*, 148, 200-205. doi:10.1016/j.carbpol.2016.04.025

Li, Q., Dunn, E. T., Grandmaison, E. W., & Goosen, M. F. (1992). Applications and properties of chitosan. *Journal of Bioactive and Compatible Polymers*, 7(4), 370-397. doi: 10.1177/088391159200700402

Mahdy Samar, M., El-Kalyoubi, M. H., Khalaf, M. M., & Abd El-Razik, M. M. (2013). Physicochemical, functional, antioxidant and antibacterial properties of chitosan extracted from shrimp wastes by microwave technique. *Annals of Agricultural Sciences*, 58(1), 33–41. doi:10.1016/j.aosas.2013.01.006

Majtán, J., Bíliková, K., Markovič, O., Gróf, J., Kogan, G., & Šimúth, J. (2007). Isolation and characterisation of chitin from bumblebee (*Bombus terrestris*). *International Journal of Biological Macromolecules*, 40, 237-241. <https://doi.org/10.1016/j.ijbiomac.2006.06.016>

Marpudi, S. L., Abirami, L., Pushkala, R., & Srividya, N. (2011). Enhancement of storage life and quality maintenance of papaya fruits using Aloe vera based antimicrobial coating. *IJBT Vol. 10*(1), [January 2011]. doi: 10.1007/s13205-011-0016-2

Martinou, A., Kafetzopoulos, D., & Bouriotis, V. (1993). Isolation of chitin deacetylase from *Mucor rouxii* by immunoaffinity chromatography. *Journal of Chromatography*, 644, 35–41. [https://doi.org/10.1016/0021-9673\(93\)80117-Q](https://doi.org/10.1016/0021-9673(93)80117-Q)

Martinou, A., Kafetzopoulos, D., & Bouriotis, V. (1995). Chitin deacetylation by enzymatic means: Monitoring of deacetylation processes. *Carbohydrate Research*, 273, 235–242. [https://doi.org/10.1016/0008-6215\(95\)00111-6](https://doi.org/10.1016/0008-6215(95)00111-6)

Mashau, M. E., Moyane, J. N., & Jideani, I. A. (2012). Assessment of post harvest losses of fruits at Tshakhuma fruit market in Limpopo Province, South Africa. *African Journal of Agricultural Research*, 7(29). <https://doi.org/10.5897/ajar12.392>

Miraftab, M., Barnabas, J., Kennedy, J. F., & Masood, R. (2011). Antimicrobial Properties of Alginate-Chitosan (Alchite) Fibers Developed for Wound Care Applications. *Journal of Industrial Textiles*. <https://doi.org/10.1177/1528083710376985>

Moradi, M., Tajik, H., Rohani, S. M. R., & Oromiehie, A. R. (2011). Effectiveness of Zataria multiflora Boiss essential oil and grape seed extract impregnated chitosan film on ready-to-eat mortadella-type sausages during refrigerated storage. *Journal of the Science of Food and Agriculture*, 91(15), 2850–2857. <https://doi.org/10.1002/jsfa.4531>

Morimoto, M., Saimoto, H., Usui, H., Okamoto, Y., Minami, S., & Shigemasa, Y. (2001). Biological activities of carbohydrate-branched chitosan derivatives. *Biomacromolecules*, 2(4), 1133-1136. doi: 10.1021/bm010063p

Mahmud, T. M. M., Eryani-Raqeeb, A. A., Omar, S. R. S., Zaki, A. R. M., & Abdul-Rahman, A. E. (2008). Effects of Different Concentrations and Applications of Calcium on Storage Life and Physicochemical Characteristics of Papaya (*Carica Papaya L.*). *American Journal of Agricultural and Biological Sciences*, 3(3), 526–533. <https://doi.org/10.3844/ajabssp.2008.526.533>

Mukdisari, Y., Suketi, K., & Widodo, W. D. (2016). Fruit Coating with Chitosan and Beeswax to Increase Papaya Shelf Life. *Journal of Tropical Crop Science*, 3(3), 93–98. <https://doi.org/10.29244/jtcs.3.3.93-98>

Neto, J. R., Copes, F., Chevallier, P., Vieira, R. S., Da Silva, J., Mantovani, D., & Beppu, M. M. (2022). Polysaccharide-based layer-by-layer nanoarchitectonics with sulfated chitosan for tuning anti-thrombogenic properties. *Colloids and Surfaces B: Biointerfaces*, 213, 112359. <https://doi.org/10.1016/j.colsurfb.2022.112359>

Niu, B., Shao, P., Chen, H., & Sun, P. (2019). Structural and physicochemical characterisation of novel hydrophobic packaging films based on pullulan derivatives for fruits preservation. *Carbohydrate Polymers*, 208, 276-284. <https://doi.org/10.1016/j.carbpol.2018.12.074>

No, H. K., Lee, S. Y., Park, N. Y., & Meyers, S. P. (2003). Comparison of Physicochemical, Binding, and Antibacterial Properties of Chitosans Prepared without and with Deproteinization Process. *Journal of Agricultural and Food Chemistry*, 51(26), 7659–7663. <https://doi.org/10.1021/jf030226w>

Novita, M., Satriana, Martunis, S., Rohaya, E., Hasmarita, & Sudirman. (2012). Pengaruh pelapisan chitosan terhadap sifat fisik dan kimia tomat segar (*Lycopersicum pyriforme*) pada berbagai tingkat kematangan. *Jurnal Teknologi Industri Pertanian Indonesia*, 4, 1-8. doi: 10.21776/ub.jtipi.2012.004.01.1

Odhong, C., Wahome, R. G., Vaarst, M., Nalubwama, S. W., Kiggundu, M., Halb, N., & Githigia, S. (2014). In vitro anthelmintic effects of crude aqueous extracts of *Tephrosia vogelii*, *tephrosia villosa* and *Carica papaya* leaves and seeds. *African Journal of Biotechnology*, 13(52), 4667-4672. doi: 10.5897/AJB2014.14097

Olcum, M., Tastan, B., Ercan, I., Eltutan, I. B., & Genc, S. (2020). Inhibitory effects of phytochemicals on NLRP3 inflammasome activation: A review. *Phytomedicine*, 75, 153238. <https://doi.org/10.1016/j.phymed.2020.153238>

Paull, R. E., Nishijima, W. T., Reyes, M. G., & Cavaletto, C. G. (1997). Post-harvest handling and losses during marketing of papaya (*Carica papaya* L.). *Post-harvest Biology and Technology*, 11(3), 165–179. [https://doi.org/10.1016/s0925-5214\(97\)00028-8](https://doi.org/10.1016/s0925-5214(97)00028-8)

Prajapati, H., Patil, R., & Shukla, Y. (2017). Studies on Biochemical Changes and Changes in Cell Wall Degrading Enzymes in Papaya Fruit Inoculated with *Colletotrichum demetium*. *International Journal of Current Microbiology and Applied Sciences*. <https://doi.org/10.20546/ijcmas.2017.607.232>

Prakash, J., Das, S. P., Bhattacharjee, T., & Singh, N. P. (2014). Studies on effect of pollarding in papaya. *Indian Journal of Horticulture*, 71(3), 419–20.

Prasad, K., & Paul, J. R., (2021). Post-harvest Losses of Papaya and Practice for Management. *Food and Scientific Reports*, 2(February), Pp. 7–11.

Pranoto, Y., Salokhe, V. M., & Rakshit, S. K. (2005). Physical and antibacterial properties of alginate-based edible film incorporated with garlic oil. *Food Research International*, 38(3), 267–272. doi:10.1016/j.foodres.2004.04.009

Puvvada, Y. S., Vankayalapati, S., & Sukhavasi, S. (2012). Extraction of chitin from chitosan from exoskeleton of shrimp for application in the pharmaceutical industry. *International Current Pharmaceutical Journal*, 1(9), 258–263. <https://doi.org/10.3329/icpj.v1i9.11616>

Raafat, D., von Barga, K., Haas, A., & Sahl, H. G. (2008). Insights into the mode of action of chitosan as an antibacterial compound. *Applied and Environmental Microbiology*, 74(12), 3764–3773. doi: 10.1128/aem.00453-08.

Rajestary, R., Landi, L., & Romanazzi, G. (2021). Chitosan and post-harvest decay of fresh fruit: Meta-analysis of disease control and antimicrobial and eliciting activities. *Comprehensive Reviews in Food Science and Food Safety*, 20(1), 563–582. <https://doi.org/10.1111/1541-4337.12672>

Rege, P. R., & Block, L. H. (1999). Chitosan processing: Influence of process parameters during acidic and alkaline hydrolysis and effect of the processing sequence on the resultant chitosan's properties. *Carbohydrate Research*, 321, 235–245. [https://doi.org/10.1016/S0008-6215\(99\)00172-X](https://doi.org/10.1016/S0008-6215(99)00172-X)

Renuka, V., Ravishankar, C. N., Elavarasan, K., Zynudheen, A. A., & Joseph, T. C. (2019). Production and Characterisation of Chitosan from Shrimp Shell Waste of *Parapeneopsis stylifera*. *International Journal of Current Microbiology and Applied Sciences*, 8(11), 2076–2083. <https://doi.org/10.20546/ijcmas.2019.811.240>

Rodrigues, J. P., de Souza Coelho, C. C., Soares, A. G., & Freitas-Silva, O. (2021). Current technologies to control fungal diseases in post-harvest papaya (*Carica papaya* L.). *Biocatalysis and Agricultural Biotechnology*, 36, 102128. doi:10.1016/j.bcab.2021.102128

Roller, S., & Covill, N. (1999). The antifungal properties of chitosan in laboratory media and apple juice. *International Journal of Food Microbiology*, 47(1-2), 67–77. doi:10.1016/s0168-1605(99)00006-9.

Romanazzi, G., Feliziani, E., & Sivakumar, D. (2018). Chitosan, a biopolymer with triple action on post-harvest decay of fruit and vegetables: Eliciting, antimicrobial and film-forming properties. *Frontiers in Microbiology*, 9, 2745. doi: 10.3389/fmicb.2018.02745.

Romanazzi, G., Feliziani, E., Baños, S. B., & Sivakumar, D. (2015). Shelf life extension of fresh fruit and vegetables by chitosan treatment. *Critical Reviews in Food Science and Nutrition*, 57(3), 579–601. doi:10.1080/10408398.2014.900474.

Rout, S. K. (2001). Physicochemical, functional and spectroscopic analysis of crawfish chitin and chitosan as affected by process modification (Doctoral dissertation, Louisiana State University and Agricultural & Mechanical College, USA). https://digitalcommons.lsu.edu/gradschool_disstheses/2916/

Samar, M. E., Mohamed, R. R., & Mohy Eldin, M. S. (2015). Chitosan nanoparticles for improving the oral bioavailability of curcumin. *International Journal of Biological Macromolecules*, 77, 126-133. <https://doi.org/10.1016/j.ijbiomac.2015.03.057>

Santos, C. M. dos, Abreu, C. M. P. de, Freire, J. M., Queiroz, E. de R., & Mendonça, M. M. (2014). Chemical characterisation of the flour of peel and seed from two papaya cultivars. *Food Science and Technology (Campinas)*, 34(2), 353–357. doi:10.1590/fst.2014.0048

Sajomsang, W., & Gonil, P. (2010). Preparation and characterisation of α -chitin from cicada sloughs. *Materials Science and Engineering: C*, 30(3), 357–363. <https://doi.org/10.1016/j.msec.2009.11.014>

Saran, P. L., & Choudhary, R. (2013). Drug bioavailability and traditional medicaments of commercially available Papaya – a review. *African Journal of Agricultural Research*, 8(25), 3216–23.

Saran, P. L., Choudhary, R., & Solanki, I. S. (2014). Papaya: Agri-hort system. *Agriculture Today*, 17(2), 40–1.

Saran, P. L., Choudhary, R., Solanki, I. S., & Kumar, P. R. (2015). Stamen feminization in hermaphrodite papaya: a new report. *Progressive Horticulture*, 47(1), 173–4. doi:10.5958/2249-5258.2015.00032.9.

Saran, P. L., & Choudhary, R. (2019). Advances in papaya cultivation. In *Burleigh Dodds series in agricultural science* (pp. 547–568). <https://doi.org/10.19103/as.2019.0054.15>

Sekarina, A. S., Supriyadi, N., Munawaroh, H., Susanto, E., Show, P., & Ningrum, A. (2023). Effects of edible coatings of chitosan - fish skin gelatine containing black tea extract on quality of minimally processed papaya during refrigerated storage. *Carbohydrate Polymer Technologies and Applications*, 5, 100287. <https://doi.org/10.1016/j.carpta.2023.100287>

Shah, S., & Hashmi, M. S. (2020). Chitosan–aloe vera gel coating delays post-harvest decay of mango fruit. *Horticulture Environment and Biotechnology*, 61(2), 279–289. <https://doi.org/10.1007/s13580-019-00224-7>

Singh, S. P., Kumar, S., & Tomar, M. S. (2019). Aqueous Extract of *Carica papaya* Leaf Elicits the Production of TNF- α and Modulates the Expression of cell Surface Receptors in Tumor-Associated Macrophages. *Bioscience Biotechnology Research Communications*, 12(4), 1115–1122. <https://doi.org/10.21786/bbrc/12.4/35>

Soares, C.G., do Prado, S.B.R., Andrade, S.C.S., & Fabi, J.P. (2021). Systems biology applied to the study of papaya fruit ripening: The influence of ethylene on pulp softening. *Cells*, 10(9), 2339. <https://doi.org/10.3390/cells10092339>

Souza, A.F., Da Silva, W.B., Gonçalves, Y.S., Da Silva, M.G., De Oliveira, J.G., (2014). Ripening physiology of varieties of papaya commercialised in Brazil. *Revista Brasileira De Fruticultura*. 36, 318–328. <https://doi.org/10.1590/0100-2945-215/13>.

Ssekatawa, K., Byarugaba, D. K., Wampande, E. M., Moja, T. N., Nxumalo, E., Maaza, M., Kirabira, J. B., (2021). Isolation and characterisation of chitosan from Ugandan edible mushrooms, Nile perch scales and banana weevils for biomedical applications. *Scientific Reports*, 11(1). doi:10.1038/s41598-021-81880-7

Struszczyk, H., Ginalska, G., Błażewicz, M., Mielcarek, A., & Lewandowska, K. (2002). Chitosan as a biomaterial: studies on its safety and hemostatic potential. *Journal of Biomedical Materials Research Part B: Applied Biomaterials*, 63(4), 438-447. <https://doi.org/10.1002/jbm.10259>

Sundara, R. G., Aruchami, M., & Gowri, N. (1982). Natural deacetylation of chitin to chitosan in the abdominal cuticle of the physogastric queen of *Macrotermes estherae*. In S. Tokura & S. Hirano (Eds.), *Proceedings Second International Conference Chitin/Chitosan* (pp. 12–14). Tottori, Japan: Japanese Society Chitin.

Sudarshan, N., Hoover, D. G., & Knorr, D. (1992). Antibacterial action of chitosan. *Food Biotechnology*, 6(3), 257–272. <https://doi.org/10.1080/08905439209549838>

Szymańska, E., & Winnicka, K. (2015). Stability of Chitosan—A Challenge for Pharmaceutical and Biomedical Applications. *Marine Drugs*, 13(4), 1819–1846. <https://doi.org/10.3390/md13041819>

Tan, C. X., Tan, S. T., & Tan, S. S. (2020). An overview of papaya seed oil extraction methods. *International Journal of Food Science & Technology*, 55(4), 1506–1514. doi:10.1111/ijfs.14431

Tabassum, N., & Khan, M. A. (2020). Modified atmosphere packaging of fresh-cut papaya using alginate based edible coating: Quality evaluation and shelf life study. *Scientia Horticulturae*, 259, 108853. doi:10.1016/j.scienta.2019.108853

Tolaimate, A., Desbrières, J., Rhazi, M., Alagui, A., Vincendon, M., & Vottero, P. (2000). On the influence of deacetylation process on the physicochemical characteristics of chitosan from squid chitin. *Polymer*, 41, 2463–2469. [https://doi.org/10.1016/S0032-3861\(99\)00400-0](https://doi.org/10.1016/S0032-3861(99)00400-0)

Torlak, E., & Sert, D. (2013). Antibacterial effectiveness of chitosan–propolis coated polypropylene films against foodborne pathogens. *International Journal of Biological Macromolecules*, 60(1), 52-55. <https://doi.org/10.1016/j.ijbiomac.2013.05.013>

Tridge.com, (2021). Fresh Papaya production and top producing countries - Tridge. Retrieved from <https://www.tridge.com/intelligences/papaya/production>

USFDA, (2013). GRAS notice inventory. GRN No. 397. Retrieved from <https://www.fda.gov/food/generally-recognized-safe-gras/gras-notice-inventory>.

Ushakumari, U. N., & Ramanujan, R. (2012). Astaxanthin from shrimp shell waste. *International Journal of Pharmaceutical Chemistry Research*, 1(3), 1-6. doi: 10.7439/ijpcr.v1i3.124.

Vargas, M., Pastor, C. B., Chiralt, A., & McClements, D. J. (2008). Recent Advances in Edible Coatings for Fresh and Minimally Processed Fruits. *Critical Reviews in Food Science and Nutrition*, 48(6), 496–511. <https://doi.org/10.1080/10408390701537344>

Vilaplana, R., Chicaiza, G., Vaca, C., & Valencia-Chamorro, S. (2019). Combination of hot water treatment and chitosan coating to control anthracnose in papaya (*Carica papaya* L.) during the post-harvest period. *Crop Protection*, 105007. doi:10.1016/j.cropro.2019.105007.

Vij, T., & Prashar, Y. (2015). A review on medicinal properties of *Carica papaya* Linn. *Asian Pacific Journal of Tropical Disease*, 5(1), 1–6. [https://doi.org/10.1016/s2222-1808\(14\)60617-4](https://doi.org/10.1016/s2222-1808(14)60617-4)

Wang, X., Del Mar Contreras, M., Xu, D., Xing, C., Wang, L., & Yang, D. (2020). Different distribution of free and bound phenolic compounds affects the oxidative stability of tea seed oil: A novel perspective on lipid antioxidation. *Lebensmittel-Wissenschaft & Technologie*, 129, 109389. <https://doi.org/10.1016/j.lwt.2020.109389>

Wardana, A. A., Koga, A., Tanaka, F., & Tanaka, F. (2021). Antifungal features and properties of chitosan/sandalwood oil Pickering emulsion coating stabilised by appropriate cellulose nanofiber dosage for fresh fruit application. *Scientific Reports*, 11, 1-15. <https://doi.org/10.1038/s41598-021-98074-w>

Yadav, M., Kaushik, B., Rao, G. K., Srivastava, C. M., & Vaya, D. (2023). Advances and challenges in the use of chitosan and its derivatives in biomedical fields: A review. *Carbohydrate Polymer Technologies and Applications*, 5, 100323. <https://doi.org/10.1016/j.carpta.2023.100323>

Yen, M. T., Yang, J. H., & Mau, J. L. (2009). Physicochemical characterisation of chitin and chitosan from crab shells. *Carbohydrate Polymers*, 75(1), 15-21. doi: 10.1016/j.carbpol.2008.06.005.

Yilmaz, A. H. (2020). Antibacterial activity of chitosan-based systems. In *Functional Chitosan: Drug Delivery and Biomedical Applications*. 457–489 https://doi.org/10.1007/978-981-15-0263-7_15

Younes, I., & Rinaudo, M. (2015). Chitin and Chitosan Preparation from Marine Sources. Structure, Properties and Applications. *Marine Drugs*, 13(3), 1133–1174. <https://doi.org/10.3390/md13031133>

Zhang, L., Huang, C., & Zhao, H. (2019). Application of Pullulan and Chitosan Multilayer Coatings in Fresh Papayas. *Coatings*, 9(11), 745. doi:10.3390/coatings9110745

Zhou, Y., Hu, L., Chen, Y., Liao, L., Li, R., Wang, H., Mo, Y., Lin, L., & Liu, K. (2022). The combined effect of ascorbic acid and chitosan coating on post-harvest quality and cell wall

metabolism of papaya fruits. Lebensmittel-Wissenschaft & Technologie, 171, 114134. doi: 10.1016/j.lwt.2022.114134

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Ethical Statement - Studies in humans and animals

- This study does not involve any animals or human subjects.




Fig. 1: Extraction of Chitosan from Crustaceans




Fig. 2: Antibacterial activity of chitosan against gram-negative (left) and gram positive (right) bacteria





Fig. 3: Factors leading to post-harvest losses in papaya

Graphical Abstract



Graphical Abstract

Table 1: Physiochemical composition of chitosan from different source origin

Parameters	Composition Value	References
Proximate Composition		
Moisture (wt %)	0.0004 – 9.23	Renuka <i>et al.</i> , 2019; Kumari <i>et al.</i> , 2017; Isa <i>et al.</i> , 2012; Ibitoye <i>et al.</i> , 2018; Islam <i>et al.</i> , 2022; Hossain & Iqbal, 2014; Ushakumari & Ramanujan, 2012
Ash (wt %)	0.03 – 11.77	Renuka <i>et al.</i> , 2019; Kumari <i>et al.</i> , 2017; Isa <i>et al.</i> , 2012; Puvvada <i>et al.</i> , 2012; Hossain & Iqbal, 2014; Ssekatawa <i>et al.</i> , 2021
Protein (wt %)	1.99 – 48.55	Renuka <i>et al.</i> , 2019; Kumari <i>et al.</i> , 2017; Isa <i>et al.</i> , 2012; Rout, 2001
Fibre (wt %)	8.74 – 55.8	Isa <i>et al.</i> , 2012; Miraftab <i>et al.</i> , 2011, Rout, 2001
Ultimate Composition		
C (wt %)	16.23 – 40.97	Isa <i>et al.</i> , 2012; Ibitoye <i>et al.</i> , 2018; Islam <i>et al.</i> , 2022
H (wt %)	6.51	Islam <i>et al.</i> , 2022
O (wt %)	34.86	Islam <i>et al.</i> , 2022
N (wt %)	2.71 – 8.50	Isa <i>et al.</i> , 2012; Ibitoye <i>et al.</i> , 2018; Ssekatawa <i>et al.</i> , 2021; Islam <i>et al.</i> , 2022; Rout, 2001
C/N (molar ratio)	5.23 – 7.79	Isa <i>et al.</i> , 2012; Ibitoye <i>et al.</i> , 2018; Kumari <i>et al.</i> , 2017; Islam <i>et al.</i> , 2022
Techno-functional Properties		
pH	7.9 – 8.5	Renuka <i>et al.</i> , 2019; Divya <i>et al.</i> , 2014; Puvvada <i>et al.</i> , 2012
Molecular Weight (KDa)	110.64 – 1200	Renuka <i>et al.</i> , 2019; Ssekatawa <i>et al.</i> , 2021; Puvvada <i>et al.</i> , 2012; Struszczyk <i>et al.</i> , 2002
Solubility (%)	48.30 – 98.40	Renuka <i>et al.</i> , 2019; Kumari <i>et al.</i> , 2017; Ssekatawa <i>et al.</i> , 2021; Hossain & Iqbal, 2014
Water Binding Capacity (%)	138.0 – 1150.0	Renuka <i>et al.</i> , 2019; Kumari <i>et al.</i> , 2017; Hossain & Iqbal, 2014; Rout, 2001; Knorr, 1982
Fat Binding Capacity (%)	104.0 – 331.28	Renuka <i>et al.</i> , 2019; Kumari <i>et al.</i> , 2017; Hossain & Iqbal, 2014; Rout, 2001; Knorr, 1982

Table 2: Studies on effect of chitosan edible coating in management of post-harvest losses in papaya

Coating Formulations Used	Papaya Form	Storage Conditions	Added Value	Effects and Potential Findings	References
T1: Chitosan Coating: 0.75% (w/v), T2: Beeswax Coating: 6% (w/v), T3: Control	Whole papaya fruit	Stored at room temperature (25-27 °C) till maturity stage six	Beeswax emulsion	<ul style="list-style-type: none"> • Keeping quality of papaya significantly increased when coating prepared with chitosan (11.33 days after harvest) and beeswax (12.33 days after harvest) was applied as compared to the untreated control group (7.67 days after harvest). • Papaya fruits that were coated with chitosan experienced delayed maturity until stage six, resulting in an extended shelf life, likely attributed to reduced rates of respiration and transpiration. • Chitosan coated fruit had lower weight loss (6.76%) as compared to the control group (8.54%) and maintained the hardness and TSS values of papaya. • Papaya coated with chitosan had a better and smooth appearance and were free from fungi due to the antimicrobial properties of chitosan, which effectively inhibits the growth of spoilage microorganisms. 	Mukdisari <i>et al.</i> , 2016

<p>T1: Control (distilled water); T2: AsA: 1.5% (w/v) ascorbic acid; T3: CTS: 1.0% (w/v) chitosan; T4: AsA + CTS: 1.5% (w/v) ascorbic acid + 1.0% (w/v) chitosan</p>	<p>Green and mature whole papaya fruit</p>	<p>Stored at 20 °C and 80 –90% relative humidity for 0, 4, 8, 12, and 16 days</p>	<p>Ascorbic acid</p>	<ul style="list-style-type: none"> • Chitosan and ascorbic acid (AsA + CTS) combined coating significantly reduced weight loss (21.12 – 25.29%) and increased firmness (64.32 – 619.16%) during the storage of papaya as compared to control. • Decreased L*, a* and b* values were observed with the application of AsA + CTS coating, which indicated delayed fruit senescence and browning • Application of AsA + CTS increased the pH of papaya fruits by 4.36 –11.73%, reduced TSS (3.45 - 4.90%) and increased TA (12.70 - 15.80%) as compared to CTS. • AsA + CTS treatments significantly decreased the ripening index by 19.15 - 34.00% as compared to control. • Reducing sugar content decreased (5.23 – 13.79%) and soluble sugar content increased (4.06 - 4.81%) with AsA + CTS treatments during the storage days as compared to the control • AsA + CTS treatment significantly reduced 	<p>Zhou <i>et al.</i>, 2022</p>
--	--	---	----------------------	---	---------------------------------

				<p>H₂O₂ content (26.23 – 66.26%) and MDA levels (28.06 %– 43.36%) at all storage dates as compared to control</p> <ul style="list-style-type: none"> • Inhibition of softening enzyme activity and gene expression level of cell wall degrading enzymes such as CX, PG, PME, and β-GAL. 	
<p>T1- Control T2- Chitosan (1% w/v) + tuna fish gelatin solution; T3- chitosan (1% w/v) + tuna fish gelatin solution + black tea extract (5% w/v); T4- chitosan (1% w/v) + tuna fish gelatin solution + black tea extract (10% w/v); T5- chitosan (1% w/v) + tuna fish gelatin solution + black tea extract (15% w/v)</p>	Minimally processed papaya fruit	Storage durations (1, 4, 7, and 10 days)	Tuna skin gelatin and black tea extract	<ul style="list-style-type: none"> • The decrease in hardness occurred very quickly during the storage period in T1 sample (0.35 Nm on day 1 and 0.13 Nm on day 10) as compared to the T3 sample (0.19 Nm on day 1 and 0.14 Nm on day 10). • The lowest weight loss was observed in T5 samples (1.91-4.61%) as compared to T1 samples (1.88-6.02%) due to the presence of gelatin and chitosan-based coatings, which act as a barrier to water vapor in minimally processed fruits, reducing the mass loss during storage. • The increase in TSS content in T5 fruit samples was higher (8.83 °Brix) as compared to T1 (8.33 °Brix) and this could be due to the influence 	Sekarina <i>et al.</i> , 2023

				<p>of edible coating layer dissolving during the test.</p> <ul style="list-style-type: none">• The antioxidant activity was found to be slightly higher in T5 samples (59.26-72.62%) as compared to T1 samples (66.97-72%) and this may be due to the presence of tea polyphenols and chitosan as both have strong antioxidant properties especially in terms of the ability to scavenge hydroxyl radicals and reactive oxygen species.• Microbial growth in control sample was faster (7.38 cfu/ml) than in the edible coating treatment sample with the addition of black tea extract (6.97-7.42 cfu/ml) and this may be due to the presence of tea polyphenols and chitosan, which can interact with negatively charged residues on the surface of microbial cells by an electrostatic attraction so that it flocculates and adsorbs on the surface of microorganisms, then inhibits the physiological metabolism of	
--	--	--	--	--	--

				microorganisms and finally inhibits the growth of microorganisms.	
T1- Control; T2- Chitosan coating (5g/L) after treatment with hot water; T3- Chitosan coating (10g/L) after treatment with hot water; T4- Chitosan coating (15g/L) after treatment with hot water; T5- Chitosan coating (20g/L) after treatment with hot water	Whole papaya fruit	Stored for 28 days at 10 °C + 7 days of shelf-life at 20 °C (35 days)	Hot Water Treatment	<ul style="list-style-type: none"> • Gradual increase in chitosan concentration in the coating showed better preserving capabilities. • Weight loss in highest concentration of chitosan (T5) was around 5.2% whereas in the case of control (T1) it was around 8.4% hence showing a significant rate of decrease in weight loss as the concentration of coating increases. • Hue angle and colour tests indicate that chitosan coating has a better effect on retaining colour of papaya as hue angles loss was observed to be lower in (T5) as compared to (T1) • Loss of firmness and textural loss was observed to be less in (T5) as it had the highest concentration of chitosan. • Increase in Total soluble solids value (TSS) in control at 28 days was observed. In the case of T5 the increase percent was lower at 28days which 	Vilaplana <i>et al.</i> , 2019

				indicates delayed ripening. Hence increasing the desirability.	
T1- Control; T2- Chitosan coating layer: 0.5% chitosan solution(w/v); T3- Pullulan coating layer: 0.5% pullulan solution (w/v); T4- Two layer coating: 0.5% pullulan solution (w/v) + 0.5% chitosan solution (w/v); T5- Four layer coating: 0.5% pullulan solution (w/v) + 0.5% chitosan solution (w/v); T6- Six layer coating by 0.5% pullulan solution (w/v) + 0.5% chitosan solution (w/v)	Whole and fresh papaya fruits	Stored at 25 °C, 50% relative humidity up to 14 days	Pullulan coating	<ul style="list-style-type: none"> • Weight loss in control was observed to be 28.86% at 14th day whereas for two layers, four layers and six layers coating, the percentage was 15.53%, 10.41% and 13.85% respectively. Hence 4 layer coating showed the best results in preventing weight loss. • Loss of firmness was observed in control papaya as it got completely softened at 14th day. 4 layer papaya gave the best results in fruit firmness meter which was 20.2 N whereas for 5 layer and 6 layer coating it was 15.6 N and 16.9 N respectively. • Colour being the indicator which represents maturity of papaya was analysed using b* value of spectrophotometer and the control showed highest value at 14th day (51.8) on the other hand 4 layer coating (T5) showed the lowest value (37.6) which 	Zhang <i>et al.</i> , 2019

				<p>ensured the delaying of undesired ripening.</p> <ul style="list-style-type: none"> • Anti-bacterial properties of chitosan decreased the respiratory rate in which the 4 layer coating was the most stable one. • Vitamin-C content showed a significant decrease in control papaya (15.06 mg/kg) and it was highest in 4 layer coated papaya (26.17mg/kg). 	
T1- Control; T2- Chitosan coating (molecular weight: 150 kDa); T3- Chitosan coating (molecular weight: 300 kDa)	Whole papaya fruits	Stored at ambient temperature (18 - 24 °C) with air relative humidity ranging from 60-80% for 20 days	-	<ul style="list-style-type: none"> • Chitosan (150 kDa) showed higher moisture content in storage condition (8.5± 0.3%) as compared to 300 kDa chitosan (6.0± 0.1%). Hence chitosan with more molecular weight showed better response towards preventing moisture gain. • Chitosan (150 kDa) showed lower log values for mesophilic bacteria, yeasts and molds as compared to uncoated and 300 kDa chitosan as it has a less organized structure with lower crystallinity, resulting in a more homogeneous coating which is desirable since it provides a more effective barrier 	Dotto <i>et al.</i> , 2015

				<p>between the fruit and the external environment.</p> <ul style="list-style-type: none"> • Shelf life study of the samples show that the chitosan (150 kDa) is most suitable among the three formulations as it shows better anti-microbial activity and can extend the shelf life of papaya about 4 days (for mesophilic bacteria) and about 7 days (for yeasts and molds). 	
<p>T1- Control coating: 1% (w/v) chitosan; T2- Coating treatment with 1% (w/v) chitosan + 5% (w/v) ethanolic extract of propolis (EEP)</p>	<p>Whole green coloured papaya fruits</p>	<p>Stored at a temperature of $25\pm 3^{\circ}\text{C}$ with relative humidity between 65-70% for 8 to 9 days</p>	<p>Propolis Extract</p>	<ul style="list-style-type: none"> • The fruits covered with EEP + chitosan (T2) demonstrated a reduced deterioration index and infection diameter of the fungus <i>Colletotrichum gloeosporioides</i>, as compared to the control (T1) papayas. • The control fruits showed a rapid decay during 4th day of storage hence increasing their deterioration index. Treated fruits were damage free till 4th day and then at the end of storage period the T2 showed only moderate damage. • The treated fruits presented a higher content of anthocyanins (1.4 mg/100g) in 	<p>Barrera <i>et al.</i>, 2015</p>

				<p>comparison with the control (0.9 mg/100g) indicating a possible retardation of maturity.</p> <ul style="list-style-type: none"> • The treated fruits demonstrated a higher concentration of chlorophyll (3.2 - 4.5 mg/g) at the end of storage, compared to the control fruits (4.7 - 7.2 mg/g), confirming that the fruits covered with EEP + chitosan presented a slower pigmentation metabolism. 	
<p>T1- Control (Uncoated); T2- Chitosan-Oxidized Starch coating (1:3 w/w): chitosan solution (1% w/v) + oxidized starch slurry (3.5% w/v)</p>	<p>Ripe yellow coloured whole papaya fruit</p>	<p>Stored at room temperature (25 ± 1 °C) for 15 days</p>	<p>Oxidized starch</p>	<ul style="list-style-type: none"> • Coating has a significant effect on the papaya fruit in terms of weight loss because results suggest that there was no significant weight change in coated papaya during the 15 day period but the uncoated papaya showed an increasing trend in weight loss percentage. • Coated papaya (T2) showed a firmness reduction of 47.3% whereas the uncoated papaya (T1) showed 92.02% firmness loss. • The ripening stage of uncoated papaya was attained at Day 5 but in the case of coated papaya the same stage was attained at Day 15, 	<p>Escamilla-García <i>et al.</i>, 2018</p>

				<p>indicating delay in ripening.</p> <ul style="list-style-type: none"> • Log value for total coliforms, yeasts, mesophilic bacteria and fungi showed a distinguishable lead in case of uncoated papaya whereas slower growth of these microorganisms were observed in coated papaya. 	
<p>T1- Control; T2- Chitosan 0.5% (w/v); T3- Chitosan 1.0% (w/v); T4- Chitosan 1.5% (w/v); T5- Chitosan 2.0% (w/v); T6- Chitosan 2.5% (w/v); T7- Chitosan 3.0% (w/v);</p>	<p>Fresh whole papaya fruit</p>	<p>Stored under two conditions: cold storage ($12\pm 1^{\circ}\text{C}$) for 23 days, ambient temperature ($28\pm 1^{\circ}\text{C}$) for 9 days</p>	<p>–</p>	<ul style="list-style-type: none"> • T1 experienced maximum weight loss (9.90% at ambient storage and 8.97% at cold storage) during 7 and 18 days of storage respectively, while T7 lost minimum weight (5.1% at ambient and 4.5% at cold storage) during 9 and 23 days of storage • Papaya fruits treated with chitosan were recorded upto 48.15% lesser change in colour than untreated fruits was evident from the colour (L, a and b) values at the end of storage. • The moisture content of T1 decreased at faster rate and recorded 86.23% (ambient storage) and 85.00% (cold storage) during 7 and 18 days of storage, respectively while 	<p>Bhanushree <i>et al.</i>, 2018</p>

				<p>moisture content of T7 registered slower decrease and registered highest moisture content of 87.23 % (ambient storage) and 88.25% in cold storage during 9 and 23 days of storage, respectively.</p> <ul style="list-style-type: none">• T1 fruits recorded lesser sensory scores while T7 fruits had higher score of overall acceptability of 7.92 (ambient storage) during 7 days of storage and 8.85 (cold storage) during 18 days of storage.• T7 fruits showed an extended storage period up to 9 days (ambient storage) and 23 days (cold storage) as compared to T1 fruits (7 days at ambient and 18 days at cold storage) due to the delay in the ripening process.	
--	--	--	--	---	--