

Evidence for Homodimerization of the c-Fos Transcription Factor in Live Cells Revealed by Fluorescence Microscopy and Computer Modeling

Nikoletta Szalóki,^a Jan Wolfgang Krieger,^b István Komáromi,^{c,d} Katalin Tóth,^b György Vámosi^{a,e}

Department of Biophysics and Cell Biology, Research Center for Molecular Medicine, University of Debrecen, Debrecen, Hungary^a; German Cancer Research Center (DKFZ), Biophysics of Macromolecules, Heidelberg, Germany^b; HAS-UD Vascular Biology, Thrombosis and Hemostasis Research Group, Hungarian Academy of Sciences, Debrecen, Hungary^c; Division of Clinical Laboratory Science, Department of Laboratory Medicine, Faculty of Medicine, University of Debrecen, Debrecen, Hungary^d; Sanford Burnham Prebys Medical Discovery Institute at Lake Nona, Orlando, Florida, USA^e

The c-Fos and c-Jun transcription factors, members of the activator protein 1 (AP-1) complex, form heterodimers and bind to DNA via a basic leucine zipper and regulate the cell cycle, apoptosis, differentiation, etc. Purified c-Jun leucine zipper fragments could also form stable homodimers, whereas c-Fos leucine zipper homodimers were found to be much less stable in earlier *in vitro* studies. The importance of c-Fos overexpression in tumors and the controversy in the literature concerning c-Fos homodimerization prompted us to investigate Fos homodimerization. Förster resonance energy transfer (FRET) and molecular brightness analysis of fluorescence correlation spectroscopy data from live HeLa cells transfected with fluorescent-protein-tagged c-Fos indicated that c-Fos formed homodimers. We developed a method to determine the absolute concentrations of transfected and endogenous c-Fos and c-Jun, which allowed us to determine dissociation constants of c-Fos homodimers ($K_d = 6.7 \pm 1.7 \mu\text{M}$) and c-Fos–c-Jun heterodimers (on the order of 10 to 100 nM) from FRET titrations. Imaging fluorescence cross-correlation spectroscopy (SPIM-FCCS) and molecular dynamics modeling confirmed that c-Fos homodimers were stably associated and could bind to the chromatin. Our results establish c-Fos homodimers as a novel form of the AP-1 complex that may be an autonomous transcription factor in c-Fos-overexpressing tissues and could contribute to tumor development.

Activator protein 1 (AP-1) is a transcriptional regulator composed of members of the Fos, Jun, and ATF families of DNA-binding proteins (1, 2). c-Fos and c-Jun regulate a variety of processes, including proliferation, differentiation, apoptosis, and oncogenesis (3). They function as dimers binding to the promoter/enhancer regions of numerous mammalian genes (4). Their DNA-binding domain is composed of a leucine zipper promoting dimerization and a basic region that binds with high affinity to a specific 8-bp-long DNA sequence (5, 6).

In addition to forming stable heterodimers with c-Fos (7–9), c-Jun can also homodimerize, as revealed *in vitro* by electrophoretic mobility shift assay (EMSA) (8), and bind to DNA as a homodimer, although with lower affinity than the heterodimer (8, 10). In contrast, the c-Fos homodimer was found to be unstable *in vitro*, and thus, c-Fos has been thought to interact with DNA only by forming heterodimers with c-Jun (9, 11, 12). The instability of the c-Fos dimer is thought to be due to repulsion between its negatively charged residues in the leucine zipper (6). The wild-type c-Fos zipper showed no homoassociation at a concentration of 0.1 μM according to EMSA (13). O’Shea and coworkers estimated the K_d of the c-Fos leucine zipper homodimer to be 3.2 and 5.6 μM at 0 and 25°C, implying that the failure of others to detect c-Fos dimerization was probably due to low protein concentrations (14). It was shown by EMSA that a single amino acid change in the leucine zipper is sufficient to allow a truncated c-Fos protein to homodimerize and bind to its DNA response element (15). Melting temperature analyses of different leucine zipper dimers revealed that thermal stability increases from c-Fos–c-Fos through c-Fos–c-Jun to c-Jun–c-Jun (16).

c-Fos expression and activation can be induced by growth factors, cytokines, or neurotransmitters via G-protein-coupled re-

ceptor-, mitogen-activated protein kinase-, cyclic AMP-, or Ca^{2+} -dependent signaling pathways (17–19). c-Fos overexpression occurs in several pathological conditions, which can have both proliferative and antiproliferative effects. c-Fos was overexpressed in some tamoxifen-resistant human breast tumors (20) and highly overexpressed in malignant oral tissues (21). It could also contribute to hepatocarcinogenesis (22). In a murine skin carcinogenesis model, c-Fos was shown to be required for malignant tumor conversion (23). c-Fos can be upregulated via the thyroid hormone nuclear receptor $\alpha 1$, which is a tumor inducer in intestinal tumorigenesis (24). Conversely, c-Fos overexpression inhibited cell cycle progression and stimulated cell death in hepatocytes (25). It also activated apoptosis in colorectal carcinoma cells in a p53-dependent manner (26).

Because c-Fos, but not c-Jun, is overexpressed in many different types of tumors, we were interested whether c-Fos at higher concentrations could form stable homodimers and bind to DNA

Received 9 April 2015 Returned for modification 13 May 2015

Accepted 20 August 2015

Accepted manuscript posted online 24 August 2015

Citation Szalóki N, Krieger JW, Komáromi I, Tóth K, Vámosi G. 2015. Evidence for homodimerization of the c-Fos transcription factor in live cells revealed by fluorescence microscopy and computer modeling. *Mol Cell Biol* 35:3785–3798. doi:10.1128/MCB.00346-15.

Address correspondence to György Vámosi, vamosig@med.unideb.hu.

K.T. and G.V. are equal senior authors.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/MCB.00346-15>.

Copyright © 2015, American Society for Microbiology. All Rights Reserved.




FIG 1 Schematic drawing of c-Fos, its mutant forms, and c-Jun. From the top: full-length Fos with fluorescent protein tag at the C terminus, C-terminally truncated Fos²¹⁵ and Fos^{ΔΔ} where the DNA-binding and dimerization domains were deleted, and Jun. Pink denotes the DNA-binding domain, yellow denotes the leucine zipper, and the dotted line denotes the linker between Fos/Jun and the fluorescent-protein tag (ECFP, EYFP, EGFP, or mRFP1).

in live cells. Förster resonance energy transfer (FRET) can be used to assess distances between two fluorophores in the range of 2 to 10 nm (27, 28), whereas fluorescence cross-correlation spectroscopy (FCCS) can demonstrate the comobility of two molecules (29–31). Using these methods, we previously demonstrated heterodimerization and chromatin binding of c-Fos and c-Jun and described the conformation of their complex in live cells (7, 32). It was shown in our lab (German Cancer Research Center) by imaging FCCS that mobility and protein-protein interaction maps of c-Fos and c-Jun were correlated (33).

Here we performed FRET measurements of fluorescent protein-tagged c-Fos molecules by confocal microscopy and flow cytometry to examine whether c-Fos could form homodimers. We developed a method combining fluorescence correlation spectroscopy (FCS) and immunofluorescence to assess the concentrations of both fluorescently labeled and unlabeled endogenous c-Fos and c-Jun in cells. This allowed us to determine the K_d of c-Fos homodimers and c-Fos–c-Jun heterodimers in live HeLa cells by FRET titrations. We found that the K_d of the c-Fos homodimer is more than 1 order of magnitude higher than that of the heterodimer. To our knowledge, this is the first report on the determination of the K_d of transcription factors from FRET titrations in live cells. Imaging FCCS measurements revealed codiffusion of stable c-Fos homodimers and their binding to chromatin. Our molecular dynamics (MD) simulations support the notion that Fos homodimers can form, bind to DNA, and remain stable over the time span of the simulation (500 ns). This novel homodimeric form of c-Fos may act as an autonomous transcriptional regulator.

MATERIALS AND METHODS

Cell culture, plasmid construction, and transfection of HeLa cells. Cell culture, plasmid construction, and transfection have been described elsewhere (34). For detailed information on these procedures and the plasmids and proteins used in this study, see the supplemental material and Fig. 1.

Confocal microscopic and flow cytometric FRET analyses. Confocal microscopic images were collected with a Zeiss LSM 510 microscope. Flow cytometric measurements were carried out with a Becton Dickinson FACSAria III instrument. For details of data acquisition and FRET analysis on a pixel-by-pixel or cell-by-cell basis, which have been described earlier (34), see the supplemental material.

FCS and calibration of fluorescence intensity as a measure of absolute concentration. In FCS (35), molecules diffuse across a subfemtoliter ($<1 \mu\text{m}^3$) detection volume defined by a focused laser beam. This causes

fluorescence fluctuations, which are analyzed to derive dynamic parameters of the molecules studied. FCS measurements were performed with a modified Olympus FluoView 1000 confocal microscope mounted on an inverted IX-81 stand with an UPlanAPO 60 \times numerical aperture 1.2 water immersion objective. The FCS extension (Steinbeis Transfer Unit for Biophysical Analytics, Heidelberg, Germany) equipped with two avalanche photodiodes (Perkin-Elmer, Wellesley, MA) is attached to the confocal scanning unit. Fluorescence of enhanced green fluorescent protein (EGFP) was excited by the 488-nm line of an Ar ion laser and detected at 500 to 550 nm. To allow FCS measurements at high EGFP concentrations (up to 25 μM), laser illumination was dimmed by a neutral-density filter (optical density of 1) to yield $P = 0.2 \mu\text{W}$ at the sample. FCS measurements of live HeLa cells were performed with eight-well chambered cover glass plates (Nunc). Points for FCS measurements were selected from confocal images. In each sample, ~ 30 cells were measured at room temperature and six 8-s runs per cell were recorded. Fluorescence autocorrelation functions (ACFs) were calculated online by an ALV-5000E hardware correlator card (ALV Laser, Langen, Germany). Autocorrelation curves were fitted to a two-component three-dimensional (3D) diffusion model with triplet correction by the program QuickFit 3.0 (36) as follows:

$$G(\tau) = \frac{1 - T + T e^{-\tau/\tau_{tr}}}{1 - T} \frac{1}{N} \left[r_1 \left(1 + \frac{\tau}{\tau_1} \right)^{-1} \left(1 + \frac{\tau}{S^2 \tau_1} \right)^{-1/2} + r_2 \left(1 + \frac{\tau}{\tau_2} \right)^{-1} \left(1 + \frac{\tau}{S^2 \tau_2} \right)^{-1/2} \right] \quad (1)$$

where τ is the lag time, T denotes the triplet fraction, τ_{tr} is the triplet correlation time, τ_1 and τ_2 are the diffusion times of the fast and slow species (average dwell times of molecules in the detection volume), r_1 and $r_2 = 1 - r_1$ are the fractional amplitudes of the two components, N is the average number of molecules in the detection volume, and S is the aspect ratio of the ellipsoidal detection volume. To assess the aggregation state of EGFP-labeled proteins, the molecular brightness or fluorescence per particle, F/N , was calculated and compared with that of monomeric EGFP.

To facilitate K_d determinations, we developed a method to assess absolute concentrations from fluorescence intensity similar to that described in reference 37. In the first step, we determined the detection volume, V_{eff} , of the microscope by using a 130 nM Alexa Fluor 488 solution as a standard. From its ACF, τ_d and S were determined by fitting and the lateral radius, ω_{xy} , and the axial radius, ω_z , were calculated with the following equation:

$$\omega_{xy} = \sqrt{4D\tau_d}, \quad \omega_z = \omega_{xy} S, \quad (2)$$

where τ_d is the measured diffusion time of the dye, which measures mobility and is inversely proportional to the diffusion coefficient and $D = 435 \mu\text{m}^2/\text{s}$ is the diffusion coefficient of Alexa Fluor 488 at 22.5°C (38). The effective detection volume is calculated with the following equation:

$$V_{\text{eff}} = \pi^3 \omega_{xy}^2 \omega_z \quad (3)$$

From autocorrelation curves with EGFP, the particle numbers, N , were determined. From these, molar concentrations, c , were calculated as follows:

$$c = N / (N_A V_{\text{eff}}) \quad (4)$$

where N_A is Avogadro's number. Before every FCS measurement, the fluorescence intensity, F , was measured with the imaging detector (photomultiplier tube) of the confocal microscope at the site of FCS measurement; thus, c -versus- F calibration lines were generated (see Fig. 6B).

To facilitate the comparison of measurements on different days, we used fluorescent beads. The fluorescence intensity of 6- μm green calibration beads (beads with 1% relative intensity from the InSpeck green microscope image intensity calibration kit [Molecular Probes, Life Technologies, Inc.]) was used to normalize EGFP fluorescence. Confocal sections in the equatorial plane of the beads were recorded on the same day as the FCS calibration with identical instrument settings. The average fluorescence intensity per pixel in the central area of the beads was obtained.

With the calibration curve (see Fig. 6B, triangles) the local intensity of the bead at its center corresponded to an EGFP concentration of $c_{\text{bead unit}}^{\text{confocal}} = \sim 15.4 \pm 0.7 \mu\text{M}$ (average \pm standard error of the mean [SEM] of five experiments).

We could also transfer the concentration calibration to flow cytometric measurements. In the microscopic calibration described above, the local fluorescence intensities at a pixel of the sample and the bead are compared. In contrast, in flow cytometry, the total intensity of the whole cell and the bead is measured. Therefore, the ratio of the cellular and bead volumes had to be taken into account. The volume of HeLa nuclei (where Fos and Jun are localized) is ~ 13.6 times larger than that of the beads, as determined by confocal microscopic 3D sectioning and the Imaris software (Bitplane AG, Zurich, Switzerland). In addition, the different detection efficiencies of the spectra of the bead and EGFP arising from different band-pass filters in the flow cytometer and the confocal microscope resulted in a factor of 13. Taking these factors into account, the total intensity of a bead corresponded to an EGFP concentration of $c_{\text{bead unit}}^{\text{flow cyt}} = \sim 1.2 \mu\text{M}$ (distributed in a HeLa cell nucleus) in flow cytometric experiments. If the localization of the protein is not perfectly nuclear, we can correct for this as well. From confocal microscopic sectioning, we determined the nuclear and cytoplasmic fractions of Fos²¹⁵-EGFP (the protein we used for FRET titrations), which were $85\% \pm 5\%$ and $15\% \pm 5\%$, independent of the expression level (see Fig. S3 in the supplemental material). With this correction, a bead unit of $\sim 1.0 \pm 0.1 \mu\text{M}$ was used to calculate the nuclear concentration of Fos²¹⁵-EGFP in flow cytometric experiments.

Determination of the absolute concentration of endogenously expressed Fos and Jun. With regular immunofluorescence assays, only the relative amounts of endogenous and transfected proteins can be assessed. By knowing the absolute concentration of the transfected proteins, the endogenous concentration can also be determined. Therefore, we combined the immunofluorescence assay with the results of FCS-based EGFP concentration calibration to assess the endogenous concentrations of Fos and Jun in HeLa cells.

Immunofluorescence labeling was carried out as follows. After being washed three times with PBS, cells were fixed with 3.7% formaldehyde (4°C, 10 min), permeabilized with 0.25% Triton and 0.1% Tween-Tris-buffered saline (TBS) at room temperature for 30 min, and blocked with 2% bovine serum albumin with 0.1% Tween-TBS at room temperature for 30 min. Cells were incubated with a mouse anti-c-Fos (Merck, Whitehouse Station, NJ) or anti-c-Jun (Millipore, Billerica, MA) monoclonal antibody at 20 $\mu\text{g/ml}$ for 1 h at room temperature and then incubated with the secondary polyclonal antibody NL-637-DAMIG (R&D Systems, Minneapolis, MN) at 50 $\mu\text{g/ml}$ for 1 h at room temperature in the dark. Between consecutive steps, cells were washed three times with PBS. Titration of antibodies showed that these concentrations were close to saturating values (see Fig. S4A to C in the supplemental material).

Flow cytometric measurements were performed on a FACSAria III flow cytometer. The green EGFP signal (I^{green}) was excited at 488 nm, and emission was detected at 515 to 545 nm, while the red NL637 signal (I^{red}) was excited at 633 nm and emission was detected through a 655-nm low-pass filter.

The amount of Fos-EGFP was determined by comparing its I^{green} fluorescence signal to that of the green bead used for concentration calibration. The red signal, I^{red} , of the NL637-DAMIG antibody used for immunofluorescence labeling is proportional to the total amount of Fos, i.e., the endogenous Fos in the nontransfected sample and the endogenous Fos plus Fos-EGFP in the transfected one. The concentrations of endogenous Fos ($c_{\text{Fos-endogenous}}$) and Fos-EGFP ($c_{\text{Fos-EGFP}}$) were calculated from the green and red signals with transfected and nontransfected cells as follows:

$$c_{\text{Fos-EGFP}} = \frac{I_{\text{transfected}}^{\text{green}}}{I_{\text{bead}}^{\text{green}}} \times c_{\text{bead unit}}^{\text{flow cyt}} \quad (5)$$

$$\frac{c_{\text{Fos-EGFP}} + (c_{\text{Fos-endogenous}})}{(c_{\text{Fos-endogenous}})} = \frac{I_{\text{transfected}}^{\text{red}}}{I_{\text{nontransfected}}^{\text{red}}} \quad (6)$$

$$(c_{\text{Fos-endogenous}}) = \frac{c_{\text{Fos-EGFP}}}{\left(\frac{I_{\text{transfected}}^{\text{red}}}{I_{\text{nontransfected}}^{\text{red}}}\right) - 1} = \frac{\left(\frac{I_{\text{transfected}}^{\text{green}}}{I_{\text{bead}}^{\text{green}}}\right) \times c_{\text{bead unit}}}{\left(\frac{I_{\text{transfected}}^{\text{red}}}{I_{\text{nontransfected}}^{\text{red}}}\right) - 1} \quad (7)$$

with $c_{\text{bead unit}}^{\text{flow cyt}} = \sim 1.2 \mu\text{M}$ in our measurements. I^{red} signals were collected from fixed cells (required by the immunofluorescence labeling protocol), whereas I^{green} signals were measured in nonfixed live cells (from the same transfected population) to avoid deterioration of EGFP fluorescence because of fixation. The concentrations of endogenous and EGFP-tagged Jun were determined by the same principle.

Calculation of dissociation equilibria from FRET data. To assess the K_d s of Fos homodimers and Fos-Jun heterodimers, we carried out FRET titration experiments. In the derivation of dissociation equilibria, we assumed that the heterodimer was more stable than the Fos homodimer. Therefore, in the case of Fos-Jun association, we neglected the presence of Fos homodimers (assuming that they were not present at lower concentrations). The law of mass action for heterodimer formation is calculated with the following equation:

$$[F][J]/[FJ] = K_d^{\text{FJ}} \quad (8)$$

where brackets denote concentrations of free monomers and heterodimers and K_d^{FJ} is the dissociation constant of the heterodimeric complex. The total concentration of Fos or Jun can be written as the sum of the concentrations of free monomers F and J and heterodimers FJ as follows:

$$[F]_t = [F] + [FJ]; [J]_t = [J] + [FJ] \quad (9)$$

The concentration of the heterodimer is calculated with the following equation:

$$[FJ] = \frac{1}{2} \left\{ [F]_t + [J]_t + K_d^{\text{FJ}} - \sqrt{[F]_t^2 - 2[F]_t \times ([J]_t - K_d^{\text{FJ}}) + ([J]_t + K_d^{\text{FJ}})^2} \right\} \quad (10)$$

The measured FRET efficiency of EGFP- and mRFP1-labeled proteins is an average value stemming from FRET-producing and non-FRET-producing donor molecules. Free donors or donors associated with endogenous unlabeled protein give zero FRET. Only donors forming a complex with an acceptor make a positive contribution to FRET (Fig. 2). Thus, we need to calculate the concentration of Fos-Jun dimers labeled with both a donor and an acceptor. The total concentrations of Fos and Jun are calculated as follows:

$$[F]_t = [F_D]_t + [F_e]_t \text{ and } [J]_t = [J_A]_t + [J_e]_t \quad (11)$$

where the indexes *D*, *A*, and *e* refer to donor-tagged, acceptor-tagged, and endogenous molecules. The fractions of donor-tagged Fos (p_D) and acceptor-tagged Jun (p_A) are calculated as follows:

$$p_D = \frac{[F_D]_t}{[F_D]_t + [F_e]_t} \text{ and } p_A = \frac{[J_A]_t}{[J_A]_t + [J_e]_t} \quad (12)$$

The concentration of doubly labeled Fos_D-Jun_A dimers is calculated as follows:

$$[F_D J_A] = [FJ] \times p_D \times p_A = [FJ] \times \frac{[F_D]_t}{[F_D]_t + [F_e]_t} \times \frac{[J_A]_t}{[J_A]_t + [J_e]_t} \quad (13)$$

We denote the FRET efficiency in the complex of a single donor-tagged Fos and an acceptor-tagged Jun by E_0 . The measured apparent FRET efficiency, E_{meas} , can be written as follows:

$$E_{\text{meas}} = \frac{[F_D J_A]}{[F_D]_t} \times E_0 + \frac{[F_D]_t - [F_D J_A]}{[F_D]_t} \times 0 = \frac{[F_D J_A]}{[F_D]_t} \times E_0 \quad (14)$$

where $[F_D J_A]$ is the concentration of complexes of donor-tagged Fos with acceptor-tagged Jun, and $[F_D]_t$ is the total concentration of donor-tagged Fos (without respect to being monomeric or in a Fos-Jun complex).

$[F_D]_t - [F_D]_A$ is the concentration of donor-tagged Fos not complexed with acceptor-tagged Jun (Fos-EGFP in monomeric form or complexed with endogenous Jun), contributing zero FRET efficiency. By introducing the acceptor-to-donor expression ratio $N_A/N_D = [J_A]/[F_D]$ (see equation S8 in the supplemental material) and combining equations 10, 13, and 14, the measured FRET efficiency can be expressed as follows:

$$E_{\text{meas}} = \frac{[F]_t + [J]_t + K_d^{FJ}}{\sqrt{[F]_t^2 - 2[F]_t([J]_t - K_d^{FJ}) + ([J]_t + K_d^{FJ})^2}} \times [F_D]_t \frac{N_A}{N_D} E_0 \quad (15)$$

By substituting equation 11 and the expression for the N_A/N_D ratio into equation 15, we get the formula used in the nonlinear fit (see equation S12 in the supplemental material) with variables $[F_D]_t$ and N_A/N_D .

To calculate Fos-Fos equilibria, we have to take Fos-Jun formation into account as well. Since the heterodimer is more stable, we make the simplifying assumption that all of the Jun molecules present are in complex with Fos at the high Fos concentrations where Fos homodimerization takes place, leaving no free Jun. We can write the following equilibrium equations:

$$\begin{aligned} [F][F]/[FF] &= K_d^{FF}, \\ [J]_t &= [FJ], \text{ and } [F]_t = [F] + [FJ] + 2[FF] \end{aligned} \quad (16)$$

where K_d^{FF} is the dissociation constant of the Fos homodimer and FF and FJ denote the homo- and heterodimer. The amount of Fos homodimer can be expressed as follows:

$$[FF] = \frac{1}{8} \left\{ 4[F]_t - 4[J]_t + K_d^{FF} - \sqrt{8K_d^{FF}[F]_t - 8K_d^{FF}[J]_t + (K_d^{FF})^2} \right\} \quad (17)$$

Fos molecules can be labeled with a donor or an acceptor or can be unlabeled, and only homodimers containing both a donor and an acceptor will produce FRET (Fig. 2). The fraction of such double-labeled pairs follows a multinomial distribution and is calculated as follows:

$$p_{D,A} = 2p_D p_A \quad (18)$$

where $p_D = [F_D]/([F_D]_t + [F_A]_t + [F_e]_t)$ and $p_A = [F_A]/([F_D]_t + [F_A]_t + [F_e]_t)$ are the donor- and acceptor-tagged fractions of Fos. The measured FRET efficiency equals the following:

$$E_{\text{meas}} = \frac{[F_D]_t [F_A]_t}{[F_D]_t} \times E_0 = \frac{[FF] p_{D,A}}{[F_D]_t} \times E_0 \quad (19)$$

Combining equations 17 and 19 (see also equation S14 in the supplemental material) yields the following equation:

$$E_{\text{meas}} = \frac{1}{4} \left(4[F]_t - 4[J]_t + K_d^{FF} - \sqrt{8K_d^{FF}[F]_t - 8K_d^{FF}[J]_t + (K_d^{FF})^2} \right) \frac{[F_D]_t \frac{N_A}{N_D} E_0}{[F]_t} \quad (20)$$

with

$$[F]_t = [F_D]_t + [F_A]_t + [F_e]_t = [F_D]_t \left(1 + \frac{N_A}{N_D} \right) + [F_e]_t \quad (21)$$

where $[F]_t$ is the total Fos concentration (including donor-tagged, acceptor-tagged, and endogenous Fos, without respect to a monomeric or dimeric state) and E_0 is the FRET efficiency of a donor-acceptor pair (this may be different from the E_0 of the heterodimer). These equations were used to determine the K_d s of dimers from flow cytometric FRET experiments by nonlinear regression using Prism (GraphPad Software, Inc). In the analysis we also considered the presence of dark acceptor species due to imperfect maturation and a cytoplasmic fraction of Fos




FIG 2 Possible combinations of fluorescently tagged and endogenous Fos and Jun. (A) In the monomer-heterodimer equilibrium, fluorescently tagged and endogenous, unlabeled Fos and Jun molecules participate. The three species containing a donor tag contribute to the measured value of FRET efficiency, E_{meas} , i.e., the doubly labeled heterodimer having a FRET efficiency of E_0 and donor-labeled Fos in complex with endogenous Jun or present as a monomer; the latter two species are characterized by zero FRET efficiency. The fractions of the different heterodimers follow a multinomial distribution. E_{meas} is a weighted average of the species-specific E values (see equation S12 in the supplemental material). (B) In the monomer-homodimer equilibrium, donor-tagged, acceptor-tagged, and endogenous Fos and endogenous Jun participate. Four heterodimeric species and the donor-tagged monomer contribute to E_{meas} (see equation S14 in the supplemental material).

(see the supplemental material). These factors influence E_0 but not the value of K_d .

SPIM-FCCS. Single-plane illumination (light sheet) microscopy-based fluorescence cross-correlation spectroscopy (SPIM-FCCS) measurements were performed with an in-house-built SPIM setup based on the design described in references 39 and 40. Data were analyzed with the software QuickFit 3.0. For details of the experimental setup and analysis, see the supplemental material.

MD simulation of Fos-Jun and Fos-Fos complexes. Two systems were submitted to MD simulation. The first one was constructed from the Fos (139-198)–Jun (257-313) protein fragments associated with the DNA fragment as deposited in the Protein Data Bank (41). It was completed by adding the missing hydrogen atoms and closing the N- and C-terminal residues by the acetyl and N-methyl groups, respectively. The second system, Fos (139-198)–Fos (139-198), was obtained from the first one by using the Jun fragment as a template in a proper position for homology modeling of the Fos protein. Each of these systems was put into a dodecahedral box, solvated by the TIP3P explicit water model, and neutralized by Na^+ ions, and further, Na^+ and Cl^- ions were added to set the ionic strength to 0.15 M. They were then minimized, slowly heated to 310 K, and after an 80-ns equilibration period, submitted to 500-ns constant particle number (123,888 and 123,870 for the Fos-Jun and Fos-Fos systems, respectively), constant-pressure (10^5 Pa), constant-temperature (310 K) production dynamics. For the simulations, the AMBER99SB force field (42) and periodic boundary condition were used. Short-range electrostatic and van der Waals interactions were calculated explicitly within a 1-nm cutoff. For the long-range electrostatic interactions, the particle mesh Ewald method (43) was applied. A Berendsen barostat and thermostat (44) were used during this simulation. For the simulations, the GROMACS packages were used (45, 46).

As controls, MD simulations of the Leu zipper region only of the

c-Fos–c-Jun and c-Fos–c-Fos dimeric structures were also carried out with the setup protocol detailed above. The Leu zipper region we considered consisted of amino acid residues 275 to 313 and 160 to 198 of the c-Jun and c-Fos fragments, respectively. Simulations were completed for both the wild-type Leu zipper regions and the corresponding structures by applying Leu280Asp and Leu294Asp virtual mutations in c-Jun protein fragments and Leu165Asp and Leu179Asp mutations in c-Fos protein fragments.

RESULTS

FRET microscopy implies Fos homodimerization. FRET is the radiationless transfer of energy from a donor fluorophore to a nearby acceptor (27, 28), which is often used to assess molecular distances. To measure the association of c-Fos molecules (referred to as Fos in the following sections), we used Fos and the C-terminally truncated mutant form Fos²¹⁵ tagged with enhanced cyan fluorescent protein (ECFP) (donor) or enhanced yellow fluorescent protein (EYFP) (acceptor) in confocal microscopic FRET experiments. Fos²¹⁵ was prepared (32) to bring the fluorescent-protein-labeled C termini of Fos and Jun to similar distances from the leucine zipper to enhance FRET (Fig. 1). Images of donor, transfer, and acceptor signals were recorded, and FRET efficiencies (E values) of labeled proteins, as well as acceptor-to-donor molecular (N_A/N_D) ratios were calculated on a pixel-by-pixel or cell-by-cell basis.

Pixel-by-pixel FRET efficiency maps and histograms of representative cells are shown in Fig. 3. Cells cotransfected with full-length Fos-ECFP–Fos-EYFP (top row) yielded a mean FRET efficiency of $E = 5.0\% \pm 0.5\%$ (SEM; $n \approx 30$ cells); for Fos²¹⁵-ECFP–Fos²¹⁵-EYFP (second row; $n \approx 30$), it was $10.0\% \pm 0.5\%$. The higher E value of the truncated Fos²¹⁵ molecules is probably due to the reduced distance between the shorter C-terminal regions (Fig. 1). We also measured the FRET efficiencies of the Fos-ECFP–Jun-EYFP and Fos²¹⁵-ECFP–Jun-EYFP samples (third and fourth rows; $n \approx 30$), which were $7.9\% \pm 0.4\%$ and $15.0\% \pm 1.1\%$. For these pairs, we have shown heterodimer formation earlier by FCCS and FRET (7, 32, 34). Both Fos and Jun molecules showed strong nuclear enrichment. The negative control (ECFP and EYFP coexpressed as separate proteins) and the positive control (ECFP-EYFP fusion protein) had mean E values of $2.8\% \pm 0.4\%$ and $48.6\% \pm 0.8\%$ (fifth and sixth rows; $n \approx 20$). These proteins had a diffuse distribution in the whole cell. The FRET efficiency of the Fos-Fos (or Fos²¹⁵-Fos²¹⁵) samples was lower than that of the Fos-Jun and Fos²¹⁵-Jun heterodimers but significantly higher than that of the negative control, indicating that Fos formed homodimers in these cells.

Dimer formation depends on the concentrations of the interacting partners; therefore, we analyzed FRET in cells expressing various amounts of the proteins. We calculated average intensities in the whole nucleus for Fos or Jun and in the whole cell for the positive and negative controls in single cells and determined FRET on a cell-by-cell basis. This method allowed a rapid analysis of several hundred cells. Figure 4A and B show the mean E value as a function of the acceptor-to-donor molecular N_A/N_D ratio for the Fos²¹⁵-ECFP–Fos²¹⁵-EYFP and Fos-ECFP–Fos-EYFP samples. Data were grouped into three subsets according to donor concentrations based on the fluorescence intensity of the donor (low, medium, high). For both protein pairs, E increases from low to high donor concentrations, in accordance with the higher probability of complex formation. E also increases with increasing N_A/N_D ratios because more acceptor-tagged Fos molecules are

available for donor-tagged ones to form a complex. The plateau or, in its absence, the average of the highest E values is presented in Fig. 4C, which shows that at higher Fos concentrations, the extent of homodimerization increased as expected. Similar to the pixel-by-pixel analysis, the mean FRET efficiencies of the Fos-Fos and Fos²¹⁵-Fos²¹⁵ dimers are between those of the negative control and the respective Fos-Jun or Fos²¹⁵-Jun dimers.

The measured E values depend on the FRET efficiency E_0 in a single donor-acceptor complex determined by the dye-to-dye distance and orientation and on the fraction of donors forming dimers with an acceptor. The length of the Fos²¹⁵ molecule downstream of the dimerization domain is similar to that of Jun; thus, the dye-to-dye distances in the Fos²¹⁵-Fos²¹⁵ and Fos²¹⁵-Jun complexes should be similar. However, in the case of Fos homodimers, only complexes of donor- and acceptor-tagged proteins yield FRET (Fig. 2). The measured mean E value is a weighted average of non-FRET-ting and FRET-ting dimers. Thus, the mean FRET efficiency of the homodimer is expected to be lower than that of the heterodimer. This was taken into account in our subsequent analyses.

Calibration of fluorescence intensity to measure absolute EGFP concentrations. The above-described FRET titration curves demonstrated that FRET efficiency can be used to monitor the extent of homo- and heteroassociations quantitatively in our system. The stability of a complex is characterized by its dissociation constant, K_d . In Materials and Methods, we outlined a method to determine the K_d of interacting proteins in live cells from FRET titration curves. This requires knowledge of the absolute concentrations of all of the interacting molecules, i.e., the transfected fluorescent and endogenous nonfluorescent ones. In subsequent measurements, we used the EGFP-mRFP1 dye pair because of the higher photostability of EGFP than ECFP. First, we developed a calibration method to determine the concentration of fluorescent proteins. Confocal images of cells expressing free EGFP were taken, and ACF curves were recorded at selected points of the images (see Fig. 6A). From ACF curves, local dye concentrations were determined by nonlinear fitting, yielding a calibration curve of EGFP concentration versus fluorescence intensity per pixel, $c(F)$ (see Fig. 6B) (37). To make the concentration calibration portable and facilitate the comparison of measurements on different days, we normalized EGFP fluorescence by using a fluorescent bead as an intensity standard. The EGFP concentration corresponding to one bead unit was $\sim 15.4 \pm 0.7 \mu\text{M}$ for our confocal microscope and $1.0 \pm 0.1 \mu\text{M}$ for the flow cytometer. For the latter calculation, we took the nucleus-to-bead volume ratio (13:1) and the nuclear fraction of Fos²¹⁵-EGFP (85%) into account. In that way, the molar concentration of EGFP-tagged protein could be assessed by simply comparing its intensity to that of the bead measured on the same day in the microscopic or flow cytometric setup without having to repeat the FCS calibration.

Determination of the absolute concentrations of endogenous and transfected Fos and Jun. To calculate the K_d of dimers, we also need to know the amount of endogenous Fos and Jun, since they can also form dimers with each other or with their fluorescent counterparts. First, we detected the green fluorescence signal of Fos-EGFP in transfected cells and compared it to that of the calibration bead to determine the absolute concentration of transfected protein (equation 5 in Materials and Methods; see Fig. S4D to G in the supplemental material). Then we used immunofluorescence labeling paired with far red channel flow cytometry




FIG 3 Subcellular pixel-by-pixel analysis of dimerization by confocal microscopic FRET on HeLa cells. ECFP (donor channel) was excited at 458 nm and detected at 475 to 525 nm. In the transfer channel, excitation was at 458 nm and detection was at 530 to 600 nm. EYFP (acceptor channel) was excited at 514 nm and detected at 530 to 600 nm. Full-length Fos-ECFP–Fos-EYFP (top row), Fos²¹⁵-ECFP–Fos²¹⁵-EYFP (second row), Fos-ECFP–Jun-EYFP (third row), and Fos²¹⁵-ECFP–Jun-EYFP (fourth row) showed nuclear localization. The negative control, ECFP and EYFP expressed independently, and the positive control, the ECFP-EYFP fusion protein (fifth and sixth rows), were evenly distributed in the whole cell. FRET efficiency (E) was calculated in each pixel. Histograms show the statistics of the subcellular distribution of E .

to detect the total Fos pool in nontransfected and Fos-EGFP-transfected samples. The immunofluorescence signal of the nontransfected sample is proportional to the endogenous Fos concentration, while that of the transfected one corresponds to the sum of the endogenous and transfected amounts. Thus, using the known concentration of Fos-EGFP, we deduced the average concentrations of endogenous Fos (113 ± 11 nM) and Jun (94 ± 10 nM) in HeLa cells (equation 7 in Materials and Methods). Our procedure combining immunofluorescence and EGFP-tagged protein expression can be generally used to assess the absolute

concentration of any endogenously expressed nonfluorescent protein.

Determination of the dissociation constant of Fos-Jun heterodimers and Fos homodimers in live cells with flow cytometric FRET data. We wanted to determine the dissociation constants of homo- and heterodimers from FRET titrations. Therefore, we derived the expressions of FRET efficiency in terms of the concentrations of donor-tagged, acceptor-tagged, and unlabeled endogenous proteins of interest and the K_d s (equations 15, 20, and 21; see equations S12 and S14 in the supplemental material). Flow




FIG 4 Cell-by-cell analysis of dimerization by confocal microscopic FRET. (A, B) FRET efficiencies of donor (ECFP)- and acceptor (EYFP)-tagged Fos²¹⁵ or full-length Fos molecules as a function of the acceptor-to-donor molecular ratio (N_A/N_D). Data from 300 cells were grouped into three subsets as a function of donor intensity (low, <800; medium, 800 to 1,200; high, >1,200 [ar-

bitrary units]). Cellular data were binned in 0.25-wide intervals of the N_A/N_D values to reduce data scatter. FRET efficiencies increased with increasing N_A/N_D ratios. (C) Saturation values of FRET efficiencies at high N_A/N_D ratios (>0.95). ECFP-EYFP fusion protein served as a positive control, and independently expressed ECFP and EYFP served as a negative control. The FRET data of the Fos-Jun and Fos²¹⁵-Jun pairs were previously published in reference 34.

cytometric FRET experiments were carried out with large cell populations expressing EGFP- and mRFP1-tagged proteins at various concentrations. Cell-by-cell FRET efficiency (E) values, donor concentrations $[F_D]_t$ (derived from the FRET-corrected donor intensity $[I_D]$ by comparison to beads), and acceptor-to-donor molecular ratios (N_A/N_D) were determined. To characterize heterodimers, cells were cotransfected with Fos²¹⁵-EGFP-Jun-mRFP1 or Jun-EGFP-Fos²¹⁵-mRFP1 pairs. The extent of association depends on the concentrations of both molecules. To create functions with a single variable, cells were grouped into classes with approximately constant N_A/N_D ratios. E values of selected N_A/N_D ratio groups were plotted as a function of the donor concentration (Fig. 5A and C). The FRET efficiency increases with increasing N_A/N_D ratios, as expected. Figure 5B and D display the theoretical values of the maximal FRET efficiencies at various N_A/N_D ratios (see equation S13 in the supplemental material), which could be measured if all possible donor-acceptor complexes were formed (and association were complete). The function increases linearly up to N_A/N_D ratio of 1, where it reaches E_0 , the FRET efficiency of a single donor-acceptor pair, and remains constant at higher N_A/N_D ratios. E -versus- N_A/N_D data were fitted to the model function describing the heterodimerization process (see equation S12 in the supplemental material), taking the presence of fluorescent-protein-tagged and unlabeled endogenous Fos and Jun into account (Fig. 2A). The apparent K_d values derived from the fits varied with various N_A/N_D ratios for both pairs between 10 and 370 nM. We received lower apparent K_d s where Fos was present in excess (larger Fos/Jun ratios) (Fig. 5E).

Fos homodimerization was studied by measuring the FRET efficiency of Fos²¹⁵-EGFP and Fos²¹⁵-mRFP1 (Fig. 5F). Here, homodimers can contain donor-acceptor-, donor-donor-, and acceptor-acceptor-tagged protein pairs (plus dimers containing endogenous Fos) (Fig. 2B); thus, the dependence of E on the N_A/N_D ratio is different from that for heterodimerization (Fig. 5F). The theoretical E_{max} value approximates the value of E_0 asymptotically as N_A/N_D tends to infinity (see equation S15 in the supplemental material). Curves were fitted by taking into account both the homoassociation of Fos (in all combinations of donor-tagged, acceptor-tagged, and endogenous molecules) and its heteroassociation with endogenous Jun (see equation S14 in the supplemental material). In the fits, the K_d and E_0 values were linked for data sets with different N_A/N_D ratios, yielding a K_d of $6.7 \pm 1.7 \mu\text{M}$ and an E_0 of $9.5\% \pm 0.8\%$ for the homodimerization process. When different N_A/N_D groups were fitted independently, K_d and E_0 ranged from 5.4 to 9.7 μM and from 9.1 to 11.9%, respectively. As expected, the K_d of the Jun-Fos²¹⁵ heterodimer is much lower than that of the Fos homodimer, which means that Fos homodimerization will be significant only in the case of its selective overexpression. At equal Fos and Jun concentrations, the formation of Fos-Jun heterodimers is more probable than the formation of Fos homodimers.

bitrary units]). Cellular data were binned in 0.25-wide intervals of the N_A/N_D values to reduce data scatter. FRET efficiencies increased with increasing N_A/N_D ratios. (C) Saturation values of FRET efficiencies at high N_A/N_D ratios (>0.95). ECFP-EYFP fusion protein served as a positive control, and independently expressed ECFP and EYFP served as a negative control. The FRET data of the Fos-Jun and Fos²¹⁵-Jun pairs were previously published in reference 34.




FIG 5 Determination of the dissociation coefficients of Fos-Jun heterodimers and Fos-Fos homodimers from flow cytometric FRET titrations. (A, C) FRET efficiency measured in cells cotransfected with Fos²¹⁵-EGFP–Jun-mRFP1 and Jun-EGFP–Fos²¹⁵-mRFP1 and plotted as a function of the donor-tagged Fos²¹⁵ or Jun concentration. Data were grouped according to acceptor-to-donor molecular ratios (N_A/N_D) and fitted as described elsewhere (see equation S12 in the supplemental material) (solid lines), yielding the K_d value of the heterodimers and the FRET efficiency (E_0) of individual donor-acceptor pairs. Endogenous Fos and Jun were also taken into account. (B, D) The solid lines represent the maximal theoretically attainable E values at different N_A/N_D ratios (assuming E_0 values of 15 and 14.1% based on the fits) when all available Jun-mRFP1 molecules are engaged in heterodimers with Fos; the marked points correspond to the experimental N_A/N_D ratios (see equation S13 in the supplemental material). (E) Dependence of the K_d values from the fits on the Fos/Jun ratio. (F) FRET efficiency of Fos²¹⁵-EGFP–Fos²¹⁵-mRFP1 homodimers as a function of the donor-tagged Fos²¹⁵ concentration with K_d and E_0 yielded from a linked fit (see equation S14 in the supplemental material). (G) Maximal attainable FRET efficiencies at different N_A/N_D ratios (assuming an E_0 value of 9.47% based on the fit) when all Fos molecules form homodimers (see equation S15 in the supplemental material).

Fluorescence brightness and slow diffusion indicate Fos homoassociation and DNA binding. FCS was used not only for concentration calibration but also as an additional tool to probe Fos homoassociation. Diffusing particle concentration-versus-fluorescence intensity curves were generated for EGFP, Fos^{ΔΔ}-EGFP (a mutant form lacking the DNA-binding and dimerization domains, see Fig. 1), full-length Fos-EGFP, and Fos-EGFP–Jun-

mRFP1 samples (Fig. 6B). The slopes of the fitted straight lines for EGFP and Fos^{ΔΔ}-EGFP were 30.0 and 30.3 (nM/intensity unit). The similar slopes indicate that equal intensities of EGFP or Fos^{ΔΔ}-EGFP correspond to equal particle numbers, suggesting that this mutant form contains one fluorophore per particle; i.e., it is monomeric. In contrast, full-length Fos-EGFP yielded a slope of 11.5, which is less than half of the previous values. Thus, an equal




FIG 6 FCS-based concentration calibration and brightness analysis. (A) The EGFP concentration in HeLa cells was determined from the amplitude of the ACF. The curve was fitted to a two-component free-diffusion model with triplet correction. (B) Diffusing particle concentration ($1/G_0$) as a function of the fluorescence intensity per pixel of EGFP, Fos $\Delta\Delta$ -EGFP, Fos-EGFP, and Fos-EGFP coexpressed with Jun-mRFP1. Data were fitted with straight lines by Deming regression. (C) Fluorescence per particle or molecular brightness values characterizing the aggregation state plotted as a function of the concentration of the EGFP tag. Symbols are the same as in panel B. (D) Normalized ACFs fitted to a two-component free-diffusion model. (E) Diffusion constants and fractions of the second, slow component derived from the fits (n , number of cells).

intensity of Fos-EGFP corresponds to a little less than half the particle concentration of the monomeric proteins, implying the formation of Fos homodimers. In fact, this curve is not expected to be linear in the low-concentration range because of the monomer-dimer transition. The slope of the Fos-EGFP–Jun-mRFP1 sample (with an average Jun/Fos ratio of 0.7) is between the monomeric and dimeric slopes. Therefore, Fos-EGFP is partially complexed with Jun-mRFP1, where the brightness of EGFP is similar to that of monomers, whereas the rest of Fos-EGFP may form homodimers. We also analyzed the specific particle brightness, F/N , defined as the ratio of the fluorescence intensity, F , to the number of particles, N . This parameter characterizes the association state of a labeled protein and is proportional to the number of fluorophores in a jointly diffusing complex. F/N -versus-EGFP concentration values are shown in Fig. 6C. Fos-EGFP is brighter than EGFP, Fos $\Delta\Delta$ -EGFP, or Fos-EGFP–Jun-mRFP1, corroborating the

conclusion that Fos-EGFP is homodimerized when there is not enough Jun present. The brightness of Fos-EGFP increases with its concentration, indicating that dimerization is enhanced at higher concentrations, whereas the brightness of EGFP or Fos $\Delta\Delta$ -EGFP does not vary with its concentration, just as expected for monomers.

From the ACF curves, we also determined molecular diffusion properties (Fig. 6D). ACF curves from the EGFP, Fos-EGFP, Fos $\Delta\Delta$ -EGFP, and Fos-EGFP–Jun-mRFP1 samples were fitted to a model assuming fast, freely diffusing and slowly moving components (7). Diffusion coefficients and the fractions of the species are shown in (Fig. 6E). The average fraction of the slow components was 0.35 ± 0.14 for Fos-EGFP expressed alone and 0.38 ± 0.10 for Fos-EGFP when coexpressed with Jun-mRFP1, whereas it was only 0.19 ± 0.12 for the nonbinding Fos $\Delta\Delta$ -EGFP mutant form and 0.07 ± 0.05 for EGFP. The similarly increased slow fractions of Fos-EGFP expressed alone or together with Jun-mRFP1 suggest

that Fos can bind to chromatin not only as a heterodimer but also as a homodimer.

SPIM-FCCS confirms stable homoassociation and chromatin binding of Fos proteins. We used FCCS, the two-color version of FCS, to characterize the comobility of dimer-forming Fos molecules. In FCCS, the ACF and cross-correlation function (CCF) from two molecular species tagged with different colors are determined. A nonzero CCF amplitude indicates that the molecules are moving together. The ratio of the CCF and ACF amplitudes from a double-labeled sample is proportional to the fraction of molecules forming a complex. FCCS measurements were performed by SPIM with an electron-multiplying charge-coupled device camera as a sensor, which allows simultaneous measurements at many pixels in a cell. This improves the statistics and provides two-dimensional interaction and mobility maps (40).

Measurements were carried out with cells cotransfected with the following protein combinations: Fos²¹⁵-EGFP-Fos²¹⁵-mRFP1, Fos²¹⁵-EGFP-Jun-mRFP1, Fos^{ΔΔ}-EGFP-Fos²¹⁵-mRFP1 (negative control), and EGFP-P30-mRFP1 (positive control, EGFP and mRFP1 connected by a 30-residue-long polyproline linker). Cells expressing about equal amounts of the green and red fluorophores were selected from the concentration range used in FRET experiments (0.3 to 10 μM). Figure 7A shows typical correlation curves obtained from the four samples. Amplitudes of the curves were low because of the high concentration required to see Fos homodimerization, but our data showed that quantitative measurements were possible even in this concentration range. At each pixel, we performed a global FCCS fit to the green and red ACF curves and the CCF (40) (see equations S16 to S18 in the supplemental material). The fit functions were parameterized by the concentrations of three diffusing species (green only, red only, green-red [GR] dimers), which were linked over all three curves. We assumed a two-component diffusion model for the ACF curves and a one-component model for the CCF. Diffusion coefficients were not linked. Figure 7B shows examples of the maps and histograms of the relative GR dimer concentration $c_{GR}/(c_{G-only} + c_{R-only} + c_{GR})$ obtained from these fits. Figure 7C shows the statistics from $n > 10$ cells as the average \pm the standard deviation (SD) of the medians extracted from the pixel distributions in each cell. As expected, the negative control had the smallest apparent dimer fraction and the positive control had the largest one, defining the dynamic range of the measurements (0.06 to 0.32). The upper limit is < 1 because of the imperfect overlap of the green and red detection volumes, partial photobleaching of the dyes, and imperfect dye maturation/folding, producing green-only and red-only species besides doubly labeled molecules. The Fos-Jun heterodimer showed a large apparent dimer fraction (0.22 ± 0.07), whereas the value from the Fos²¹⁵ homodimer was slightly smaller (0.16 ± 0.05) but still significantly larger than that of the negative control.

We also analyzed the protein mobility of the GR dimers that could be extracted from the fits of the CCFs. We used a single-component fit; a second component could not be fitted to the CCFs of the Fos-EGFP-Jun-mRFP1 and Fos²¹⁵-EGFP-Fos²¹⁵-mRFP1 samples, suggesting the presence of only a single slow species. The CCFs of the negative control could not be fitted reliably because of their very low amplitude. The diffusion coefficient (D_{cross}) of the EGFP-P30-mRFP1 fusion protein was $\sim 4.3 \mu\text{m}^2/\text{s}$ (the mean value of two components present for this protein), whereas the average diffusion coefficients of the Fos²¹⁵-Fos²¹⁵ and

Fos-Jun dimers were much lower, ~ 0.3 and $0.4 \mu\text{m}^2/\text{s}$ (Fig. 7D). The presence of a single slow component for Fos homodimers and Fos-Jun heterodimers indicated that these complexes could bind to slowly moving nuclear components, supposedly the chromatin.

Fos-Jun and Fos-Fos complexes form stable dimers, as revealed by MD modeling. On the basis of the distance-related data from FRET measurements, we performed MD modeling to verify the stable formation of Fos homodimers. In Fig. 8, the first frames from the production dynamic trajectories are presented. During the 500-ns simulation interval (see Movies S5 and S6 in the supplemental material), not only the Fos-Jun heterodimer but also the Fos-Fos homodimer remained bound to DNA and the dimeric structures (coiled coil motifs) stayed strongly associated by their leucine zipper regions. It is noteworthy that the Fos-Fos homodimer exists despite the net negative charge of the leucine zipper regions. This indicates that in the dimeric form, hydrophobic interactions play a crucial role and electrostatic forces are largely shielded by counterions. While H bonds and even salt bridges can contribute to the stabilization of the dimeric structure as well, the H-bond networks between protein chains have variability in their connection patterns, which further supports the importance of hydrophobic interactions in the dimeric structures.

Simulations carried out solely for the leucine zipper region of c-Jun-c-Fos and c-Fos-c-Fos dimers indicated stable structures with coiled-coil motifs (Fig. 8C and D). These results are both in good accordance with former simulations carried out for the c-Jun-c-Fos leucine zipper region (47) and underline again the role of hydrophobic forces even in the stability of the c-Fos-c-Fos homodimer. However, when Leu-Asp virtual mutations were introduced into these dimeric structures (as a negative control), the contact between the corresponding regions of helices was either weakened (c-Jun-c-Fos) or even destroyed (c-Fos-c-Fos), as demonstrated in Fig. 8E and F.

DISCUSSION

Homodimer formation of short fragments, mainly the leucine zippers, of Fos proteins has been studied earlier. However, *in vitro* studies reported low stabilities of the homodimer, and it was assumed that it could not be present in live cells. By combining FRET, FCS, and imaging FCCS, we demonstrated that Fos proteins formed homodimers in live cells and presented a method of calculating their dissociation constant. The K_d of Fos homodimers in HeLa cells was $6.7 \pm 1.7 \mu\text{M}$, which is on the same order of magnitude as the value of $5.6 \mu\text{M}$ determined for its isolated leucine zippers *in vitro* by circular dichroism (14). Values reported for the heterodimers of the isolated leucine zippers (10, 48, 49) or longer polypeptides (50) *in vitro* varied between 1 and 140 nM. For the Fos-Jun heterodimer, we found a K_d range of 10 to 370 nM in live cells, which depended on the Fos/Julun ratio and on putting the donor and acceptor tags on one or the other protein. The variation of the K_d with different Fos/Julun ratios may be caused by the formation of Jun homodimers, which could interfere with the heterodimerization process. At lower Fos/Julun ratios, when there is excess Jun present, the relative amount of Jun homodimers is expected to be higher; thus, less free Jun is available and the heterodimerization process could shift toward higher concentrations (Fig. 5E), resulting in a higher apparent K_d . At higher Fos/Julun ratios, where the Jun homodimer is expected to be less abundant, we got a K_d of < 100 nM for the heterodimer, in agreement with earlier *in vitro* results. The shift between the Fos-Jun and Jun-Fos




FIG 7 SPIM-FCCS data analysis showing codiffusion and DNA binding of Fos homodimers. (A) ACFs and CCFs from SPIM-FCCS measurements. Green, EGFP ACF; red, mRFP1 ACF. Solid lines indicate the experimental data, whereas dashed lines are fits assuming two diffusing components (ACF curves) or one component (CCFs). The red horizontal line is the cross talk-corrected red ACF amplitude, and the blue horizontal line is the level of cross-correlation due to cross talk. Cross-correlation above this value is due to the codiffusion of green and red molecules. (B) The first two columns are fluorescence intensity maps of EGFP or mRFP1 from a selected cell. The third column is a map of the fraction of GR dimers among all of the molecules detected [$c_{GR}/(c_{G-only} + c_{R-only} + c_{GR})$], determined from the fits, and the histograms show their distributions. (C and D) Average fractions of GR dimers (C) and diffusion coefficients, D_{cross} (D), from the cross-correlation fits (mean \pm SD; $n > 20$ for each sample). Fits were carried out on a pixel-by-pixel basis, and the median of the respective parameter from each cell was then averaged. ***, $P < 0.0001$ (t test).




FIG 8 Both Fos-Jun and Fos-Fos complexes form stable dimers and bind to DNA. (A and B) MD simulations were carried out on small Fos-Jun (A) and Fos-Fos (B) fragments bound to the DNA fragment. Ribbon representation (colored cyan) was applied for the helical secondary structure of the Fos protein fragment (A, B). The atomic details of constituent residues are shown by stick representation with the C, H, N, O, and S atoms in gray, white, blue, red, and yellow, respectively. For the Jun fragment (A) or the second Fos fragment (B), a solvent-excluded surface representation was applied by using the above-described color codes. (C to F) Visual representation of trajectories from MD simulations of the Leu zipper region of the Jun-Fos (C, E) and Fos-Fos (D, F) dimeric structures. Wild-type protein fragments (C, D) and virtually mutated (Leu280Asp and Leu294Asp in c-Jun and Leu165Asp and Leu179Asp in c-Fos) fragments (E, F) were considered. Mutant residues are shown by stick representation with the color scheme of the atoms as above. Jun is represented by the orange helix, and Fos is represented by the green and yellow helices. From each 500-ns dynamic trajectory, 100 frames were saved equidistantly and superimposed (after removal of rotation and translation). Wild-type protein fragments (C, D) demonstrate stable coiled-coil motifs with relatively small fluctuations. The mutations in the Fos-Jun fragment (E) resulted in a somewhat distorted structure and larger fluctuations, indicating weakening of the interaction between the monomers. This is even more strongly expressed in the mutant dimeric Fos-Fos fragment (F), where the hydrophobic interaction between regions affected by the mutations is completely destroyed.

curves in Fig. 5E might be due to dark states (51) and incomplete maturation of mRFPI, resulting in an error in the acceptor-to-donor ratios. Furthermore, the autofluorescence intensity of HeLa cells in the green channel corresponds to the specific intensity of ~ 50 nM EGFP; therefore, the signal-to-noise ratio in the concentration range of the K_d is lower than in the case of the homodimer, making the K_d for the heterodimers less accurate.

Several groups have used FRET to determine the K_d s of isolated proteins (50, 52). Other groups have used microscopic FRET to determine K_d s in cells by utilizing prior estimates of protein copy numbers per cell (53) or applying *in vitro* concentration calibration with purified proteins (54). Here we present a method to calculate K_d values based on FRET titrations after concentration calibration by FCS, where the whole procedure was carried out with live cells. None of the earlier studies took into account the presence of endogenous, unlabeled proteins. With our method, the absolute concentrations of both overexpressed fluorescent and endogenous nonfluorescent proteins were determined and included

in dissociation equilibria. Our concentration calibration method is transferable to measurements performed with different instruments or on different days with fluorescent beads as a standard. The procedure can be generally used to determine K_d values and absolute concentrations of proteins in live cells.

FRET reveals that a certain fraction of molecules is colocalized within Förster distance. We used FCS to assess the codiffusion of molecules, which is a direct indication of stable interaction. Our molecular brightness analysis of FCS data indicated that Fos-EGFP, when expressed alone, had a higher ($\geq 2\times$) molecular brightness than its dimerization- and DNA-binding-deficient Fos $^{\Delta\Delta}$ mutant form or the free EGFP dye. This corroborated that at a concentration of a few micromolar, Fos was present mainly as a homodimer, which was stable at least for a few tens of milliseconds (the mean dwell time of particles in the focal volume setting the upper limit of the time scales observed in our FCS experiments). When fitted with slow and fast diffusion components, the slow fraction of Fos was about the same whether expressed alone

or together with Jun; in contrast, the slow fraction of the Fos^{ΔΔ} mutant form was significantly smaller, hinting at DNA binding of wild-type Fos either as a homodimer or as a heterodimer. The presence of a very small apparent slow fraction in the case of lone EGFP and Fos^{ΔΔ} is probably due to molecular crowding in the nucleus leading to anomalous subdiffusion (37). This makes the autocorrelation curves less steep than for free diffusion, mimicking the presence of a second, slowly moving component with a longer diffusion time.

SPIM-FCCS allowed us to confirm the presence, visualize the distribution, and characterize the intranuclear mobility of Fos homodimers. These were stable for at least a few hundred milliseconds, the time window defined by the cross-correlation diffusion time. Their diffusion coefficient derived from the cross-correlation curve was $\sim 0.3 \mu\text{m}^2/\text{s}$, similar to that of Fos-Jun heterodimers ($0.4 \mu\text{m}^2/\text{s}$). The measured diffusion coefficients are similar to those determined by confocal FCCS for the same proteins (7) and to values observed for other chromatin-binding proteins, e.g., nuclear receptors (55, 56) or HP1 α (57).

Our MD modeling simulations also supported the possibility of homodimerization, showing that homodimers stayed together for the duration of the simulation.

The existence of stable Fos homodimers capable of chromatin binding brings up the possibility that they act as transcriptional regulators and may explain the importance of Fos overexpression in oncogenesis. Various complexes of different Fos and Jun variants occur in different cell types, contributing to cell proliferation or apoptosis (58–60). It is not clear yet whether the Fos homodimer could function as an autonomous transcription factor or, alternatively, occupy the binding sites of AP-1 heterodimers, preventing their normal function and interfering with their proliferative or antiproliferative effects.

ACKNOWLEDGMENTS

We thank Gabriele Müller for preparing the EGFP-P30-mRFP1 construct, László Nagy, Jörg Langowski, and Matt Peloquin for useful comments on the manuscript, and Edina Nagy for excellent technical assistance.

This work was supported by grants from the Hungarian Scientific Research Fund (OTKA K103965 to G.V.); the Magyar Ösztöndíj Bizottság and the Deutscher Akademischer Austauschdienst (MÖB/21-1/2013 to G.V. and K.T.); TÁMOP-4.2.2.A-11/1/KONV-2012-0023 VÉD-ELEM implemented through the New Hungary Development Plan cofinanced by the European Social Fund and the European Regional Development Fund (to G.V.); the European Union and the State of Hungary, cofinanced by the European Social Fund in the framework of the TÁMOP-4.2.4.A/2-11/1-2012-0001 National Excellence Program (to G.V.); and the Deutsche Forschungsgemeinschaft and the Heidelberg Graduate School for Mathematical and Computational Methods for the Sciences (HGS MathComp; GSC 220) in the framework of the German Excellency Initiative (to J.W.K.).

REFERENCES

- Angel P, Karin M. 1991. The role of Jun, Fos and the AP-1 complex in cell-proliferation and transformation. *Biochim Biophys Acta* 1072: 129–157.
- Zenz R, Eferl R, Scheinecker C, Redlich K, Smolen J, Schonhaler HB, Kenner L, Tschachler E, Wagner EF. 2008. Activator protein 1 (Fos/Jun) functions in inflammatory bone and skin disease. *Arthritis Res Ther* 10:201.
- Shaulian E, Karin M. 2001. AP-1 in cell proliferation and survival. *Oncogene* 20:2390–2400. <http://dx.doi.org/10.1038/sj.onc.1204383>.
- Curran T, Franza BR, Jr. 1988. Fos and jun: the AP-1 connection. *Cell* 55:395–397. [http://dx.doi.org/10.1016/0092-8674\(88\)90024-4](http://dx.doi.org/10.1016/0092-8674(88)90024-4).
- Hurst HC. 1995. Transcription factors 1: bZIP proteins. *Protein Profile* 2:101–168.
- Porte D, Oertel-Buchheit P, John M, Granger-Schnarr M, Schnarr M. 1997. DNA binding and transactivation properties of Fos variants with homodimerization capacity. *Nucleic Acids Res* 25:3026–3033. <http://dx.doi.org/10.1093/nar/25.15.3026>.
- Baudendistel M, Müller G, Waldeck W, Angel P, Langowski J. 2005. Two-hybrid fluorescence cross-correlation spectroscopy detects protein-protein interactions in vivo. *Chemphyschem* 6:984–990. <http://dx.doi.org/10.1002/cphc.200400639>.
- Nakabeppu Y, Ryder K, Nathans D. 1988. DNA binding activities of three murine Jun proteins: stimulation by Fos. *Cell* 55:907–915. [http://dx.doi.org/10.1016/0092-8674\(88\)90146-8](http://dx.doi.org/10.1016/0092-8674(88)90146-8).
- Smeal T, Angel P, Meek J, Karin M. 1989. Different requirements for formation of Jun:Jun and Jun:Fos complexes. *Genes Dev* 3:2091–2100. <http://dx.doi.org/10.1101/gad.3.12b.2091>.
- John M, Leppik R, Busch SJ, Granger-Schnarr M, Schnarr M. 1996. DNA binding of Jun and Fos bZip domains: homodimers and heterodimers induce a DNA conformational change in solution. *Nucleic Acids Res* 24:4487–4494. <http://dx.doi.org/10.1093/nar/24.22.4487>.
- Chinenov Y, Kerppola TK. 2001. Close encounters of many kinds: Fos-Jun interactions that mediate transcription regulatory specificity. *Oncogene* 20:2438–2452. <http://dx.doi.org/10.1038/sj.onc.1204385>.
- Curran T. 1992. Fos and Jun: oncogenic transcription factors. *Tohoku J Exp Med* 168:169–174. <http://dx.doi.org/10.1620/tjem.168.169>.
- Porte D, Oertel-Buchheit P, Granger-Schnarr M, Schnarr M. 1995. Fos leucine zipper variants with increased association capacity. *J Biol Chem* 270:22721–22730. <http://dx.doi.org/10.1074/jbc.270.39.22721>.
- O'Shea EK, Rutkowski R, Stafford Iii WF, Kim PS. 1989. Preferential heterodimer formation by isolated leucine zippers from Fos and Jun. *Science* 245:646–648. <http://dx.doi.org/10.1126/science.2503872>.
- Nicklin MJH, Casari G. 1991. A single site mutation in a truncated Fos protein allows it to interact with the TRE in vitro. *Oncogene* 6:173–179.
- Mason JM, Schmitz MA, Müller KM, Arndt KM. 2006. Semirational design of Jun-Fos coiled coils with increased affinity: universal implications for leucine zipper prediction and design. *Proc Natl Acad Sci U S A* 103:8989–8994. <http://dx.doi.org/10.1073/pnas.0509880103>.
- Greenberg ME, Ziff EB. 1984. Stimulation of 3T3 cells induces transcription of the c-fos proto-oncogene. *Nature* 311:433–438. <http://dx.doi.org/10.1038/311433a0>.
- Piechaczyk M, Blanchard JM. 1994. c-fos proto-oncogene regulation and function. *Crit Rev Oncol Hematol* 17:93–131. [http://dx.doi.org/10.1016/1040-8428\(94\)90021-3](http://dx.doi.org/10.1016/1040-8428(94)90021-3).
- Sheng M, Thompson MA, Greenberg ME. 1991. CREB: A Ca²⁺-regulated transcription factor phosphorylated by calmodulin-dependent kinases. *Science* 252:1427–1430. <http://dx.doi.org/10.1126/science.1646483>.
- Johnston SRD, Lu B, Scott GK, Kushner PJ, Smith IE, Dowsett M, Benz CC. 1999. Increased activator protein-1 DNA binding and c-Jun NH₂-terminal kinase activity in human breast tumors with acquired tamoxifen resistance. *Clin Cancer Res* 5:251–256.
- Mishra A, Bharti AC, Saluja D, Das BC. 2010. Transactivation and expression patterns of Jun and Fos/AP-1 super-family proteins in human oral cancer. *Int J Cancer* 126:819–829.
- Güller M, Toulbi-Abed K, Legrand A, Michel L, Mauviel A, Bernuau D, Daniel F. 2008. c-Fos overexpression increases the proliferation of human hepatocytes by stabilizing nuclear cyclin D1. *World J Gastroenterol* 14:6339–6346. <http://dx.doi.org/10.3748/wjg.14.6339>.
- Saez E, Rutberg SE, Mueller E, Oppenheim H, Smoluk J, Yuspa SH, Spiegelman BM. 1995. c-fos is required for malignant progression of skin tumors. *Cell* 82:721–732. [http://dx.doi.org/10.1016/0092-8674\(95\)90469-7](http://dx.doi.org/10.1016/0092-8674(95)90469-7).
- Kress E, Skah S, Sirakov M, Nadjar J, Gadot N, Scoazec J, Samarut J, Plateroti M. 2010. Cooperation between the thyroid hormone receptor TR α 1 and the WNT pathway in the induction of intestinal tumorigenesis. *Gastroenterology* 138:1863–1874.e1861. <http://dx.doi.org/10.1053/j.gastro.2010.01.041>.
- Mikula M, Gotzmann J, Fischer AN, Wolschek MF, Thallinger C, Schulte-Hermann R, Beug H, Mikulits W. 2003. The proto-oncoprotein c-Fos negatively regulates hepatocellular tumorigenesis. *Oncogene* 22: 6725–6738. <http://dx.doi.org/10.1038/sj.onc.1206781>.
- Preston GA, Lyon TT, Yin Y, Lang JE, Solomon G, Annab L, Srinivasan

- DG, Alcorta DA, Barrett JC. 1996. Induction of apoptosis by c-Fos protein. *Mol Cell Biol* 16:211–218.
27. Clegg RM. 1995. Fluorescence resonance energy transfer. *Curr Opin Biotechnol* 6:103–110. [http://dx.doi.org/10.1016/0958-1669\(95\)80016-6](http://dx.doi.org/10.1016/0958-1669(95)80016-6).
 28. Jares-Erijman EA, Jovin TM. 2003. FRET imaging. *Nat Biotechnol* 21:1387–1395. <http://dx.doi.org/10.1038/nbt896>.
 29. Bacia K, Kim SA, Schwille P. 2006. Fluorescence cross-correlation spectroscopy in living cells. *Nat Methods* 3:83–89. <http://dx.doi.org/10.1038/nmeth822>.
 30. Liu P, Sudhaharan T, Koh RML, Hwang LC, Ahmed S, Maruyama IN, Wohland T. 2007. Investigation of the dimerization of proteins from the epidermal growth factor receptor family by single wavelength fluorescence cross-correlation spectroscopy. *Biophys J* 93:684–698. <http://dx.doi.org/10.1529/biophysj.106.102087>.
 31. Tian Y, Martinez MM, Pappas D. 2011. Fluorescence correlation spectroscopy: a review of biochemical and microfluidic applications. *Appl Spectrosc* 65:115A–124A.
 32. Vámosi G, Baudendistel N, Von Der Lieth CW, Szalóki N, Mocsar G, Müller G, Brázda P, Waldeck W, Damjanovich S, Langowski J, Tóth K. 2008. Conformation of the c-Fos/c-Jun complex in vivo: a combined FRET, FCCS, and MD-modeling study. *Biophys J* 94:2859–2868. <http://dx.doi.org/10.1529/biophysj.107.120766>.
 33. Pernuš A, Langowski J. 2015. Imaging Fos-Jun transcription factor mobility and interaction in live cells by single plane illumination-fluorescence cross correlation spectroscopy. *PLoS One* 10:e0123070. <http://dx.doi.org/10.1371/journal.pone.0123070>.
 34. Szalóki N, Doan-Xuan QM, Szöllosi J, Tóth K, Vámosi G, Bacsó Z. 2013. High throughput FRET analysis of protein-protein interactions by slide-based imaging laser scanning cytometry. *Cytometry A* 83:818–829. <http://dx.doi.org/10.1002/cyto.a.22315>.
 35. Haustein E, Schwille P. 2007. Fluorescence correlation spectroscopy: novel variations of an established technique. *Annu Rev Biophys Biomol Struct* 36:151–169. <http://dx.doi.org/10.1146/annurev.biophys.36.040306.132612>.
 36. Krieger JW, Langowski J. 2015. QuickFit 3.0 (compiled: 8 January 2015, SVN: 3697): a data evaluation application for biophysics. <http://www.dkfz.de/Macromol/quickfit/>. Accessed 8 January 2015.
 37. Weidemann T, Wachsmuth M, Knoch TA, Müller G, Waldeck W, Langowski J. 2003. Counting nucleosomes in living cells with a combination of fluorescence correlation spectroscopy and confocal imaging. *J Mol Biol* 334:229–240. <http://dx.doi.org/10.1016/j.jmb.2003.08.063>.
 38. Petrásek Z, Schwille P. 2008. Precise measurement of diffusion coefficients using scanning fluorescence correlation spectroscopy. *Biophys J* 94:1437–1448. <http://dx.doi.org/10.1529/biophysj.107.108811>.
 39. Greger K, Swoger J, Stelzer EH. 2007. Basic building units and properties of a fluorescence single plane illumination microscope. *Rev Sci Instrum* 78:023705. <http://dx.doi.org/10.1063/1.2428277>.
 40. Krieger JW, Singh AP, Garbe CS, Wohland T, Langowski J. 2014. Dual-color fluorescence cross-correlation spectroscopy on a single plane illumination microscope (SPIM-FCCS). *Opt Express* 22:2358–2375. <http://dx.doi.org/10.1364/OE.22.002358>.
 41. Glover JNM, Harrison SC. 1995. Crystal structure of the heterodimeric bZIP transcription factor c-Fos–c-Jun bound to DNA. *Nature* 373:257–261. <http://dx.doi.org/10.1038/373257a0>.
 42. Hornak V, Abel R, Okur A, Strockbine B, Roitberg A, Simmerling C. 2006. Comparison of multiple amber force fields and development of improved protein backbone parameters. *Proteins* 65:712–725. <http://dx.doi.org/10.1002/prot.21123>.
 43. Darden T, York D, Pedersen L. 1993. Particle mesh Ewald: an N-log(N) method for Ewald sums in large systems. *J Chem Phys* 98:10089–10092. <http://dx.doi.org/10.1063/1.464397>.
 44. Berendsen HJC, Postma JPM, Van Gunsteren WF, Dinola A, Haak JR. 1984. Molecular dynamics with coupling to an external bath. *J Chem Phys* 81:3684–3690. <http://dx.doi.org/10.1063/1.448118>.
 45. Hess B, Kutzner C, Van Der Spoel D, Lindahl E. 2008. GROMACS 4: algorithms for highly efficient, load-balanced, and scalable molecular simulation. *J Chem Theory Comput* 4:435–447. <http://dx.doi.org/10.1021/ct700301q>.
 46. Pronk S, Páll S, Schulz R, Larsson P, Bjelkmar P, Apostolov R, Shirts MR, Smith JC, Kasson PM, Van Der Spoel D, Hess B, Lindahl E. 2013. GROMACS 4.5: a high-throughput and highly parallel open source molecular simulation toolkit. *Bioinformatics* 29:845–854. <http://dx.doi.org/10.1093/bioinformatics/btt055>.
 47. Zuo ZL, Gandhi NS, Mancera RL. 2010. Calculations of the free energy of interaction of the c-Fos–c-Jun coiled coil: effects of the solvation model and the inclusion of polarization effects. *J Chem Inf Model* 50:2201–2212. <http://dx.doi.org/10.1021/ci100321h>.
 48. Heuer KH, Mackay JP, Podzebenko P, Bains NPS, Weiss AS, King GF, Easterbrook-Smith SB. 1996. Development of a sensitive peptide-based immunoassay: application to detection of the Jun and Fos oncoproteins. *Biochemistry* 35:9069–9075. <http://dx.doi.org/10.1021/bi952817o>.
 49. Pernelle C. 1993. An efficient screening assay for the rapid and precise determination of affinities between leucine zipper domains. *Biochemistry* 32:11682–11687. <http://dx.doi.org/10.1021/bi00094a026>.
 50. Patel LR, Curran T, Kerppola TK. 1994. Energy transfer analysis of fos-jun dimerization and DNA binding. *Proc Natl Acad Sci U S A* 91:7360–7364. <http://dx.doi.org/10.1073/pnas.91.15.7360>.
 51. Hendrix J, Flors C, Dedecker P, Hofkens J, Engelborghs Y. 2008. Dark states in monomeric red fluorescent proteins studied by fluorescence correlation and single molecule spectroscopy. *Biophys J* 94:4103–4113. <http://dx.doi.org/10.1529/biophysj.107.123596>.
 52. Song Y, Madahar V, Liao J. 2011. Development of FRET assay into quantitative and high-throughput screening technology platforms for protein-protein interactions. *Ann Biomed Eng* 39:1224–1234. <http://dx.doi.org/10.1007/s10439-010-0225-x>.
 53. Mehta K, Hoppe AD, Kainkaryam R, Woolf PJ, Linderman JJ. 2009. A computational approach to inferring cellular protein-binding affinities from quantitative fluorescence resonance energy transfer imaging. *Proteomics* 9:5371–5383. <http://dx.doi.org/10.1002/pmic.200800494>.
 54. Erickson MG, Alseikhan BA, Peterson BZ, Yue DT. 2001. Preassociation of calmodulin with voltage-gated Ca(2+) channels revealed by FRET in single living cells. *Neuron* 31:973–985. [http://dx.doi.org/10.1016/S0896-6273\(01\)00438-X](http://dx.doi.org/10.1016/S0896-6273(01)00438-X).
 55. Brázda P, Krieger J, Daniel B, Jonas D, Szekeres T, Langowski J, Tóth K, Nagy L, Vámosi G. 2014. Ligand binding shifts highly mobile retinoid X receptor to the chromatin-bound state in a coactivator-dependent manner, as revealed by single-cell imaging. *Mol Cell Biol* 34:1234–1245. <http://dx.doi.org/10.1128/MCB.01097-13>.
 56. Brázda P, Szekeres T, Bravics B, Tóth K, Vámosi G, Nagy L. 2011. Live-cell fluorescence correlation spectroscopy dissects the role of coregulator exchange and chromatin binding in retinoic acid receptor mobility. *J Cell Sci* 124:3631–3642. <http://dx.doi.org/10.1242/jcs.086082>.
 57. Capoulade J, Wachsmuth M, Hufnagel L, Knop M. 2011. Quantitative fluorescence imaging of protein diffusion and interaction in living cells. *Nat Biotechnol* 29:835–839. <http://dx.doi.org/10.1038/nbt.1928>.
 58. Ameyar M, Wisniewska M, Weitzman JB. 2003. A role for AP-1 in apoptosis: the case for and against. *Biochimie* 85:747–752. <http://dx.doi.org/10.1016/j.biochi.2003.09.006>.
 59. Eferl R, Wagner EF. 2003. AP-1: a double-edged sword in tumorigenesis. *Nat Rev Cancer* 3:859–868. <http://dx.doi.org/10.1038/nrc1209>.
 60. Shaulian E, Karin M. 2002. AP-1 as a regulator of cell life and death. *Nat Cell Biol* 4:E131–E136. <http://dx.doi.org/10.1038/ncb0502-e131>.

SUPPLEMENTAL MATERIAL

for

Evidence for homodimerization of the c-Fos transcription factor in live cells revealed by fluorescence microscopy and computer modeling

Nikoletta Szalóki, Jan Wolfgang Krieger, István Komáromi, Katalin Tóth and György Vámosi

Mol. Cell Biol. 2015

Cell culture and transfection of HeLa cells

Adherent HeLa cells were grown in a 5% CO₂ humidified atmosphere at 37°C and passaged in RPMI 1640 supplemented with 10% FBS, 2 mM glutamine, 100 U/l penicillin and 100 mg/l streptomycin without phenol red. For measurements with confocal microscopy, HeLa cells were plated 48 h before the measurement in 8-well chambered Ibidi coverglasses (Ibidi GmbH, Planegg/Martinsried, Germany) at a concentration of 15,000 cells in 300 µl medium per well. For FACS, they were plated in 24-well cell culture plates (Greiner Bio-One GmbH, Frickenhausen, Germany) using 50,000 cells in 500 µl per well. For transfection 25 µl of FBS-free RPMI, 2 µl FuGene HD reagent (Promega, Madison, Wisconsin, USA) and 0.8 µg DNA were mixed, incubated for 20 minutes at room temperature, then 5 µl of this mix was added to each well. For SPIM-FCCS 100,000 cells were plated in 35-mm Petri dishes. 90 µl FBS-free RPMI was mixed with 0.1 to 6 µg of plasmid DNA and 4 µl FuGene HD reagent and the total amount of the mix was used for transfection in a small Petri dish. Small pieces of No.3 cover glass (approximately 5×8 mm², thoroughly cleaned and sterilized) were put into the Petri dishes before seeding the cells. These were later mounted in the SPIM.

Plasmid construction

Experiments were carried out using HeLa cells transfected with fluorescent proteins. The ECFP, EYFP pair was used for FRET measurements with confocal microscopy, and the EGFP, mRFP1 pair for flow cytometric FRET and SPIM-FCCS experiments. The expression vectors pSV-c-Fos-ECFP, pSV-c-Fos-EGFP, pSV-c-Fos-EYFP, pSV-c-Fos-mRFP1 and the positive controls, pSV-ECFP-EYFP and pSV-EGFP-mRFP1 (coding for the fusion of the two fluorescent proteins connected by a RNPPVAT linker) were constructed using a multi-step cloning strategy (1). Fos²¹⁵ is a truncated version of full-length c-Fos, where the last 165 amino acids have been removed (2). Cells co-transfected with ECFP and EYFP or with EGFP and mRFP1 plasmids served as negative controls. Cells transfected with ECFP, EGFP, EYFP, mRFP1 alone were used for the determination of spectral cross-talk of the dyes between detection channels. As a biological negative control, a deletion mutant of c-Fos lacking the DNA-binding and dimerization domains was used (1). For the SPIM-FCCS measurements an EGFP-P30-mRFP1 construct containing a 30 amino acid long linear proline linker was inserted to reduce FRET and to increase the size of the product; addition of the linker increased the diffusion time of the molecule making it more suitable for camera-based SPIM-FCCS measurement, which has a time resolution of ~0.5 ms.

Calculation of FRET efficiency from confocal images

FRET measurements between ECFP as donor and EYFP as acceptor labels were performed on a Zeiss LSM 510 confocal microscope (Carl Zeiss, Jena, Germany) as described earlier (3). Images were collected in three channels, I_1 (donor signal), I_2 (FRET signal) and I_3 (acceptor signal), which contain different contributions from donor and acceptor emission (see Eq. 1). The donor signal was excited at 458 nm and emission was detected

between 475-525 nm (I_1), while the FRET signal was also excited at 458 nm and detected between 530-600 nm (I_2). The acceptor signal was excited at 514 nm and detected between 530-600 nm (I_3). The “Multi Track” option of the data acquisition software was used switching laser illumination line by line to minimize spectral crosstalk between the channels. These three fluorescence signals can be expressed in terms of contributions due to direct excitation of the donor and the acceptor, and the sensitized emission of the acceptor:

$$\begin{aligned} I_1(458, 475-525) &= I_D(1-E) + I_A S_4 + I_D E \alpha \frac{S_4}{S_2} \\ I_2(458, 530-600) &= I_D(1-E) S_1 + I_A S_2 + I_D E \alpha \\ I_3(514, 530-600) &= I_A \end{aligned} \quad (S1)$$

Intensities I_{1-3} were background-corrected by subtracting mean intensities of non-transfected cells. I_D is the unquenched pure donor intensity in channel 1, I_A is the pure acceptor signal in channel 3 and E is the FRET efficiency. The spectral crosstalk factor S_1 was calculated using cells expressing ECFP alone, whereas S_2 and S_4 were measured from cells expressing EYFP only:

$$S_1 = \frac{I_2^D}{I_1^D}, \quad S_2 = \frac{I_2^A}{I_3^A}, \quad S_4 = \frac{I_1^A}{I_3^A}. \quad (S2)$$

The term describing sensitized acceptor emission (acceptor emission excited via FRET from the donor) is $I_D E \alpha$. The factor α is the ratio of the signal from excited EYFP molecules detected in channel 2 to the signal of an equal number of excited ECFP molecules in channel 1. To determine α , we used cells transfected with the fusion protein ECFP-EYFP, in which donor and acceptor are expressed at a one-to-one ratio. For this sample α can be written as (3):

$$\alpha = \frac{I_A S_2 \frac{\epsilon_{458}^D}{\epsilon_{458}^A}}{I_D \frac{\epsilon_{458}^D}{\epsilon_{458}^A}}. \quad (S3)$$

The term $\frac{\epsilon_{458}^D}{\epsilon_{458}^A} = 4.43$ is the ratio of the extinction coefficients of ECFP and EYFP at the wavelength used for exciting the I_1 and I_2 signals. Equations S1 and S3 compose a set of four equations with four unknowns (I_D , I_A , E , and α). The solution for α , using the I_1 - I_3 intensities measured for the ECFP-EYFP fusion, is:

$$\alpha = \frac{S_2 \left(I_1 S_1 - I_2 + \left(1 + \frac{\epsilon_{458}^D}{\epsilon_{458}^A} \right) I_3 (S_2 - S_1 S_4) \right)}{I_1 S_2 - I_2 S_4}. \quad (S4)$$

The average α value was used for subsequent FRET analyses of all the other doubly transfected samples. The FRET efficiency for the doubly labeled samples was expressed as

$$E = \frac{S_2 (I_2 - I_1 S_1 - I_3 S_2 + I_3 S_1 S_4)}{\alpha (I_1 S_2 - I_2 S_4) + S_2 (I_2 - I_1 S_1 - I_3 S_2 + I_3 S_1 S_4)}. \quad (\text{S5})$$

The total unquenched donor fluorescence (I_D), can be calculated as:

$$I_D = \frac{\frac{\epsilon_{458}^D}{\epsilon_{458}^A} I_3 (I_1 S_2 - I_2 S_4)}{I_1 S_1 - I_2 + \left(1 + \frac{\epsilon_{458}^D}{\epsilon_{458}^A}\right) I_3 (S_2 - S_1 S_4)}. \quad (\text{S6})$$

Because there was no donor crosstalk to the acceptor channel, the pure acceptor emission is $I_A = I_3$. We calculated the acceptor-to-donor molecular ratios for doubly labeled samples. First, the acceptor/donor intensity ratio Q was calculated with the ECFP-EYFP fusion sample, for which the number of donor and acceptor molecules are equal:

$$Q = \frac{I_A}{I_D} \quad (\text{S7})$$

Then, for any doubly labeled sample, the acceptor-to-donor molecular ratio is:

$$\frac{N_A}{N_D} = \frac{I_A}{I_D Q} \quad (\text{S8})$$

FRET analysis was carried out on a cell-by-cell or pixel-by-pixel basis. For selecting regions of interest (ROIs) and calculating the mean cellular fluorescence intensities in each channel, the LSM data acquisition software was employed. FRET calculations with the mean intensities were performed in Microsoft Excel using the above equations.

For pixel-by-pixel FRET calculations, the RiFRET plugin (4) of the ImageJ software was applied using the same equations. Images were background-corrected and pixels below the threshold ($\sim 3 \times$ the background) were not considered. Images were low-pass filtered by Gaussian blurring (with an $e^{-1/2}$ radius of 2 pixels) and corrected for pixel shift between the channels. FRET histograms and FRET efficiency maps were created in Matlab environment.

Flow cytometric FRET

Flow cytometric FRET measurements using EGFP (donor) and mRFP1 (acceptor) fluorophores were carried out on a FACSARIA III instrument (Becton Dickinson, San Jose, CA). The laser lines of this instrument are optimal for these dyes. For gating out debris and apoptotic cells, forward and side scatter signals were applied. FRET analysis follows the same overall procedure as that described for confocal microscopy, with appropriate modifications due to the different spectral properties of the used fluorescent proteins. Donor and transfer signals were excited with a 488-nm solid state laser and detected using 515-545 nm band pass and 561 nm long pass filters, whereas the acceptor signal was excited by a 561-nm solid state laser and measured via a 561 nm long pass filter. The factor S_4 was negligible thanks to the non-excitability of mRFP1 at 488 nm. Equations analogous to Eqs. S1 can be written for I_1 , I_2 and I_3 , with the simplification that $S_4=0$. For FRET calculations from flow cytometric data the Reflex software developed at our institute was employed (5).

To determine α , the sample expressing the EGFP-mRFP1 fusion protein was used. From Eqs. S1 and S3 α can be expressed with the following simplified formula:

$$\alpha = \frac{I_1 S_1 - I_2 + \left(1 + \frac{\epsilon_{488}^D}{\epsilon_{488}^A}\right) I_3 S_2}{I_1}. \quad (\text{S9})$$

For the FRET efficiency calculation of all doubly labeled samples the previously determined factors S_1 (derived from the EGFP-alone donor sample), S_2 (derived from the mRFP1-alone acceptor sample) and α were used, and E was expressed as:

$$E = \frac{S_2 (I_2 - I_1 S_1 - I_3 S_2)}{\alpha (I_1 S_2) + S_2 (I_2 - I_1 S_1 - I_3 S_2)}. \quad (\text{S10})$$


I_A is equal to I_3 , and the total unquenched donor fluorescence (I_D) can be calculated as:

$$I_D = \frac{\alpha (I_1 S_2) + S_2 (I_2 - I_1 S_1 - I_3 S_2)}{\alpha S_2}. \quad (\text{S11})$$

N_A/N_D values were determined according to Eqs. S7 and S8 the same way as for confocal microscopy.


Detector responses

Detector response curves were measured to check linearity, which is important for quantitative analyses. For the BD FACSAria III flow cytometer a series of calibration beads with factory defined relative fluorescence intensities (InSpeck™ Green (505/515) Microscope Image Intensity Calibration Kit, 6 μm) were used, which showed good linearity (Suppl. Fig. 1). For the concentration calculation we used the “1%” bead (having a factory provided 1.2% relative intensity).



Supplementary Figure 1. Detector response of FACSAria flow cytometer

The response curve of the PMT detector of the Olympus FV1000 confocal microscope was recorded by using a dilution series of Alexa 488 solutions (Suppl. Fig. 2). The response had also a good linearity.



Supplementary Figure 2. Detector response of Olympus FV1000 confocal microscope

Model functions for Fos hetero- and homodimerization

In the main text the dependence of the measured FRET efficiency E_{meas} on the concentrations of donor- and acceptor-tagged as well as endogenous Fos and Jun was derived. The exact model functions used for nonlinear fitting are given below. For Fos-Jun heterodimerization, combining equations 11 and 15 from the main text yields the expression for the measured value of E_{meas} as a function of donor concentration $[F_D]_t$ and acceptor-to-donor ratio N_A/N_D :

$$E_{meas} = \frac{[F_D]_t \left(1 + \frac{N_A}{N_D} \right) + [F_e]_t + [J_e]_t + K_d^{FJ} - \sqrt{\left([F_D]_t + [F_e]_t \right)^2 - 2 \left([F_D]_t + [F_e]_t \right) \left([F_D]_t \frac{N_A}{N_D} + [J_e]_t - K_d^{FJ} \right) + \left([F_D]_t \frac{N_A}{N_D} + [J_e]_t + K_d^{FJ} \right)^2}}{2 \left([F_D]_t + [F_e]_t \right) \left([F_D]_t \frac{N_A}{N_D} + [J_e]_t \right)} \times$$

$$\times [F_D]_t \frac{N_A}{N_D} E_0 \quad (S12)$$

where $[F_D]_t$, and $[F_e]_t$ are the total concentrations of donor-tagged and endogenous Fos (without respect to monomeric or dimeric state), $[J_A]_t$ and $[J_e]_t$ refer to acceptor-tagged and endogenous Jun, $N_A/N_D = [J_A]_t / [F_D]_t$ is the acceptor-to-donor expression ratio (determined according to Eq. S8 in the FRET analysis), K_d^{FJ} is the dissociation constant of the heterodimeric complex and E_0 is the FRET efficiency between a single donor-acceptor pair.

To get the expected maximal FRET efficiency E_{max} at a given N_A/N_D ratio, we can calculate the limit of E_{meas} as $[F_D]_t$ tends to infinity; in this case all possible Fos-Jun dimers are formed; there is no free Jun if Fos is in excess ($N_A/N_D < 1$), and vice versa, there is no free Fos if Jun is in excess ($N_A/N_D > 1$). By setting the endogenous expressions and K_d to zero (these will vanish beside $[F_D]_t$ as $[F_D]_t \rightarrow \infty$), we get the following expression:

$$E_{\max} = \frac{[F_D]_t \left(1 + \frac{N_A}{N_D}\right) - \sqrt{[F_D]_t^2 - 2[F_D]_t^2 \frac{N_A}{N_D} + \left([F_D]_t \frac{N_A}{N_D}\right)^2}}{2[F_D]_t} \times E_0 = \frac{\left(1 + \frac{N_A}{N_D}\right) - \sqrt{\left(1 - \frac{N_A}{N_D}\right)^2}}{2} =$$

$$= \begin{cases} \frac{N_A}{N_D} \times E_0 & \text{if } \frac{N_A}{N_D} < 1 \\ E_0 & \text{if } \frac{N_A}{N_D} > 1 \end{cases} \quad (\text{S13})$$

This function is shown in Fig. 5B and D in the main text; it is linearly increasing until N_A/N_D reaches 1, and then remains constant with a value of E_0 .

For the Fos homodimerization process, substitution of Eq. 21 into Eq. 20 from the main text yields the following expression for the measured FRET efficiency:

$$E_{\text{meas}} = \frac{1}{4} \left(4 \left([F_D]_t \left(1 + \frac{N_A}{N_D}\right) + [F_e]_t \right) - 4[J_e]_t + K_d^{FF} - \sqrt{K_d^{FF}} \sqrt{8 \left([F_D]_t \left(1 + \frac{N_A}{N_D}\right) + [F_e]_t \right) - 8[J_e]_t + K_d^{FF}} \right) \times$$

$$\times \frac{[F_D]_t \frac{N_A}{N_D} E_0}{\left([F_D]_t \left(1 + \frac{N_A}{N_D}\right) + [F_e]_t \right)^2} \quad (\text{S14})$$

where $[F_D]_t$, $[F_A]_t$ and $[F_e]_t$ are the total concentrations of donor-tagged, acceptor-tagged and endogenous Fos (without respect to monomeric or dimeric state), $[J_e]_t$ is the total concentration of endogenous Jun, $N_A/N_D = [F_A]_t/[F_D]_t$ is the acceptor-to-donor expression ratio, K_d^{FF} is the dissociation constant of Fos molecules, and E_0 is the actual FRET efficiency between a single donor-acceptor pair. In the above equation we assumed that all Jun molecules were in complex with Fos (because Fos is in excess).

To get the expected maximal FRET efficiency E_{\max} at a given N_A/N_D ratio for the homodimerization process, we can calculate the limit of E_{meas} as $[F_D]_t$ tends to infinity; in this case all possible Fos homodimers are formed, there is no free Fos monomer. By setting the endogenous expressions and K_d^{FF} to zero (these will vanish beside $[F_D]_t$ as $[F_D]_t \rightarrow \infty$), we get the following expression:

$$E_{\text{meas}} = \frac{N_A/N_D}{1 + N_A/N_D} \times E_0 \quad (\text{S15})$$

This function is shown in Fig. 5G in the main text.

Nonlinear fitting was carried out by using the software Graphpad Prism 5 (GraphPad Software, Inc., La Jolla, CA). In the fits the running variable was $[F_D]_t$, while the other variable, N_A/N_D was kept constant by grouping cells into classes according to N_A/N_D values. The fit parameters yielded by the fits were E_0 and K_d .


Possible effect of imperfect dye maturation or dark states on FRET analysis and K_d determination

Imperfect dye maturation may produce green-only and red-only Fos-Fos (or Fos-Jun) complexes besides doubly labeled green-red complexes. Dark mRFP1 species have been described by Hendrix and coworkers (6). We calculated a model function including a dark acceptor fraction for fitting the E_{meas} vs. donor concentration graphs (such as Fig. 5 in the main text) for homodimerization. The fit resulted a different E_0 (FRET between a single donor and acceptor), but identical K_d for the Fos homodimer as without a dark fraction. When assuming e.g. 33.3% dark component for mRFP1, the E_0 FRET efficiency changed 1.5 fold, from 9.14% to 13.17% (at $N_A/N_D=1.15$). We measure a lower E_{meas} FRET efficiency because some acceptors engaged in donor-acceptor complexes are dark, and these complexes contribute zero FRET. We can also consider this situation as a lower average FRET efficiency between a donor and an acceptor (a weighted average of the real E_0 and 0 values, weighted by the fractions of fluorescent and dark acceptors).

The reason why the presence of a dark acceptor fraction does not change the K_d is the following. From the EGFP-mRFP1 fusion sample we determine the red:green fluorescence intensity ratio for a 1:1 acceptor-to-donor expression ratio. If a Fos-EGFP-Fos-mRFP1 co-transfected sample has the same red:green intensity ratio as the EGFP-mRFP1 fusion sample, its ratio of acceptor-tagged to donor-tagged molecules is also 1:1 (just some acceptors don't fluoresce). Therefore, the total (donor-tagged+acceptor-tagged) Fos concentration is indeed $2\times$ the concentration of donor-tagged Fos. Thus, total Fos concentrations are always determined accurately, and dissociation equilibria are analyzed correctly.

Determination of the cytoplasmic and nuclear concentrations of Fos²¹⁵-EGFP in HeLa cells

We estimated the fraction of the integrated cytoplasmic fluorescence intensity compared to the total intensity of the whole cell from confocal z-stacks by using the Imaris software. The cytoplasmic fraction of Fos²¹⁵-EGFP is constant with some random variation, $15\pm 5\%$ (s.d.), and the nuclear fraction is correspondingly $85\pm 5\%$. These relative fractions do not vary with the expression level: the total cellular fluorescence (gained by integration of pixel intensities of z-stacks) is proportional to the nuclear concentration (presented as the average fluorescence intensity per pixel in the nucleus) (Suppl. Fig. 3).



Supplementary Figure 3. Relationship between total cellular and total nuclear fluorescence intensities

We also checked whether the presence of the cytoplasmic fraction affects the FRET-derived K_d and E_0 . We tested a model function considering a constant cytoplasmic fraction (15%) of donor and acceptor-tagged Fos²¹⁵ (contributing zero FRET). First, the actual nuclear concentrations are lower by 15%, therefore, the K_d is also

reduced by this amount (this was taken into account in our calculations). Second, the cytoplasmic concentration is only 13% of the nuclear concentration, thus, most of the cytoplasmic Fos²¹⁵ molecules are assumedly monomeric, contributing zero FRET. When assuming 15% cytoplasmic fraction in the model, the derived E_0 value occurring between a single donor-acceptor pair is 15% higher as compared to the case of assuming no cytoplasmic fraction. The reason is that if 15% of all donors are surely monomeric (those being in the cytoplasm), then the donors engaged in a donor-acceptor complex must have a higher FRET efficiency to produce the measured E value.

Single Plane Illumination Microscopy – fluorescence cross-correlation spectroscopy (SPIM-FCCS)

SPIM-FCCS measurements were performed on an in-house built selective plane illumination microscope setup based on the design described in (7, 8). In brief, a blue 491 nm and a green 561 nm (both from Calypso/Jive Lasers, Cobolt AB, Solna, Sweden) laser beam were expanded with distinct beam expanders and then combined by a dichroic mirror into a dual color excitation beam. Then the laser beam was relayed onto a cylindrical lens followed by a projection objective, which formed an approximately 1.3 μm thick light sheet ($1/e^2$ half-width) illuminating the cells. Samples were mounted into the sample chamber hanging from above at an angle slightly below 45° to the light sheet. The chamber was filled with Hanks' solution. The fluorescence light was collected with a 60 \times /NA 1 water dipping objective (CFI Apo-W NIR 60 \times /1.0W, Nikon, Hamburg). The sample fluorescence was split into two color channels using an image splitter optics (DualView DV2, Photometrics, Tucson), and imaged onto an EMCCD-camera (iXon X3 860, Andor, Belfast) which has a pixel-size of 400 \times 400 nm² in the object plane. A daily alignment and focus volume calibration of the setup were performed as detailed in (8) using 100-nm multi-colored fluorescent microspheres (T7279 Microspheres, Life Technologies GmbH, Darmstadt, Germany) and a 170 bp long double-stranded DNA labeled with Alexa-488 and Alexa-594 on opposing ends. The shift between the green and red foci was regularly below 100 nm, so these shifts did not have to be incorporated into the fitting models (8), whereas the different focus sizes were taken into account. The cross-talk of EGFP fluorescence to the red channel, $\kappa_{gr}=I_{red}/I_{green}$, was measured with cells expressing c-Fos-EGFP only, yielding a value of 3.35%, which was incorporated in the fit model.

Data acquisition and analysis were performed using our own software QuickFit 3.0 as follows (for details, see (9)). For each measurement, ~ 100 thousand frames were acquired at a repetition time of $\sim 530 \mu\text{s}$ (exposure time: $\sim 470 \mu\text{s}$) and an EM-Gain setting of 300. Background acquired without illumination was subtracted from the frames. Photobleaching was corrected on a per-pixel basis (8, 10). Then the autocorrelation and cross-correlation curves of the two color channels (green: 500-550 nm, red: >600 nm) were calculated. We found that binning the intensities for 2 \times 2 pixels (resulting a virtual pixel size of 800 \times 800 nm²) significantly improved the quality of correlation curves (at the expense of losing some resolution), thus, we used binned data to calculate correlation curves. Finally, a global fit to the two auto- and the cross-correlation functions was performed for each (binned) pixel. We used the model described by equations 17-19 in (8) with two diffusing components for the green and red autocorrelation functions and one component for the cross-correlation function. A second component could not be fitted well to the cross-correlation curves. The following equations were used:


$$g_{gg}(\tau) = \frac{\eta_g^2 c_A + \eta_g^2 c_{AB}}{\eta_g^2 \cdot (c_A + c_{AB})^2} \cdot G_{gg}^G(\tau) \quad (\text{S16})$$

$$g_{rr}(\tau) = \frac{\eta_r^2 \cdot [c_B + c_{AB}] + \kappa_{gr}^2 \eta_g^2 \cdot [c_A + c_{AB}] + 2\kappa_{gr} \eta_r \eta_g c_{AB}}{(\kappa_{gr} \eta_g c_A + (\eta_r + \kappa_{gr} \eta_g) \cdot c_{AB} + \eta_r c_B)^2} \cdot G_{rr}^R(\tau) \quad (\text{S17})$$

$$g_{gr}(\tau) = g_{rg}(\tau) = \frac{\eta_g \eta_r c_{AB} + \kappa_{gr} \eta_g \eta_r c_A + \kappa_{gr} \eta_g^2 \cdot c_{AB}}{(\eta_g c_A + \eta_g c_{AB}) \cdot (\kappa_{gr} \eta_g c_A + (\eta_r + \kappa_{gr} \eta_g) \cdot c_{AB} + \eta_r c_B)} \cdot G_{gr}^{GR}(\tau) \quad (\text{S18})$$

where η_g and η_r are the molecular brightness values of fluorophore A in the green channel and fluorophore B in the red channel, which can be estimated as a ratio of fluorescence intensity in a pixel, divided by number of particles in the detection volume, $G(\tau)$ factors are the non-normalized auto- and cross-correlation functions accounting for the diffusive motion of the appropriate species (8), and c_A , c_B and c_{AB} are the concentrations of green-only and red-only molecules and green-red complexes, respectively. The diffusion parameters in the three correlation functions were not linked for the Fos-Fos and Fos-Jun samples, whereas they were linked for the EGFP-P30-mRFP1 sample.

Titration of antibodies and expression of endogenous and transfected Fos and Jun



Supplementary Figure 4. Measurement of expression of endogenous and transfected Fos and Jun by flow cytometry

(A-C) Titration curves of NL-637-DAMIG secondary antibody, anti-c-Jun and anti-c-Fos primary antibodies. In panels (D-G) the primary antibodies were used at 0.02 mg/ml, and the secondary antibody at 0.05 mg/ml, which

are close to saturating conditions. Nonspecific binding of the primary antibody is low according to the fast saturation of the primary antibody titrations. (D) Fluorescence intensity distribution (red channel) of immunolabeled Fos in cells transfected with Fos-EGFP and in non-transfected cells. Endogenous Fos and Fos-EGFP were targeted by mouse anti-Fos monoclonal antibody, followed by staining with NL-637-tagged donkey anti-mouse polyclonal antibody. The control sample was stained with the secondary antibody alone; this signal was used for background correction. (E) Fluorescence intensity in the green channel from live cells transfected with Fos-EGFP. For background correction the autofluorescence of non-transfected cells was measured. Cells were not fixed to avoid degradation of EGFP fluorescence. (F-G) Analogous measurements for Jun. The average intensities of fluorescence histograms were used to assess the average cellular concentration of endogenous Fos and Jun as described in the main text. Because the antibodies were used close to saturating conditions, it may be expected that relative changes in immunofluorescence intensity faithfully reflect differences in protein expression levels.

Supplementary movies 5 and 6

Movie representation of the trajectory frames (saved at every 800 ps) from the 500 ns simulations carried out for the Fos-Jun (Fos-Jun_500_mov, supplemental movie 2) and Fos-Fos (Fos-Fos_500_mov, supplemental movie 3) dimers complexed with the DNA fragment. The solvent molecules, the ions and the hydrogens are not shown for clarity. The movies were prepared by the VMD 1.9.2 software package. For the DNA fragment "paperchain", whereas for the Fos and Jun fragments "new cartoon" representation were applied. The atomic details of the fragments are shown using "licorice" drawing method.

REFERENCES

1. **Baudendistel M, Müller G, Waldeck W, Angel P, Langowski J.** 2005. Two-hybrid fluorescence cross-correlation spectroscopy detects protein-protein interactions in vivo. *ChemPhysChem* **6**:984-990.
2. **Vámosi G, Baudendistel N, Von Der Lieth CW, Szalóki N, Mocsár G, Müller G, Brázda P, Waldeck W, Damjanovich S, Langowski J, Tóth K.** 2008. Conformation of the c-Fos/c-Jun complex in vivo: A combined FRET, FCCS, and MD-modeling study. *Biophysical Journal* **94**:2859-2868.
3. **Szalóki N, Doan-Xuan QM, Szöllosi J, Tóth K, Vámosi G, Bacsó Z.** 2013. High throughput FRET analysis of protein-protein interactions by slide-based imaging laser scanning cytometry. *Cytometry Part A* **83**:818-829.
4. **Roszik J, Lisboa D, Szöllosi J, Vereb G.** 2009. Evaluation of intensity-based ratiometric FRET in image cytometry - Approaches and a software solution. *Cytometry Part A* **75**:761-767.
5. **Szentesi G, Vereb G, Horváth G, Bodnár A, Fábíán Á, Matkó J, Gáspár R, Damjanovich S, Mátyus L, Jenei A.** 2005. Computer program for analyzing donor photobleaching FRET image series. *Cytometry Part A* **67**:119-128.
6. **Hendrix J, Flors C, Dedecker P, Hofkens J, Engelborghs Y.** 2008. Dark states in monomeric red fluorescent proteins studied by fluorescence correlation and single molecule spectroscopy. *Biophysical Journal* **94**:4103-4113.
7. **Greger K, Swoger J, Stelzer EHK.** 2007. Basic building units and properties of a fluorescence single plane illumination microscope. *Review of Scientific Instruments* **78**.
8. **Krieger JW, Singh AP, Garbe CS, Wohland T, Langowski J.** 2014. Dual-color fluorescence cross-correlation spectroscopy on a single plane illumination microscope (SPIM-FCCS). *Optics Express* **22**:2358-2375.
9. **Krieger JW, Langowski J.** 2015. QuickFit 3.0 (compiled:2015/01/08, SVN: 3697): A data evaluation application for biophysics. <http://www.dkfz.de/Macromol/quickfit/>. Accessed
10. **Ries J, Chiantia S, Schwille P.** 2009. Accurate determination of membrane dynamics with line-scan FCS. *Biophysical Journal* **96**:1999-2008.