

Review

Legume Protein Extracts: The Relevance of Physical Processing in the Context of Structural, Techno-Functional and Nutritional Aspects of Food Development

Chaima Neji ¹, Jyoti Semwal ^{1,2}, Mohammad Hassan Kamani ³, Endre Máthé ¹ and Péter Sipos ^{1,*}

¹ Faculty of Agricultural and Food Sciences and Environmental Management, Institute of Nutrition, University of Debrecen, Böszörményi út 138, 4032 Debrecen, Hungary

² Department of Grain Science and Technology, CSIR-Central Food Technological Research Institute, Mysore 570020, Karnataka, India

³ Food Chemistry and Technology Department, Teagasc Food Research Centre, Moorepark, P61 C996 Fermoy, County Cork, Ireland

* Correspondence: siposp@agr.unideb.hu

Abstract: Legumes are sustainable protein-rich crops with numerous industrial food applications, which give them the potential of a functional food ingredient. Legume proteins have appreciable techno-functional properties (e.g., emulsification, foaming, water absorption), which could be affected along with its digestibility during processing. Extraction and isolation of legumes' protein content makes their use more efficient; however, exposure to the conditions of further use (such as temperature and pressure) results in, and significantly increases, changes in the structural, and therefore functional and nutritional, properties. The present review focuses on the quality of legume protein concentrates and their changes under the influence of different physical processing treatments and highlights the effect of processing techniques on the structural, functional, and some of the nutritional, properties of legume proteins.

Keywords: legume protein; protein extraction; techno-functional and nutritional properties of protein isolates

Citation: Neji, C.; Semwal, J.; Kamani, M.H.; Máthé, E.; Sipos, P. Legume Protein Extracts: The Relevance of Physical Processing in the Context of Structural, Techno-Functional and Nutritional Aspects of Food Development. *Processes* **2022**, *10*, 2586. <https://doi.org/10.3390/pr10122586>

Academic Editor: Vlad Mureşan

Received: 29 October 2022

Accepted: 24 November 2022

Published: 4 December 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

With the increase in population and rising awareness of consumption of quality proteinaceous foods, researches for finding sustainable high-quality protein sources are proceeding apace [1–3]. Quality of a protein ingredient is generally described by its ability to meet the requirements of essential amino acids and physiological needs of the organism [4]. However, in this era of climate change, the sustainability features of the proteins have gained even more importance. Hence, alternative proteins such as cultured meat, microbial protein, edible fungi, microalgae, and insect proteins are being explored as new and sustainable sources [2,5]. Although several novel alternatives are available in the current market, plant-based proteins are still being preferred, owing to advantages such as having better consumer acceptability that is related to many subjective reasons, such as taste and health, familiarity, attitudes, food neophobia and social norms [6].

On the other hand, plant sources seem to be more reliable and sustainable protein compared to animal products [7]. Recently, legumes have received much interest owing to their lower cost of production, easy accessibility, health benefits, and their nitrogen-fixing ability in soil [8–11]. They grow globally in diverse agroclimatic zones, from alpine and arctic regions to the equatorial tropics [12]. They can be used as sources of food, flavouring plants, fodder and green manure plants or other industrial raw materials, such as oils, biofuel, timber trees, gums, dyes, and insecticides (Figure 1). Therefore, legumes

can be envisioned as super crops, meeting the challenges of climatic changes and the increasing population demands.




Figure 1. Multi-functional utilization of legumes. The scheme was constructed based on [13–19].

Various studies have brought about evidence that indicates the great potential of legumes as a source for providing high quality proteins. Furthermore, this potential highly depends on vital factors such as bioavailability, digestibility, amino acid profile and the possible anti-nutrient content of food/protein extract matrices [20–22]. Raw legumes generally have a low degree of bioavailability, and consequently poor nutritional value [23]. Compared with animal proteins, legume proteins are absorbed at slower rates, which might explain their lower dietary utilisation [24]. Therefore, it is of paramount importance to improve their nutritional quality. The nutritional quality is linked to the chemical composition (including protein/peptide fractions and their proportion; anti-nutrient content, etc.). In particular, from an industrial point of view, the antinutrient is one of the factors that can be influenced by the extraction of protein or by physical, chemical, or biological processes [25–27].

During the extraction processes of legume protein, the structural and functional properties of proteins could be affected due to the intense pH or temperature conditions [25,28]. These events might include physical processing steps such as dehulling, milling, isolation (by air classification, electrostatic separation, hydrocyclone or solvent extraction) and drying (freeze-, spray-, oven- or vacuum-drying) [29–31]. From the industrial perspective, extraction techniques are a part of processing of commercial protein powder [30].

During the manufacturing of protein-containing foods, proteins may be exposed to several processing steps, such as soaking, cooking, baking, drying, pressure-cooking, autoclaving, extrusion and microwaving [23]. The study of changes in the structure and behavior of proteins as a result of the industrially applied physical processes highlights the techno-functional and nutritional parameters that must be carefully considered during

product development [32]. Hence, from industrial and consumer viewpoints, understanding protein's behaviour within the food system is absolutely crucial for any food product development.

Therefore, the current review focuses on presenting the effects of extraction, thermal and non-thermal physical processes affecting the structural, functional, and nutritional characteristics of legume protein(s) isolates, which can give streamlined indications to improve the processing technologies to obtain properly balanced protein-containing foods.

2. Legume Proteins

2.1. Classification and Composition

Legumes are principally stored in the embryonic cotyledons [33]. The protein content in legumes ranges from 13 to 30 g/100 g dry matter. Some commonly consumed legumes with respective protein contents are pinto beans (15.41%), navy beans (14.98%), black beans (15.24%), cowpea (13.22%), kidney bean (15.35%), chickpea (14.53%), lentils (17.86%), lupin (25.85%) and soybean (28.62%) [34].

Legume proteins can be classified by their sequence, structure, function, conserved motifs, biological activity, solubility, and sedimentation coefficient [35] (Figure 2). According to their biological activities, proteins are primarily of two types: (i) storage proteins, which account for approximately 70% of total seed nitrogen, and (ii) enzymatic, regulatory, and structural proteins, which perform normal cellular activities, including the synthesis of storage proteins [36]. Further, the storage proteins can be classified into seed storage proteins and vegetative storage proteins. The vegetative storage proteins are the proteins that accumulate in vegetative tissues, such as leaves, stems, depending on plant species. It serves as a temporal reservoir of amino acids for use in subsequent phases of growth and development [37]. According to the Osborne classification scheme, legume proteins can be classified based on their solubility and can be fractionated as globulins (70%), albumins (10–20%), glutelins [38] and prolamines [39,40] (Figure 2).




Figure 2. Classification of legume proteins [35–37,40–43].

Overall, all legume flours are dominated by albumin and globulin proteins [42,44]. The globulin is the predominant protein fraction in legumes such as soybean, green gram bean, kidney bean and pea [45–47], which can be further classified based on the sedimentation coefficient: legumins (11S), vicilin (7S) and convicilin (Figure 2). Vicilins are found in various legumes, such as kidney bean, mung bean, red bean, cowpea and bambara

groundnut, while chickpeas are high in legumins. Pea and faba bean proteins comprise of both vicilins and legumins [47–49].

Since legume seeds contain more globulins than prolamin, they are rich in lysine and tryptophan [44,50] but contain less methionine and cysteine [50]. The nature of the protein depends on their amino acids, which are the basic building blocks of a protein. Amino acids are considered as the indicators of nutritional qualities and functionalities of proteins. It is an important parameter to consider when they are used to improve the nutritional value of a food products [25,51]. There are 20 amino acids, which dictate the quality and nature of protein [52]. Overall, glutamic acid and aspartic acid amino acids are the most abundant amino acids in legumes. However, tryptophan, methionine and cysteine were found less abundantly. Legumes are a poor source of essential sulphur-containing amino acids [53,54]. Table 1 shows the amino acid composition of different legumes. It is interesting to note that almost all the legumes have similar essential amino acid ratios, as recommended by the FAO [55].

Table 1. Amino acid content of various legumes and the recommended amino acid scoring patterns for adults (g/100 g protein) [55–66].

Amino Acids /Legumes	Chickpea	Soybean	Lentil	Bean	Faba Bean	Lupin	Cowpea	Pea	FAO
Alanine	4.5–5.2	3.615–4.7	4.2–4.7	3.07–4.89	3.97–4.15	3.17–3.8	4.2–4.67	3.54–5.2	NR *
Arginine	8–9.2	6.18–8.3	7.6–7.8	5.29–6.08	8.96–9.46	8.51–14.1	6.66–7.69	4–8.6	NR
Aspartic acid	10.2–12.1	7.13–11.8	11.8–13.7	9.02–12.94	9.28–10.77	8.2–15.1	10.8–11.44	8.06–12.37	NR
Cysteine	0.4–1.7	1.5–2.065	0.7–0.9	0–0.94	0.85–1.33	1.12–1.72	0.28–0.32	0.35–1.8	NR
Glutamic acid	16.5–17.8	9.115–18.2	21.4–21.5	11.42–17.06	15.67–16.51	19.19–22.15	17.2–18.54	8.53–19.7	NR
Glycine	3.4–4.3	3.71–4.4	3.6	3.19–4.55	3.95–4.73	3.83–4.77	3.8–4.48	3.87–5.27	NR
<u>Histidine</u>	2.7–3.2	2.4–3	2.2–2.5	2.59–3.42	2.41–2.61	2.31–2.95	3.06–3.19	1.92–2.94	1.6
<u>Isoleucine</u>	4.1–5.2	4.2–5.9	3.8–4.1	3.42–5.21	3.67–3.94	2.89–4.62	3.75–3.84	3.09–4.5	3
<u>Leucine</u>	7.7–9.5	7.095–7.9	7.8	6.72–8.46	6.57–7.47	5.83–7.3	7.65–7.7	6.7–7.84	6.1
<u>Lysine</u>	6.7–7.8	6–6.58	7–7.3	4.91–6.48	5.97–7.08	4.35–4.92	5.74–7.5	3.41–8.1	4.8
<u>Methionine</u>	0.8–1.6	1.1–2.72	0.8	0.72–1.76	0.52–1.06	0.35–0.7	1.46–2.11	0.72–1.6	NR
<u>Phenylalanine</u>	5–6.2	3.88–5.8	4.5–5	4.48–5.91	3.98–4.19	3.42–4	5.75–5.51	4.02–5.2	NR
Proline	3.5–4.4	3.63–5.3	3.5–4.9	2.95–6.49	3.86–4.27	4.4–5.72	4–5.91	2.11–5	NR
Serine	3.3–5.6	5.5–6.375	3.5–5.2	4.59–6.9	4.28–4.76	4.31–5.98	4.5–5.6	2.93–5.71	NR
<u>Threonine</u>	2.7–3.9	2.68–4.1	3–3.5	3.17–4.72	2.96–3.4	2.9–5.02	3.8–4.1	1.55–4.46	2.5
<u>Tryptophan</u>	0.6–1.4	1–7.64	0.7–1.2	1.11–1.18	0.85–0.87	0.49–1	0.7–1.11	0.61–3	0.66
Tyrosine	2.6–3.1	4–4.14	3.2–3.3	2.75–5.25	2.59–2.78	3.11–5.1	2.92–4.04	3.17–3.7	NR
<u>Valine</u>	3.9–5.2	4.4–5.245	4.5–5	4.58–5.38	3.41–4.31	2.46–4.2	4.68–5.1	3.97–5.11	4

* No recommended score. Underline labelling indicates essential amino acids.

2.2. Protein Quality of Legumes

The quality of legume protein is linked to the higher levels of albumin and globulin fractions, owing to the high amount of essential amino acid lysine [45]. The differences in the amino acid composition of these fractions could have an effect on protein digestibility. The studies suggested that globulin has higher digestibility compared to the albumin fraction in lentil and horse gram, which was associated with less cystine content, and therefore less disulfide bonds [67]. In contrast, a negative correlation between globulin content and its *in vitro* protein digestibility of legumes such as garden pea, grass pea, soybean and lentil was related to their resistance to gastric digestion [68].

However, globulin is hydrolysed into peptides and amino acids during the intestinal phase [68]. For the garden pea, grass pea and lentil, large-size peptides (6–20 kDa) were detected at the end of the gastric phase, but in the intestinal digests, fragments up to 6 kDa were found, except in the case of soybean, where peptides were below 4 kDa [68]. Similarly, cowpea globulins were poorly digested by pepsin, but not by serine proteases [69].

In the case of albumin, a high amount of aromatic (tyrosine, phenylalanine and tryptophan) and β -branched (threonine, valine and isoleucine) amino acids did reduce the digestibility. This was attributed to the restricted enzyme accessibility due to aggregate formation by hydrophobic linkages [67]. Another possible reason for the low digestibility of albumin could be its compact quaternary structure, which is known to impart low digestibility [70]. These studies suggested that the quality of legume proteins could be varied predominantly depending on their amino acid profile, hydrophobic interactions, and the generated protein fractions.

2.3. Extraction Techniques

In order to increase the protein content and to reduce the impact of anti-nutrients, the proteins are extracted, including isolates [25] and concentrates [39]. The extraction process depends on many factors, such as pH, temperature, particle size, ionic strength, type of salt used and solvent-to-flour ratio [71]. During the selection of an appropriate extraction technique for protein, the efficiency of the process (i.e., recovery rate), and the characteristics of the final product (i.e., nutritional quality) should be also considered. For example, the extraction process could involve intense conditions, such as very high/low pH and/or temperature, which could induce damage to the proteins [72]. Additionally, the extraction process must be energy efficient and environmentally friendly [73]. Therefore, all the mentioned factors should be considered before selecting a suitable extraction method.

The extraction can be performed in both dry and wet conditions [71]. Dry extraction consists of milling followed by the fractionation of flour particles into its constituents such as starch, fiber, or protein [74]. Protein enrichment by dry-based extraction technique is typically based on differences in size and density of legume flour [74]. Factors such as starch granule size, uniformity and its interaction with protein molecules (as they dictate the grain hardness) are important parameters affecting its efficacy [75].

For instance, the strong adherence of large amounts of insoluble protein to starch granules might cause difficulty during the separation process [76]. Starch granules in legumes range from 4 to 85 μm [77], whereas specifically for the pulse sub-group it is about 20 μm [78]. Pulses contain a uniform size of starch particles, which makes them suitable raw material for dry fractionation [78]. For pulses, the fractionation of different components can be performed by air classification [71], electrostatic separation, and sieving (sifting) technique [78]. Among these, air classification is the commonly used separation technique. It separates the light fine fraction (protein) from the heavy coarse fraction (starch) [42] (Figure 3).




Figure 3. Legume protein extraction methods [29,44,75,79–88].

Seed properties including hardness, particle density, starch granule size, fat content and flour dispersibility influence the protein content during air classification [89]. Therefore, this technique is useful particularly for the production of protein-rich meals and flours (protein content up to about 60%) from non-oilseed legumes like peas [90]. For instance, protein extracted from legumes with low lipid content such as lentil and chickpea exhibited higher protein recovery rate and yield compared to other legumes such as broad and kidney bean [91]. However, it is difficult to fractionate legumes with high oil content by air classification [82]. In such seeds, oil may promote the cohesion between powder particles through viscous and capillary bridging, which negatively affects the dispersion of individual particles in air [74]. Hence, the air classification technique for lipid-containing legumes might produce protein fractions with less purity [73]. Additionally, oil might soften the grain, giving it a ductile behaviour, and thus increasing milling energy demand [74]. Therefore, dry fractionation of protein is an appropriate technology only for the low lipid legumes. For oil-rich legumes, defatting prior to dry fractionation would be a straightforward approach to obtain higher protein content and purity [74,92].

Furthermore, the difference in the particle size of protein and starch affects the separation process. It was demonstrated that the smaller protein-rich fragments could be separated efficiently [89]. However, the protein separation efficiency reduces by the decrease in size of starch or the increasing ratio of damaged or broken starch granules. As a consequence, their differentiation from protein bodies become difficult [81].

Unlike air classification, the electrostatic separation, which is one of the eco-innovative technologies [93] is based on differences in dielectric properties of flour particles instead of their size and density [94]. However, this difference does not have any advantage on the purity of plant protein isolate/concentrates, including legumes [29]. Based on the methods used to impart surface charge, three main types of electrostatic separators are the induction charging, corona discharge, and tribocharging, based on the methods used to impart surface charge [29].

Another approach for production of protein ingredient is wet extraction (Figure 3). It is an efficient method, which can increase protein content up to 90% and 70% in isolates

and concentrates, respectively [71]. This method employs different solvents such as alkaline, acidic, salty aqueous solutions or water as extraction media followed by isoelectric precipitation (IEP) or membrane separation [42,71,95].

In general, to evaluate the separation technique efficiency, the purity and yield of the protein isolate/concentrate should be considered. The protein yield depends on the extraction method and conditions (i.e., particle size of the flour, solubilising agent, pH of solubilisation and precipitation [96], nature of the legume [25,97] and variety [95], classifier wheel speed [98,99], etc.). Many researchers compared the purity of legume proteins that was extracted with different dry, wet and hybrid fractionation methods [29,100]. The purity of protein isolate obtained from dry fractionation is lower than from wet extraction. While higher than 80% protein concentration can be achieved by wet process, dry fractionation results only 54% protein concentration in the case of lupin [73], however, further treatments (e.g., electrostatic separation) can improve efficiency [101]. Dry technique preserves the native state of the protein (i.e., structure and function) [82,89,95] and consumes lower energy and water, and retains the native structure and function of components [78,81,82].

Several modifications can be made to the basic steps of the protein extraction process to enhance its efficiency. For an instance, flowability-aiding agents such as fused silica particles can be added during air classification. Research in [74] showed an increase in the yield of the lupine protein-rich fraction with the addition of flowability aids. Additionally, repetition of the extraction cycle enhanced the separation efficiency as reported by [78]. Researches have showed that the combination of protein extraction technologies could improve its separation efficiency [29]. For instance, the combination of electrostatic separation with air classification could increase the purity of protein isolate compared to air classification alone [92,102]. First, the air classification could remove starch, and subsequent electrostatic separation separated the protein and fiber into oppositely charged fractions [92,102].

However, the combination of dry fractionation with aqueous could be used to obtain fractions with higher protein contents [73,103]. Thus far, several novel techniques, such as direct-assisted extraction, enzymatic-assisted extraction, cavitation-assisted extraction (e.g., ultrasound-assisted extraction, hydrodynamic cavitation extraction), supercritical extraction, pulsed electric field, liquid biphasic flotation, hybrid extraction processes and microwave-assisted extraction are being explored to increase the efficiency of protein extraction, specifically from food waste [72,104]. A recent study demonstrated that the intervention of ultrasound technology during the extraction of pea protein enhanced its extraction efficiency in short time with low water consumption [105]. Other studies also showed similar outcomes for the extraction yield of soy protein using ultrasonication [85,106], and lupin [107]. The application of the high pressure homogenisation treatment improved extraction yields of soybean protein up to 82%, which was attributed to the reduction in particle size and cell disruption [84].

In addition to the enhance extraction efficiency, the combination of extraction techniques can enhance the functional and nutritional proprieties of the protein isolate. The intervention of ultrasonication during extraction of lupin protein reduced the toxicity and improved its amino acid score [107]. Interestingly, the extraction of pea protein with ultrasound-assisted alkali extraction method improved its functional properties (solubility, water/oil-holding capacity, foaming/emulsifying capacity, stability, and gel formation capacity) with strong antioxidant activities [105]. Furthermore, the application of the alkaline extraction, before the first and the second enzymatic hydrolysis, enhanced the yield of peanut protein extract and their richness in small peptides [83]. Likewise, the extraction of protein from heat-denatured soy meal using protease pre-hydrolysis followed by sub-critical water treatment improved its hydrophobic amino acids composition, surface hydrophobicity, interfacial adsorption in addition to an improvement of emulsifying ability and physical stability of emulsion. However, it decreased the protein purity with prolonged hydrolysis [86].

3. Processing Affects Structural and Functional Properties of Proteins

During food processing, the variation of conditions (temperature, timing, pressure, pH, etc.) can lead to changes in the structural (Table 2) and consequently the functional properties of proteins (Table 3). In an ideal situation, once the protein is produced in the cytoplasm, its folding will be aided by chaperons and post-translational modifications to develop the structural conformation compatible with its own function. After functional accomplishment, the proteins will become ultimately degraded into amino acids that can be used for the next protein synthesis. This ongoing saga of proteins is regulated by multiple cellular mechanisms at relatively high energetic costs. Extracting proteins from their natural life cycle will render them non-functional and stable, to some extent, so that such a fact stands at the basis of food preservation. The food-specific stability of proteins is paramount to manufacturers but for consumers the importance would fall onto the reutilization of amino acids from proteins. It remains therefore a constant challenge how to match the protein stability and amino acids availability in the case of every foodstuff containing protein extract.

Table 2. Effect of processing on the structure of legume protein isolate/concentrate.

Physical Process	Effect	Protein Extract	References
Heat treatment	Unfolding and denaturation	Cowpea protein isolates	[108]
		Soybean protein isolate	[109]
	Denaturation and/or subsequent aggregation	Kidney, red and mung beans protein isolates	[110]
	Alteration in secondary and tertiary conformational		
	No dissociation in the protein subunit	Mung bean protein isolate	[30]
Increased α -helix decrease in β -sheets	Soybean Protein Isolate	[111]	
Extrusion treatment	Denaturation	Mung bean protein isolate	[112]
	Increase degree of aggregation and crosslinking		
	Increase in the proportion of β -turn structure	Soy protein concentrate	[113]
		Pea protein isolate	[114]
	Decrease of α -helix and β -sheet	Pea protein isolate	[115]
	Increase in α -helix	Mung bean protein isolate	[112].
	Decrease in β -sheet content		
High pressure	Unfolding /denaturation/ aggregation	Red kidney bean protein isolate	[116]
		Cowpea protein isolates	[108]
		Yellow field pea protein isolate	[117]
	Changes in the secondary structure (β -sheets)	Red kidney bean protein isolate	[118]
Cold plasma	Compact tertiary structure	Grass pea protein isolate	[119]
	Higher ordered secondary structure		
	Dissociation of globulins	Soy protein isolate	[120]
	Oxidation /alteration of the secondary and tertiary structures		
	Alteration of the secondary structures	Peanut protein Isolate	[121]
Irradiation	Modification of the secondary and tertiary structure	Red kidney bean phytohemagglutinin (lectin)	[122]
	Insoluble amorphous aggregates and partially unfolded	Jack bean Concanavalin A	[123]

Table 3. Effect of processing on the functional properties of legume protein isolate/concentrate.

Properties	Treatment	Effect	References
Water-holding capacity	Heat treatment	Increase	[124]
	High hydrostatic pressure	Increase	[108,118]
	Cold plasma	Increase	[121,125]
	Irradiation (X-ray irradiation)	Decrease	[126]
Oil-holding capacity	Heat treatment	Increase	[127]
	Cold plasma	Increase	[121,125]
Protein solubility	Heat treatment	Increase/Decrease Factors: pH, Temperature	[30,127]
	Extrusion	Decrease	[113,114]
	High hydrostatic pressure	Increase/Decrease Factor: Protein isolate's nature, Pressure	[116,128,129]
	Cold plasma	Increase	[119,121,125]
	Irradiation (Gamma/ X-ray irradiation)	Decrease	[126]
Surface hydrophobicity	Heat treatment	Increase	[130]
	Extrusion	Increase/Decrease Factor: Feed moisture, Temperature	[112,113]
	High hydrostatic pressure	Increase	[131]
	Heat treatment	Increase/Decrease Factor: pH	[30,110,124,132]
Emulsifying activity and stability	Extrusion	Stability: Increase	[113,114]
	High hydrostatic pressure	Increase/Decrease Factor: Pressure	[116,118,129]
	Cold plasma	Increase	[120,121]
	Irradiation (Gamma/ Electron beam irradiation)	Increase	[126,133]
	Extrusion	Decrease	[114]
	High hydrostatic pressure	Increase	[129]
	Cold plasma	Increase	[120,121]
	Irradiation (X-ray irradiation)	Decrease	[126]
Gelling capacity	Heat treatment	Increase	[30]
	High hydrostatic pressure	Increase	[108,134,135]
	Cold plasma	Increase	[136]

3.1. Effect of Thermal Processing

3.1.1. Effect of Heat Treatment

Thermal processing is one of the most common processing technologies employed in the food industry. It seems to be the most promising processing method, with a great influence on the structure of proteins [137].

Generally, an increase in temperature may lead to rearrangement and partial unfolding of protein molecules [127]. Several studies showed that thermal processing could result in protein aggregation due to disulfide bond formation [108,109,130,132]. This processing method could induce conformational changes in the cowpea, pea and soybean protein isolates when heated at 90 °C, 80–100 °C, and 90 °C and 120 °C, respectively [108,109,132]. The authors suggested that the high temperature could denature the protein isolate as observed in case of soybean protein [109]. Likewise, 95 °C heat processing for 30

min led to extensive denaturation and/or subsequent aggregation accompanied by decrease in free SH and alteration in secondary and tertiary conformational structure in kidney, red and mung beans protein isolates [110].

However, oven drying (50 °C for 24 h) of mung bean protein isolate showed no dissociation in the protein subunit, suggesting its high thermal stability [30]. The secondary structure of the protein mainly consists of α -helix, β -sheets, strands and turns. The α -helix is present in relatively lower proportion [48]. During processing, the amount of each secondary structure present in protein depends on time, temperature, and intensity of heat processing applied [111]. For instance, heat processing of soybean protein isolate increased α -helix and an overall decrease was observed in β -sheets when compared with non-treated counterpart [111]. The secondary structure of protein might also get affected during formulation and long-term storage [127].

Additionally, studies demonstrated that the surface hydrophobicity and hydrophobic interactions increased after thermal processing [109,130]. Researches showed that the β -sheet exhibited a significant inverse correlation with the surface hydrophobicity of soybean protein isolate when heated above 90 °C [111]. Likewise, a negative correlation between solubility and protein denaturation was reported, as well as the high solubility indicating a high ratio of native proteins and low denaturation [124,138].

Since functional properties are directly linked to the structural characteristics of protein, thermal processing might alter the functionality of protein [124]. For instance, the size of soluble aggregates can influence the functional property of the protein. The heat processed soybean protein isolate with a large proportion of higher than 1000 kDa protein aggregates promoted foaming property 2% concentration. However, when the total amount of protein exceeded 60%, those large protein aggregates promoted emulsification. On the contrary, the medium-size aggregates (670 to 1000 kDa) seemed to have a complementary or supportive role in stabilizing emulsion systems [130].

Heat processing can ameliorate various functionalities such as emulsifying properties [124]. Comparing freeze and convective drying at 40 °C and 50 °C, convective drying had higher values of emulsifying activity indices and emulsifying stability indices compared to the freeze-drying extracts [124]. The study conducted on vacuum-dried lentil protein isolate (at 60 °C and 85 kPa for 48 h) revealed that the processing reduced the solubility with high level of denaturation, which resulted in poor gelling ability when compared to the spray and freeze-dried lentil protein isolate powders. This finding corroborated with the solubility results, since a low concentration of highly soluble protein could make an acceptable gel [138]. Another study revealed that the heat processing of kidney, red and mung beans protein isolates at 95 °C for 30 min improved their solubility and emulsifying activity [110].

For soybean protein isolates, heat treatment using a water bath (90 °C) and autoclave (120 °C) for 15 min denatured the protein isolate, leading to facilitated droplet breakup and decreased droplet size. This treatment was inclined to adsorb on the surface of oil droplets during emulsion storage, which provides a thick and viscoelastic interfacial layer [109]. Another study revealed that heating during spray-drying causes the partial unfolding of globular protein, leading to the exposure of hydrophobic amino acid residues towards the external environment, which increases the adsorption at the oil-water interface and subsequently improved emulsion activity index and stability [30].

Heat processing conditions such as temperature, duration, pH, influence protein structure in different ways, and consequently, their functional properties. For instance, heat pre-treatment (at pH 7.0) enhanced the emulsion-forming ability of isolated pea protein, which led to smaller-sized emulsified oil droplets. On the contrary, pre-treated samples at pH 3.0, 5.0, and 7.0 affected negatively the foaming ability [132].

The difference in thermal stability of proteins depends on species and thermal stability too. For instance, the gelling ability of the globular proteins of pea, soybean and lupin are different, which was due to the variation in their thermal stability. This phenomenon is linked to their structural variations such as amino acid composition, molecular weight,

protein subunits and surface hydrophobicity [139]. The difference in the amino acid composition of the protein sub-fraction could reflect in different conformations, which makes it impossible to generalise the type and amount of bonds, secondary, tertiary or quaternary structures present in protein [137]. However, the proportion and the amount of a particular sub-fraction can affect the final functional characteristics. For instance, the pre-heated mixture of pea vicilin and globulins increased final moduli (elastic (G') and loss (G'') modulus) of the acid gels, whereas legumin-enriched samples displayed low gelling properties [140].

3.1.2. The Effects of Extrusion Treatment

Extrusion is the most versatile processing technology used in the food industry [141]. It is defined as a high-temperature and short-time process [142]. It is a multi-functional and thermal/mechanical process [143] that enables the control of different processes such as mixing, heating, cooking, shearing, and shaping of products [141]. It can be used to produce pre-cooked and dehydrated foods [144] and has the ability to develop products with better nutritional, functional and sensory characteristics [141]. Several products are made using extrusion technology such as breakfast cereals, savory snacks, crispy flat-bread, pre-cooked flours, cereal-based baby food, and textural proteins [141,145]. Currently, many studies are being focussed on the production of legume protein meat extenders and meat analogues using extrusion technology [112,146].

During extrusion treatment, the free sulfhydryl content, particle size, secondary structure, and tertiary structure of protein isolate might change, influencing the functional properties of the legume protein isolate [114,115]. Further, the extrusion could influence the interaction of legume proteins with non-protein matrix, such as microstructure, solubility and emulsifying capacity of mixtures of starch–pea protein isolate [147], soy protein concentrates–rice [148], soy protein isolate–corn [149] and isolated/textured soybean proteins–pectin [150]. These interactions could be affected by the barrel temperature, screw speed, feed rate and feed moisture [146].

The hydrophobic interactions, hydrogen bonds, disulfide bonds and their interactions are the most critical factors that collectively hold the protein structure [114,151]. This structural assembly within protein is destroyed due to the triple action of high temperature, pressure and shear force, ultimately resulting in protein depolymerisation. These depolymerised proteins recombine under specific temperature and moisture content conditions to form protein aggregates [114].

Among all factors, the feed moisture plays a major role during the extrusion process. For instance, denaturation and formation of mung bean protein aggregates could be controlled by manipulating feed moisture content during extrusion [112]. An extrusion process with low feed moisture content of 30% resulted in extensive protein denaturation, resulting in an increased degree of aggregation and crosslinking [112]. However, similar effects were observed for soy protein concentrate at lower moisture content (18 and 25%) [113]. According to these authors, further increase in feed moisture content to 30%, could cause an increase in the proportion of β -turn structure [113]. For pea protein isolate, the reduction of β -sheet and α -helices after extrusion under 150 °C and 26% of moisture feed was associated with the changes in the secondary structure [115]. Similar reduction in the β -sheet structure of extruded pea protein isolate was observed at 30% moisture content [114]. The decrease in β -sheet and α -helical structures were related to the increase in protein aggregates [115]. Therefore, it can be interpreted that the changes in the secondary structure during extrusion depend on the protein source as well as on the feed moisture content.

Increasing moisture content could increase the interactions between disulfide bonds with hydrogen bonds and hydrophobic interactions, which reduces the degree of aggregation [142]. The feed moisture content ~49% promoted partial protein unfolding and formation of small aggregates. This partial unfolding could be manifested by increase in α -helix and disulfide bond content and a decrease in β -sheet content [112]. However, studies

showed that the high-moisture extrusion could result in the denaturation of protein concentrate [152]. A complete denaturation of mung bean protein isolate was reported along with formation of large protein aggregates when extruded at 60% feed moisture content [112]. A reduction in the α -helices and β -turn was also noticed as this moisture content [114]. Hence, the feed moisture content could directly influence the structural organisation of protein, which might further depend on factors such as temperature.

During the extrusion process, high temperature and shearing causes protein dissociation inside of the equipment; water loss and aggregation after the molten mass exits the die and a sudden drop in temperature. This aggregation usually decreases protein solubility, and hence influences hydrodynamic properties, which are related to the formation of supra-molecular assemblies [153]. Studies suggested that the temperature and ingredients during extrusion could affect the pea protein solubility [154]. Extrusion of pea protein isolate at different moisture feed (30, 40, 50, and 60%) and temperature (60 °C, 100 °C, 120 °C, 140 °C, and 175 °C) resulted in protein aggregation, which reduced its solubility [114]. Similar effects were observed for soy protein after processing at 30, 60, 80, 115, and 130 °C with 35% of moisture feed [153]. However, for pea protein isolates, a higher feed moisture content of 55% reduced its solubility [154].

Extrusion cooking at 130–170 °C with 26–35% moisture, reduced solubility of pea protein isolate [115]. Likewise, at 18 and 25% moisture content and 110, 130 and 160 °C temperatures, extrusion cooking decreased the protein solubility of soy protein concentrate compared to the native protein [113]. However, increasing temperature from 110 to 160 °C at 18 and/or 25 % moisture content gradually enhanced the solubility of soy protein [113]. The decrease in the solubility might be related to the formation of heat-induced insoluble aggregates due to the interaction between hydrophobic groups of unfolded protein structures [113]. On the contrary, extrusion of mung bean protein at 144.57 °C with 49.3% feed moisture content improved its solubility [112]. The partial denaturation and formation of small-sized protein aggregates along with increased interactive surface region between protein and water molecules can positively affect the solubilisation of protein, thus improving the overall solubility [112]. Therefore, it can be concluded that the solubility of protein isolate depends on the moisture feed and temperature, which can influence the aggregate size or nature, and ultimately the bioavailability of amino acids from proteins.

The structural modifications during the extrusion process can affect the protein solubility and hence its hydrodynamic properties such as water absorption capacity, emulsification and foaming properties. For example, a higher water absorption capacity of isolated soy protein and mung bean protein was attributed to the higher amount of protein solubility [155]. Similarly, the protein denaturation of the extruded faba bean concentrate (high extrusion moisture content) was translated by a high-water binding capacity [152]. In addition to the solubility, the hydrophilic–hydrophobic amino acids and polar/non-polar amino acids can serve as a useful indicator of the functionality of protein, such as water and fat absorption, and surfactant properties [30,155]. The study showed that the temperature at the last heating zone of extrusion equipment and moisture content significantly affected the water binding capacity of the meat analogues, which was in negative relation with temperature (120 °C to 140 °C) [152].

The foaming properties of a protein can be characterized by measuring the foaming capacity and foaming stability [156]. Foaming capacity is affected by the solubility of the protein isolate [114]. Compared with native pea protein, the foaming capacity of extrudate of pea protein hydrated at 30, 40, 50 and 60% moisture content reduced due to the decrease in its solubility. However, the extrudate with 30% moisture content demonstrated the highest foaming stability [114]. This finding is correlated to the irreversible thermal denaturing of proteins, which might result in the unfolding of the peptide chain. The unfolding phenomenon led to the structural rearrangement within the machine barrel to form pro-

tein aggregates. These aggregates are adsorbed on the air–water interface to create a network structure, thus improving the stability of the foam. Additionally, the insoluble protein particles play a favourable role in the stability of the foam [114].

Moisture content seems to have a lubricant role during the extrusion process. At 30 and 40% moisture content, protein is denatured by raising the temperature from 60 to 175 °C, which exposes hidden hydrophobic groups, thus increasing the emulsifying activity index and emulsifying stability index [114]. In contrast, the increase in the extrusion temperature from 110 to 130 °C decreases the emulsifying stability of extruded soy protein emulsion, which could be related to the formation of larger particles when the extrusion temperature increases to 130 °C [113].

Compared with soy protein concentrate extruded at 25% of feed moisture, soy protein concentrate extruded at 18% feed formed unstable emulsion that can be related to the particle droplet size in emulsions. The emulsions fabricated by extruded soy protein concentrate at 25% feed moisture had small droplets size which promotes more stability against coalescence and flocculation [113]. Similarly, the droplet size of the emulsions produced by soy protein was reduced after extrusion treatment at 30, 60, 80, 115, and 130 °C and 35% of moisture feed [153].

The surface hydrophobicity is correlated to functional properties of protein such as emulsification and foaming [112]. It is a physicochemical property that greatly determines the tendency of protein molecules to aggregate and thus to lose solubility [157]. The soy protein concentrate extruded at 110, 130 and 160 °C with 18 and 25% moisture contents exhibiting lower surface hydrophobicity compared with the untreated one. This was attributed to the protein–protein interaction inside the extruder and the formation of protein aggregates, which hide the hydrophobic sites on the surface of protein and inhibit its accessibility [113]. In contrast, a slight increase in the soy protein concentrate’s superficial hydrophobicity was reported after extrusion with single-screw extruder at 30, 60, 80, 115, and 130 °C and 35% moisture [153]. For mung bean proteins, the increase in feed moisture content during extrusion effectively increased surface hydrophobicity by denaturing them and exposing more hydrophobic domains, consequently making the protein structure more flexible [112]. Moreover, the increase in the moisture content from 18 to 25% during extrusion-enhanced surface hydrophobicity, which was related to the formation of larger protein aggregates [113].

3.2. Effect of Non-Thermal Technologies

3.2.1. Effect of High-Pressure Treatment

High-pressure processing (HPP) is an interesting alternative to traditional processing [158]. It is a non-thermal technology that has several applications including the modification of proteins structure [159]. High hydrostatic pressure (HHP), high-pressure homogenization (HPH) and ultra-high-pressure homogenization are improved forms of HPP [160].

HPP induces gradual unfolding and denaturation in the structure of protein molecules [108,116]. This effect was observed when the yellow field pea protein isolate was treated under 600 MPa [117]. Interestingly, HPP at similar pressure (400 and 600 MPa) completely denatured soybean protein isolates [131]. Indeed, for the same protein isolate a greater extent of protein denaturation, aggregation, and network formation occurred with increasing pressure level (250–550 MPa for 15 min), due to protein tertiary and quaternary conformation changes. Pressure treatments above 350 MPa induced protein denaturation and subsequent gel structure formation [135]. Another study conducted on red kidney bean protein isolate demonstrated that the HPP had little effect on its structure when treated below 600 MPa. However, the increase in pressure up to 600 MPa led to the aggregation of protein molecules/monomers [118]. Likewise, HPH could cause unfolding of lentil proteins at 50–150 MPa [129]. On the contrary, a study conducted on lentil protein

isolate showed that HPP could not significantly alter protein structure, even when exposed at high level of 600 MPa for 15 min. The authors suggested that HPP might not have affected covalent bonds, and thus, the primary protein structure did not alter after the processing [161].

Since HPP can alter protein structure, it can induce several subsequent changes in its physico-chemical properties, such as hydration, hydrophobicity, and hydrophilicity [159]. Some of the researches employ solubility and surface hydrophobicity as the indices of degree of denaturation caused by high pressure; [131,162] reported that HPP (200–600 MPa at pH 8) increased surface hydrophobicity and protein aggregation accompanied with a decrease in free SH and partial unfolding of 7S and 11S fractions of soybean protein isolate. Interestingly, HPP at pH 3 resulted in partial denaturation of protein, leading to decreased thermal stability, and increase in protein solubility [131].

Protein solubility is a critical factor in the hydrolytic yield of proteins that may get affected by HPP. HPP of lentil protein concentrates at 100, 200 and 300 MPa exhibited similar soluble protein content as control. However, increase in pressure (400 and 500 MPa) significantly reduced the soluble protein content, probably due to the formation of insoluble protein aggregates [128]. On the contrary, the solubility of red kidney bean protein isolate was significantly improved at 400 MPa or higher, possibly due to formation of soluble aggregate from insoluble precipitate [116]. A study conducted on lentil protein isolate demonstrated that the HPH at 100 MPa could increase its solubility as a consequence of protein unfolding and increased solvent–protein interactions. However, HPH at 150 MPa significantly decreased the solubility, probably as a consequence of the over-processing effect of pressure [129].

In the case of water-holding capacity and gelation property, HHP (200, 400 and 600 MPa) was found to be more efficient than thermal processing in cowpea protein [108]. HHP led to the generation of compact aggregates at high protein concentrations (10.5–12.0 g/100 g), which would have increased the ability of protein to interact through hydrophobic interactions, ultimately producing more elastic gels [134]. HHP aids the production of gels at lower temperatures, which make cowpea proteins useful to texturize hot-serving foodstuff [134].

Emulsification property is another functional property of the proteins which might be affected by HPP [116]. Emulsification is generally known as dispersing one liquid in another immiscible liquid [163]. Factors such as surface hydrophobicity, partial denaturation and disordered structure can affect the potential of protein for the adsorption at the oil–water interface [164]. HPP of kidney bean protein isolate at 200 and 400 MPa significantly increased its emulsifying activity and stability index [116]. Another study showed similar effects of HPP on soybean protein, particularly on β -7S and A-11S polypeptides. However, the authors suggested that the alterations in emulsification properties could be pH dependent [164]. Since the pH determines the net protein charge, especially below and above the isoelectric point, it might influence the adsorption phenomenon [117]. The HPP at neutral pH could improve the emulsifying activity of soy proteins isolate [162]. However, HHP of yellow field pea protein isolate at pH 3.0 improved emulsion quality with oil droplet size of 26–68 μ m, along with higher foaming capacity [117]. Apart from pH, intensity of pressure can also influence the functional properties of legume proteins. A study conducted on red kidney bean protein isolate revealed that the functional properties such as water-holding capacity, and emulsifying increased with the increase in pressure intensity from 200 to 600 MPa, unlike the foaming properties that decreased [118].

3.2.2. Cold Plasma

Generally, cold plasma is being employed for improving antimicrobial activity, structural alterations, surface decontamination, etc. At present, the application of cold plasma is being expanded by combining it with other technologies such as nanotechnology, pulsed electric field, pulsed light and ultrasound [165]. Cold plasma is a non-thermal pro-

cessing technology [136,166], which utilizes energetic and reactive gases [167]. In this technology, a simple gas (such as air or nitrogen) may be used, or the system may rely on a mixture of noble gases, such as helium, argon, or neon [168]. Through ionization of inducer gas with electrical discharges, cold plasma consisting of electrons, ions, neutral particles, and free radicals can be obtained [120].

Studies demonstrated that cold plasma could change the secondary and tertiary structure of grass pea protein isolate, imparting more thermodynamically stable structure. This was attributed to the increase in hydrogen bonding in the α -helix and β -sheet structures [119]. However, another study reported that the atmospheric cold plasma induced reactive oxygen species-mediated oxidation of soy protein isolate, which ultimately altered its secondary and tertiary structures [120]. For peanut protein isolate, the primary structures were unaltered under cold plasma treatment, unlike the secondary structure [121]. The modification of the secondary structure manifested in an increase in the content of β -sheets and random coil and a decrease in the content of α -helix and β -turn [121].

Atmospheric pressure plasma could improve the functional properties of pea protein in term of solubility [125]. However, the combination of the thermal and plasma-induced pH conditions lead to an increase in protein solubility to 327% [125]. Concomitant to the previous mentioned study, cold plasma processing of grass pea protein isolate for 60 s, increased the protein solubility. It was ascribed to the reduction in protein particle size and the increase in protein surface charge [119]. Similarly, the atmosphere cold plasma treatment improved the solubility of peanut protein isolate [121,169]. The maximum of solubility was reached after 7 min of treatment, with a 12.17% increase comparing with untreated samples [169].

Additionally, the unfolding of protein structure, exposure of active sites, and binding of water micelles to protein molecules could improve the water-holding capacity and emulsion stability of the protein [121]. Another study conducted by [136] showed an improvement in gelling capacity of cold plasma-treated pea protein. Their finding indicated that native pea protein concentrate could not form gel at 90 °C, while the cold plasma could induce a better gelling property. The gels from the processed sample exhibited homogeneous three-dimensional network structure with interconnected macropores. However, gels prepared from processed samples at 80 and 90 °C possessed good mechanical strength and viscoelasticity along with high water-holding capacity [136].

Another study showed that the cold plasma processing could improve the interfacial and emulsifying properties of grass pea protein isolate in terms of thermodynamic stability of protein on interface and globulin dissociation. It also increased oil-droplet surface electrical charge [119]. For peanut protein isolate, the improvement of the emulsifying properties was related to the increased random coils [121]. The intensity of changes in the functional properties of protein seems to be time dependent. For instance, cold plasma processing of soybean protein isolate for 1 min improved its emulsifying and foaming properties. However, prolonged processing for 5 and 10 min led to the subsequent reduction in both interfacial characteristics [120].

3.2.3. Irradiation

Irradiation is a highly efficient and low cost technology with the advantages of short processing time [105]. Gamma irradiation, X-rays and electron beam are the most commonly used radiation treatments in food industries. The term 'food irradiation' is used to describe a process where food is exposed to ionizing energy, just utilizing gamma photons, machine-generated X-rays or accelerated electrons [170]. In case of γ -irradiation, the rays are emitted from the radioisotopes ^{60}Co and ^{137}Cs , or high energy electrons; however, X-rays are produced by machine sources [171]. Depending on the absorbed radiation dose, irradiation can reduce the storage losses, extend the product shelf life and improve microbiological and parasitological safety of foods [171]. Despite being declared as safe and nutritionally adequate by numerous international expert groups, the commercial use

of irradiation is still limited [170]. Therefore, there is limited literature available on the effect of γ -radiation on the properties of protein, specifically legume proteins.

The treatment of soy protein isolate by γ -irradiation caused the disruption of the ordered structure and induced degradation, cross-linking, and aggregation of the polypeptide chains [172]. Likewise, a high dose of γ -irradiation of jack bean Concanavalin A (lectin) reduces its β -rich secondary structures. This indicates an accumulation of completely unfolded and fragmented peptides that have a tendency to form insoluble amorphous aggregates [123]. Similarly, γ -irradiation of red kidney bean phytohemagglutinin resulted in complete destruction of secondary and tertiary structure at a dose superior or equal to 10 kGy [122].

Furthermore, studies showed that irradiation treatments, such as γ -irradiation, X-rays and electron beam, could influence the functional properties of legume proteins. The treatment of soy proteins concentrates with γ - (1 kGy intensity) and X-ray irradiation for 15 min enhanced the emulsification activity and foaming capacity; however, it decreased the protein solubility [126]. Similarly, the electron beam irradiation treatment of pea protein hydrolysates increased its emulsifying properties with a maximum effect at 50 kGy. Additionally, both foaming capacity and stability increased with the increase in irradiation dose [133]. A reduction in the water-binding capacity of the gamma-irradiated soy protein concentrate was experienced, which was attributed to the modification in the secondary and tertiary structures [126].

4. Effect of Extraction and Processing on the Nutritional Characteristics

The nutritional value of plant-based food depends on the amount of proteins, the specific distribution of the amino acids, and, most importantly, the bioavailability of amino acids [20,23]. The latter is determined by digestibility, that is measured to assess the susceptibility of protein to proteolysis. Therefore, highly digestible proteins can provide more amounts of amino acids for absorption due to their higher degree of hydrolysis, thereby providing better nutritional value [173].

In general, legume proteins have low in-situ digestibility, which could be related to both external and internal factors [174]. The physical inaccessibility due to the entrapment in, for example, intact cell structures, the presence of antinutritional factors [174] and the protein interactions with other compounds such as carbohydrates, lipids and especially anti-nutritional factors (anti-nutritional proteins (trypsin inhibitors, lectins, etc.) and the anti-nutritional chemicals (tannins, phytates, and polyphenols)) are among the external factors [23]. These factors could be influenced by the extraction and/or during other processing steps of legumes grains (Figure 4).




Figure 4. Effect of extraction and processing techniques on the digestibility of the protein isolate/concentrate [20,21,30,49,116,120,133,137,161,174–182].

4.1. Effect of Extraction

The extraction methods, can influence the *in vitro* protein digestibility [22]. The changes in the protein digestibility could be related to the modifications in the structure during processing. The literature suggests an increase in protein digestibility after their extraction, which is related to the changes in exposure of peptide bonds and increase in accessibility to susceptible protein sites [182]. The higher digestibility of the lentil protein isolate in comparison with lentil flour might be attributed to variations in the processing such as thermal treatment and spray drying for isolate preparation [182]. However, the authors emphasized that this difference in digestibility might largely depends on the reduction of anti-nutritional factors [182]. Hence, the extraction of protein can significantly influence its digestibility. For instance, protein extracted from African mesquite bean using isoelectric precipitation exhibited low digestibility compared to micellization precipitation. This was suggested to be the consequence of the high pH achieved by sodium hydroxide, which resulted into amino acid cross-linking, and subsequent formation of lysinoalanine from lysine and dehydroalanine via serine and cysteine degradation [22].

Furthermore, approaches including digestibility model systems and quantification methods, can offer further details on the digestibility of protein isolates [182].

Additionally, protein extraction has been found to reduce the content of anti-nutritional factors that might subsequently lead to improved digestibility [22,183]. The isolation techniques such as alkaline extraction, iso-electric precipitation and ultrafiltration seemed to lower trypsin inhibitor in lentil protein concentrates and isolates [182,183]. Moreover, the size of the protein bodies seems to be associated with its digestibility. The study showed that the protein extracted from lentils using isoelectric precipitation had a lower amount of larger molar mass proteins (>699 kDa), compared with the dry-milled flour, that is often less digestible and may be often associated with intolerance and/or allergy [184].

4.2. Effect of Processing

4.2.1. Effect of Thermal Processing

Heat Treatment

Proteins vary in proteolytic susceptibility and hence their digestibility, owing to the differences in their origin, nature and structure [23]. The main structural component of legume protein is β -sheets and the literature showed its strong inverse correlation with protein digestibility [185]. The disappearance of intramolecular β -sheets could lead to the increase in protein digestibility since it influences the accessibility of enzymes for digestion [137]. Further, the unfolding of the globular protein subunits could enhance its digestibility as it might expose the otherwise buried hydrophobic groups to pepsin [186]. A study conducted on vicilin-rich protein isolated from *Phaseolus* legumes (kidney bean, red bean and mung bean) showed that the thermal treatment improved their nutritional property, which was largely dependent on their amino acid composition and protein aggregation. The heat-induced protein aggregation varied with the type of protein isolate, and the formation of disulphide bonds.

The nutritional relevance of the legume protein such as the protein-specific amino acid content could also be influenced by the thermal processing [49,187]. The roasting of bambara groundnut protein isolate reduced its protein content, which was attributed to the protein aggregation. The larger protein aggregates can be removed during centrifugation before isoelectric precipitation step [49]. The thermal processing seemed to improve the nutritional properties of *Phaseolus* legumes and promoted protein aggregation [187]. Another important factor affecting the nutritional property would be the anti-nutritional factors, which could be reduced by using thermal processing [21]. The thermal treatment of commercial soybean protein products could inactivate up to 80% of trypsin inhibitory activity [21]. However, in case of lentil and faba bean protein concentrates, the thermal treatment at 95 °C for 15 min could lead to reduction in trypsin inhibitor activity up to 86% and 78%, respectively, when compared with untreated samples [178]. Interestingly, there is a possibility that thermal and/or alkaline processing led to the formation of compounds such as Maillard reaction compounds, D-amino acids and/or lysinoalanine, which reduced the protein digestibility by up to 28% in rats and pigs [21].

Extrusion Treatment

The primary reason for the improvement in digestibility upon extrusion has been linked to the inactivation of anti-nutritional compounds and to the thermal denaturation of proteins due to high temperatures and shearing [180,188]. The reduction in anti-nutritional factors was noticed during the processing of protein concentrates and extruded products [180,181].

The extrusion process significantly improves the *in vitro* protein digestibility of lupin protein extrudates compared to that of raw material mixture [180]. Similarly, the extrusion cooking of soybean protein concentrate significantly improved the *in vitro* protein and

starch digestibilities and significantly reduced some anti-nutritional factors of the formulated meals [181]. Partial denaturation of mung bean protein isolate improved its digestibility due to the increased α -helix along with reduced β -sheet content, while stronger gel-forming behaviour was also experienced due to the increased disulfide bond content, frequency-independent viscoelasticity and shear thinning effect [30]. However, another study on the mice demonstrated that the extrusion process did not decrease the protein digestibility of soy protein isolate, and hence could not affect the growth rate of male and female mice [189].

Barrel temperature and feed moisture are two other factors, which can affect the protein digestibility during the extrusion process. The increase in barrel temperature was found to decrease the *in vitro* protein digestibility of lupin significantly, which could be due to non-enzymatic browning reactions and thermal crosslinking [180]. On the other hand, feed moisture content is an essential factor that should be adjusted during the extrusion of protein ingredients to achieve desirable quality and product performance as a vegetarian-based meat extender [30]. In particular, increasing water feed from 40% to 55% did not show significant difference in *in vitro* protein digestibility, whereas the sample with 68% water feed exhibited significantly higher *in vitro* protein digestibility. At higher water feeds (~60%), protein aggregation could have been reduced, which enhanced protein digestibility [180].

Concerning the amino acid composition, the amounts of sulfur-containing amino acids, including cystine and methionine of plant proteins such as soybean and pea tend to increase after extrusion, while the amounts of acid-stable amino acids of texturized vegetable proteins tend to decrease [155]. In contrast, [154] found that the extrusion had no effect on the degree of hydrolysis and amino acid composition indicating that the thermal and mechanical energy during extrusion did not cause the formation of peptide bonds or the degradation of amino acids due to the Maillard reactions.

4.2.2. Effect of Non-Thermal Processing

The Effect of High Hydrostatic Pressure Processing

High hydrostatic pressure (HHP) involves absorption of water and induction of heat within the protein, which result in changes in its conformation, thereby impairing its susceptibility to proteolysis [190]. The changes in protein structure during HHP, such as denaturation and formation of random coil, can have a positive effect on protein digestibility [137]. However, the aggregation and Maillard reactions during HHP can lead to a decrease in digestibility [137,174].

The pressure-induced gels (600 MPa/5 °C/4 min) exhibited more accessibility to gastric proteolysis than heat-induced gels [177]. For lentil protein, pressure treatment (300 MPa for 15 min) increased the enzyme accessibility to cleavage sites of protein, which improved the degree of hydrolysis and their antioxidant activity [161]. In contrast, pressure treatment of red kidney bean protein isolate with 200 MPa or above significantly decreased its digestibility. The decrease in digestibility was due to the protein unfolding or denaturation and protein aggregation but it was noticed only when the pressure was increased above 200 MPa and the incubation time was long (120 min) [116].

Cold Plasma

The digestibility and bioavailability of the amino acids is related to the nutritional properties of the protein since their degradation; fragmentation or cross-linking reactions can influence the digestibility of proteins [176]. Cold plasma treatment may trigger changes in the amino acid side chains, such as the -SH groups [119,121]. In general, these changes are manifested by a reduction in the free -SH groups due to oxidation of proteins [120]. The aromatic and sulfur-containing amino acid side chains are known to be particularly susceptible to oxidation, given the extreme sensitivity of cysteine residues

[119,121,191]. Similar reduction in cysteine was reported in grass pea protein isolate, probably due to the oxidative effect of ozone produced during plasma discharge [119]. Plasma treatment of pea protein isolate resulted in alteration in fluorescence spectra of tryptophan, which was attributed to the oxidation and the fluorescence quenching [125]. Similarly, cold plasma treatment of pea protein concentrate isolate decreased its fluorescence intensity, which was related to tryptophan oxidation due to the elimination of hydrogen atom on its indole ring by radicals [136]. The cold plasma induced-oxidation of sulfur-containing amino acid residues, such as cysteine and methionine resulted in the degradation of sulfhydryl groups [120].

During cold plasma treatment, the oxidation induced loss of amino acids that can lead to a reduced nutritional quality of protein [176]. However, the effect of the oxidation is dose-dependent [179]. Mild oxidation might enhance the functionality and might not have a significant impact on the nutritional quality of proteinaceous foods since the extent of the loss of essential amino acids is limited [179]. Several studies reported the reduction of allergenicity of proteins after the application of non-thermal treatments due to the changes in the antibody's binding ability [192]. Among these treatments, the application of cold plasma was successful in inactivating allergens in most proteins [176]. For soybean protein isolate, the application of atmospheric cold plasma at 120 Hz for 5 min decreased the level of IgE-binding by up to 75% compared to the control [120]. Another study stated that the cold atmospheric pressure plasma was the most effective in the reduction of Gly m5 immunoreactivity comparing with pulsed ultraviolet light and gamma-irradiation (3–100 kGy) [175].

Irradiation

Similar to the functional properties, the effect of irradiation on nutritional aspect of protein concentrates is less studied area and limited to γ -irradiation and electron beam. The literature suggested that irradiation levels up to 10 kGy could be an effective measure in deactivating anti-nutrients such as protease inhibitors, lectin, phytic acid, non-starch polysaccharides and oligosaccharides without altering the nutritional quality of the food [178]. Higher than 25 kGy, γ -irradiation treatment of soy protein isolate reduced its allergenic proteins [175]. Moreover, 30–50 kGy irradiation reduced the potential of red kidney bean lectin to agglutinate rabbit blood erythrocytes [122]. However, [20] reported that the γ -irradiation negatively affected protein digestibility, which could be due to the destruction of aromatic and sulfur containing amino acids. On the other hand, the electron beam irradiation influences protein structure, which could be manifested in the form of improved antioxidant activity pea protein hydrolysates [133]. Despite these positive changes of irradiation, the legislation has allowed 10 kGy as maximum dose limit of γ -irradiation for specific commercial food items [175].

5. Conclusions

Due to the rapidly growing numbers of the world population, the consumer-specific health awareness, the scientific knowledge advancement in the field of nutrition and improvement of the food manufacturing concepts and technologies, it clearly indicates that the proteins are once again back in the forefront of research. On a larger scale, we also have to address the global warming-induced changes to crop production and the declining health conditions of populations. Moreover, there are other contrasting and even moral issues to consider related to the relevance of proteins in the food chain, including those of food waste, the declining popularity of animal foodstuffs and the increasing relevance of a plant-based diet, all of which is more specific to developed countries. Conversely, in developing countries, food security and safety emerge as the major restrictive factors that must be faced and through multiple solutions ensure the sustainability of social and economic progress. We also have to admit that the focus on diets advocating for proteins being the most critical macronutrients, is shifting towards the bioavailability of essential and non-essential amino acids [193]. The utmost significance of some amino

acids for vital functions is obtaining increasing scientific evidences. For example, amino acids such as glutamine and glutamate were proposed to interfere with the onset of autism spectrum disorder [194], but the glutamate can also regulate the glutamylation of microtubules affecting cytoskeletal functions and cancer metastasis [195].

Legumes are one of the most important sustainable sources of protein, but, by any means, attention should be paid to their digestibility, knowing that several endogenous and exogenous factors have negative influences. The prime examples of these factors are the interaction of proteins with starch and lipids or the damaging effect of the harsh extraction conditions (pH and temperature). Nevertheless, physical food processing techniques, such as wet or dry heat treatment, irradiation or high-pressure treatment, could produce protein extract matrices that would confer additional nutritional benefits regarding the delivery of amino acids, dietary fibers and polyphenols.

Author Contributions: Conceptualization, P.S. and C.N.; writing—original draft preparation, C.N. and J.S.; writing—review and editing, P.S., M.H.K. and E.M.; supervision, P.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. FAO. Available online: <https://www.fao.org/news/story/en/item/35571/icode/> (accessed on 14 September 2022).
2. Anzani, C.; Boukid, F.; Drummond, L.; Mullen, A.M.; Álvarez, C. Optimising the use of proteins from rich meat co-products and non-meat alternatives: Nutritional, technological and allergenicity challenges. *Food Res. Int.* **2020**, *137*, 109575. <https://doi.org/10.1016/j.foodres.2020.109575>.
3. Ingerslev, A.K.; Rasmussen, L.; Zhou, P.; Nørgaard, J.V.; Theil, P.K.; Jensen, S.K.; Lærke, H.N. Effects of dairy and plant protein on growth and growth biomarkers in a piglet model. *Food Funct.* **2021**, *12*, 11625–11640. <https://doi.org/10.1039/d1fo02092g>.
4. Kurpad, A.V. Protein: Quality and Sources. In *Encyclopedia of Human Nutrition*, 3rd ed.; Caballero, B., Ed.; Elsevier Ltd.: Amsterdam, The Netherlands, 2013; pp. 123–130.
5. Zhang, C.; Guan, X.; Yu, S.; Zhou, J.; Chen, J. Production of meat alternatives using live cells, cultures and plant proteins. *Curr. Opin. Food Sci.* **2022**, *43*, 43–52. <https://doi.org/10.1016/j.cofs.2021.11.002>.
6. Onwezen, M.C.; Bouwman, E.P.; Reinders, M.J.; Dagevos, H. A systematic review on consumer acceptance of alternative proteins: Pulses, algae, insects, plant-based meat alternatives, and cultured meat. *Appetite* **2021**, *159*, 105058. <https://doi.org/10.1016/j.appet.2020.105058>.
7. Sabaté, J.; Soret, S. Sustainability of plant-based diets: Back to the future. *Am. J. Clin. Nutr.* **2014**, *100*, 476S–482S. <https://doi.org/10.3945/ajcn.113.071522>.
8. Duranti, M. Grain legume proteins and nutraceutical properties. *Fitoterapia* **2006**, *77*, 67–82. <https://doi.org/10.1016/j.fitote.2005.11.008>.
9. Sim, S.Y.J.; SRV, A.; Chiang, J.H.; Henry, C.J. Plant Proteins for Future Foods: A Roadmap. *Foods* **2021**, *10*, 1967. <https://doi.org/10.3390/foods10081967>.
10. Hertzler, S.R.; Lieblein-Boff, J.C.; Weiler, M.; Allgeier, C. Plant Proteins: Assessing Their Nutritional Quality and Effects on Health and Physical Function. *Nutrients* **2020**, *12*, 3704. <https://doi.org/10.3390/nu12123704>.
11. Gutiérrez-Urbe, J.A.; Guajardo-Flores, D.; López-Barrios, L. Legumes in the Diet. In *Encyclopedia of Food and Health*, 1st ed.; Caballero, B., Finglas, P.M.; Toldra, F., Eds.; Academic Press: Oxford, UK, 2016; pp. 539–543.
12. Hasanuzzaman, M. *Legume Crops: Prospects*; BoD—Books on Demand: London, UK, 2020; p. 182.
13. Simpson, M.G. 8—Diversity and Classification of Flowering Plants: Eudicots. In *Plant Systematics*, 2nd ed.; Simpson, M.G., Ed.; Elsevier Inc.: New York, NY, USA, 2010; pp. 275–448.
14. Singhee, D. Review on Natural Dyes for Textiles from Wastes. In *Chemistry and Technology of Natural and Synthetic Dyes and Pigments*; IntechOpen: Rijeka, Croatia, 2020.
15. Semba, R.D.; Ramsing, R.; Rahman, N.; Kraemer, K.; Bloem, M.W. Legumes as a sustainable source of protein in human diets. *Glob. Food Secur.* **2021**, *28*, 100520. <https://doi.org/10.1016/j.gfs.2021.100520>.
16. Kelemu, S.; Cardona, C.; Segura, G. Antimicrobial and insecticidal protein isolated from seeds of *Clitoria ternatea*, a tropical forage legume. *Plant Physiol. Biochem.* **2004**, *42*, 867–873. <https://doi.org/10.1016/j.plaphy.2004.10.013>.
17. Somers, D.A.; Samac, D.A.; Olhoft, P.M. Recent Advances in Legume Transformation. *Plant Physiol.* **2003**, *131*, 892–899. <https://doi.org/10.1104/pp.102.017681>.

18. National Research Council. *Tropical Legumes: Resources for the Future*; The Minerva Group, Ed.; The Minerva Group: New York, NY, USA; Hong Kong, China, 2002; p. 340.
19. Gresshoff, P.M.; Hayashi, S.; Biswas, B.; Mirzaei, S.; Indrasumunar, A.; Reid, D.; Samuel, S.; Tollenaere, A.; van Hameren, B.; Hastwell, A.; et al. The value of biodiversity in legume symbiotic nitrogen fixation and nodulation for biofuel and food production. *J. Plant Physiol.* **2015**, *172*, 128–136. <https://doi.org/10.1016/j.jplph.2014.05.013>.
20. Sá, A.G.A.; Moreno, Y.M.F.; Carciofi, B.A.M. Food processing for the improvement of plant proteins digestibility. *Crit. Rev. Food Sci. Nutr.* **2020**, *60*, 3367–3386. <https://doi.org/10.1080/10408398.2019.1688249>.
21. Sarwar Gilani, G.; Wu Xiao, C.; Cockell, K.A. Impact of Antinutritional Factors in Food Proteins on the Digestibility of Protein and the Bioavailability of Amino Acids and on Protein Quality. *Br. J. Nutr.* **2012**, *108*, S315–S332. <https://doi.org/10.1017/S0007114512002371>.
22. Falade, K.O.; Akeem, S.A. Physicochemical properties, protein digestibility and thermal stability of processed African mesquite bean (*Prosopis africana*) flours and protein isolates. *Food Meas.* **2020**, *14*, 1481–1496. <https://doi.org/10.1007/s11694-020-00398-0>.
23. Drulyte, D.; Orlien, V. The Effect of Processing on Digestion of Legume Proteins. *Foods* **2019**, *8*, 224. <https://doi.org/10.3390/foods8060224>.
24. Rubio, L.A.; Clemente, A. In vivo (rat) and in vitro (Caco-2 cells) absorption of amino acids from legume protein isolates as compared to lactalbumin or casein. *Arch. Anim. Nutr.* **2009**, *63*, 413–426. <https://doi.org/10.1080/17450390903169237>.
25. Arntfield, S.D.; Maskus, H.D. 9—Peas and other legume proteins. In *Handbook of Food Proteins?* 1st ed.; Phillips, G.O., Williams, P.A., Eds.; Elsevier Ltd.: London, UK, 2011; pp. 233–266.
26. Abbas, Y.; Ahmad, A. Impact of processing on nutritional and antinutritional factors of legumes: a review. *Desalination* **2018**, *2*, 199–215.
27. Patterson, C.A.; Curran, J.; Der, T. Effect of Processing on Antinutrient Compounds in Pulses. *Cereal Chem.* **2017**, *94*, 2–10. <https://doi.org/10.1094/CCHEM-05-16-0144-FL>.
28. Tiwari, B.K.; Gowen, A.; McKenna, B. *Pulse Foods: Processing, Quality and Nutraceutical Applications*, 2nd ed.; Academic Press: Cambridge, MA, USA, 2020; pp. 536.
29. Zhu, H.; Tang, H.; Cheng, Y.; Li, Z.; Tong, L. Electrostatic separation technology for obtaining plant protein concentrates: A review. *Trends Food Sci. Technol.* **2021**, *113*, 66–76. <https://doi.org/10.1016/j.tifs.2021.04.044>.
30. Brishti, F.H.; Chay, S.Y.; Muhammad, K.; Ismail-Fitry, M.R.; Zarei, M.; Karthikeyan, S.; Saari, N. Effects of drying techniques on the physicochemical, functional, thermal, structural and rheological properties of mung bean (*Vigna radiata*) protein isolate powder. *Food Res. Int.* **2020**, *138*, 109783. <https://doi.org/10.1016/j.foodres.2020.109783>.
31. Cruz Solorio, A.; Garín Aguilar, M.; Valencia del Toro, G. Nutritional and functional properties of protein concentrate and protein isolates of foods. In *Science within Food: Up-to-Date Advances on Research and Educational Ideas*, 1st ed.; Méndez-Vilas, A., Ed.; FORMATEX: Badajoz, Spain, 2020; pp. 176.
32. Schefer, S.; Oest, M.; Rohn, S. Interactions between Phenolic Acids, Proteins, and Carbohydrates—Influence on Dough and Bread Properties. *Foods* **2021**, *10*, 2798. <https://doi.org/10.3390/foods1011279>
33. Gallardo; Thompson; Burstin Reserve accumulation in legume seeds. *Comptes Rendus Biol.* **2008**, *331*, 755–762. <https://doi.org/10.1016/j.crv.2008.07.017>.
34. Rebello; Greenway; Finley A review of the nutritional value of legumes and their effects on obesity and its related co-morbidities. *Obes. Rev.* **2014**, *15*, 392–407. <https://doi.org/10.1111/obr.12144>.
35. Mani-López, E.; Palou, E.; López-Malo, A. Legume proteins, peptides, water extracts, and crude protein extracts as antifungals for food applications. *Trends Food Sci. Technol.* **2021**, *112*, 16–24. <https://doi.org/10.1016/j.tifs.2021.03.035>.
36. Allen, L.H. Legumes. In *Encyclopedia of Human Nutrition*, 3rd ed.; Caballero, B., Ed.; Elsevier Ltd.: London, UK, 2013; pp. 74–79.
37. Fujiwara, T.; Nambara, E.; Yamagishi, K.; Goto, D.B.; Naito, S. Storage Proteins. *Arab. Book* **2002**, *1*, e0020. <https://doi.org/10.1199/tab.0020>.
38. Belitz, H.; Grosch, W.; Schieberle, P. *Food Chemistry*, 3rd ed.; Translation from the Fifth German Edition; Springer: Berlin/Heidelberg, Germany, 2004; p. 1070.
39. Alting, A.C.; van de Velde, F. 9—Proteins as clean label ingredients in foods and beverages. In *Natural Food Additives, Ingredients and Flavourings*, 1st ed.; Baines, D., Seal, R., Eds.; Elsevier Ltd.: London, UK, 2012; pp. 197–211.
40. Goyal, A.K. (Ed.). *Grain Legumes*; BoD Books on Demand: Zagreb, Croatia, 2016; p. 188.
41. Moreno-Valdespino, C.A.; Luna-Vital, D.; Camacho-Ruiz, R.M.; Mojica, L. Bioactive proteins and phytochemicals from legumes: Mechanisms of action preventing obesity and type-2 diabetes. *Food Res. Int.* **2020**, *130*, 108905. <https://doi.org/10.1016/j.foodres.2019.108905>.
42. Boye, J.; Zare, F.; Pletch, A. Pulse proteins: Processing, characterization, functional properties and applications in food and feed. *Food Res. Int.* **2010**, *43*, 414–431. <https://doi.org/10.1016/j.foodres.2009.09.003>.
43. Vitale, A.; Bollini, R. Legume storage Proteins. In *Seed Development and Germination*, 2nd ed.; Kigel, J., Ed.; Routledge: New York, NY, USA, 2017; pp. 73–95.
44. Stone, A.K.; Nosworthy, M.G.; Chiremba, C.; House, J.D.; Nickerson, M.T. A comparative study of the functionality and protein quality of a variety of legume and cereal flours. *Cereal Chem.* **2019**, *96*, 1159–1169. <https://doi.org/10.1002/cche.10226>.

45. Sashikala, V.B.; Sreerama, Y.N.; Pratape, V.M.; Narasimha, H.V. Effect of thermal processing on protein solubility of green gram (*Phaseolus aureus*) legume cultivars. *J Food Sci Technol* **2013**, *52*, 1552–1560. <https://doi.org/10.1007/s13197-013-1149-x>.
46. Alonso, R.; Grant, G.; Dewey, P.; Marzo, F. Nutritional Assessment in Vitro and in Vivo of Raw and Extruded Peas (*Pisum sativum* L.). *J. Agric. Food Chem.* **2000**, *48*, 2286–2290. <https://doi.org/10.1021/jf000095o>.
47. Wrigley, C.W. *Encyclopedia of Food Grains*, 2nd ed.; Elsevier Science: San Diego, CA, USA, 2015; p. 1976.
48. Shevkani, K.; Singh, N.; Chen, Y.; Kaur, A.; Yu, L. Pulse proteins: Secondary structure, functionality and applications. *J Food Sci Technol* **2019**, *56*, 2787–2798. <https://doi.org/10.1007/s13197-019-03723-8>.
49. Diedericks, C.F.; Venema, P.; Mubaiwa, J.; Jideani, V.A.; van der Linden, E. Effect of processing on the microstructure and composition of Bambara groundnut (*Vigna subterranea* (L.) Verdc.) seeds, flour and protein isolates. *Food Hydrocoll.* **2020**, *108*, 106031. <https://doi.org/10.1016/j.foodhyd.2020.106031>.
50. Kigel, J. *Seed Development and Germination*, 1st ed.; Routledge: New York, NY, USA, 2017; p. 872.
51. Keskin, S.O.; Ali, T.M.; Ahmed, J.; Shaikh, M.; Siddiq, M.; Uebersax, M.A. Physico-chemical and functional properties of legume protein, starch, and dietary fiber—A review. *Legume Sci.* **2022**, *4*, 117. <https://doi.org/10.1002/leg3.117>.
52. Lopez, M.J.; Mohiuddin, S.S. *Biochemistry, Essential Amino Acids*. StatPearls Publishing, Ed.; StatPearls Publishing: Treasure Island, FL, USA, 2022. <https://www.statpearls.com/ArticleLibrary/viewarticle/36202>
53. Baptista; Pinho; Pinto; Casal; Mota; Ferreira Characterization of protein and fat composition of seeds from common beans (*Phaseolus vulgaris* L.), cowpea (*Vigna unguiculata* L. Walp) and bambara groundnuts (*Vigna subterranea* L. Verdc) from Mozambique. *Food Meas.* **2016**, *11*, 442–450. <https://doi.org/10.1007/s11694-016-9412-2>.
54. Margier, M.; Georgé, S.; Hafnaoui, N.; Remond, D.; Nowicki, M.; Du Chaffaut, L.; Amiot, M.; Reboul, E. Nutritional Composition and Bioactive Content of Legumes: Characterization of Pulses Frequently Consumed in France and Effect of the Cooking Method. *Nutrients* **2018**, *10*, 1668. <https://doi.org/10.3390/nu10111668>.
55. *FAO Dietary Protein Quality Evaluation in Human Nutrition: Report of an FAO Expert Consultation*; Food and Agriculture Organization of the United Nations: Rome, Italy, 2013; p. I.
56. Iqbal, A.; Khalil, I.A.; Ateeq, N.; Sayyar Khan, M. Nutritional quality of important food legumes. *Food Chem.* **2006**, *97*, 331–335. <https://doi.org/10.1016/j.foodchem.2005.05.011>.
57. Yao, D.N.; Kouassi, K.N.; Erba, D.; Scazzina, F.; Pellegrini, N.; Casiraghi, M.C. Nutritive Evaluation of the Bambara Groundnut Ci12 Landrace [*Vigna subterranea* (L.) Verdc. (Fabaceae)] Produced in Côte d’Ivoire. *Int. J. Mol. Sci.* **2015**, *16*, 21428–21441. <https://doi.org/10.3390/ijms160921428>.
58. Khattab, R.Y.; Arntfield, S.D.; Nyachoti, C.M. Nutritional quality of legume seeds as affected by some physical treatments, Part 1: Protein quality evaluation. *Food Sci. Technol.* **2009**, *42*, 1107–1112. <https://doi.org/10.1016/j.lwt.2009.02.008>.
59. Jahreis, G.; Brese, M.; Leiterer, M.; Schäfer, U.; Böhm, V.; Jena Legume flours: Nutritionally important sources of protein and dietary fiber. *Sci. Res.* **2015**, *63*, 36–42.
60. Liu, B.; Peng, J.; Zhang, S.; Zou, B.; Zhong, G. Chemical composition, in vitro starch digestibility and amino acid analysis of a underexplored kidney bean (*Phaseolus vulgaris* L.) grown in Chongqing, China. *Int. J. Food Sci. Technol.* **2013**, *48*, 527–532. <https://doi.org/10.1111/j.1365-2621.2012.03214.x>.
61. Fouad, A.A.; Rehab, F.M.A. Effect of germination time on proximate analysis, bioactive compounds and antioxidant activity of lentil (*Lens culinaris* Medik.) Sprouts. *Acta Sci. Polonorum. Technol. Aliment.* **2015**, *14*, 233–246. <https://doi.org/10.17306/J.AFS.2015.3.25>.
62. Aremu, M.O.; Stephen Audu, S.; Lyambee Gav, B. Comparative Review of Crude Protein and Amino Acids of Leguminous Seeds Grown in Nigeria. *Int. J. Sci.* **2017**, *3*, 88–97. <https://doi.org/10.18483/ijSci.1390>.
63. Arowora, K.A.; Popoola, C.A.; Yakubu, O.E.; Ugwuoke, K.C.; Aliyu, F. Proximate Composition and Amino Acids Contents in Selected Legumes and Hura crepitans seed in Wukari Taraba State. *Covenant J. Phys. Life Sci.* **2020**, *8*, 2354–3574. <https://doi.org/10.47231/SFXI7117>.
64. Palander, S.; Laurinen, P.; Perttilä, S.; Valaja, J.; Partanen, K. Protein and amino acid digestibility and metabolizable energy value of pea (*Pisum sativum*), faba bean (*Vicia faba*) and lupin (*Lupinus angustifolius*) seeds for turkeys of different age. *Anim. Feed Sci. Technol.* **2006**, *127*, 89–100. <https://doi.org/10.1016/j.anifeedsci.2005.07.003>.
65. Lásztity, R.; Khalil, M.M.; Haraszi, R.; Baticz, O.; Tömösközi, S. Isolation, functional properties and potential use of protein preparations from lupin. *Food Nahr.* **2001**, *45*, 396–398. [https://doi.org/10.1002/1521-3803\(20011001\)45:6<396::AID-FOOD396>3.0.CO;2-C](https://doi.org/10.1002/1521-3803(20011001)45:6<396::AID-FOOD396>3.0.CO;2-C).
66. Martineau-Côté, D.; Achouri, A.; Karboune, S.; L’Hocine, L. Faba Bean: An Untapped Source of Quality Plant Proteins and Bioactives. *Nutrients* **2022**, *14*, 1541. <https://doi.org/10.3390/nu14081541>.
67. Ghumman, A.; Kaur, A.; Singh, N. Functionality and digestibility of albumins and globulins from lentil and horse gram and their effect on starch rheology. *Food Hydrocoll.* **2016**, *61*, 843–850. <https://doi.org/10.1016/j.foodhyd.2016.07.013>.
68. Santos-Hernández, M.; Alfieri, F.; Gallo, V.; Miralles, B.; Masi, P.; Romano, A.; Ferranti, P.; Recio, I. Compared digestibility of plant protein isolates by using the INFOGEST digestion protocol. *Food Res. Int.* **2020**, *137*, 109708. <https://doi.org/10.1016/j.foodres.2020.109708>.

69. Araújo, A.H.; Cardoso, P.C.B.; Pereira, R.A.; Lima, L.M.; Oliveira, A.S.; Miranda, M.R.A.; Xavier-Filho, J.; Sales, M.P. In vitro digestibility of globulins from cowpea (*Vigna unguiculata*) and xerophitic algaroba (*Prosopis juliflora*) seeds by mammalian digestive proteinases: A comparative study. *Food Chem.* **2002**, *78*, 143–147. [https://doi.org/10.1016/S0308-8146\(01\)00391-0](https://doi.org/10.1016/S0308-8146(01)00391-0).
70. Teka, T.A.; Retta, N.; Bultosa, G.; Admassu, H.; Astatkie, T. Protein fractions, in vitro protein digestibility and amino acid composition of select cowpea varieties grown in Ethiopia. *Food Biosci.* **2020**, *36*, 100634. <https://doi.org/10.1016/j.fbio.2020.100634>.
71. Singhal, A.; Karaca, A.C.; Tyler, R.; Nickerson, M. *Pulse Proteins: From Processing to Structure-Function Relationships*; IntechOpen: Rijeka, Croatia, 2016; p. 188.
72. Prandi, B.; Zurlini, C.; Maria, C.I.; Cutroneo, S.; Di Massimo, M.; Bondi, M.; Brutti, A.; Sforza, S.; Tedeschi, T. Targeting the Nutritional Value of Proteins From Legumes By-Products Through Mild Extraction Technologies. *Front. Nutr.* **2021**, *8*, 695793. <https://doi.org/10.3389/fnut.2021.695793>.
73. Berghout, J.A.M.; Pelgrom, P.J.M.; Schutyser, M.A.I.; Boom, R.M.; van der Goot, A.J. Sustainability assessment of oilseed fractionation processes: A case study on lupin seeds. *J. Food Eng.* **2015**, *150*, 117–124. <https://doi.org/10.1016/j.jfoodeng.2014.11.005>.
74. Pelgrom, P.J.M.; Berghout, J.A.M.; van der Goot, A.J.; Boom, R.M.; Schutyser, M.A.I. Preparation of functional lupine protein fractions by dry separation. *Food Sci. Technol.* **2014**, *59*, 680–688. <https://doi.org/10.1016/j.lwt.2014.06.007>.
75. Schutyser, M.A.I.; Pelgrom, P.J.M.; van der Goot, A.J.; Boom, R.M. Dry fractionation for sustainable production of functional legume protein concentrates. *Trends Food Sci. Technol.* **2015**, *45*, 327–335. <https://doi.org/10.1016/j.tifs.2015.04.013>.
76. Czuchajowska, Z. Opportunities for pulses in food and non-food industrial processes by fractionation. In *Linking Research and Marketing Opportunities for Pulses in the 21st Century: Proceedings of the Third International Food Legumes Research Conference*, 3rd ed.; Springer Science & Business Media, Ed.; Springer Science & Business Media: Berlin/Heidelberg, Germany, 2012; p. 707.
77. Do, D.T.; Singh, J. Legume Microstructure. In *Encyclopedia of Food Chemistry*; Melton, L., Shahidi, F., Varelis, P., Eds.; Elsevier Inc: Amsterdam, The Netherlands, 2019; pp. 15–21.
78. Schutyser, M.A.I.; van der Goot, A.J. The potential of dry fractionation processes for sustainable plant protein production. *Trends Food Sci. Technol.* **2011**, *22*, 154–164. <https://doi.org/10.1016/j.tifs.2010.11.006>.
79. Tabtabaei, S.; Konakbayeva, D.; Rajabzadeh, A.R.; Legge, R.L. Functional properties of navy bean (*Phaseolus vulgaris*) protein concentrates obtained by pneumatic tribo-electrostatic separation. *Food Chem.* **2019**, *283*, 101–110. <https://doi.org/10.1016/j.foodchem.2019.01.031>.
80. Bayram, M.; Alameen, A. Protein Extraction Techniques from Cereals and Legumes. In Proceedings of the International Gap Agriculture and Livestock Congress, Sanliurfa, Turkey, 25–27 April 2018; pp. 948–953.
81. Pelgrom; Vissers; Boom; Schutyser Dry fractionation for production of functional pea protein concentrates. *Food Res. Int.* **2013**, *53*, 232–239. <https://doi.org/10.1016/j.foodres.2013.05.004>.
82. Assatory, A.; Vitelli, M.; Rajabzadeh, A.R.; Legge, R.L. Dry fractionation methods for plant protein, starch and fiber enrichment: A review. *Trends Food Sci. Technol.* **2019**, *86*, 340–351. <https://doi.org/10.1016/j.tifs.2019.02.006>.
83. Jiang, L.; Hua, D.; Wang, Z.; Xu, S. Aqueous enzymatic extraction of peanut oil and protein hydrolysates. *Food Bioprod. Process.* **2010**, *88*, 233–238. <https://doi.org/10.1016/j.fbp.2009.08.002>.
84. Preece, K.E.; Hooshyar, N.; Krijgsman, A.J.; Fryer, P.J.; Zuidam, N.J. Intensification of protein extraction from soybean processing materials using hydrodynamic cavitation. *Innov. Food Sci. Emerg. Technol.* **2017**, *41*, 47–55. <https://doi.org/10.1016/j.ifset.2017.01.002>.
85. Preece, K.E.; Hooshyar, N.; Krijgsman, A.; Fryer, P.J.; Zuidam, N.J. Intensified soy protein extraction by ultrasound. *Chem. Eng. Processing* **2017**, *113*, 94–101. <https://doi.org/10.1016/j.cep.2016.09.003>.
86. Lu, W.; Chen, X.; Wang, J.; Yang, X.; Qi, J. Enzyme-assisted subcritical water extraction and characterization of soy protein from heat-denatured meal. *J. Food Eng.* **2016**, *169*, 250–258. <https://doi.org/10.1016/j.jfoodeng.2015.09.006>.
87. Tanger, C.; Engel, J.; Kulozik, U. Influence of extraction conditions on the conformational alteration of pea protein extracted from pea flour. *Food Hydrocoll.* **2020**, *107*, 105949. <https://doi.org/10.1016/j.foodhyd.2020.105949>.
88. Klupšaitė, D.; Juodeikienė, G. Legume: Composition, protein extraction and functional properties. A review. *Chem. Technol.* **2015**, *66*, 5–12. <https://doi.org/10.5755/j01.ct.66.1.12355>.
89. Pelgrom, P.J.M.; Boom, R.M.; Schutyser, M.A.I. Method Development to Increase Protein Enrichment During Dry Fractionation of Starch-Rich Legumes. *Food Bioprocess Technol.* **2015**, *8*, 1495–1502. <https://doi.org/10.1007/s11947-015-1513-0>.
90. Dijkstra, D.S.; Linnemann, A.R.; van Boekel, T.A.J.S. Towards Sustainable Production of Protein-Rich Foods: Appraisal of Eight Crops for Western Europe. PART II: Analysis of the Technological Aspects of the Production Chain. *Crit. Rev. Food Sci. Nutr.* **2003**, *43*, 481–506. <https://doi.org/10.1080/10408690390246332>.
91. Qayyum, M.M.N.; Butt, F.M.; Anjum, H.; Nawaz composition analysis of some selected legumes for protein isolates recovery. *J. Anim. Plant Sci.* **2012**, *22*, 1156–1162.
92. Pelgrom, P.J.M.; Wang, J.; Boom, R.M.; Schutyser, M.A.I. Pre- and post-treatment enhance the protein enrichment from milling and air classification of legumes. *J. Food Eng.* **2015**, *155*, 53–61. <https://doi.org/10.1016/j.jfoodeng.2015.01.005>.
93. Pojić, M.; Mišan, A.; Tiwari, B. Eco-innovative technologies for extraction of proteins for human consumption from renewable protein sources of plant origin. *Trends Food Sci. Technol.* **2018**, *75*, 93–104.
94. Khazaei; Subedi; Nickerson; Martínez-Villaluenga; Frias; Vandenberg Seed Protein of Lentils: Current Status, Progress, and Food Applications. *Foods* **2019**, *8*, 391. <https://doi.org/10.3390/foods8090391>.

95. Ferreira, K.C.; Bento, J.A.C.; Caliari, M.; Bassinello, P.Z.; Berrios, J.D.J. Dry bean proteins: Extraction methods, functionality, and application in products for human consumption. *Cereal Chem.* **2022**, *99*, 67–77. <https://doi.org/10.1002/cche.10514>.
96. Sandberg, A. Developing Functional Ingredients: A Case Study of Pea Protein. In *Functional Foods—Concept to Product*, 2nd ed.; Woodhead Publishing: Sawston, UK, 2011; p. 1.
97. Boye, J.I.; Aksay, S.; Roufik, S.; Ribéreau, S.; Mondor, M.; Farnworth, E.; Rajamohamed, S.H. Comparison of the functional properties of pea, chickpea and lentil protein concentrates processed using ultrafiltration and isoelectric precipitation techniques. *Food Res. Int.* **2010**, *43*, 537–546. <https://doi.org/10.1016/j.foodres.2009.07.021>.
98. Wang, J.; Zhao, J.; de Wit, M.; Boom, R.M.; Schutyser, M.A.I. Lupine protein enrichment by milling and electrostatic separation. *Innov. Food Sci. Emerg. Technol.* **2016**, *33*, 596–602. <https://doi.org/10.1016/j.ifset.2015.12.020>.
99. Xing, Q.; de Wit, M.; Kyriakopoulou, K.; Boom, R.M.; Schutyser, M.A.I. Protein enrichment of defatted soybean flour by fine milling and electrostatic separation. *Innov. Food Sci. Emerg. Technol.* **2018**, *50*, 42–49. <https://doi.org/10.1016/j.ifset.2018.08.014>.
100. Allotey, D.K.; Kwofie, E.M.; Adewale, P.; Lam, E.; Ngadi, M. A meta-analysis of pulse-protein extraction technologies: Impact on recovery and purity. *J Food Eng* **2022**, *327*, 111048. <https://doi.org/10.1016/j.jfoodeng.2022.111048>.
101. Parvathy Shanthakumar; Klepacka, J.; Bains, A.; Chawla, P.; Sanju Bala Dhull; Najda, A. The Current Situation of Pea Protein and Its Application in the Food Industry. *Molecules* **2022**, *27*, 5354. <https://doi.org/10.3390/molecules27165354>.
102. Xing, Q.; Utami, D.P.; Demattey, M.B.; Kyriakopoulou, K.; de Wit, M.; Boom, R.M.; Schutyser, M.A.I. A two-step air classification and electrostatic separation process for protein enrichment of starch-containing legumes. *Innov. Food Sci. Emerg. Technol.* **2020**, *66*, 102480. <https://doi.org/10.1016/j.ifset.2020.102480>.
103. Pelgrom, P.J.M.; Boom, R.M.; Schutyser, M.A.I. Functional analysis of mildly refined fractions from yellow pea. *Food Hydrocoll.* **2015**, *44*, 12–22. <https://doi.org/10.1016/j.foodhyd.2014.09.001>.
104. Kamal, H.; Le, C.F.; Salter, A.M.; Ali, A. Extraction of protein from food waste: An overview of current status and opportunities. *Compr. Rev. Food Sci. Food Saf.* **2021**, *20*, 2455–2475. <https://doi.org/10.1111/1541-4337.12739>.
105. Wang, F.; Zhang, Y.; Xu, L.; Ma, H. An efficient ultrasound-assisted extraction method of pea protein and its effect on protein functional properties and biological activities. *LWT Food Sci. Technol.* **2020**, *127*, 109348.
106. Fukase, H.; Ohdaira, E.; Masuzawa, N.; Ide, M. Effect of Ultrasound in Soybean Protein Extraction. *Jpn. J. Appl. Phys.* **1994**, *33*, 3088–3090. <https://doi.org/10.1143/JJAP.33.3088>.
107. Aguilar-Acosta, L.A.; Serna-Saldivar, S.O.; Rodríguez-Rodríguez, J.; Escalante-Aburto, A.; Chuck-Hernández, C. Effect of Ultrasound Application on Protein Yield and Fate of Alkaloids during Lupin Alkaline Extraction Process. *Biomolecules* **2020**, *10*, 292. <https://doi.org/10.3390/biom10020292>.
108. Peyrano, F.; Speroni, F.; Avanza, M.V. Physicochemical and functional properties of cowpea protein isolates treated with temperature or high hydrostatic pressure. *Innov. Food Sci. Emerg. Technol.* **2016**, *33*, 38–46. <https://doi.org/10.1016/j.ifset.2015.10.014>.
109. Li, Q.; Zheng, J.; Ge, G.; Zhao, M.; Sun, W. Impact of heating treatments on physical stability and lipid-protein co-oxidation in oil-in-water emulsion prepared with soy protein isolates. *Food Hydrocoll.* **2020**, *100*, 105167. <https://doi.org/10.1016/j.foodhyd.2019.06.012>.
110. Tang, C.; Sun, X.; Yin, S. Physicochemical, functional and structural properties of vicilin-rich protein isolates from three Phaseolus legumes: Effect of heat treatment. *Food Hydrocoll.* **2009**, *23*, 1771–1778. <https://doi.org/10.1016/j.foodhyd.2009.03.008>.
111. Wang, Z.; Li, Y.; Jiang, L.; Qi, B.; Zhou, L. Relationship between Secondary Structure and Surface Hydrophobicity of Soybean Protein Isolate Subjected to Heat Treatment. *J. Chem.* **2014**, *2014*, 475389. <https://doi.org/10.1155/2014/475389>.
112. Brishti, F.H.; Chay, S.Y.; Muhammad, K.; Rashedi Ismail-Fitry, M.; Zarei, M.; Karthikeyan, S.; Caballero-Briones, F.; Saari, N. Structural and rheological changes of texturized mung bean protein induced by feed moisture during extrusion. *Food Chem.* **2021**, *344*, 128643. <https://doi.org/10.1016/j.foodchem.2020.128643>.
113. Mozafarpour, R.; Koocheki, A.; Milani, E.; Varidi, M. Extruded soy protein as a novel emulsifier: Structure, interfacial activity and emulsifying property. *Food Hydrocoll.* **2019**, *93*, 361–373. <https://doi.org/10.1016/j.foodhyd.2019.02.036>.
114. Zhang, B.; Kang, X.; Cheng, Y.; Cui, B.; Abd El-Aty, A.M. Impact of high moisture contents on the structure and functional properties of pea protein isolate during extrusion. *Food Hydrocoll.* **2022**, *127*, 107508. <https://doi.org/10.1016/j.foodhyd.2022.107508>.
115. Beck, S.M.; Knoerzer, K.; Arcot, J. Effect of low moisture extrusion on a pea protein isolate's expansion, solubility, molecular weight distribution and secondary structure as determined by Fourier Transform Infrared Spectroscopy (FTIR). *J. Food Eng.* **2017**, *214*, 166–174. <https://doi.org/10.1016/j.jfoodeng.2017.06.037>.
116. Yin, S.; Tang, C.; Wen, Q.; Yang, X.; Li, L. Functional properties and in vitro trypsin digestibility of red kidney bean (*Phaseolus vulgaris* L.) protein isolate: Effect of high-pressure treatment. *Food Chem.* **2008**, *110*, 938–945. <https://doi.org/10.1016/j.foodchem.2008.02.090>.
117. Chao, D.; Jung, S.; Aluko, R.E. Physicochemical and functional properties of high pressure-treated isolated pea protein. *Innov. Food Sci. Emerg. Technol.* **2018**, *45*, 179–185. <https://doi.org/10.1016/j.ifset.2017.10.014>.
118. Ahmed, J.; Al-Ruwaih, N.; Mulla, M.; Rahman, M.H. Effect of high pressure treatment on functional, rheological and structural properties of kidney bean protein isolate. *Food Sci. Technol.* **2018**, *91*, 191–197. <https://doi.org/10.1016/j.lwt.2018.01.054>.
119. Mehr, H.M.; Koocheki, A. Effect of atmospheric cold plasma on structure, interfacial and emulsifying properties of Grass pea (*Lathyrus sativus* L.) protein isolate. *Food Hydrocoll.* **2020**, *106*, 105899. <https://doi.org/10.1016/j.foodhyd.2020.105899>.

120. Zhang, Q.; Cheng, Z.; Zhang, J.; Nasiru, M.M.; Wang, Y.; Fu, L. Atmospheric cold plasma treatment of soybean protein isolate: Insights into the structural, physicochemical, and allergenic characteristics. *J. Food Sci.* **2021**, *86*, 68–77. <https://doi.org/10.1111/1750-3841.15556>.
121. Ji; Dong; Han; Li; Chen; Li; Chen Effects of Dielectric Barrier Discharge (DBD) Cold Plasma Treatment on Physicochemical and Functional Properties of Peanut Protein. *Food Bioprocess Technol.* **2017**, *11*, 344–354. <https://doi.org/10.1007/s11947-017-2015-z>.
122. Mallikarjunan, N.; Marathe, S.; Rajalakshmi, D.; Mahesh, S.; Jamdar, S.N.; Sharma, A. Effect of ionizing radiation on structural and functional attributes of red kidney bean (*Phaseolus vulgaris* L.) lectin. *Food Sci. Technol.* **2014**, *59*, 300–307. <https://doi.org/10.1016/j.lwt.2014.04.052>.
123. Vaz, A.F.M.; Souza, M.P.; Medeiros, P.L.; Melo, A.M.M.A.; Silva-Lucca, R.A.; Santana, L.A.; Oliva, M.L.V.; Perez, K.R.; Cuccovia, I.M.; Correia, M.T.S. Low-dose gamma irradiation of food protein increases its allergenicity in a chronic oral challenge. *Food Chem. Toxicol.* **2013**, *51*, 46–52. <https://doi.org/10.1016/j.fct.2012.09.011>.
124. Ghribi, A.M.; Gafsi, I.M.; Blecker, C.; Danthine, S.; Attia, H.; Besbes, S. Effect of drying methods on physico-chemical and functional properties of chickpea protein concentrates. *J. Food Eng.* **2015**, *165*, 179–188. <https://doi.org/10.1016/j.jfoodeng.2015.06.021>.
125. Bußler, S.; Steins, V.; Ehlbeck, J.; Schlüter, O. Impact of thermal treatment versus cold atmospheric plasma processing on the techno-functional protein properties from *Pisum sativum* ‘Salamanca’. *J. Food Eng.* **2015**, *167*, 166–174. <https://doi.org/10.1016/j.jfoodeng.2015.05.036>.
126. Maurya, A.; Mhatre, S.; Kalha, R.; Dabade, A. Effect of gamma radiation treatment on physico-chemical and functional properties of soya proteins isolate. *Ann. Food Sci. Technol.* **2020**, *21*, 274–285 Available online: https://afst.valahia.ro/wp-content/uploads/2022/09/I.4_Maurya.pdf (accessed on 25 September 2022).
127. Ngui, S.P.; Nyobe, C.E.; Bassogog, C.B.B.; Tang, E.N.; Minka, S.R.; Mune, M.A.M. Influence of pH and temperature on the physicochemical and functional properties of Bambara bean protein isolate. *Heliyon* **2021**, *7*, e07824. <https://doi.org/10.1016/j.heliyon.2021.e07824>.
128. Garcia-Mora, P.; Peñas, E.; Frias, J.; Gomez, R.; Martinez-Villaluenga, C. High-pressure improves enzymatic proteolysis and the release of peptides with angiotensin I converting enzyme inhibitory and antioxidant activities from lentil proteins. *Food Chem.* **2015**, *171*, 224–232. <https://doi.org/10.1016/j.foodchem.2014.08.116>.
129. Saricaoglu, F.T. Application of high-pressure homogenization (HPH) to modify functional, structural and rheological properties of lentil (*Lens culinaris*) proteins. *Int. J. Biol. Macromol.* **2020**, *144*, 760–769. <https://doi.org/10.1016/j.ijbiomac.2019.11.034>.
130. Guo, F.; Xiong, Y.L.; Qin, F.; Jian, H.; Huang, X.; Chen, J. Surface Properties of Heat-Induced Soluble Soy Protein Aggregates of Different Molecular Masses. *J. Food Sci.* **2015**, *80*, C279–C287. <https://doi.org/10.1111/1750-3841.12761>.
131. Puppo, C.; Chapleau, N.; Speroni, F.; de Lamballerie-Anton, M.; Michel, F.; Añón, C.; Anton, M. Physicochemical Modifications of High-Pressure-Treated Soybean Protein Isolates. *J. Agric. Food Chem.* **2004**, *52*, 1564–1571. <https://doi.org/10.1021/jf034813t>.
132. Chao, D.; Aluko, R.E. Modification of the structural, emulsifying, and foaming properties of an isolated pea protein by thermal pretreatment. *CYTA: J. Food* **2018**, *16*, 357–366. <https://doi.org/10.1080/19476337.2017.1406536>.
133. Wang, L.; Zhang, X.; Liu, F.; Ding, Y.; Wang, R.; Luo, X.; Li, Y.; Chen, Z. Study of the functional properties and anti-oxidant activity of pea protein irradiated by electron beam. *Innov. Food Sci. Emerg. Technol.* **2017**, *41*, 124–129. <https://doi.org/10.1016/j.ifset.2017.01.005>.
134. Peyrano, F.; de Lamballerie, M.; Avanza, M.V.; Speroni, F. Rheological characterization of the thermal gelation of cowpea protein isolates: Effect of pretreatments with high hydrostatic pressure or calcium addition. *Food Sci. Technol.* **2019**, *115*, 108472. <https://doi.org/10.1016/j.lwt.2019.108472>.
135. Sim, S.Y.J.; Karwe, M.V.; Moraru, C.I. High pressure structuring of pea protein concentrates. *J. Food Process Eng.* **2019**, *42*, e13261. <https://doi.org/10.1111/jfpe.13261>.
136. Zhang, S.; Huang, W.; Feizollahi, E.; Roopesh, M.S.; Chen, L. Improvement of pea protein gelation at reduced temperature by atmospheric cold plasma and the gelling mechanism study. *Innov. Food Sci. Emerg. Technol.* **2021**, *67*, 102567. <https://doi.org/10.1016/j.ifset.2020.102567>.
137. Salazar-Villanea, S.; Hendriks, W.H.; Bruininx, E.M.A.M.; Gruppen, H.; van der Poel, A.F.B. Protein structural changes during processing of vegetable feed ingredients used in swine diets: Implications for nutritional value. *Nutr. Res. Rev.* **2016**, *29*, 126–141. <https://doi.org/10.1017/S0954422416000056>.
138. Joshi, M.; Adhikari, B.; Aldred, P.; Panozzo, J.; Kasapis, S. Physicochemical and functional properties of lentil protein isolates prepared by different drying methods. *Food Chem.* **2011**, *129*, 1513–1522. <https://doi.org/10.1016/j.foodchem.2011.05.131>.
139. Al-Ali, H.A.; Shah, U.; Hackett, M.J.; Gulzar, M.; Karakyriakos, E.; Johnson, S.K. Technological strategies to improve gelation properties of legume proteins with the focus on lupin. *Innov. Food Sci. Emerg. Technol.* **2021**, *68*, 102634. <https://doi.org/10.1016/j.ifset.2021.102634>.
140. Messian, J.; Chihi, M.L.; Sok, N.; Saurel, R. Effect of globular pea proteins fractionation on their heat-induced aggregation and acid cold-set gelation. *Food Hydrocoll.* **2015**, *46*, 233–243. <https://doi.org/10.1016/j.foodhyd.2014.11.025>.
141. Dalbhagat, C.G.; Mahato, D.K.; Mishra, H.N. Effect of extrusion processing on physicochemical, functional and nutritional characteristics of rice and rice-based products: A review. *Trends Food Sci. Technol.* **2019**, *85*, 226–240. <https://doi.org/10.1016/j.tifs.2019.01.001>.

142. Chen, F.L.; Wei, Y.M.; Zhang, B. Chemical cross-linking and molecular aggregation of soybean protein during extrusion cooking at low and high moisture content. *Food Sci. Technol.* **2011**, *44*, 957–962. <https://doi.org/10.1016/j.lwt.2010.12.008>.
143. Singh, S.; Gamlath, S.; Wakeling, L. Nutritional aspects of food extrusion: A review. *Int. J. Food Sci. Technol.* **2007**, *42*, 916–929. <https://doi.org/10.1111/j.1365-2621.2006.01309.x>.
144. Ojokoh, A.O.; Yimin, W. Effect of Fermentation on Chemical Composition and Nutritional Quality of Extruded and Fermented Soya Products. *Int. J. Food Eng.* **2011**, *7*, 1857. <https://doi.org/10.2202/1556-3758.1857>.
145. Alam, M.S.; Kaur, J.; Khaira, H.; Gupta, K. Extrusion and Extruded Products: Changes in Quality Attributes as Affected by Extrusion Process Parameters: A Review. *Crit. Rev. Food Sci. Nutr.* **2016**, *56*, 445–473. <https://doi.org/10.1080/10408398.2013.779568>.
146. Samard, S.; Gu, B.; Ryu, G.-H. Effects of extrusion types, screw speed and addition of wheat gluten on physicochemical characteristics and cooking stability of meat analogues. *J. Sci. Food Agric.* **2019**, *99*, 4922–4931. <https://doi.org/10.1002/jsfa.9722>.
147. Philipp, C.; Oey, I.; Silcock, P.; Beck, S.M.; Buckow, R. Impact of protein content on physical and microstructural properties of extruded rice starch-pea protein snacks. *J. Food Eng.* **2017**, *212*, 165–173. <https://doi.org/10.1016/j.jfoodeng.2017.05.024>.
148. Phongthai, S.; D'Amico, S.; Schoenlechner, R.; Homthawornchoo, W.; Rawdkuen, S. Effects of protein enrichment on the properties of rice flour based gluten-free pasta. *Food Sci. Technol.* **2017**, *80*, 378–385. <https://doi.org/10.1016/j.lwt.2017.02.044>.
149. Yu, L.; Ramaswamy, H.S.; Boye, J. Protein rich extruded products prepared from soy protein isolate-corn flour blends. *Food Sci. Technol.* **2013**, *50*, 279–289. <https://doi.org/10.1016/j.lwt.2012.05.012>.
150. Bueno, A.S.; Pereira, C.M.; Menegassi, B.; Arêas, J.A.G.; Castro, I.A. Effect of extrusion on the emulsifying properties of soybean proteins and pectin mixtures modelled by response surface methodology. *J. Food Eng.* **2009**, *90*, 504–510. <https://doi.org/10.1016/j.jfoodeng.2008.07.028>.
151. Chen, Q.; Zhang, J.; Zhang, Y.; Meng, S.; Wang, Q. Rheological properties of pea protein isolate-amylose/amylopectin mixtures and the application in the high-moisture extruded meat substitutes. *Food Hydrocoll.* **2021**, *117*, 106732. <https://doi.org/10.1016/j.foodhyd.2021.106732>.
152. Carmo, C.S.D.; Knutsen, S.H.; Malizia, G.; Dessev, T.; Geny, A.; Zobel, H.; Myhrer, K.S.; Varela, P.; Sahlstrøm, S. Meat analogues from a faba bean concentrate can be generated by high moisture extrusion. *Future Foods A Dedic. J. Sustain. Food Sci.* **2021**, *3*, 100014. <https://doi.org/10.1016/j.fufo.2021.100014>.
153. Silva, A.C.C.; Arêas, E.P.G.; Silva, M.A.; Arêas, J.A.G. Effects of Extrusion on the Emulsifying Properties of Rumen and Soy Protein. *Food Biophys.* **2010**, *5*, 94–102. <https://doi.org/10.1007/s11483-010-9149-0>.
154. Osen, R.; Toelstede, S.; Eisner, P.; Schweiggert-Weisz, U. Effect of high moisture extrusion cooking on protein-protein interactions of pea (*Pisum sativum* L.) protein isolates. *Int. J. Food Sci. Technol.* **2015**, *50*, 1390–1396. <https://doi.org/10.1111/ijfs.12783>.
155. Samard, S.; Ryu, G. Physicochemical and functional characteristics of plant protein-based meat analogs. *J. Food Process. Preserv.* **2019**, *43*, e14123. <https://doi.org/10.1111/jfpp.14123>.
156. Ma, K.K.; Greis, M.; Lu, J.; Nolden, A.A.; McClements, D.J.; Kinchla, A.J. Functional Performance of Plant Proteins. *Foods* **2022**, *11*, 594.
157. Wagner, J.R.; Sorgentini, D.A.; Añón, M.C. Relation between Solubility and Surface Hydrophobicity as an Indicator of Modifications during Preparation Processes of Commercial and Laboratory-Prepared Soy Protein Isolates. *J. Agric. Food Chem.* **2000**, *48*, 3159–3165. <https://doi.org/10.1021/jf990823b>.
158. Gharibzahedi, S.M.T.; Smith, B. Effects of high hydrostatic pressure on the quality and functionality of protein isolates, concentrates, and hydrolysates derived from pulse legumes: A review. *Trends Food Sci. Technol.* **2021**, *107*, 466–479. <https://doi.org/10.1016/j.tifs.2020.11.016>.
159. Queirós, R.P.; Saraiva, J.A.; da Silva, J.A.L. Tailoring structure and technological properties of plant proteins using high hydrostatic pressure. *Crit. Rev. Food Sci. Nutr.* **2018**, *58*, 1538–1556. <https://doi.org/10.1080/10408398.2016.1271770>.
160. Patrignani, F.; Lanciotti, R. Applications of high and ultra high pressure homogenization for food safety. *Front. Microbiol.* **2016**, *7*, 1132. <https://doi.org/10.3389/fmicb.2016.01132>.
161. Ahmed, J.; Mulla, M.; Al-Ruwaih, N.; Arfat, Y.A. Effect of high-pressure treatment prior to enzymatic hydrolysis on rheological, thermal, and antioxidant properties of lentil protein isolate. *Legum. Sci.* **2019**, *1*, e10. <https://doi.org/10.1002/leg3.10>.
162. Molina, E.; Papadopoulou, A.; Ledward, D.A. Emulsifying properties of high pressure treated soy protein isolate and 7S and 11S globulins. *Food Hydrocoll.* **2001**, *15*, 263. [https://doi.org/10.1016/s0268-005x\(01\)00023-6](https://doi.org/10.1016/s0268-005x(01)00023-6).
163. Ishak, N.F.; Hashim, N.A.; Othman, M.H.D.; Monash, P.; Zuki, F.M. Recent progress in the hydrophilic modification of alumina membranes for protein separation and purification. *Ceram. Int.* **2017**, *43*, 915–925. <https://doi.org/10.1016/j.ceramint.2016.10.044>.
164. Puppo, M.C.; Speroni, F.; Chapleau, N.; de Lamballerie, M.; Añón, M.C.; Anton, M. Effect of high-pressure treatment on emulsifying properties of soybean proteins. *Food Hydrocoll.* **2005**, *19*, 289–296. <https://doi.org/10.1016/j.foodhyd.2004.07.001>.
165. Ucar, Y.; Ceylan, Z.; Durmus, M.; Tomar, O.; Cetinkaya, T. Application of cold plasma technology in the food industry and its combination with other emerging technologies. *Trends Food Sci. Technol.* **2021**, *114*, 355–371. <https://doi.org/10.1016/j.tifs.2021.06.004>.
166. Saremnezhad, S.; Soltani, M.; Faraji, A.; Hayaloglu, A.A. Chemical changes of food constituents during cold plasma processing: A review. *Food Res. Int.* **2021**, *147*, 110552. <https://doi.org/10.1016/j.foodres.2021.110552>.

167. Niemira, B.A. Cold Plasma Decontamination of Foods. *Annu. Rev. Food Sci. Technol.* **2012**, *3*, 125–142. <https://doi.org/10.1146/annurev-food-022811-101132>.
168. Niemira, B.A. Chapter 18—Decontamination of Foods by Cold Plasma. In *Emerging Technologies for Food Processing*, 2nd ed.; Sun, D., Ed.; Elsevier Ltd.: London, UK, 2015; pp. 327–333.
169. Ji, Y.; Wang, Z.; Pei, F.; Shi, J.; Li, J.; Gunosewoyo, H.; Yang, F.; Tang, J.; Xie, X.; Yu, L. Introducing nitrogen atoms to amidoalkylindoles: Potent and selective cannabinoid type 2 receptor agonists with improved aqueous solubility. *Med. Chem. Commun.* **2019**, *10*, 2131–2139. <https://doi.org/10.1039/c9md00411d>.
170. Farkas, J.; Mohácsi-Farkas, C. History and future of food irradiation. *Trends Food Sci. Technol.* **2011**, *22*, 121–126. <https://doi.org/10.1016/j.tifs.2010.04.002>.
171. Farkas, J. Irradiation for better foods. *Trends Food Sci. Technol.* **2006**, *17*, 148–152. <https://doi.org/10.1016/j.tifs.2005.12.003>.
172. Lee, M.; Lee, S.; Song, K.B. Effect of γ -irradiation on the physicochemical properties of soy protein isolate films. *Radiat Phys Chem* **2005**, *72*, 35–40. <https://doi.org/10.1016/j.radphyschem.2004.01.006>.
173. Singh Pulses: An overview. *J. Food Sci. Technol.* **2017**, *54*, 853–857. <https://doi.org/10.1007/s13197-017-2537-4>.
174. Joye, I. Protein Digestibility of Cereal Products. *Foods* **2019**, *8*, 199. <https://doi.org/10.3390/foods8060199>.
175. Meinschmidt, P.; Ueberham, E.; Lehmann, J.; Reineke, K.; Schlüter, O.; Schweiggert-Weisz, U.; Eisner, P. The effects of pulsed ultraviolet light, cold atmospheric pressure plasma, and gamma-irradiation on the immunoreactivity of soy protein isolate. *Innov. Food Sci. Emerg. Technol.* **2016**, *38*, 374–383. <https://doi.org/10.1016/j.ifset.2016.06.007>.
176. Basak, S.; Annapure, U.S. Recent trends in the application of cold plasma for the modification of plant proteins—A review. *Future Foods : A Dedic. J. Sustain. Food Sci.* **2022**, *5*, 100119. <https://doi.org/10.1016/j.fufo.2022.100119>.
177. Hall, A.E.; Moraru, C.I. Effect of High Pressure Processing and heat treatment on in vitro digestibility and trypsin inhibitor activity in lentil and faba bean protein concentrates. *Food Sci. Technol.* **2021**, *152*, 112342. <https://doi.org/10.1016/j.lwt.2021.112342>.
178. Siddhuraju; Makkar; Becker the effect of ionising radiation on antinutritional factors and the nutritional value of plant materials with reference to human and animal food. *Food Chem.* **2002**, *78*, 187. [https://doi.org/10.1016/s0308-8146\(01\)00398-3](https://doi.org/10.1016/s0308-8146(01)00398-3).
179. Xiong; Guo Animal and Plant Protein Oxidation: Chemical and Functional Property Significance. *Foods* **2020**, *10*, 40. <https://doi.org/10.3390/foods10010040>.
180. Palanisamy, M.; Franke, K.; Berger, R.G.; Heinz, V.; Töpfl, S. High moisture extrusion of lupin protein: Influence of extrusion parameters on extruder responses and product properties. *J. Sci. Food Agric.* **2018**, *99*, 2175–2185. <https://doi.org/10.1002/jsfa.9410>.
181. Omosebi, M.O.; Osundahunsi, O.F.; Fagbemi, T.N. Effect of extrusion on protein quality, antinutritional factors, and digestibility of complementary diet from quality protein maize and soybean protein concentrate. *J. Food Biochem.* **2018**, *42*, e12508. <https://doi.org/10.1111/jfbc.12508>.
182. Joehnke, M.S.; Jeske, S.; Ispiryani, L.; Zannini, E.; Arendt, E.K.; Bez, J.; Sørensen, J.C.; Petersen, I.L. Nutritional and anti-nutritional properties of lentil (*Lens culinaris*) protein isolates prepared by pilot-scale processing. *Food Chem. X* **2021**, *9*, 100112. <https://doi.org/10.1016/j.fochx.2020.100112>.
183. Barbana, C.; Boye, J.I. In vitro protein digestibility and physico-chemical properties of flours and protein concentrates from two varieties of lentil (*Lens culinaris*). *Food Funct.* **2013**, *4*, 310–321. <https://doi.org/10.1039/c2fo30204g>.
184. Aryee, A.N.A.; Boye, J.I. Comparative Study of the Effects of Processing on the Nutritional, Physicochemical and Functional Properties of Lentil. *J. Food Process. Preserv.* **2017**, *41*, e12824. <https://doi.org/10.1111/jfpp.12824>.
185. Carbonaro; Maselli; Nucara Relationship between digestibility and secondary structure of raw and thermally treated legume proteins: A Fourier transform infrared (FT-IR) spectroscopic study. *Amino Acids* **2011**, *43*, 911–921. <https://doi.org/10.1007/s00726-011-1151-4>.
186. Laguna, L.; Picouet, P.; Guàrdia, M.D.; Renard, C.M.G.C.; Sarkar, A. In vitro gastrointestinal digestion of pea protein isolate as a function of pH, food matrices, autoclaving, high-pressure and re-heat treatments. *Food Sci. Technol.* **2017**, *84*, 511–519. <https://doi.org/10.1016/j.lwt.2017.06.021>.
187. Tang, C.; Chen, L.; Ma, C. Thermal aggregation, amino acid composition and in vitro digestibility of vicilin-rich protein isolates from three Phaseolus legumes: A comparative study. *Food Chem.* **2009**, *113*, 957–963. <https://doi.org/10.1016/j.foodchem.2008.08.038>.
188. Gulati, P.; Brahma, S.; Rose, D.J. Chapter 13—Impacts of extrusion processing on nutritional components in cereals and legumes: Carbohydrates, proteins, lipids, vitamins, and minerals. In *Extrusion Cooking*, 2nd ed.; Ganjyal, G.M., Ed.; Elsevier Inc.: Edinburgh, UK, 2020; pp. 415–443.
189. MacDonald; Pryzbyszewski; Hsieh Soy Protein Isolate Extruded with High Moisture Retains High Nutritional Quality. *J. Agric. Food Chem.* **2009**, *57*, 3550–3555. <https://doi.org/10.1021/jf803435x>.
190. Lee, H.; Ha, M.J.; Shahbaz, H.M.; Kim, J.U.; Jang, H.; Park, J. High hydrostatic pressure treatment for manufacturing of red bean powder: A comparison with the thermal treatment. *J. Food Eng.* **2018**, *238*, 141–147. <https://doi.org/10.1016/j.jfoodeng.2018.06.016>.

191. Takai, E.; Kitamura, T.; Kuwabara, J.; Ikawa, S.; Yoshizawa, S.; Shiraki, K.; Kawasaki, H.; Arakawa, R.; Kitano, K. Chemical modification of amino acids by atmospheric-pressure cold plasma in aqueous solution. *J. Phys. D Appl. Phys.* **2014**, *47*, 285403. <https://doi.org/10.1088/0022-3727/47/28/285403>.
192. Esteghlal, S.; Gahruie, H.H.; Niakousari, M.; Barba, F.J.; Bekhit, A.E.; Mallikarjunan, K.; Roohinejad, S. Bridging the Knowledge Gap for the Impact of Non-Thermal Processing on Proteins and Amino Acids. *Foods* **2019**, *8*, 262. <https://doi.org/10.3390/foods8070262>.
193. Mariotti, F.; Gardner, C.D. Dietary Protein and Amino Acids in Vegetarian Diets—A Review. *Nutrients* **2019**, *11*, 2661. <https://doi.org/10.3390/nu1112661>.
194. Kawada, K.; Kuramoto, N.; Mimori, S. Possibility that the Onset of Autism Spectrum Disorder is Induced by Failure of the Glutamine-Glutamate Cycle. *Curr. Mol. Pharmacol.* **2021**, *14*, 170–174. <https://doi.org/10.2174/1874467213666200319125109>.
195. Strzyz, P. Forcing MT glutamylation. *Nat. Rev. Mol. Cell Biol.* **2021**, *22*, 509. <https://doi.org/10.1038/s41580-021-00391-5>