

Hemolysed blood elicits - calcium antagonist and high CO₂ reversible - constrictions via elevation of Ca²⁺ in isolated cerebral arteries

Peter Cseplo^{1,2}, Zoltan Vamos¹, Orsolya Torok¹, Ivan Ivic¹, Attila Toth³, Andras Buki⁴ and Akos Koller^{1,5,6}

¹ Department of Pathophysiology and Gerontology, University of Pecs, Medical School, and Szentagothai Research Centre, Pecs, Hungary

² Department of Central Anesthesiology and Intensive Therapy, Petz Aladar County Teaching Hospital, Győr, Hungary

³ Institute of Cardiology, Division of Clinical Physiology, Medical and Health Science Centre, University of Debrecen, Hungary

⁴ Department of Neurosurgery, University of Pecs, Medical School, Pecs, Hungary

⁵ Institute of Natural Sciences, University of Physical Education, Budapest, Hungary

⁶ Department of Physiology, New York Medical College, Valhalla, NY, USA

Correspondence:

Akos Koller, MD, PhD.

Department of Physiology, New York Medical College, Valhalla, NY 10595, USA

and Institute of Natural Sciences, University of Physical Education, Budapest, Hungary

E-mail: koller@nymc.edu or koller.akos@tf.hu

Running title: Perivascular blood constricts cerebral arteries

CONTACT INFORMATION

Peter Cseplo^{1,2} [+36702505028, cseplopeti@gmail.com; fax: +3696507902]

¹ Department of Pathophysiology and Gerontology, University of Pecs, Medical School, and Szentagothai Research Centre, Pecs, Hungary, [7624, Pecs, Szigeti street 12, HUNGARY]

² Department of Central Anesthesiology and Intensive Therapy, Petz Aladar County Teaching Hospital, Győr, Hungary [9024, Győr, Vasvári Pal street 2-4, HUNGARY]

Zoltan Vamos¹ [+36303573557, azozoka@gmail.com; fax: +3672536247]

¹ Department of Pathophysiology and Gerontology, University of Pecs, Medical School, and Szentagothai Research Centre, Pecs, Hungary, [7624, Pecs, Szigeti street 12, HUNGARY]

Orsolya Torok¹ [+36304298827, torokorsi55@gmail.com; fax: +3672536247]

¹ Department of Pathophysiology and Gerontology, University of Pecs, Medical School, and Szentagothai Research Centre, Pecs, Hungary, [7624, Pecs, Szigeti street 12, HUNGARY]

Ivan Ivic¹ [+385911814211, ivic.ivan@gmail.com; fax: +3672536247]

¹ Department of Pathophysiology and Gerontology, University of Pecs, Medical School, and Szentagothai Research Centre, Pecs, Hungary, [7624, Pecs, Szigeti street 12, HUNGARY]

Attila Toth³ [+3652411600, atitoth@dote.hu; fax: +3652255978 extension 56869]

³ Institute of Cardiology, Division of Clinical Physiology, Medical and Health Science Centre, University of Debrecen, Hungary [4032, Debrecen, Moricz Zs krt. 22, HUNGARY]

Andras Buki⁴ [+3672535932, buki.andras@pte.hu; fax: +3672535931]

⁴ Department of Neurosurgery, University of Pecs, Medical School, Pecs, Hungary [7623, Pecs, Ret street 2, HUNGARY]

Akos Koller^{1,5,6} [+36709020681, koller.akos@tf.hu; fax: +3613566337]

¹ Department of Pathophysiology and Gerontology, University of Pecs, Medical School, and Szentagothai Research Centre, Pecs, Hungary, [7624, Pecs, Szigeti street 12, HUNGARY]

⁵ Institute of Natural Sciences, University of Physical Education, Budapest, Hungary [1123, Budapest, Alkotas street 44, HUNGARY]

⁶ Department of Physiology, New York Medical College, Valhalla, NY, USA [40 Sunshine Cottage Rd, Valhalla, NY 10595, USA]

Keywords: SUBARACHNOID HEMORRHAGE, IN VITRO STUDIES, VASCULAR INJURY, TRAUMATIC BRAIN INJURY, CBF AUTOREGULATION

Abstract

During acute subarachnoid hemorrhage blood is hemolysed, which is followed by a significant cerebrovascular spasm resulting in serious clinical condition. Interestingly, however the direct vasomotor effect of perivascular hemolysed blood (HB) has not yet been characterized preventing the assessment of contribution of vasoconstrictor mechanisms deriving from brain tissue and/or blood and development of possible treatments. We hypothesized that perivascular HB reduces the diameter of the cerebral arteries (BA: basilar artery; MCA: middle cerebral artery) via elevating vascular tissue $[Ca^{2+}]_i$ level. Vasomotor responses were measured by videomicroscopy and intracellular Ca^{2+} by the fura2-AM ratiometric method. Adding HB to the vessel chamber reduced the diameter significantly (BA: from $264 \pm 7 \mu m$ to $164 \pm 11 \mu m$; $-24 \pm 3 \%$ of PD; MCA: from $187 \pm 14 \mu m$ to $155 \pm 14 \mu m$), which was reversed to control level by wash-out of HB. Potassium chloride (KCl), HB, serum, hemolysed red blood cell (RBC), plasma and platelet suspension (PLTs), elicited significant constrictions of isolated basilar arteries. There was a significant increase in K^+ concentration in hemolysed HB (7.02 ± 0.22 mmol/L) compared to whole blood (6.20 ± 0.01 mmol/L). Before HB, acetylcholine (ACh), sodium-nitroprussid (SNP), nifedipin, and CO_2 elicited substantial dilations in cerebral arteries. In contrast, in the presence of HB dilations to ACh, SNP, decreased, but not to nifedipine and CO_2 . After washout of HB, NO-mediated dilations remained significantly reduced compared to control. HB significantly increased the ratiometric Ca-signal, which returned to control level after washout.

In conclusion, perivascular hemolysed blood elicits significant - nifedipine and high CO_2 reversible - constrictions of isolated basilar and middle cerebral arteries, primarily via increasing intracellular Ca^{2+} , findings, which can contribute to the refinement of local treatment of subarachnoid hemorrhage.

Introduction

Clinical and experimental studies showed that acute subarachnoid hemorrhage due to traumatic brain injury or stroke¹ is followed by serious local vasospasm², which can severely reduce regional cerebral blood flow, with the consequent loss of brain function. Proper resistance (i.e. diameter) of cerebral arteries plays an important role in maintaining continuous blood supply of the brain to preserve its functions.^{1, 3, 4} Disturbances of cerebrovascular autoregulation³ may occur as a result of intracranial hemorrhage and brain injury, as well. The hemolysed blood (HB) then can affect local tissues (neurons, glia cells, vascular cells, etc.), but primarily impairs the regulation of cerebrovascular tone endangering maintenance of normal flow to brain and thus functions.^{5, 6, 7, 8} Increased vascular contractility to hemolysed blood may be attributed to endothelial dysfunction and/or increased contractility of vascular smooth muscle.⁹ However, in such conditions several cell types can be involved in the pathological regulation of vascular resistance in addition to HB. Earlier experiments showed that purified hemoglobin induces vasoconstriction¹⁰ of cerebral vessels, which was explained by its nitric oxide scavenging effect.¹¹ However, the vasoconstrictor effects of perivascular hemolysed blood (HB), which is present in vivo during hemorrhage and traumatic brain injury have not yet been fully characterized.¹² We hypothesized that perivascular HB reduces the diameter of the cerebral vessels via elevating $[Ca^{2+}]_i$ levels.

In order to test this hypothesis we have utilized isolated basilar and middle cerebral arteries of rat, known to be importantly involved in the blood supply of brain and allowing us to single out the vasomotor effects of hemolysed blood without the interference of other mechanisms associated with tissue hemorrhage and traumatic brain injury.

Materials and Methods

Animals

For these experiments ~2 months-old (250 ± 50 g) male Wistar rats (CrI:WI, Charles River Hungary Kft; n=6-12 in each group) were used. Animals were housed on a 12h light/dark cycle and were ad-libitum fed on standard rat chow and free access to tap water. All experiments and interventions were undertaken according to the general rules and special approval of the University of Pecs Ethical Committee for the Protection of Animals in Research (BA 02/2000-8/2008), in accordance with the directives of the National Ethical Council for Animal Research and those of the EU Directive (2010/63/EU), in accordance with the ARRIVE guidelines.

Isolation of cerebral arteries and measurements of diameter

Cerebral vessels were isolated as previously described.¹³⁻¹⁵ In brief, animals were anesthetized by ether and decapitated according to Institutional Animal Care and Use Committee of University of Pecs, Medical School, Pecs, Hungary. The brains were immediately removed and placed in Krebs' buffer. Basilar arteries (BA) and middle cerebral arteries (MCA) were isolated from the brain of each animal. Segments of the BAs and MCAs were isolated using microsurgery instruments. Both ends of the arteries were mounted onto two glass micropipettes in a vessel chamber and pressurized to 80 mmHg with zero flow. The hydrodynamic resistances of the micropipettes were matched. Inflow and outflow pressures were controlled and measured by a pressure servo-control system (Living Systems Instrumentation, Burlington, VT, USA). Inner vascular diameter was measured with a video-micrometer system and continuously recorded using a computerized data acquisition system (LabChart 7 pro by PowerLab, ADInstruments, Australia). All vessels were allowed to stabilize for 60 min in oxygenated (21% O₂; 5% CO₂;

74% N₂) Krebs' buffer (at 37°C). After the equilibration period, during which spontaneous myogenic tone developed (measured as a basal diameter; BD), and the vascular responses were assessed, as reported previously.¹⁵⁻¹⁷ At the end of each experiment the passive diameters (PD) of the vessels were measured at 80 mmHg intraluminal pressure in the presence of Ca²⁺-free Krebs' buffer containing the L-type Ca²⁺ channel inhibitor nifedipine (10⁻⁴ mol/L) to achieve maximal vasodilatation.

Administration of Vasoactive Agents and Inhibitors

The vasomotor effect of perivascular blood was investigated by adding autologous hemolysed whole blood (HB) directly into the vessel chamber. Hemolysed whole blood (200 µL) was prepared by osmolysis from 40 µL whole blood (B) and 160 µL bidestillated water (DW) at ratio B:DW=1:4. In other series of experiments vasomotor function of cerebral arteries were studied in response to blood components, such as blood serum, blood plasma, hemolysed red blood cell (RBC), platelet concentrate (PLTc), platelet suspension (PLTs) and purified hemoglobin (Hgb).

For testing the receptor-independent vasoconstriction 60 mmol/L KCl was used. To assess endothelial function, was tested by vascular responses to acetylcholine (ACh, 10⁻⁴ mol/L), whereas that of smooth muscle by sodium nitroprusside (SNP; 10⁻⁴ mol/L) and the L-type Ca²⁺ channel inhibitor nifedipine (10⁻⁶ mol/L), which was also used to assess the passive diameter (PD) of arteries (10⁻⁴ mol/L).

To assess the vasodilator effect of carbon dioxide (CO₂), normal (5% CO₂; 21% O₂; 74% N₂) and elevated CO₂ (15% CO₂; 21% O₂; 64% N₂) gas mixture were used to bubble Krebs' buffer (for 5 minutes; at 37°C; n=10) in vessel chamber. All drugs were purchased from Sigma Aldrich (Budapest, Hungary).

Potassium concentration was measured by Nova Biomedical pHox plus blood gas analyzer (Massachusetts, USA).

Assessment of intravascular calcium ion level

As described previously¹⁹ changes in intracellular Ca^{2+} -ion concentration were assessed with ratiometric (R) calcium-measurement at the wavelength of 340 nm and 380 nm using 5-Oxazolecarboxylic acid, 2-(6-(bis(2-((acetyloxy)methoxy)-2-oxoethyl)amino)-5-(2-(2-(bis(2-((acetyloxy)methoxy)-2-oxoethyl)amino)-5-methylphenoxy)ethoxy)-2-benzofuranyl)-, (acetyloxy)methyl ester (Fura2-AM; Invitrogen, Life Technologies, Budapest, Hungary) fluorescent dyes.^{20, 21} The physiological Krebs solution was supplemented with Fura2-AM (5 $\mu\text{mol/L}$) fluorescent Ca^{2+} indicator dye and BSA (bovine serum albumin; 1%) for 60 min during which spontaneous myogenic tone developed. We have used fluorescent microscope to measure intracellular Ca^{2+} concentrations by an IncyteIm2 instrument (Intracellular Imaging Inc, Cincinnati, OH, USA) by recording images (cut off >510 nm) excited alternatively by 340 and 380 nm wavelengths. Images were recorded every 4 s and evaluated offline. Arterial Ca^{2+} concentrations were detected by calculating ratios (R) between averaged signal intensity at 340 and 380 nm excitation in the whole arterial segment.

Statistical Analysis

Experimental results are presented as mean \pm S.E.M. Data are expressed as either micrometer or percentage of basal [BD%] and passive diameter [PD%]. The changes in ratiometric intracellular calcium measurements are indicated either as ratio (R) or as a delta ratio (ΔR). Statistical analysis was performed by one-way ANOVA (Holm-Sidak method) or Student's t-test as appropriate by SPSS 11.0 for Windows software. P-values <0.05 were considered to be statistically significant. Figures were made by SigmaPlot 11.0 for Windows software..

Results

Effect of perivascular hemolysed blood and its components on the diameter of cerebral arteries

The basal diameter of BA was 264 ± 7 μm and MCA was 185 ± 15 μm in the presence of 80 mmHg intraluminal pressure, whereas the passive diameter of BA was 392 ± 8 μm and MCA was 282 ± 10 μm . Summary data (Fig. 1) shows that HB elicited significant constrictions of BA (top, $164 \pm 11 \mu\text{m}$, -23.9 ± 3 of PD%) and also in MCA (bottom; $155 \pm 14 \mu\text{m}$, -11.4 ± 0.8 of PD%).

Importantly, after wash-out of HB the basal diameters of cerebral arteries reached level (BA: 288 ± 12 μm ; MCA: 195 ± 12 μm).

Page 10 of 27

Figure 2 shows that KCl (control: 255 ± 18 μm , KCl: 170 ± 20 μm ; -21 ± 2 of PD%), and HB elicited constrictions of cerebral arteries and at the same time there was a significant increase in K^+ concentration in hemolysed blood (HB 7.02 ± 0.22 mmol/L) compared to whole blood (6.20 ± 0.01 mmol/L).

Figure 3 shows summary data of diameter changes [PD%] of BA in response to hemolysed blood (HB), blood serum, hemolysed red blood cell (RBC), blood plasma, platelet suspension (PLTs), platelet concentration (PLTc) and hemoglobin (Hgb). HB (control: 264 ± 7 μm , HB: $164 \pm 11 \mu\text{m}$, -23.9 ± 3 of PD%), Blood serum (control: 246 ± 8 μm , serum: 170 ± 6 μm ; -19 ± 0.9 of PD%), the hemolysed red blood cell (RBC), (control: 217 ± 9 μm , RBC: 166 ± 6 μm ; -14 ± 1 of PD%), blood plasma (control: 258 ± 7 μm , plasma: 226 ± 7 μm ; -7.7 ± 0.5 of PD%), platelet suspension (PLTs), (control: 191 ± 15 μm , PLTs: 165 ± 16 μm ; -7.5 ± 2 of PD%). Whereas, hemoglobin (Hgb) (control: 263 ± 16 μm ; Hgb 10^{-12} M: 263 ± 15 μm ; -0.27 ± 2.11 of PD%; Hgb 10^{-6} M: 274 ± 21 μm ; -0.12 ± 1.91 of PD%) and platelet concentrate (PLTc) (control: 188 ± 11 μm , PLTc: 185 ± 12 μm ; -0.8 ± 0.9 of PD%) did not affect the diameter in the present experimental conditions.

Changes in agonist-induced vasomotor responses to presence of HB

Responses were measured before (control), in the presence of HB and after wash-out of HB. Summary data shows in Figure 4 that in control, the ACh-induced dilations were 19.9 ± 4.6 (% of basal diameter, BD%), presence of HB significantly decreased the dilation to 7.4 ± 1.4 of BD% and after wash-out HB it remained at 5.7 ± 1.7 of BD%). As Figure 4 shows, dilations to SNP in control were 26 ± 2.6 of BD%, which was reduced to 11.8 ± 1.7 of BD% by HB and remained at 13.9 ± 2.2 of BD%. In contrast nifedipine-induced dilations were not significantly affected by HB:

re 32.6 ± 5.1 of BD%, during HB 28.7 ± 3.5 of BD%, after wash-out the HB 30 ± 2.3 of BD%.

Reversal of HB-induced cerebrovascular constrictions in the presence of high CO₂ and nifedipine

Dilations to increased level of CO₂ were measured before (control), during (HB) and after (wash-out) of hemolysed blood (Fig. 4). High CO₂ elicited significant dilations in control (25.7 ± 2.7 of BD%), which did not change significantly in the presence of HB (29.5 ± 1.7 of BD%), or after wash-out HB (27 ± 3 of BD%). Similarly, nifedipine-induced dilations (control: 32.6 ± 5.1 of BD%) were not affected by the presence of HB (28.7 ± 3.5 of BD%) or after wash-out of HB (30.1 ± 2.3 of BD%, respectively).

Changes in vascular [Ca²⁺]_i in response to HB

Summary data (Fig. 5) shows that perivascular HB elicited increases in the ratiometric (R) Ca signal in a concentration-dependent manner (by 20 μ L steps from 0 μ L up to 200 μ L), indicating increase in intravascular [Ca²⁺]_i concentrations. In control conditions, before administration of HB the ratio was 1.118 ± 0.043 ; 100 μ L HB it increased to 1.352 ± 0.019 (Δ ratio= 0.154 ± 0.013) and 200 μ L HB it significantly increased to 1.397 ± 0.016 (Δ ratio= 0.211 ± 0.022), respectively. The

HB-induced constrictions of basilar arteries paralleled with the increases in intracellular Ca^{2+} concentration. After wash-out the ratio significantly decreased (1.076 ± 0.069 ; $\Delta\text{ratio} = -0.293 \pm 0.079$) resulting in dilation.

Discussion

The salient findings of the present study are: 1) perivascular hemolysed blood elicited substantial constrictions of isolated basilar and middle cerebral arteries, 2) which corresponded with increases in vascular wall Ca^{2+} , and could be reversed by the calcium-channel antagonist nifedipine and increased level of CO_2 . In addition it reduced agonists-induced dilations

Page 12 of 27

HB elicits vasoconstriction both in basilar and middle cerebral arteries

In all experiments vessels developed myogenic tone^{3, 13, 16, 17} (passive diameters vs. basal diameters), thus vasomotor capacity of both basilar (BA) and middle cerebral arteries (MCA) could be observed in the presence of optimal tone, without the use of pre-constrictor, which could interfere with cellular vasomotor mechanisms. The data show that addition of HB to the chamber caused significant constrictions in basilar arteries. Interestingly, after washout of HB, basal diameter returned to the control level (Fig. 1). Importantly smaller intracerebral arteries (middle cerebral artery) are also responded with constriction to HB. It is likely that even smaller arterial vessels are affected by HB as previous studies showed that myogenic tone of pial vessels were impaired even after washout of blood²². Nevertheless, HB may elicit vasomotor responses, which are region specific.

Potential mechanisms of reversal of HB-induced constrictions by high pCO₂

Interestingly, data reported in the literature regarding the effect and mediation of pCO₂-induced dilations of cerebral vessels are not unequivocal. For example, the nature of response (dilation or constriction) varied depending on the experimental conditions. The potential effect of changes in pH was supported by some, but refuted by other studies.²³⁻²⁵ There were studies suggesting endothelial^{26, 27} and nitric oxide mediations²⁸, and role for arachidonic acid metabolites^{27, 29}, SK_{Ca}/IK_{Ca} channels^{27, 30} and also changes in vascular cell membrane polarization.^{31, 32} Because of

mentioned we felt it is important to establish the effects of pCO₂ on the vasomotor tone of isolated cerebral arteries, especially in the presence of HB; a condition in which the presence of in vivo confounding factors can be excluded. The finding that vasoconstrictor effect of HB can be reversed by wash-out of blood or decreasing intracellular Ca²⁺ concentration using locally applied Ca-channel antagonists, or increase locally perivascular pCO₂ suggest a key role for intracellular Ca²⁺ level rather than to calcium sensitivity. Also, it seems that high pCO₂ is powerful enough to overcome any constrictor mechanisms or factors operating during hemolysis of blood. We believe that extrapolating these experimental findings to clinical conditions may open up novel therapeutic avenues for subarachnoid hemorrhage especially the powerful effect of perivascular application of high pCO₂ should be explored and documented.

Proposed mechanisms of action of hemolysed blood (HB) and blood component-induced constrictions of cerebral arteries

Blood contains myriad of vasoactive components^{9, 11, 12, 33} thus future studies need to single out the mechanisms finally leading to constriction. For example, blood serum via activating coagulation cascade may contain eicosanoids³⁴/prostanoids³⁵, low molecule weight peptides (endothelin-1³⁶) and thrombin.^{37, 38} Blood plasma circulating with inactive coagulation factors has

less vasoactive properties than serum, but containing fibrinogen³⁹⁻⁴¹ or plasma protein⁴² may result in vasoconstriction. Interestingly, while others^{10, 11} demonstrated that hemoglobin causes vasoconstriction, we could not confirm it in isolated basilar arteries. Hemolysed red blood cell suspension induces vasoconstriction, which could be explained -in part- by released hemoglobin and bilirubin oxidation products¹¹ and potassium¹² ions (see Fig. 2) derived from de-compartmentalized RBC. Interestingly, while platelet concentration had no vasoconstrictor effect, platelet suspension elicited vasoconstriction (also see Fig. 3), likely due to release of thromboxane-A₂ from platelets.⁴³ In addition to these mechanisms, we propose a po

Page 14 of 27

for high K⁺ in HB-induced constrictions. During hemolysis high amount of K⁺ is released from red blood cells, which can reach a constrictor level. Perivascular application of KCl shows (Fig. 2) that it can elicit substantial constrictions, similar to that of HB.

The finding that HB impaired the endothelium and smooth muscle nitric oxide pathways (ACh, SNP; Fig. 4) suggest in the presence of HB high level of K⁺ effects directly the smooth muscle eliciting depolarization²² and thus increases Ca²⁺ level and similar level of constriction as exogenous KCl. Nevertheless, it is a clinically relevant finding that washing out of blood reversed the constrictions (Fig. 1).

Effect of HB on agonists-induced dilations

Many previous studies^{3, 44, 45} established that endothelium-derived factors are important in the modulation of vasomotor tone of cerebral arteries. Previous data showed that oxyhemoglobin induces significant constriction of cerebral arteries which was explained – in part - by binding nitric oxide (NO).^{10, 11} On the other hand hemoglobin may act directly on smooth muscle cell by activating tyrosin-kinase thus inactivating voltage dependent potassium channels (K_{v1.5}).²² In the presence of HB (Fig. 4), ACh- and SNP-induced dilations mechanisms were significantly reduced which, after washout of HB remained impair, suggesting that HB affects both endothelial

an smooth muscle NO-related mediations, which remain impaired even after washout of HB. These findings suggest that although HB-induced constrictions can be reversed, some of the important vasomotor mechanisms remain impaired, which may have clinical significance.

HB increases the level of vascular wall $[Ca^{2+}]_i$

Summary data (Fig. 5) shows that HB, in a concentration-dependent manner (by 20 μ L steps from 0 μ L up to 200 μ L) increased the ratiometric (R) Ca signal indicating increase in

ar $[Ca^{2+}]_i$ concentration¹⁹⁻²¹. Since we have found that HB elicited constrictions of basilar arteries, we hypothesized that regardless of proximal signaling pathways, HB by increasing the intravascular Ca^{2+} level, results in constrictions. Interestingly, wash-out of HB, significantly decreased the $[Ca^{2+}]_i$ reaching the control level. The findings regarding the parallel changes in the vascular $[Ca^{2+}]_i$ and the diameter suggests that the final signaling mechanism by which HB elicits constriction of cerebral arteries is an elevation of smooth muscle intracellular Ca^{2+} concentration.

Clinical importance

Searching for effective pharmaceutical treatments to improve cerebral blood flow in diseased conditions, such as hemorrhagic stroke⁴⁶ or traumatic brain injury (TBI)⁴⁷⁻⁴⁹ is an ongoing clinical effort. In these conditions the resistance of cerebral vessel greatly increases reducing the regional blood supply of brain. Our findings that direct perivascular application of HB (without traumatic brain injury, and in the absence of neural or other tissue factors) elicited substantial constrictions, which however can be reversed by local application of calcium channel antagonist or high pCO₂ suggest that they could be utilized in clinical area and may open up novel therapeutic possibilities for subarachnoid hemorrhage.

In conclusion, extravascular hemolysed blood elicits substantial constriction of cerebral arteries of different sizes by increasing the level of smooth muscle Ca^{2+} , which however could be reversed by perivascular administration of calcium antagonist and increasing CO_2 level. These findings could advance the development of novel therapies during hemorrhagic stroke, traumatic brain injury and surgery to overcome cerebrovascular spasm and thereby providing appropriate blood flow to the affected brain regions.

Page 16 of 27

Acknowledgements

We gratefully thank Professor Janos Hamar for his valuable suggestions and critical remarks. We thank Ms. Viktoria Csato for additional calcium image measurements and Istvan Zoard Batai for performing some of the vascular experiments.

Support: American Heart Association, Founders Affiliate, 0855910D, Hungarian National Science Research Fund (OTKA) K 108444; Developing Competitiveness of Universities in the South Transdanubian Region, “Identification of new biomarkers...”, SROP-4.2.2.A-11/1/KONV-2012–0017 and “Complex examination of neuropeptide...” SROP-4.2.2.A-11/1/KONV-2012-0024; SROP-4.2.4.A/2-11-1-2012-0001 “National Excellence Program”; Hungarian Hypertension Society (MHT) 2012-2015.

Disclosure/conflict of Interest

No competing financial interests exist.

References

1. Diedler, J., Sykora, M., Rupp, A., Poli, S., Karpel-Massler, G., Sakowitz, O. and Steiner, T. (2009). Impaired cerebral vasomotor activity in spontaneous intracerebral hemorrhage. *Stroke* 40, 815-819.
2. Delgado, T.J., Brismar, J. and Svendgaard, N.A. (1985). Subarachnoid haemorrhage in the rat: angiography and fluorescence microscopy of the major cerebral arteries. *Stroke* 16, 595-602.
3. Koller, A. and Toth, P. (2012). Contribution of Flow-Dependent Vasomotor Mechanisms to the Autoregulation of Cerebral Blood Flow. *Journal of vascular research* 49, 375-389.
4. Paulson, O.B., Strandgaard, S. and Edvinsson, L. (1990). Cerebral autoregulation. *Cerebrovascular and brain metabolism reviews* 2, 161-192.
5. Faraci, F.M., Baumbach, G.L. and Heistad, D.D. (1989). Myogenic mechanisms in the cerebral circulation. *J Hypertens Suppl* 7, S61-64; discussion S65.
6. Johansson, B. (1989). Myogenic tone and reactivity: definitions based on muscle physiology. *J Hypertens Suppl* 7, S5-8; discussion S9.
7. Cipolla, M.J. and Curry, A.B. (2002). Middle cerebral artery function after stroke: the threshold duration of reperfusion for myogenic activity. *Stroke* 33, 2094-2099.
8. Tamaki, K., Sadoshima, S., Baumbach, G.L., Iadecola, C., Reis, D.J. and Heistad, D.D. (1984). Evidence that disruption of the blood-brain barrier precedes reduction in cerebral blood flow in hypertensive encephalopathy. *Hypertension* 6, 175-81.
9. Sasaki, T. and Kikkawa, Y. (2013). Proposed mechanism of cerebral vasospasm: our hypothesis and current topics. *Acta Neurochir Suppl* 115, 53-56.
10. Minneci, P.C., Deans, K.J., Zhi, H., Yuen, P.S., Star, R.A., Banks, S.M., Schechter, A.N., Natanson, C., Gladwin, M.T. and Solomon, S.B. (2005). Hemolysis-associated endothelial dysfunction mediated by accelerated NO inactivation by decompartmentalized oxyhemoglobin. *J Clin Invest* 115, 3409-3417.
11. Clark, J.F. and Sharp, F.R. (2006). Bilirubin oxidation products (BOXes) and their role in cerebral vasospasm after subarachnoid hemorrhage. *J Cereb Blood Flow Metab* 26, 1223-1233.
12. Sobey, C.G. and Faraci, F.M. (1998). Subarachnoid haemorrhage: what happens to the cerebral arteries? *Clin Exp Pharmacol Physiol* 25, 867-876.
13. Toth, P., Rozsa, B., Springo, Z., Doczi, T. and Koller, A. (2011). Isolated human and rat cerebral arteries constrict to increases in flow: role of 20-HETE and TP receptors. *J Cereb Blood Flow Metab* 31, 2096-2105.
14. Ungvari, Z., Pacher, P., Kecskemeti, V. and Koller, A. (1999). Fluoxetine dilates isolated small cerebral arteries of rats and attenuates constrictions to serotonin, norepinephrine, and a voltage-dependent Ca(2+) channel opener. *Stroke* 30, 1949-1954.
15. Toth, P., Csiszar, A., Sosnowska, D., Tucsek, Z., Cseplo, P., Springo, Z., Tarantini, S., Sonntag, W.E., Ungvari, Z. and Koller, A. (2013). Treatment with the cytochrome P450 omega-hydroxylase inhibitor HET0016 attenuates cerebrovascular inflammation, oxidative stress and improves vasomotor function in spontaneously hypertensive rats. *Br J Pharmacol* 168, 1878-1888.
16. Osol, G. and Halpern, W. (1985). Myogenic properties of cerebral blood vessels from normotensive and hypertensive rats. *Am J Physiol* 249, H914-921.
17. Osol, G., Laher, I. and Cipolla, M. (1991). Protein kinase C modulates basal myogenic tone in resistance arteries from the cerebral circulation. *Circ Res* 68, 359-367.
18. Vamos, Z., Cseplo, P., Ivic, I., Matics, R., Hamar, J. and Koller, A. (2013). Age Determines the Magnitudes of Angiotensin II-Induced Contractions, mRNA, and Protein Expression of Angiotensin Type 1 Receptors in Rat Carotid Arteries. *J Gerontol A Biol Sci Med Sci*.

19. Czikora, A., Lizanecz, E., Bako, P., Rutkai, I., Ruzsnavszky, F., Magyar, J., Porszasz, R., Kark, T., Facsko, A., Papp, Z., Edes, I. and Toth, A. (2012). Structure-activity relationships of vanilloid receptor agonists for arteriolar TRPV1. *Br J Pharmacol* 165, 1801-1812.
20. Ungvari, Z., Pacher, P. and Koller, A. (2000). Serotonin reuptake inhibitor fluoxetine decreases arteriolar myogenic tone by reducing smooth muscle $[Ca^{2+}]_i$. *J Cardiovasc Pharmacol* 35, 849-854.
21. Grynkiewicz, G., Poenie, M. and Tsien, R.Y. (1985). A new generation of Ca^{2+} indicators with greatly improved fluorescence properties. *J Biol Chem* 260, 3440-3450.
22. Nystoriak, M.A., O'Connor, K.P., Sonkusare, S.K., Brayden, J.E., Nelson, M.T. and Wellman, G.C. (2011). Fundamental increase in pressure-dependent constriction of brain parenchymal arterioles from subarachnoid hemorrhage model rats due to membrane depolarization. *Am J Physiol Heart Circ Physiol* 300, H803-812.
23. Harder, D.R. and Madden, J.A. (1985). Cellular mechanism of force development in cat middle cerebral artery by reduced PCO_2 . *Pflugers Arch* 403, 402-406.
24. Toda, N., Hatano, Y. and Mori, K. (1989). Mechanisms underlying response to hypercapnia and bicarbonate of isolated dog cerebral arteries. *Am J Physiol* 257, H141-146.
25. Edvinsson, L. and Sercombe, R. (1976). Influence of pH and pCO_2 on alpha-receptor mediated contraction in brain vessels. *Acta Physiol Scand* 97, 325-331.
26. Aoyama, Y., Ueda, K., Setogawa, A. and Kawai, Y. (1999). Effects of pH on contraction and Ca^{2+} mobilization in vascular smooth muscles of the rabbit basilar artery. *Jpn J Physiol* 49, 55-62.
27. Yoon, S., Zuccarello, M. and Rapoport, R.M. (2012). pCO_2 and pH regulation of cerebral blood flow. *Front Physiol* 3, 365.
28. Kim, Y.C., Lee, S.J. and Kim, K.W. (2004). Effects of pH on vascular tone in rabbit basilar arteries. *Journal of Korean medical science* 19, 42-50.
29. Leffler, C.W., Parfenova, H., Basuroy, S., Jaggar, J.H., Umstot, E.S. and Fedinec, A.L. (2011). Hydrogen sulfide and cerebral microvascular tone in newborn pigs. *Am J Physiol Heart Circ Physiol* 300, H440-447.
30. Hannah, R.M., Dunn, K.M., Bonev, A.D. and Nelson, M.T. (2011). Endothelial SK(Ca) and IK(Ca) channels regulate brain parenchymal arteriolar diameter and cortical cerebral blood flow. *J Cereb Blood Flow Metab* 31, 1175-1186.
31. Peng, H.L., Ivarsen, A., Nilsson, H. and Aalkjaer, C. (1998). On the cellular mechanism for the effect of acidosis on vascular tone. *Acta Physiol Scand* 164, 517-525.
32. Peng, H.L., Jensen, P.E., Nilsson, H. and Aalkjaer, C. (1998). Effect of acidosis on tension and $[Ca^{2+}]_i$ in rat cerebral arteries: is there a role for membrane potential? *Am J Physiol* 274, H655-662.
33. Qureshi, A.I., Mendelow, A.D. and Hanley, D.F. (2009). Intracerebral haemorrhage. *Lancet* 373, 1632-1644.
34. Kehl, F., Cambj-Sapunar, L., Maier, K.G., Miyata, N., Kametani, S., Okamoto, H., Hudetz, A.G., Schulte, M.L., Zagorac, D., Harder, D.R. and Roman, R.J. (2002). 20-HETE contributes to the acute fall in cerebral blood flow after subarachnoid hemorrhage in the rat. *Am J Physiol Heart Circ Physiol* 282, H1556-1565.
35. Uski, T.K. and Andersson, K.E. (1984). Effects of prostanoids on isolated feline cerebral arteries. I. Characterization of the contraction-mediating receptor. *Acta Physiol Scand* 120, 131-136.
36. Peterson, E.C., Wang, Z. and Britz, G. (2011). Regulation of cerebral blood flow. *Int J Vasc Med* 2011, 823525.
37. Hirano, K. and Hirano, M. (2010). Current perspective on the role of the thrombin receptor in cerebral vasospasm after subarachnoid hemorrhage. *J Pharmacol Sci* 114, 127-133.
38. Maeda, Y., Hirano, K., Kai, Y., Hirano, M., Suzuki, S.O., Sasaki, T. and Kanaide, H. (2007). Up-regulation of proteinase-activated receptor 1 and increased contractile responses to thrombin after subarachnoid haemorrhage. *Br J Pharmacol* 152, 1131-1139.
39. Lominadze, D., Tsakadze, N., Sen, U., Falcone, J.C. and D'Souza, S.E. (2005). Fibrinogen and fragment D-induced vascular constriction. *Am J Physiol Heart Circ Physiol* 288, H1257-1264.

40. Sen, U., Tyagi, N., Patibandla, P.K., Dean, W.L., Tyagi, S.C., Roberts, A.M. and Lominadze, D. (2009). Fibrinogen-induced endothelin-1 production from endothelial cells. *Am J Physiol Cell Physiol* 296, C840-847.
41. Wurzel, M., Bacon, R.C., Kalt, R.B. and Zweifach, B.W. (1964). Vasoactive Properties of Plasma Protein Fractions. *Am J Physiol* 206, 923-925.
42. Culliver, H.A. and Penington, D.G. (1979). Mechanisms of vasomotor reactions in the use of SPPS. *Vox Sang* 36, 201-207.
43. Neppl, R.L., Lubomirov, L.T., Momotani, K., Pfitzer, G., Eto, M. and Somlyo, A.V. (2009). Thromboxane A₂-induced bi-directional regulation of cerebral arterial tone. *J Biol Chem* 284, 6348-6360.
44. Andresen, J., Shafi, N.I. and Bryan, R.M., Jr. (2006). Endothelial influences on cerebrovascular tone. *Journal of applied physiology* 100, 318-327.
45. Cosentino, F., Rubattu, S., Savoia, C., Venturelli, V., Pagannone, E. and Volpe, M. (2001). Endothelial dysfunction and stroke. *J Cardiovasc Pharmacol* 38 Suppl 2, S75-78.
46. Pool, J.L., Jacobson, S. and Fletcher, T.A. (1958). Cerebral vasospasm; clinical and experimental evidence. *Journal of the American Medical Association* 167, 1599-1601.
47. Kontos, H.A., Wei, E.P. and Povlishock, J.T. (1981). Pathophysiology of vascular consequences of experimental concussive brain injury. *Transactions of the American Clinical and Climatological Association* 92, 111-121.
48. Rosenblum, W.I., Povlishock, J.T., Wei, E.P., Kontos, H.A. and Nelson, G.H. (1987). Ultrastructural studies of pial vascular endothelium following damage resulting in loss of endothelium-dependent relaxation. *Stroke* 18, 927-931.
49. Baranova, A.I., Wei, E.P., Ueda, Y., Sholley, M.M., Kontos, H.A. and Povlishock, J.T. (2008). Cerebral vascular responsiveness after experimental traumatic brain injury: the beneficial effects of delayed hypothermia combined with superoxide dismutase administration. *J Neurosurg* 109, 502-509.

Figure Legends:





FIG. 1. Summary data of changes in diameter (μm) of basilar arteries BA (top, from $264 \pm 7 \mu\text{m}$ to $164 \pm 11 \mu\text{m}$, -23.9 ± 3 [PD%]; $n=12$) and middle cerebral arteries MCA (bottom, from $185 \pm 15 \mu\text{m}$ to $155 \pm 14 \mu\text{m}$, -11.4 ± 0.8 [PD%]; $n=6$) in response to hemolysed blood (HB). Data are mean \pm S.E.M. (* $p < 0.05$ between either HB and control or HB and wash-out; # $p < 0.05$ between control and passive diameter).




FIG. 2. Summary data of changes in diameter ([μm] on axis Y1) of basilar arteries (BA) in response to hemolysed blood (HB) or potassium chloride (KCl). Summary data of changes in K⁺ concentration [mmol/L] in whole blood (K⁺c; 6.20±0.01 mmol/L) and in hemolysed blood (K⁺HB; 7.02±0.22 mmol/L) on axis Y2. Data are mean ± S.E.M. (* p<0.05 between either control and HB or control and KCl, n=9-12 in each group; # p<0.05 between K⁺c and K⁺HB; n=9).




FIG. 3. Summary data of changes in diameter ([PD%]; % of passive diameter) of basilar arteries (BA) in response to hemolysed blood (HB), blood serum, hemolysed red blood cell (RBC), blood plasma, platelet suspension (PLTs), platelet concentration (PLTc) and hemoglobin (Hgb). Data are mean \pm S.E.M. (* $p < 0.05$, $n = 9-12$ in each group). KCl (control: 255 ± 18 μm , KCl: 170 ± 20 μm ; -21 ± 2 [PD%]), HB (control: 264 ± 7 μm , HB: 164 ± 11 μm , -23.9 ± 3 [PD%]), Blood serum (control: 246 ± 8 μm , serum: 170 ± 6 μm ; -19 ± 0.9 [PD%]), the hemolysed red blood cell (RBC), (control: 217 ± 9 μm , RBC: 166 ± 6 μm ; -14 ± 1 [PD%]), blood plasma (control: 258 ± 7 μm , plasma: 226 ± 7 μm ; -7.7 ± 0.5 [PD%]), platelet suspension (PLTs), (control: 191 ± 15 μm , PLTs: 165 ± 16 μm ; -7.5 ± 2 [PD%]) caused significant vasoconstriction. However, hemoglobin (Hgb) (control: 263 ± 16 μm ; Hgb 10^{-12} M: 263 ± 15 μm ; -0.27 ± 2.11 [PD%]; Hgb 10^{-6} M: 274 ± 21 μm ; -0.12 ± 1.91

[PD%]) and platelet concentrate (PLTc) (control: $188 \pm 11 \mu\text{m}$, PLTc: $185 \pm 12 \mu\text{m}$; -0.8 ± 0.9 [PD%]) did not elicit significant vasoconstriction.




FIG. 4. Summary data of changes in diameter of isolated basilar arteries (BA). (% of basal diameter at 80 mmHg; [BD%]) in response to ACh, SNP, nifedipine and CO₂ before (control), during (HB) and after (wash-out) of hemolysed blood (HB). Data are mean \pm S.E.M. (* $p < 0.05$ between either control and HB or control and wash-out; $n = 10-12$ in each group).




FIG. 5. Summary data of ratiometric (R) changes indicating the level of intravascular $[Ca^{2+}]_i$ of basilar arteries in response to increased concentrations of hemolysed blood (HB). Data are mean \pm S.E.M. (* $p < 0.05$; $n = 18$).