



# **Hermite–Hadamard-type inequalities for generalized convex functions**

**doktori (Ph.D.) értekezés**

**Bessenyei Mihály**

**DEBRECENI EGYETEM  
TERMÉSZETTUDOMÁNYI KAR**

**Debrecen, 2004**



Ezen értekezést a Debreceni Egyetem Matematika doktori program Analízis alprogramja keretében készítettem 2000-2003 között és ezúton nyújtom be a Debreceni Egyetem doktori (Ph.D.) fokozatának elnyerése céljából.

Debrecen, 2004. december 27.

.....  
Bessenyei Mihály  
jelölt

Tanúsítom, hogy Bessenyei Mihály doktorjelölt 2000-2003 között a fent nevezett doktori alprogram keretében irányításommal végezte munkáját. Az értekezésben foglaltak a jelölt önálló munkáján alapulnak, az eredményekhez önálló alkotó tevékenységével meghatározó módon járult hozzá. Az értekezés elfogadását javaslom.

Debrecen, 2004. december 27.

.....  
Dr. Páles Zsolt  
témavezető



*“Laus viventi Deo!”*

## **Acknowledgements**

First I thank my parents and my family for providing the possibility of suitable education and peaceful background to my studies, my work and my life. I thank them and also my friends for their frequent encouragement and support.

I wish to express my deep gratitude to my supervisor, DR. ZSOLT PÁLES for the outstanding guidance and support during my PhD years and also till now. His invaluable advice and excellent ideas exhaustively influenced both the contents and the form of the present dissertation. I thank him for accepting me as a Ph.D. student and conducting my researches.

I thank the leader of the Ph.D. school, DR. ZOLTÁN DARÓCZY as well as DR. LÁSZLÓ LOSONCZI for their confidence in me and providing me the opportunity to study and do research work as a Ph.D. fellow. I thank them the kind support and help they presented in various ways during my studies.

I also wish to express my gratitude to DR. GYULA MAKSA, the leader of the department, and all of my colleagues for the continuous encouragement over the years and sharing their knowledge and love of the field of Analysis with me.

My special thanks go to ANDREA GILÁNYINÉ PÁKOZDY for reviewing and correcting the text of this dissertation. Without her valuable comments and advice my co-author were Miss Prints...

At last, but not at least I express my thanks to GÁBOR DEÁK, EDIT KUN-SZABÓNÉ DANCS, DR. JÓZSEF SZILASI and all my former teachers who deeply influenced my choice of profession for arousing my interest towards the beauty of mathematics.



## Contents

Introduction	1
Chapter 1. Polynomial convexity	5
1.1. Orthogonal polynomials and basic quadrature formulae	5
1.2. An approximation theorem	12
1.3. Hermite–Hadamard-type inequalities	14
1.4. Applications	21
Chapter 2. Generalized 2-convexity	25
2.1. Characterizations via generalized lines	25
2.2. Connection with standard convexity	34
2.3. Hermite–Hadamard-type inequalities	36
2.4. Applications	38
Chapter 3. Generalized convexity induced by Tchebychev systems	41
3.1. Characterizations and regularity properties	41
3.2. Moment spaces induced by Tchebychev systems	44
3.3. Hermite–Hadamard-type inequalities	46
3.4. An alternative approach in a particular case	51
Chapter 4. Characterizations via Hermite–Hadamard inequalities	57
4.1. Further properties of generalized lines	57
4.2. Hermite–Hadamard-type inequalities and $(\omega_1, \omega_2)$ -convexity	60
Summary	65
Összefoglaló	73
Bibliography	81



## Introduction

Let  $I$  be a real interval, that is, a nonempty, connected and bounded subset of  $\mathbb{R}$ . An  $n$ -dimensional *Tchebychev system* on  $I$  consists of a set of real valued continuous functions  $\omega_1, \dots, \omega_n$  and is determined by the property that each  $n$  points of  $I \times \mathbb{R}$  with distinct first coordinates can uniquely be interpolated by a linear combination of the functions. More precisely, we have the following

**DEFINITION.** *Let  $I \subset \mathbb{R}$  be a real interval and  $\omega_1, \dots, \omega_n : I \rightarrow \mathbb{R}$  be continuous functions. Denote the column vector whose components are  $\omega_1, \dots, \omega_n$  in turn by  $\boldsymbol{\omega}$ , that is,  $\boldsymbol{\omega} := (\omega_1, \dots, \omega_n)$ . We say that  $\boldsymbol{\omega}$  is a Tchebychev system over  $I$  if, for all elements  $x_1 < \dots < x_n$  of  $I$ , the following inequality holds:*

$$\begin{vmatrix} \boldsymbol{\omega}(x_1) & \cdots & \boldsymbol{\omega}(x_n) \end{vmatrix} > 0.$$

In fact, it suffices to assume that the determinant above is nonvanishing whenever the arguments  $x_1, \dots, x_n$  are pairwise distinct points of the domain. Indeed, Bolzano's theorem guarantees that its sign is constant if the arguments are supposed to be in an increasing order, hence the components  $\omega_1, \dots, \omega_n$  can always be rearranged such that  $\boldsymbol{\omega}$  fulfills the requirement of the definition. However, considering Tchebychev systems as vectors of functions instead of sets of functions is widely accepted in the technical literature and also turns out to be very convenient in our investigations.

Without claiming completeness, let us list some important and classical examples of Tchebychev systems. In each example  $\boldsymbol{\omega}$  is defined on an arbitrary  $I \subset \mathbb{R}$  except for the last one where  $I \subset ] -\frac{\pi}{2}, \frac{\pi}{2}[$ .

- polynomial system:  $\boldsymbol{\omega}(x) := (1, x, \dots, x^n)$
- exponential system:  $\boldsymbol{\omega}(x) := (1, \exp x, \dots, \exp nx)$
- hyperbolic system:  $\boldsymbol{\omega}(x) := (1, \cosh x, \sinh x, \dots, \cosh nx, \sinh nx)$
- trigonometric system:  $\boldsymbol{\omega}(x) := (1, \cos x, \sin x, \dots, \cos nx, \sin nx)$ .

We make no attempt here to present an exhaustive account of the theory of Tchebychev systems, just mention that, motivated by some results of A. A. Markov, the first systematic investigations of the geometric theory of Tchebychev systems were done by M. G. Krein. However, let us note that Tchebychev systems play an important role, sometimes indirectly, in numerous fields of mathematics, for example, in the theory of approximation, numerical analysis and the theory of inequalities. The books [KS66] and [Kar68] contain a rich literature and bibliography of the topics for the interested reader.

The notion of convexity can also be extended applying Tchebychev systems:

DEFINITION. Let  $\omega = (\omega_1, \dots, \omega_n)$  be a Tchebychev system over the real interval  $I$ . A function  $f : I \rightarrow \mathbb{R}$  is said to be generalized convex with respect to  $\omega$  if, for all elements  $x_0 < \dots < x_n$  of  $I$ , it satisfies the inequality

$$(-1)^n \begin{vmatrix} f(x_0) & \dots & f(x_n) \\ \omega(x_0) & \dots & \omega(x_n) \end{vmatrix} \geq 0.$$

There are other alternatives to express that  $f$  is generalized convex with respect to  $\omega$ , for example,  $f$  is *generalized  $\omega$ -convex* or simply  *$\omega$ -convex*. If the underlying  $n$ -dimensional Tchebychev system can uniquely be identified from the context, we briefly say that  $f$  is *generalized  $n$ -convex*.

If  $\omega$  is the polynomial Tchebychev system, the definition leads to the notion of higher-order monotonicity which was introduced and studied by T. Popoviciu in a sequence of papers [Pop36, Pop38b, Pop38a, Pop39b, Pop39a, Pop40c, Pop40e, Pop40a, Pop40f, Pop40d, Pop40b, Pop41, Pop42b, Pop42a, Pop42c, Pop43]. A summary of these results can be found in [Pop44] and also in [Kuc85]. For the sake of unique terminology, throughout the dissertation Popoviciu's setting is called polynomial convexity. That is, a function  $f : I \rightarrow \mathbb{R}$  is said to be *polynomially  $n$ -convex* if, for all elements  $x_0 < \dots < x_n$  of  $I$ , it satisfies the inequality

$$(-1)^n \begin{vmatrix} f(x_0) & \dots & f(x_n) \\ 1 & \dots & 1 \\ x_0 & \dots & x_n \\ \vdots & \ddots & \vdots \\ x_0^{n-1} & \dots & x_n^{n-1} \end{vmatrix} \geq 0.$$

Observe that polynomially 2-convex functions are exactly the "standard" convex ones. The case, when the "generalized" convexity notion is induced by the special two dimensional Tchebychev system  $\omega_1(x) := 1$  and  $\omega_2(x) := x$ , is termed *standard setting* and *standard convexity*, respectively.

The integral average of any standard convex function  $f : [a, b] \rightarrow \mathbb{R}$  can be estimated from the midpoint and the endpoints of the domain as follows:

$$f\left(\frac{a+b}{2}\right) \leq \int_a^b f(x)dx \leq \frac{f(a)+f(b)}{2}.$$

This is the well known Hadamard's inequality ([Had93]) or, as it is quoted for historical reasons (see [ML85] for interesting remarks), the Hermite–Hadamard-inequality.

The aim of the dissertation is to verify analogous inequalities for generalized convex functions, that is, to give lower and upper estimations for the integral average of the function using certain base points of the domain. Of course, the base points are supposed to depend only on the underlying Tchebychev system of the induced convexity.

---

For this purpose, we shall follow an inductive approach since it seems to have more advantages than the deductive one. First of all, it makes the original motivations clear; on the other hand, it allows us to use the most suitable mathematical tools. Hence sophisticated proofs that sometimes occur when using deductive approach can also be avoided.

CHAPTER 1 investigates the case of polynomial convexity. The base points of the Hermite–Hadamard-type inequalities turn out to be the roots of certain orthogonal polynomials. The main tools of the chapter are based on some methods of numerical analysis, like Gauss quadrature formula and Hermite-interpolation. A smoothing technique and two theorems of Popoviciu are also crucial.

In CHAPTER 2 we present Hermite–Hadamard-type inequalities for generalized 2-convex functions. The most important auxiliary result of the proof is a characterization theorem which, in the standard setting, reduces to the well known characterization properties of convex functions. Another theorem of the chapter establishes a tight relationship between standard and generalized 2-convexity. This result has important regularity consequences and is also essential in verifying Hermite–Hadamard-type inequalities.

The general case is studied in CHAPTER 3. The main results guarantee only the existence and also the uniqueness of the base points of the Hermite–Hadamard-type inequalities but offer no explicit formulae for determining them. The main tool of the chapter is the Krein–Markov theory of moment spaces induced by Tchebychev systems. In some special cases (when the dimension of the underlying Tchebychev systems are “small”), an elementary alternative approach is also presented.

CHAPTER 4 is devoted to showing that, at least in the two dimensional case and requiring weak regularity conditions, Hermite–Hadamard-type inequalities are not merely the consequences of generalized convexity, but they also characterize it.

Specializing the members of Tchebychev systems, several applications and examples are presented for concrete Hermite–Hadamard-type inequalities in both the cases of polynomial convexity and generalized 2-convexity. As a simple consequence, the classical Hermite–Hadamard inequality is among the corollaries in each cases, too.

The results of the dissertation can be found in [BP02, BP03, BP04, BP05, BP] and [Bes04]. In the sequel, we present them without any further references to the mentioned papers.



## CHAPTER 1

### Polynomial convexity

The main results of this chapter state Hermite–Hadamard-type inequalities for polynomially convex functions. Let us recall that a function  $f : I \rightarrow \mathbb{R}$  is said to be *polynomially  $n$ -convex* if, for all elements  $x_0 < \dots < x_n$  of  $I$ , it satisfies the inequality

$$(-1)^n \begin{vmatrix} f(x_0) & \dots & f(x_n) \\ 1 & \dots & 1 \\ x_0 & \dots & x_n \\ \vdots & \ddots & \vdots \\ x_0^{n-1} & \dots & x_n^{n-1} \end{vmatrix} \geq 0.$$

In order to determine the base points and the coefficients of the inequalities, Gauss-type quadrature formulae are applied. Then, using the remainder term of the Hermite-interpolation, the main results follow immediately for “sufficiently smooth” functions due to the next two theorems of Popoviciu:

**THEOREM A.** ([Kuc85, Theorem 1. p. 387]) *Assume that  $f : I \rightarrow \mathbb{R}$  is continuous and  $n$  times differentiable on the interior of  $I$ . Then,  $f$  is polynomially  $n$ -convex if and only if  $f^{(n)} \geq 0$  on the interior of  $I$ .*

**THEOREM B.** ([Kuc85, Theorem 1. p. 391]) *Assume that  $f : I \rightarrow \mathbb{R}$  is polynomially  $n$ -convex and  $n \geq 2$ . Then,  $f$  is  $(n - 2)$  times differentiable and  $f^{(n-2)}$  is continuous on the interior of  $I$ .*

To drop the regularity assumptions, a smoothing technique is developed that guarantees the approximation of polynomially convex functions with smooth polynomially convex ones.

#### 1.1. Orthogonal polynomials and basic quadrature formulae

In what follows,  $\rho$  denotes a positive, locally integrable function (shortly: *weight function*) on an interval  $I$ . The polynomials  $P$  and  $Q$  are said to be *orthogonal on  $[a, b] \subset I$  with respect to the weight function  $\rho$*  or simply  *$\rho$ -orthogonal on  $[a, b]$*  if

$$\langle P, Q \rangle_\rho := \int_a^b PQ\rho = 0.$$

A system of polynomials is called a  $\rho$ -orthogonal polynomial system on  $[a, b] \subset I$  if each member of the system is  $\rho$ -orthogonal to the others on  $[a, b]$ . Define the moments of  $\rho$  by the formulae

$$\mu_k := \int_a^b x^k \rho(x) dx \quad (k = 0, 1, 2, \dots).$$

Then, the  $n^{\text{th}}$  degree member of the  $\rho$ -orthogonal polynomial system on  $[a, b]$  has the following representation via the moments of  $\rho$ :

$$P_n(x) := \begin{vmatrix} 1 & \mu_0 & \cdots & \mu_{n-1} \\ x & \mu_1 & \cdots & \mu_n \\ \vdots & \vdots & \ddots & \vdots \\ x^n & \mu_n & \cdots & \mu_{2n-1} \end{vmatrix}.$$

Clearly, it suffices to show that  $P_n$  is  $\rho$ -orthogonal to the special polynomials  $1, x, \dots, x^{n-1}$ . Indeed, for  $k = 1, \dots, n$ , the first and the  $(k+1)^{\text{st}}$  columns of the determinant  $\langle P_n(x), x^{k-1} \rangle_\rho$  are linearly dependent according to the definition of the moments.

In fact, the moments and the orthogonal polynomials depend heavily on the interval  $[a, b]$ . Therefore, we use the notions  $\mu_{k;[a,b]}$  and  $P_{n;[a,b]}$  instead of  $\mu_k$  and  $P_n$  above when we want to or have to emphasize the dependence on the underlying interval.

Throughout this chapter, the following property of the roots of orthogonal polynomials plays a key role (see [Szeg39]). Let  $P_n$  denote the  $n^{\text{th}}$  degree member of the  $\rho$ -orthogonal polynomial system on  $[a, b]$ . Then,  $P_n$  has  $n$  pairwise distinct roots  $\xi_1 < \dots < \xi_n$  in  $]a, b[$ .

Let us consider the following

$$(1.1) \quad \int_a^b f \rho = \sum_{k=1}^n c_k f(\xi_k)$$

$$(1.2) \quad \int_a^b f \rho = c_0 f(a) + \sum_{k=1}^n c_k f(\xi_k)$$

$$(1.3) \quad \int_a^b f \rho = \sum_{k=1}^n c_k f(\xi_k) + c_{n+1} f(b)$$

$$(1.4) \quad \int_a^b f \rho = c_0 f(a) + \sum_{k=1}^n c_k f(\xi_k) + c_{n+1} f(b)$$

Gauss-type quadrature formulae where the coefficients and the base points are to be determined so that (1.1), (1.2), (1.3) and (1.4) be exact when  $f$  is a polynomial of degree at most  $2n-1$ ,  $2n$ ,  $2n$  and  $2n+1$ , respectively. The subsequent four theorems investigate these cases.

**THEOREM 1.1.** *Let  $P_n$  be the  $n^{\text{th}}$  degree member of the orthogonal polynomial system on  $[a, b]$  with respect to the weight function  $\rho$ . Then (1.1) is exact for polynomials  $f$  of degree at most  $2n - 1$  if and only if  $\xi_1, \dots, \xi_n$  are the zeros of  $P_n$ , and*

$$(1.5) \quad c_k = \int_a^b \frac{P_n(x)}{(x - \xi_k)P_n'(\xi_k)} \rho(x) dx.$$

Furthermore,  $\xi_1, \dots, \xi_n$  are pairwise distinct elements of  $]a, b[$ , and  $c_k \geq 0$  for all  $k = 1, \dots, n$ .

This theorem follows easily from well known results in numerical analysis (see [HH94], [Joh66], [Szeg39]). For the sake of completeness, we provide a proof.

**PROOF.** First assume that  $\xi_1, \dots, \xi_n$  are the zeros of the polynomial  $P_n$  and, for all  $k = 1, \dots, n$ , denote the primitive Lagrange-interpolation polynomials by  $L_k : [a, b] \rightarrow \mathbb{R}$ . That is,

$$L_k(x) := \begin{cases} \frac{P_n(x)}{(x - \xi_k)P_n'(\xi_k)} & \text{if } x \neq \xi_k \\ 1 & \text{if } x = \xi_k. \end{cases}$$

If  $Q$  is a polynomial of degree at most  $2n - 1$ , then, using Euclidian algorithm,  $Q$  can be written in the form  $Q = PP_n + R$  where  $\deg P, \deg R \leq n - 1$ . The inequality  $\deg P \leq n - 1$  implies the  $\rho$ -orthogonality of  $P$  and  $P_n$ :

$$\int_a^b PP_n \rho = 0.$$

On the other hand,  $\deg R \leq n - 1$  yields that  $R$  is equal to its Lagrange-interpolation polynomial:

$$R = \sum_{k=1}^n R(\xi_k)L_k.$$

Therefore, considering the definition of the coefficients  $c_1, \dots, c_n$  in formula (1.5), we obtain that

$$\begin{aligned} \int_a^b Q\rho &= \int_a^b PP_n\rho + \int_a^b R\rho = \sum_{k=1}^n R(\xi_k) \int_a^b L_k\rho \\ &= \sum_{k=1}^n c_k R(\xi_k) = \sum_{k=1}^n c_k (P(\xi_k)P_n(\xi_k) + R(\xi_k)) = \sum_{k=1}^n c_k Q(\xi_k). \end{aligned}$$

That is, the quadrature formula (1.1) is exact for polynomials of degree at most  $2n - 1$ .

Conversely, assume that (1.1) is exact for polynomials of degree at most  $2n - 1$ . Define the polynomial  $Q$  by the formula  $Q(x) := (x - \xi_1) \cdots (x - \xi_n)$  and let  $P$

be a polynomial of degree at most  $n - 1$ . Then,  $\deg PQ \leq 2n - 1$  thus

$$\int_a^b PQ\rho = c_1P(\xi_1)Q(\xi_1) + \cdots + c_nP(\xi_n)Q(\xi_n) = 0.$$

Therefore  $Q$  is  $\rho$ -orthogonal to  $P$ . The uniqueness of  $P_n$  implies that  $P_n = a_nQ$ , and  $\xi_1, \dots, \xi_n$  are the zeros of  $P_n$ . Furthermore, (1.1) is exact if we substitute  $f := L_k$  and  $f := L_k^2$ , respectively. The first substitution gives (1.5), while the second one shows the nonnegativity of  $c_k$ . For further details, consult the book [Szeg39, p. 44].  $\square$

**THEOREM 1.2.** *Let  $P_n$  be the  $n^{\text{th}}$  degree member of the orthogonal polynomial system on  $[a, b]$  with respect to the weight function  $\rho_a(x) := (x - a)\rho(x)$ . Then (1.2) is exact for polynomials  $f$  of degree at most  $2n$  if and only if  $\xi_1, \dots, \xi_n$  are the zeros of  $P_n$ , and*

$$(1.6) \quad c_0 = \frac{1}{P_n^2(a)} \int_a^b P_n^2(x)\rho(x)dx,$$

$$(1.7) \quad c_k = \frac{1}{\xi_k - a} \int_a^b \frac{(x - a)P_n(x)}{(x - \xi_k)P_n'(\xi_k)}\rho(x)dx.$$

Furthermore,  $\xi_1, \dots, \xi_n$  are pairwise distinct elements of  $]a, b[$ , and  $c_k \geq 0$  for all  $k = 0, \dots, n$ .

**PROOF.** Assume that the quadrature formula (1.2) is exact for polynomials of degree at most  $2n$ . If  $P$  is a polynomial of degree at most  $2n - 1$ , then

$$\int_a^b P\rho_a = \int_a^b (x - a)P(x)\rho(x)dx = c_1(\xi_1 - a)P(\xi_1) + \cdots + c_n(\xi_n - a)P(\xi_n).$$

Applying Theorem 1.1 to the weight function  $\rho_a$  and the coefficients

$$c_{a;k} := c_k(\xi_k - a),$$

we get that  $\xi_1, \dots, \xi_n$  are the zeros of  $P_n$  and, for all  $k = 1, \dots, n$ , the coefficients  $c_{a;k}$  can be computed using formula (1.5). Therefore,

$$c_k(\xi_k - a) = \int_a^b \frac{P_n(x)}{(x - \xi_k)P_n'(\xi_k)}\rho_a(x)dx = \int_a^b \frac{(x - a)P_n(x)}{(x - \xi_k)P_n'(\xi_k)}\rho(x)dx.$$

Substituting  $f := P_n^2$  into (1.1), we obtain that

$$c_0 = \frac{1}{P_n^2(a)} \int_a^b P_n^2\rho.$$

Thus (1.6) and (1.7) are valid, and  $c_k \geq 0$  for  $k = 0, 1, \dots, n$ .

Conversely, assume that  $\xi_1, \dots, \xi_n$  are the zeros of the orthogonal polynomial  $P_n$ , and the coefficients  $c_1, \dots, c_n$  are given by the formula (1.7). Define the

coefficient  $c_0$  by  $c_0 = \int_a^b \rho - (c_1 + \cdots + c_n)$ . If  $P$  is a polynomial of degree at most  $2n$ , then there exists a polynomial  $Q$  with  $\deg Q \leq 2n - 1$  such that

$$P(x) = (x - a)Q(x) + P(a).$$

Indeed, the polynomial  $P(x) - P(a)$  vanishes at the point  $x = a$  hence it is divisible by  $(x - a)$ . Applying Theorem 1.1 again to the weight function  $\rho_a$ ,

$$\int_a^b Q \rho_a = c_{a;1}Q(\xi_1) + \cdots + c_{a;n}Q(\xi_n)$$

holds. Thus, using the definition of  $c_0$ , the representation of the polynomial  $P$  and the quadrature formula above, we have that

$$\begin{aligned} \int_a^b P(x)\rho(x)dx &= \int_a^b ((x - a)Q(x) + P(a))\rho(x)dx \\ &= \sum_{k=1}^n c_k(\xi_k - a)Q(\xi_k) + \sum_{k=0}^n P(a)c_k \\ &= c_0P(a) + \sum_{k=1}^n c_k((\xi_k - a)Q(\xi_k) + P(a)) \\ &= c_0P(a) + \sum_{k=1}^n c_kP(\xi_k), \end{aligned}$$

which yields that the quadrature formula (1.2) is exact for polynomials of degree at most  $2n$ . Therefore, substituting  $f := P_n^2$  into (1.2), we get formula (1.6).  $\square$

**THEOREM 1.3.** *Let  $P_n$  be the  $n^{\text{th}}$  degree member of the orthogonal polynomial system on  $[a, b]$  with respect to the weight function  $\rho^b(x) := (b - x)\rho(x)$ . Then (1.3) is exact for polynomials  $f$  of degree at most  $2n$  if and only if  $\xi_1, \dots, \xi_n$  are the zeros of  $P_n$ , and*

$$(1.8) \quad c_k = \frac{1}{b - \xi_k} \int_a^b \frac{(b - x)P_n(x)}{(x - \xi_k)P_n'(\xi_k)} \rho(x)dx,$$

$$(1.9) \quad c_{n+1} = \frac{1}{P_n^2(b)} \int_a^b P_n^2(x)\rho(x)dx.$$

Furthermore,  $\xi_1, \dots, \xi_n$  are pairwise distinct elements of  $]a, b[$ , and  $c_k \geq 0$  for all  $k = 1, \dots, n + 1$ .

**HINT.** Applying a similar argument to the previous one to the weight function  $\rho^b$ , we obtain the statement of the theorem.  $\square$

**THEOREM 1.4.** *Let  $P_n$  be the  $n^{\text{th}}$  degree member of the orthogonal polynomial system on  $[a, b]$  with respect to the weight function  $\rho_a^b$ . Then (1.4) is exact for polynomials  $f$  of degree at most  $2n + 1$  if and only if  $\xi_1, \dots, \xi_n$  are the zeros of  $P_n$ , and*

$$(1.10) \quad c_0 = \frac{1}{(b-a)P_n^2(a)} \int_a^b (b-x)P_n^2(x)\rho(x)dx,$$

$$(1.11) \quad c_k = \frac{1}{(b-\xi_k)(\xi_k-a)} \int_a^b \frac{(b-x)(x-a)P_n(x)}{(x-\xi_k)P_n'(\xi_k)} \rho(x)dx,$$

$$(1.12) \quad c_{n+1} = \frac{1}{(b-a)P_n^2(b)} \int_a^b (x-a)P_n^2(x)\rho(x)dx.$$

Furthermore,  $\xi_1, \dots, \xi_n$  are pairwise distinct elements of  $]a, b[$ , and  $c_k \geq 0$  for all  $k = 0, \dots, n + 1$ .

**PROOF.** Assume that the quadrature formula (1.4) is exact for polynomials of degree at most  $2n + 1$ . If  $P$  is a polynomial of degree at most  $2n - 1$ , then

$$\begin{aligned} \int_a^b P \rho_a^b &= \int_a^b (b-x)(x-a)P(x)\rho(x)dx \\ &= c_1(b-\xi_1)(\xi_1-a)P(\xi_1) + \dots + c_n(b-\xi_n)(\xi_n-a)P(\xi_n). \end{aligned}$$

Applying Theorem 1.1 to the weight function  $\rho_a^b$  and the coefficients

$$c_{a,b;k} := c_k(b-\xi_k)(\xi_k-a),$$

we get that  $\xi_1, \dots, \xi_n$  are the zeros of  $P_n$  and, for all  $k = 1, \dots, n$ , the coefficients  $c_{a,b;k}$  can be computed using formula (1.5). Therefore,

$$\begin{aligned} c_k(b-\xi_k)(\xi_k-a) &= \int_a^b \frac{P_n(x)}{(x-\xi_k)P_n'(\xi_k)} \rho_a^b(x)dx \\ &= \int_a^b \frac{(b-x)(x-a)P_n(x)}{(x-\xi_k)P_n'(\xi_k)} \rho(x)dx. \end{aligned}$$

Substituting  $f := (b-x)P_n^2(x)$  and  $f := (x-a)P_n^2(x)$  into (1.1), we obtain that

$$\begin{aligned} c_0 &= \frac{1}{(b-a)P_n^2(a)} \int_a^b (b-x)P_n^2(x)\rho(x)dx, \\ c_{n+1} &= \frac{1}{(b-a)P_n^2(b)} \int_a^b (x-a)P_n^2(x)\rho(x)dx. \end{aligned}$$

Thus (1.10), (1.11) and (1.12) are valid, furthermore,  $c_k \geq 0$  for  $k = 0, \dots, n + 1$ .

Conversely, assume that  $\xi_1, \dots, \xi_n$  are the zeros of  $P_n$ , and the coefficients  $c_1, \dots, c_n$  are given by the formula (1.11). Define the coefficients  $c_0$  and  $c_{n+1}$  by

the equations

$$\begin{aligned}\int_a^b (b-x)\rho(x)dx &= c_0(b-a) + \sum_{k=1}^n c_k(b-\xi_k) \\ \int_a^b (x-a)\rho(x)dx &= \sum_{k=1}^n c_k(\xi_k-a) + c_{n+1}(b-a).\end{aligned}$$

If  $P$  is a polynomial of degree at most  $2n+1$ , then there exists a polynomial  $Q$  with  $\deg Q \leq 2n-1$  such that

$$(b-a)P(x) = (b-x)(x-a)Q(x) + (x-a)P(b) + (b-x)P(a).$$

Indeed, the polynomial  $(b-a)P(x) - (x-a)P(b) - (b-x)P(a)$  is divisible by  $(b-x)(x-a)$  since  $x=a$  and  $x=b$  are its roots. Applying Theorem 1.1 again,

$$\int_a^b Q\rho_a^b = c_{a,b;1}Q(\xi_1) + \cdots + c_{a,b;n}Q(\xi_n)$$

holds. Thus, using the definition of  $c_0$  and  $c_{n+1}$ , the representation of the polynomial  $P$  and the quadrature formula above, we have that

$$\begin{aligned}(b-a) \int_a^b P(x)\rho(x)dx &= \\ &= \int_a^b ((b-x)(x-a)Q(x) + (x-a)P(b) + (b-x)P(a))\rho(x)dx \\ &= \sum_{k=1}^n c_k(b-\xi_k)(\xi_k-a)Q(\xi_k) \\ &\quad + P(b) \int_a^b (x-a)\rho(x)dx + P(a) \int_a^b (b-x)\rho(x)dx \\ &= \sum_{k=1}^n c_k(b-\xi_k)(\xi_k-a)Q(\xi_k) \\ &\quad + c_0(b-a)P(a) + \sum_{k=1}^n c_k(b-\xi_k)P(a) \\ &\quad + \sum_{k=1}^n c_k(\xi_k-a)P(b) + c_{n+1}(b-a)P(b) \\ &= \sum_{k=1}^n c_k((b-\xi_k)(\xi_k-a)Q(\xi_k) + (\xi_k-a)P(b) + (b-\xi_k)P(a)) \\ &\quad + c_0(b-a)P(a) + c_{n+1}(b-a)P(b) \\ &= c_0(b-a)P(a) + \sum_{k=1}^n c_k(b-a)P(\xi_k) + c_{n+1}(b-a)P(b),\end{aligned}$$

which yields that the quadrature formula (1.4) is exact for polynomials of degree at most  $2n + 1$ . Therefore, substituting  $f := (b - x)P_n^2(x)$  and  $f := (x - a)P_n^2(x)$  into (1.4), formulae (1.10) and (1.12) follow.  $\square$

Let  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable function,  $x_1, \dots, x_n$  be pairwise distinct elements of  $[a, b]$ , and  $1 \leq r \leq n$  be a fixed integer. Denote the Hermite interpolation polynomial by  $H$  that satisfies the following conditions:

$$\begin{aligned} H(x_k) &= f(x_k) \quad (k = 1, \dots, n) \\ H'(x_k) &= f'(x_k) \quad (k = 1, \dots, r). \end{aligned}$$

We recall that  $\deg H = n + r - 1$ . From a well known result, (see [HH94, Sec. 5.3, pp. 230-231]), for all  $x \in [a, b]$  there exists  $\theta$  such that

$$(1.13) \quad f(x) - H(x) = \frac{\omega_n(x)\omega_r(x)}{(n+r)!} f^{(n+r)}(\theta),$$

where

$$\omega_k(x) = (x - x_1) \cdots (x - x_k).$$

## 1.2. An approximation theorem

It is well known that there exists a function  $\varphi$  which possesses the following properties:

- (i)  $\varphi : \mathbb{R} \rightarrow \mathbb{R}_+$  is  $\mathcal{C}^\infty$ , i. e., it is infinitely many times differentiable;
- (ii)  $\text{supp } \varphi \subset [-1, 1]$ ;
- (iii)  $\int_{\mathbb{R}} \varphi = 1$ .

Using  $\varphi$ , one can define the function  $\varphi_\varepsilon$  for all  $\varepsilon > 0$  by the formula

$$\varphi_\varepsilon(x) = \frac{1}{\varepsilon} \varphi\left(\frac{x}{\varepsilon}\right) \quad (x \in \mathbb{R}).$$

Then, as it can easily be checked,  $\varphi_\varepsilon$  satisfies the following conditions:

- (i')  $\varphi_\varepsilon : \mathbb{R} \rightarrow \mathbb{R}_+$  is  $\mathcal{C}^\infty$ ;
- (ii')  $\text{supp } \varphi_\varepsilon \subset [-\varepsilon, \varepsilon]$ ;
- (iii')  $\int_{\mathbb{R}} \varphi_\varepsilon = 1$ .

Let  $I \subset \mathbb{R}$  be a nonempty open interval,  $f : I \rightarrow \mathbb{R}$  be a continuous function, and choose  $\varepsilon > 0$ . Denote the convolution of  $f$  and  $\varphi_\varepsilon$  by  $f_\varepsilon$ , that is

$$f_\varepsilon(x) := \int_{\mathbb{R}} \bar{f}(y) \varphi_\varepsilon(x - y) dy \quad (x \in \mathbb{R})$$

where  $\bar{f}(y) = f(y)$  if  $y \in I$ , otherwise  $\bar{f}(y) = 0$ . Let us recall, that  $f_\varepsilon \rightarrow f$  uniformly as  $\varepsilon \rightarrow 0$  on each compact subinterval of  $I$ , and  $f_\varepsilon$  is infinitely many times differentiable on  $\mathbb{R}$ . These important results can be found for example in [Zei86, p. 549].

**THEOREM 1.5.** *Let  $I \subset \mathbb{R}$  be an open interval,  $f : I \rightarrow \mathbb{R}$  be a polynomially  $n$ -convex continuous function. Then, for all compact subintervals  $[a, b] \subset I$ , there exists a sequence of polynomially  $n$ -convex and  $\mathcal{C}^\infty$  functions  $(f_k)$  which converges uniformly to  $f$  on  $[a, b]$ .*

**PROOF.** Choose  $a, b \in I$  and  $\varepsilon_0 > 0$  such that the inclusion  $[a - \varepsilon_0, b + \varepsilon_0] \subset I$  hold. We show that the function  $\tau_\varepsilon f : [a, b] \rightarrow \mathbb{R}$  defined by the formula

$$\tau_\varepsilon f(x) := f(x - \varepsilon)$$

is polynomially  $n$ -convex on  $[a, b]$  for  $0 < \varepsilon < \varepsilon_0$ . Let  $a \leq x_0 < \dots < x_n \leq b$  and  $k \leq n - 1$  be fixed. By induction, we are going to verify the identity

$$(1.14) \quad \begin{vmatrix} \tau_\varepsilon f(x_0) & \cdots & \tau_\varepsilon f(x_n) \\ 1 & \cdots & 1 \\ x_0 & \cdots & x_n \\ \vdots & \ddots & \vdots \\ x_0^{k-1} & \cdots & x_n^{k-1} \\ x_0^k & \cdots & x_n^k \\ \vdots & \ddots & \vdots \\ x_0^{n-1} & \cdots & x_n^{n-1} \end{vmatrix} = \begin{vmatrix} \tau_\varepsilon f(x_0) & \cdots & \tau_\varepsilon f(x_n) \\ 1 & \cdots & 1 \\ x_0 - \varepsilon & \cdots & x_n - \varepsilon \\ \vdots & \ddots & \vdots \\ (x_0 - \varepsilon)^{k-1} & \cdots & (x_n - \varepsilon)^{k-1} \\ x_0^k & \cdots & x_n^k \\ \vdots & \ddots & \vdots \\ x_0^{n-1} & \cdots & x_n^{n-1} \end{vmatrix}.$$

If  $k = 1$ , then this equation obviously holds. Assume, for a fixed positive integer  $k \leq n - 2$ , that (1.14) remains true. The binomial theorem implies the identity

$$x^k = \binom{k}{0} \varepsilon^k + \binom{k}{1} \varepsilon^{k-1} (x - \varepsilon) + \cdots + \binom{k}{k} (x - \varepsilon)^k.$$

That is,  $(x - \varepsilon)^k$  is the linear combination of the elements  $1, x - \varepsilon, \dots, (x - \varepsilon)^k$  and  $x^k$ . Therefore, adding the adequate linear combination of the  $2^{nd}, \dots, (k + 1)^{st}$  rows to the  $(k + 2)^{nd}$  row, we arrive at the equation

$$\begin{vmatrix} \tau_\varepsilon f(x_0) & \cdots & \tau_\varepsilon f(x_n) \\ 1 & \cdots & 1 \\ x_0 - \varepsilon & \cdots & x_n - \varepsilon \\ \vdots & \ddots & \vdots \\ (x_0 - \varepsilon)^{k-1} & \cdots & (x_n - \varepsilon)^{k-1} \\ x_0^k & \cdots & x_n^k \\ x_0^{k+1} & \cdots & x_n^{k+1} \\ \vdots & \ddots & \vdots \\ x_0^{n-1} & \cdots & x_n^{n-1} \end{vmatrix} = \begin{vmatrix} \tau_\varepsilon f(x_0) & \cdots & \tau_\varepsilon f(x_n) \\ 1 & \cdots & 1 \\ x_0 - \varepsilon & \cdots & x_n - \varepsilon \\ \vdots & \ddots & \vdots \\ (x_0 - \varepsilon)^{k-1} & \cdots & (x_n - \varepsilon)^{k-1} \\ (x_0 - \varepsilon)^k & \cdots & (x_n - \varepsilon)^k \\ x_0^{k+1} & \cdots & x_n^{k+1} \\ \vdots & \ddots & \vdots \\ x_0^{n-1} & \cdots & x_n^{n-1} \end{vmatrix}.$$

Hence formula (1.14) holds for all fixed positive  $k$  whenever  $1 \leq k \leq n - 1$ . The particular case  $k = n - 1$  gives the polynomial  $n$ -convexity of  $\tau_\varepsilon f$ . Applying

integral transformation and the previous result, we get that

$$\begin{aligned}
& (-1)^n \begin{vmatrix} f_\varepsilon(x_0) & \cdots & f_\varepsilon(x_n) \\ 1 & \cdots & 1 \\ x_0 & \cdots & x_n \\ \vdots & \ddots & \vdots \\ x_0^{n-1} & \cdots & x_n^{n-1} \end{vmatrix} = \\
& = \int_{\mathbb{R}} (-1)^n \begin{vmatrix} \bar{f}(t)\varphi_\varepsilon(x_0 - t) & \cdots & \bar{f}(t)\varphi_\varepsilon(x_n - t) \\ 1 & \cdots & 1 \\ x_0 & \cdots & x_n \\ \vdots & \ddots & \vdots \\ x_0^{n-1} & \cdots & x_n^{n-1} \end{vmatrix} dt \\
& = \int_{\mathbb{R}} (-1)^n \begin{vmatrix} \bar{f}(x_0 - s) & \cdots & \bar{f}(x_n - s) \\ 1 & \cdots & 1 \\ x_0 & \cdots & x_n \\ \vdots & \ddots & \vdots \\ x_0^{n-1} & \cdots & x_n^{n-1} \end{vmatrix} \varphi_\varepsilon(s) ds \\
& = \int_{\mathbb{R}} (-1)^n \begin{vmatrix} \tau_s f(x_0) & \cdots & \tau_s f(x_n) \\ 1 & \cdots & 1 \\ x_0 & \cdots & x_n \\ \vdots & \ddots & \vdots \\ x_0^{n-1} & \cdots & x_n^{n-1} \end{vmatrix} \varphi_\varepsilon(s) ds \geq 0,
\end{aligned}$$

which shows the polynomial  $n$ -convexity of  $f_\varepsilon$  on  $[a, b]$  for  $0 < \varepsilon < \varepsilon_0$ .

To complete the proof, choose a positive integer  $n_0$  such that the relation  $\frac{1}{n_0} < \varepsilon_0$  hold. If we define  $\varepsilon_k$  and  $f_k$  by  $\varepsilon_k := \frac{1}{n_0+k}$  and  $f_k := f_{\varepsilon_k}$  for  $k \in \mathbb{N}$ , then  $0 < \varepsilon_k < \varepsilon_0$  thus  $(f_k)_{k=1}^\infty$  satisfies the requirements of the theorem.  $\square$

### 1.3. Hermite–Hadamard-type inequalities

In the sequel, we shall need two additional auxiliary results. The first one investigates the convergence properties of the roots of orthogonal polynomials.

**LEMMA 1.6.** *Let  $\rho$  be a weight function on  $[a, b]$  furthermore  $(a_j)$  be strictly monotone decreasing,  $(b_j)$  be strictly monotone increasing sequences such that  $a_j \rightarrow a$ ,  $b_j \rightarrow b$  and  $a_1 < b_1$ . Denote the roots of  $P_{m;j}$  by  $\xi_{1;j}, \dots, \xi_{m;j}$  where  $P_{m;j}$  is the  $m^{\text{th}}$  degree member of the  $\rho|_{[a_j, b_j]}$ -orthogonal polynomial system on  $[a_j, b_j]$ , and denote the roots of  $P_m$  by  $\xi_1, \dots, \xi_m$  where  $P_m$  is the  $m^{\text{th}}$  degree member of the  $\rho$ -orthogonal polynomial system on  $[a, b]$ . Then,*

$$\lim_{j \rightarrow \infty} \xi_{k;j} = \xi_k \quad (k = 1, \dots, n).$$

PROOF. Observe first that the mapping  $(a, b) \mapsto \mu_{k;[a,b]}$  is continuous, therefore  $\mu_{k;[a_j,b_j]} \rightarrow \mu_{k;[a,b]}$  hence  $P_{m;j} \rightarrow P_m$  pointwise according to the representation of orthogonal polynomials. Take  $\varepsilon > 0$  such that

$$] \xi_k - \varepsilon, \xi_k + \varepsilon [ \subset ] a, b [$$

$$] \xi_k - \varepsilon, \xi_k + \varepsilon [ \cap ] \xi_l - \varepsilon, \xi_l + \varepsilon [ = \emptyset \quad (k \neq l, k, l \in \{1, \dots, m\}).$$

The polynomial  $P_m$  changes its sign on  $] \xi_k - \varepsilon, \xi_k + \varepsilon [$  since it is of degree  $m$  and it has  $m$  pairwise distinct roots; therefore, due to the pointwise convergence,  $P_{m;j}$  also changes its sign on the same interval up to an index. That is, for sufficiently large  $j$ ,  $\xi_{k;j} \in ] \xi_k - \varepsilon, \xi_k + \varepsilon [$ .  $\square$

The other auxiliary result investigates the one-sided limits of polynomially  $n$ -convex functions at the endpoints of the domain. Let us note that its first assertion involves, in fact, two cases according to the parity of the convexity.

LEMMA 1.7. *Let  $f : [a, b] \rightarrow \mathbb{R}$  be a polynomially  $n$ -convex function. Then,*

- (i)  $(-1)^n f(a) \geq \limsup_{t \rightarrow a+0} (-1)^n f(t)$ ;
- (ii)  $f(b) \geq \limsup_{t \rightarrow b-0} f(t)$ .

PROOF. It suffices to restrict the investigations to the even case of assertion (i) only since the proofs of the other ones are completely the same. For the sake of brevity, we shall use the notation  $f_+(a) := \limsup_{t \rightarrow a+0} f(t)$ . Take the elements  $x_0 := a < x_1 := t < \dots < x_n$  of  $[a, b]$ . Then, the (even order) polynomial convexity of  $f$  implies

$$\begin{vmatrix} f(a) & f(t) & f(x_2) & \dots & f(x_n) \\ 1 & 1 & 1 & \dots & 1 \\ a & t & x_2 & \dots & x_n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a^{n-1} & t^{n-1} & x_2^{n-1} & \dots & x_n^{n-1} \end{vmatrix} \geq 0.$$

Therefore, taking the limsup as  $t \rightarrow a + 0$ , we obtain that

$$\begin{vmatrix} f(a) & f_+(a) & f(x_2) & \dots & f(x_n) \\ 1 & 1 & 1 & \dots & 1 \\ a & a & x_2 & \dots & x_n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a^{n-1} & a^{n-1} & x_2^{n-1} & \dots & x_n^{n-1} \end{vmatrix} \geq 0.$$

The adjoint determinants of the elements  $f(x_2), \dots, f(x_n)$  in the first row are equal to zero since their first and second columns coincide; on the other hand,  $f(a)$  and  $f_+(a)$  have the same (positive) Vandermonde-type adjoint determinant. Hence, applying the expansion theorem on the first row, we obtain the desired inequality

$$f(a) - f_+(a) \geq 0.$$

$\square$

The main results concern the cases of odd and even order polynomial convexity separately in the subsequent two theorems.

**THEOREM 1.8.** *Let  $\rho : [a, b] \rightarrow \mathbb{R}$  be a positive integrable function. Denote the roots of  $P_m$  by  $\xi_1, \dots, \xi_m$  where  $P_m$  is the  $m^{\text{th}}$  degree member of the orthogonal polynomial system on  $[a, b]$  with respect to the weight function  $(x - a)\rho(x)$ , and denote the roots of  $Q_m$  by  $\eta_1, \dots, \eta_m$  where  $Q_m$  is the  $m^{\text{th}}$  degree member of the orthogonal polynomial system on  $[a, b]$  with respect to the weight function  $(b - x)\rho(x)$ . Define the coefficients  $\alpha_0, \dots, \alpha_m$  and  $\beta_1, \dots, \beta_{m+1}$  by the formulae*

$$\begin{aligned}\alpha_0 &:= \frac{1}{P_m^2(a)} \int_a^b P_m^2(x)\rho(x)dx, \\ \alpha_k &:= \frac{1}{\xi_k - a} \int_a^b \frac{(x - a)P_m(x)}{(x - \xi_k)P'_m(\xi_k)} \rho(x)dx\end{aligned}$$

and

$$\begin{aligned}\beta_k &:= \frac{1}{b - \eta_k} \int_a^b \frac{(b - x)Q_m(x)}{(x - \eta_k)Q'_m(\eta_k)} \rho(x)dx, \\ \beta_{m+1} &:= \frac{1}{Q_m^2(b)} \int_a^b Q_m^2(x)\rho(x)dx.\end{aligned}$$

If a function  $f : [a, b] \rightarrow \mathbb{R}$  is polynomially  $(2m + 1)$ -convex, then it satisfies the following Hermite–Hadamard-type inequality

$$\alpha_0 f(a) + \sum_{k=1}^m \alpha_k f(\xi_k) \leq \int_a^b f \rho \leq \sum_{k=1}^m \beta_k f(\eta_k) + \beta_{m+1} f(b).$$

**PROOF.** First assume that  $f$  is  $(2m + 1)$  times differentiable. Then, according to Theorem A,  $f^{(2m+1)} \geq 0$  on  $]a, b[$ . Let  $H$  be the Hermite interpolation polynomial determined by the conditions

$$\begin{aligned}H(a) &= f(a) \\ H(\xi_k) &= f(\xi_k) \\ H'(\xi_k) &= f'(\xi_k).\end{aligned}$$

By the remainder term (1.13) of the Hermite interpolation, if  $x$  is an arbitrary element of  $]a, b[$ , then there exists  $\theta \in ]a, b[$  such that

$$f(x) - H(x) = \frac{(x - a)(x - \xi_1)^2 \cdots (x - \xi_m)^2}{(2m + 1)!} f^{(2m+1)}(\theta).$$

That is,  $f\rho \geq H\rho$  on  $[a, b]$  due to the nonnegativity of  $f^{(2m+1)}$  and the positivity of  $\rho$ . On the other hand,  $H$  is of degree  $2m$ , therefore Theorem 1.2 yields that

$$\int_a^b f\rho \geq \int_a^b H\rho = \alpha_0 H(a) + \sum_{k=1}^m \alpha_k H(\xi_k) = \alpha_0 f(a) + \sum_{k=1}^m \alpha_k f(\xi_k).$$

For the general case, let  $f$  be an arbitrary polynomially  $(2m + 1)$ -convex function. Without loss of generality we may assume that  $m \geq 1$ ; in this case,  $f$  is continuous (see Theorem B). Let  $(a_j)$  and  $(b_j)$  be sequences fulfilling the requirements of Lemma 1.6. According to Theorem 1.5, there exists a sequence of  $\mathcal{C}^\infty$ , polynomially  $(2m + 1)$ -convex functions  $(f_{i;j})$  such that  $f_{i;j} \rightarrow f$  uniformly on  $[a_j, b_j]$  as  $i \rightarrow \infty$ . Denote the roots of  $P_{m;j}$  by  $\xi_{1;j}, \dots, \xi_{m;j}$  where  $P_{m;j}$  is the  $m^{\text{th}}$  degree member of the orthogonal polynomial system on  $[a_j, b_j]$  with respect to the weight function  $(x - a)\rho(x)$ . Define the coefficients  $\alpha_{0;j}, \dots, \alpha_{m;j}$  analogously to  $\alpha_0, \dots, \alpha_m$  with the help of  $P_{m;j}$ . Then,  $\xi_{k;j} \rightarrow \xi_k$  due to Lemma 1.6, and hence  $\alpha_{k;j} \rightarrow \alpha_k$  as  $j \rightarrow \infty$ . Applying the previous step of the proof on the smooth functions  $(f_{i;j})$ , it follows that

$$\alpha_{0;j} f_{i;j}(a_j) + \sum_{k=1}^m \alpha_{k;j} f_{i;j}(\xi_{k;j}) \leq \int_{a_j}^{b_j} f_{i;j} \rho.$$

Taking the limits  $i \rightarrow \infty$  and then  $j \rightarrow \infty$ , we get the inequality

$$\alpha_0 \left( \liminf_{t \rightarrow a+0} f(t) \right) + \sum_{k=1}^m \alpha_k f(\xi_k) \leq \int_a^b f \rho.$$

This, together with Lemma 1.7, gives the left hand side inequality to be proved. The proof of the right hand side inequality is analogous, therefore it is omitted.  $\square$

The second main result offers Hermite–Hadamard-type inequalities for even-order polynomially convex functions. In this case, the symmetrical structure disappears: the lower estimation involves none of the endpoints, while the upper estimation involves both of them.

**THEOREM 1.9.** *Let  $\rho : [a, b] \rightarrow \mathbb{R}$  be a positive integrable function. Denote the roots of  $P_m$  by  $\xi_1, \dots, \xi_m$  where  $P_m$  is the  $m^{\text{th}}$  degree member of the orthogonal polynomial system on  $[a, b]$  with respect to the weight function  $\rho(x)$ , and denote the roots of  $Q_{m-1}$  by  $\eta_1, \dots, \eta_{m-1}$  where  $Q_{m-1}$  is the  $(m - 1)^{\text{st}}$  degree member of the orthogonal polynomial system on  $[a, b]$  with respect to the weight function  $(b - x)(x - a)\rho(x)$ . Define the coefficients  $\alpha_1, \dots, \alpha_m$  and  $\beta_0, \dots, \beta_{m+1}$  by the formulae*

$$\alpha_k := \int_a^b \frac{P_m(x)}{(x - \xi_k)P'_m(\xi_k)} \rho(x) dx$$

and

$$\begin{aligned} \beta_0 &= \frac{1}{(b - a)Q_{m-1}^2(a)} \int_a^b (b - x)Q_{m-1}^2(x)\rho(x) dx, \\ \beta_k &= \frac{1}{(b - \eta_k)(\xi_k - a)} \int_a^b \frac{(b - x)(x - a)Q_{m-1}(x)}{(x - \eta_k)Q'_{m-1}(\eta_k)} \rho(x) dx, \\ \beta_{m+1} &= \frac{1}{(b - a)Q_{m-1}^2(b)} \int_a^b (x - a)Q_{m-1}^2(x)\rho(x) dx. \end{aligned}$$

If a function  $f : [a, b] \rightarrow \mathbb{R}$  is polynomially  $(2m)$ -convex, then it satisfies the following Hermite–Hadamard-type inequality

$$\sum_{k=1}^m \alpha_k f(\xi_k) \leq \int_a^b f \rho \leq \beta_0 f(a) + \sum_{k=1}^{m-1} \beta_k f(\eta_k) + \beta_m f(b).$$

PROOF. First assume that  $f$  is  $n = 2m$  times differentiable. Then  $f^{(2m)} \geq 0$  on  $]a, b[$  according to Theorem B. Consider the Hermite interpolation polynomial  $H$  that interpolates the function  $f$  in the roots of  $P_m$  in the following manner:

$$\begin{aligned} H(\xi_k) &= f(\xi_k) \\ H'(\xi_k) &= f'(\xi_k). \end{aligned}$$

By the remainder term (1.13) of the Hermite interpolation, if  $x$  is an arbitrary element of  $]a, b[$ , then there exists  $\theta \in ]a, b[$  such that

$$f(x) - H(x) = \frac{(x - \xi_1)^2 \cdots (x - \xi_m)^2}{(2m)!} f^{(2m)}(\theta).$$

Hence  $f \rho \geq H \rho$  on  $[a, b]$  due to the nonnegativity of  $f^{(2m)}$  and the positivity of  $\rho$ . On the other hand,  $H$  is of degree  $2m - 1$ , therefore Theorem 1.1 yields the left hand side of the inequality to be proved:

$$\int_a^b f \rho \geq \int_a^b H \rho = \sum_{k=1}^m \alpha_k H(\xi_k) = \sum_{k=1}^m \alpha_k f(\xi_k).$$

Now consider the Hermite interpolation polynomial  $H$  that interpolates the function  $f$  in the roots of  $Q_{m-1}$  and in the endpoints of the domain in the following way:

$$\begin{aligned} H(a) &= f(a) \\ H(\eta_k) &= f(\eta_k) \\ H'(\eta_k) &= f'(\eta_k) \\ H(b) &= f(b). \end{aligned}$$

By the remainder term (1.13) of the Hermite interpolation, if  $x$  is an arbitrary element of  $]a, b[$ , then there exists  $\theta \in ]a, b[$  such that

$$f(x) - H(x) = \frac{(x - a)(x - b)(x - \eta_1)^2 \cdots (x - \eta_{m-1})^2}{(2m)!} f^{(2m)}(\theta).$$

The factors of the right hand side are nonnegative except for the factor  $(x - b)$  which is negative hence  $f \rho \leq H \rho$ . On the other hand,  $H$  is of degree  $2m - 1$ ,

therefore Theorem 1.4 yields the right hand side inequality to be proved:

$$\begin{aligned} \int_a^b f \rho \leq \int_a^b H \rho &= \beta_0 H(a) + \sum_{k=1}^{m-1} \beta_k H(\eta_k) + \beta_m H(b) \\ &= \beta_0 f(a) + \sum_{k=1}^{m-1} \beta_k f(\eta_k) + \beta_m f(b). \end{aligned}$$

From this point, an analogous argument to the corresponding part of the previous proof gives the statement of the theorem without any differentiability assumptions on the function  $f$ .  $\square$

Specializing the weight function  $\rho \equiv 1$ , the roots of the inequalities can be obtained as convex combinations of the endpoints of the domain. The coefficients of the convex combinations are the roots of certain orthogonal polynomials on  $[0, 1]$  in both cases. Observe that interchanging the role of the endpoints in any side of the inequality concerning the odd order case, we obtain the other side of the inequality.

**THEOREM 1.10.** *Let, for  $m \geq 0$ , the polynomial  $P_m$  be defined by the formula*

$$P_m(x) := \begin{vmatrix} 1 & \frac{1}{2} & \cdots & \frac{1}{m+1} \\ x & \frac{1}{3} & \cdots & \frac{1}{m+2} \\ \vdots & \vdots & \ddots & \vdots \\ x^m & \frac{1}{m+2} & \cdots & \frac{1}{2m+1} \end{vmatrix}.$$

*Then,  $P_m$  has  $m$  pairwise distinct roots  $\lambda_1, \dots, \lambda_m$  in  $]0, 1[$ . Define the coefficients  $\alpha_0, \dots, \alpha_m$  by*

$$\begin{aligned} \alpha_0 &:= \frac{1}{P_m^2(0)} \int_0^1 P_m^2(x) dx, \\ \alpha_k &:= \frac{1}{\lambda_k} \int_0^1 \frac{x P_m(x)}{(x - \lambda_k) P_m'(\lambda_k)} dx. \end{aligned}$$

*If a function  $f : [a, b] \rightarrow \mathbb{R}$  is polynomially  $(2m + 1)$ -convex, then it satisfies the following Hermite–Hadamard-type inequality*

$$\begin{aligned} \alpha_0 f(a) + \sum_{k=1}^m \alpha_k f((1 - \lambda_k)a + \lambda_k b) &\leq \frac{1}{b - a} \int_a^b f(x) dx \\ &\leq \sum_{k=1}^m \alpha_k f(\lambda_k a + (1 - \lambda_k)b) + \alpha_0 f(b). \end{aligned}$$

**PROOF.** Apply Theorem 1.8 in the particular setting when  $a := 0$ ,  $b := 1$  and the weight function is  $\rho \equiv 1$ . Then, as simple calculations show,  $P_m$  is exactly the  $m^{\text{th}}$  degree member of the orthogonal polynomial system on  $[0, 1]$  with respect to the weight function  $\rho(x) = x$  (see the beginning of this chapter). Therefore,  $P_m$

has  $m$  pairwise distinct roots  $0 < \lambda_1 < \dots < \lambda_m < 1$ , indeed. Moreover, the coefficients  $\alpha_0, \dots, \alpha_m$  have the form above. Define the function  $F : [0, 1] \rightarrow \mathbb{R}$  by the formula

$$F(t) := f((1-t)a + tb).$$

It is easy to check that  $F$  is polynomially  $(2m+1)$ -convex on the interval  $[0, 1]$ . Hence, applying Theorem 1.8 and the previous observations, it follows that

$$\begin{aligned} \int_0^1 F(t) dt &\geq \alpha_0 F(0) + \sum_{k=1}^m \alpha_k F(\lambda_k) \\ &= \alpha_0 f(a) + \sum_{k=1}^m \alpha_k f((1-\lambda_k)a + \lambda_k b). \end{aligned}$$

On the other hand, to complete the proof of the left hand side inequality, observe that

$$\frac{1}{b-a} \int_a^b f(x) dx = \int_0^1 F(t) dt.$$

For verifying the right hand side one, define the function  $\varphi : [a, b] \rightarrow \mathbb{R}$  by the formula

$$\varphi(x) := -f(a+b-x).$$

Then,  $\varphi$  is polynomially  $(2m+1)$ -convex on  $[a, b]$ . The previous inequality applied on  $\varphi$  gives the upper estimation of the Hermite–Hadamard-type inequality for  $f$ .  $\square$

**THEOREM 1.11.** *Let, for  $m \geq 1$ , the polynomials  $P_m$  and  $Q_{m-1}$  be defined by the formulae*

$$P_m(x) := \begin{vmatrix} 1 & 1 & \dots & \frac{1}{m} \\ x & \frac{1}{2} & \dots & \frac{1}{m+1} \\ \vdots & \vdots & \ddots & \vdots \\ x^m & \frac{1}{m+1} & \dots & \frac{1}{2m} \end{vmatrix},$$

$$Q_{m-1}(x) := \begin{vmatrix} 1 & & \frac{1}{2 \cdot 3} & \dots & \frac{1}{m(m+1)} \\ x & & \frac{1}{3 \cdot 4} & \dots & \frac{1}{(m+1)(m+2)} \\ \vdots & & \vdots & \ddots & \vdots \\ x^{m-1} & & \frac{1}{(m+1)(m+2)} & \dots & \frac{1}{(2m-1)2m} \end{vmatrix}.$$

*Then,  $P_m$  has  $m$  pairwise distinct roots  $\lambda_1, \dots, \lambda_m$  in  $]0, 1[$  and  $Q_{m-1}$  has  $m-1$  pairwise distinct roots  $\mu_1, \dots, \mu_{m-1}$  in  $]0, 1[$ , respectively. Define the coefficients  $\alpha_1, \dots, \alpha_m$  and  $\beta_0, \dots, \beta_m$  by*

$$\alpha_k := \int_0^1 \frac{P_m(x)}{(x-\lambda_k)P'_m(\lambda_k)} dx$$

and

$$\begin{aligned}\beta_0 &:= \frac{1}{Q_{m-1}^2(0)} \int_0^1 (1-x)Q_{m-1}^2(x)dx, \\ \beta_k &:= \frac{1}{(1-\mu_k)\mu_k} \int_0^1 \frac{x(1-x)Q_{m-1}(x)}{(x-\mu_k)Q'_{m-1}(\mu_k)}dx, \\ \beta_m &:= \frac{1}{Q_{m-1}^2(1)} \int_0^1 xQ_{m-1}^2(x)dx.\end{aligned}$$

If a function  $f : [a, b] \rightarrow \mathbb{R}$  is polynomially  $(2m)$ -convex, then it satisfies the following Hermite–Hadamard-type inequality

$$\begin{aligned}\sum_{k=1}^m \alpha_k f((1-\lambda_k)a + \lambda_k b) &\leq \frac{1}{b-a} \int_a^b f(x)dx \\ &\leq \beta_0 f(a) + \sum_{k=1}^{m-1} \beta_k f((1-\mu_k)a + \mu_k b) + \beta_m f(b).\end{aligned}$$

PROOF. Substitute  $a := 0$ ,  $b := 1$  and  $\rho \equiv 1$  into Theorem 1.9. Then,  $P_m$  is exactly the  $m^{\text{th}}$  degree member of the orthogonal polynomial system on the interval  $[0, 1]$  with respect to the weight function  $\rho(x) = 1$ ; similarly,  $Q_{m-1}$  is the  $(m-1)^{\text{st}}$  degree member of the orthogonal polynomial system on the interval  $[0, 1]$  with respect to the weight function  $\rho(x) = (1-x)x$ . Therefore,  $Q_m$  has  $m$  pairwise distinct roots  $0 < \lambda_1 < \dots < \lambda_m < 1$  and  $Q_{m-1}$  has  $m-1$  pairwise distinct roots  $0 < \mu_1 < \dots < \mu_{m-1} < 1$ , indeed. Moreover, the coefficients  $\alpha_1, \dots, \alpha_m$  and  $\beta_0, \dots, \beta_m$  have the form above. To complete the proof, apply Theorem 1.9 on the function  $F : [0, 1] \rightarrow \mathbb{R}$  defined by the formula

$$F(t) := f((1-t)a + tb).$$

□

### 1.4. Applications

In the particular setting when  $m = 1$ , Theorem 1.10 reduces to the classical Hermite–Hadamard inequality:

COROLLARY 1.12. *If  $f : [a, b] \rightarrow \mathbb{R}$  is a polynomially 2-convex (i.e. convex) function, then the following inequalities hold*

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{f(a) + f(b)}{2}.$$

In the subsequent corollaries we present Hermite–Hadamard-type inequalities in those cases when the roots of the polynomials in Theorem 1.10 and Theorem 1.11 can explicitly be computed.

**COROLLARY 1.13.** *If  $f : [a, b] \rightarrow \mathbb{R}$  is a polynomially 3-convex function, then the following inequalities hold*

$$\frac{1}{4}f(a) + \frac{3}{4}f\left(\frac{a+2b}{3}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{3}{4}f\left(\frac{2a+b}{3}\right) + \frac{1}{4}f(b).$$

**COROLLARY 1.14.** *If  $f : [a, b] \rightarrow \mathbb{R}$  is a polynomially 4-convex function, then the following inequalities hold*

$$\begin{aligned} & \frac{1}{2}f\left(\frac{3+\sqrt{3}}{6}a + \frac{3-\sqrt{3}}{6}b\right) + \frac{1}{2}f\left(\frac{3-\sqrt{3}}{6}a + \frac{3+\sqrt{3}}{6}b\right) \\ & \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{1}{6}f(a) + \frac{2}{3}f\left(\frac{a+b}{2}\right) + \frac{1}{6}f(b). \end{aligned}$$

**COROLLARY 1.15.** *If  $f : [a, b] \rightarrow \mathbb{R}$  is a polynomially 5-convex function, then the following inequalities hold*

$$\begin{aligned} & \frac{1}{9}f(a) + \frac{16+\sqrt{6}}{36}f\left(\frac{4+\sqrt{6}}{10}a + \frac{6-\sqrt{6}}{10}b\right) \\ & + \frac{16-\sqrt{6}}{36}f\left(\frac{4-\sqrt{6}}{10}a + \frac{6+\sqrt{6}}{10}b\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \\ & \leq \frac{16-\sqrt{6}}{36}f\left(\frac{6+\sqrt{6}}{10}a + \frac{4-\sqrt{6}}{10}b\right) \\ & + \frac{16+\sqrt{6}}{36}f\left(\frac{6-\sqrt{6}}{10}a + \frac{4+\sqrt{6}}{10}b\right) + \frac{1}{9}f(b). \end{aligned}$$

In some other cases analogous statements can be formulated applying Theorem 1.11. For simplicity, instead of writing down these corollaries explicitly, we shall present a list which contains the roots of  $P_n$  (denoted by  $\lambda_k$ ), and the coefficients  $\alpha_k$  for the left hand side inequality furthermore the roots of  $Q_n$  (denoted by  $\mu_k$ ), and the coefficients  $\beta_k$  for the right hand side inequality, respectively.

**Case  $n = 6$**

The roots of  $P_3$ :

$$\frac{5-\sqrt{15}}{10}, \quad \frac{1}{2}, \quad \frac{5+\sqrt{15}}{10};$$

the corresponding coefficients:

$$\frac{5}{18}, \quad \frac{4}{9}, \quad \frac{5}{18}.$$

The roots of  $Q_2$ :

$$\frac{5 - \sqrt{5}}{10}, \quad \frac{5 + \sqrt{5}}{10};$$

the corresponding coefficients:

$$\frac{1}{12}, \quad \frac{5}{12}, \quad \frac{5}{12}, \quad \frac{1}{12}.$$

**Case  $n = 8$**

The roots of  $P_4$ :

$$\begin{aligned} \frac{1}{2} - \frac{\sqrt{525 + 70\sqrt{30}}}{70}, \quad \frac{1}{2} - \frac{\sqrt{525 - 70\sqrt{30}}}{70}, \\ \frac{1}{2} + \frac{\sqrt{525 - 70\sqrt{30}}}{70}, \quad \frac{1}{2} + \frac{\sqrt{525 + 70\sqrt{30}}}{70}; \end{aligned}$$

the corresponding coefficients:

$$\frac{1}{4} - \frac{\sqrt{30}}{72}, \quad \frac{1}{4} + \frac{\sqrt{30}}{72}, \quad \frac{1}{4} + \frac{\sqrt{30}}{72}, \quad \frac{1}{4} - \frac{\sqrt{30}}{72}.$$

The roots of  $Q_3$ :

$$\frac{1}{2} - \frac{\sqrt{21}}{14}, \quad \frac{1}{2}, \quad \frac{1}{2} + \frac{\sqrt{21}}{14};$$

the corresponding coefficients:

$$\frac{1}{20}, \quad \frac{49}{180}, \quad \frac{16}{45}, \quad \frac{49}{180}, \quad \frac{1}{20}.$$

**Case  $n = 10$**

The roots of  $P_5$ :

$$\begin{aligned} \frac{1}{2} - \frac{\sqrt{245 + 14\sqrt{70}}}{42}, \quad \frac{1}{2} - \frac{\sqrt{245 - 14\sqrt{70}}}{42}, \\ \frac{1}{2}, \quad \frac{1}{2} + \frac{\sqrt{245 - 14\sqrt{70}}}{42}, \quad \frac{1}{2} + \frac{\sqrt{245 + 14\sqrt{70}}}{42}; \end{aligned}$$

the corresponding coefficients:

$$\frac{322 - 13\sqrt{70}}{1800}, \quad \frac{322 + 13\sqrt{70}}{1800}, \quad \frac{64}{225}, \quad \frac{322 + 13\sqrt{70}}{1800}, \quad \frac{322 - 13\sqrt{70}}{1800}.$$

The roots of  $Q_4$ :

$$\begin{aligned} \frac{1}{2} - \frac{\sqrt{147 + 42\sqrt{7}}}{42}, \quad \frac{1}{2} - \frac{\sqrt{147 - 42\sqrt{7}}}{42}, \\ \frac{1}{2} + \frac{\sqrt{147 - 42\sqrt{7}}}{42}, \quad \frac{1}{2} + \frac{\sqrt{147 + 42\sqrt{7}}}{42}; \end{aligned}$$

the corresponding coefficients:

$$\frac{1}{30}, \quad \frac{14 - \sqrt{7}}{60}, \quad \frac{14 + \sqrt{7}}{60}, \quad \frac{14 + \sqrt{7}}{60}, \quad \frac{14 - \sqrt{7}}{60}, \quad \frac{1}{30}.$$

**Case  $n = 12$  (right hand side inequality)**

The roots of  $Q_5$ :

$$\frac{1}{2} - \frac{\sqrt{495 + 66\sqrt{15}}}{66}, \quad \frac{1}{2} - \frac{\sqrt{495 - 66\sqrt{15}}}{66},$$

$$\frac{1}{2}, \quad \frac{1}{2} + \frac{\sqrt{495 - 66\sqrt{15}}}{66}, \quad \frac{1}{2} + \frac{\sqrt{495 + 66\sqrt{15}}}{66};$$

the corresponding coefficients:

$$\frac{1}{42}, \quad \frac{124 - 7\sqrt{15}}{700}, \quad \frac{124 + 7\sqrt{15}}{700}, \quad \frac{128}{525},$$

$$\frac{124 + 7\sqrt{15}}{700}, \quad \frac{124 - 7\sqrt{15}}{700}, \quad \frac{1}{42}.$$

During the investigations of the higher-order cases above, we can use the symmetry of the roots of the orthogonal polynomials with respect to  $1/2$ , and therefore the calculations lead to solving linear or quadratic equations. The first case where “casus irreducibilis” appears is  $n = 7$ ; similarly, this is the reason for presenting only the right hand side inequality for polynomially 12-convex functions.

## Generalized 2-convexity

In terms of geometry, the Tchebychev property of a two dimensional system can equivalently be formulated: the linear combinations of the members of the system (shortly: *generalized lines*) are continuous furthermore any two points of the plain with distinct first coordinates can be connected by a unique generalized line. That is, generalized lines have the most important properties of affine functions. These properties turn out to be so strong that most of the classical results of standard convexity can be generalized for this setting.

First we investigate some basic properties of generalized lines of two dimensional Tchebychev systems. Then the most important tool of the chapter, a characterization theorem is proved for generalized 2-convex functions. Two consequences of this theorem, namely the existence of generalized support lines and the property of generalized chords are crucial to verify Hermite–Hadamard-type inequalities. Another result states a tight connection between standard and  $(\omega_1, \omega_2)$ -convexity, and also guarantees the integrability of  $(\omega_1, \omega_2)$ -convex functions. Some classical results of the theory of convex functions, like their representation and stability are also generalized for this setting.

### 2.1. Characterizations via generalized lines

Let us recall that  $(\omega_1, \omega_2)$  is said to be a *Tchebychev system* over an interval  $I$  if  $\omega_1, \omega_2 : I \rightarrow \mathbb{R}$  are continuous functions and, for all elements  $x < y$  of  $I$ ,

$$\begin{vmatrix} \omega_1(x) & \omega_1(y) \\ \omega_2(x) & \omega_2(y) \end{vmatrix} > 0.$$

Some concrete examples on Tchebychev systems are presented in the last section of the chapter. Given a Tchebychev system  $(\omega_1, \omega_2)$ , a function  $f : I \rightarrow \mathbb{R}$  is called *generalized convex with respect to  $(\omega_1, \omega_2)$*  or shortly: *generalized 2-convex* if, for all elements  $x < y < z$  of  $I$ , it satisfies the inequality

$$\begin{vmatrix} f(x) & f(y) & f(z) \\ \omega_1(x) & \omega_1(y) & \omega_1(z) \\ \omega_2(x) & \omega_2(y) & \omega_2(z) \end{vmatrix} \geq 0.$$

Clearly, in the standard setting this definition reduces to the notion of (ordinary) convexity. Let  $(\omega_1, \omega_2)$  be a Tchebychev system on an interval  $I$ , and denote the set of all linear combinations of the functions  $\omega_1$  and  $\omega_2$  by  $\mathcal{L}(\omega_1, \omega_2)$ . We say that

a function  $\omega : I \rightarrow \mathbb{R}$  is a *generalized line* if it belongs to the linear hull  $\mathcal{L}(\omega_1, \omega_2)$ . The properties of generalized lines play the key role in our further investigations; first we need the following simple but useful ones.

LEMMA 2.1. *Let  $(\omega_1, \omega_2)$  be a Tchebychev system over an interval  $I$ . Then, two different generalized lines of  $\mathcal{L}(\omega_1, \omega_2)$  have at most one common point; moreover, if two different generalized lines have the same value at some  $\xi \in I^\circ$ , then the difference of the lines is positive on one side of  $\xi$  while negative on the other side of  $\xi$ . In particular,  $\omega_1$  and  $\omega_2$  have at most one zero; moreover, if  $\omega_1$  (resp.,  $\omega_2$ ) vanishes at some  $\xi \in I^\circ$ , then  $\omega_1$  is positive on one side of  $\xi$  while negative on the other.*

PROOF. Due to the linear structure of  $\mathcal{L}(\omega_1, \omega_2)$ , without loss of generality we may assume that one of the lines is the constant zero line. Then, the other generalized line  $\omega$  has the representation  $\alpha\omega_1 + \beta\omega_2$ , with  $\alpha^2 + \beta^2 > 0$ .

The first assertion of the theorem is equivalent to the property that  $\omega$  has at most one zero. To show this, assume indirectly that  $\omega(\xi)$  and  $\omega(\eta)$  equal zero for  $\xi \neq \eta$ ; that is,

$$\begin{aligned}\alpha\omega_1(\xi) + \beta\omega_2(\xi) &= 0 \\ \alpha\omega_1(\eta) + \beta\omega_2(\eta) &= 0.\end{aligned}$$

By the Tchebychev property of  $(\omega_1, \omega_2)$ , the base determinant of the system is nonvanishing, therefore the system has only trivial solutions  $\alpha = 0$  and  $\beta = 0$  which contradicts the property  $\alpha^2 + \beta^2 > 0$ .

An equivalent formulation of the second assertion is the following: if  $\omega(\xi) = 0$  for some interior point  $\xi$ , then  $\omega > 0$  on one side of  $\xi$  while  $\omega < 0$  on the other. If this is not true, then, according to the previous result and Bolzano's theorem,  $\omega$  is strictly positive (or negative) on both sides of  $\xi$ . For simplicity, assume that  $\omega(t) > 0$  for  $t \neq \xi$ . Define the generalized line  $\omega^*$  by  $\omega^* := -\beta\omega_1 + \alpha\omega_2$ . Then,  $(\omega, \omega^*)$  is also a Tchebychev system: if  $x < y$  are elements of  $I$ , then

$$\begin{aligned}\begin{vmatrix} \omega(x) & \omega(\xi) \\ \omega^*(x) & \omega^*(y) \end{vmatrix} &= \begin{vmatrix} \alpha & \beta \\ -\beta & \alpha \end{vmatrix} \cdot \begin{vmatrix} \omega_1(x) & \omega_1(y) \\ \omega_2(x) & \omega_2(y) \end{vmatrix} \\ &= (\alpha^2 + \beta^2) \begin{vmatrix} \omega_1(x) & \omega_1(y) \\ \omega_2(x) & \omega_2(y) \end{vmatrix} > 0.\end{aligned}$$

Therefore, taking the elements  $x < \xi < y$  of  $I$ , we arrive at the inequalities

$$\begin{aligned}0 &< \begin{vmatrix} \omega(x) & \omega(\xi) \\ \omega^*(x) & \omega^*(\xi) \end{vmatrix} = \omega(x)\omega^*(\xi) \\ 0 &< \begin{vmatrix} \omega(\xi) & \omega(y) \\ \omega^*(\xi) & \omega^*(y) \end{vmatrix} = -\omega(y)\omega^*(\xi),\end{aligned}$$

which yields the contradiction that  $\omega^*(\xi)$  is simultaneously positive and negative.

For the last assertion, notice that  $\omega_1$ ,  $\omega_2$  and the constant zero functions are special generalized lines and apply the previous part of the theorem.  $\square$

The most important property of  $\mathcal{L}(\omega_1, \omega_2)$  guarantees the existence of a generalized line “parallel” to the constant zero function, which itself is a generalized line, too (see below). Moreover, as it can also be shown,  $\mathcal{L}(\omega_1, \omega_2)$  fulfills the axioms of hyperbolic geometry.

**LEMMA 2.2.** *If  $(\omega_1, \omega_2)$  is a Tchebychev system on an interval  $I$ , then there exists  $\omega \in \mathcal{L}(\omega_1, \omega_2)$  such that  $\omega$  is positive on  $I^\circ$ .*

**PROOF.** If  $\omega_1$  has no zero in  $I^\circ$ , then  $\omega := \omega_1$  or  $\omega := -\omega_1$  (according to the sign of  $\omega_1$ ) will do. Suppose that  $\omega_1(\xi) = 0$  for some  $\xi \in I^\circ$ . Due to Lemma 2.1, without loss of generality we may assume that

$$\begin{aligned}\omega_1(x) &< 0 & (x < \xi, x \in I) \\ \omega_1(y) &> 0 & (y > \xi, y \in I).\end{aligned}$$

Choose the elements  $x < \xi < y$  of  $I$ . The Tchebychev property of  $(\omega_1, \omega_2)$  and the negativity of  $\omega_1(x)\omega_2(y)$  implies the inequality

$$\frac{\omega_2(y)}{\omega_1(y)} < \frac{\omega_2(x)}{\omega_1(x)}.$$

Hence

$$(2.1) \quad \alpha := \sup_{y > \xi} \left[ \frac{\omega_2(y)}{\omega_1(y)} \right] \leq \inf_{x < \xi} \left[ \frac{\omega_2(x)}{\omega_1(x)} \right];$$

moreover, both sides are real numbers. We show that the generalized line defined by  $\omega := \alpha\omega_1 - \omega_2$  is positive on the interior of  $I$ .

First observe that  $\omega$  takes positive value at the point  $\xi$ . Indeed, by the definition of  $\omega$  we have  $\omega(\xi) := \alpha\omega_1(\xi) - \omega_2(\xi) = -\omega_2(\xi)$ ; on the other hand, for  $y > \xi$ , the positivity of  $\omega_1(y)$  and the Tchebychev property of  $(\omega_1, \omega_2)$  yields  $-\omega_2(\xi) > 0$ .

If  $y > \xi$ , then the definition of  $\alpha$  implies

$$\alpha \geq \frac{\omega_2(y)}{\omega_1(y)};$$

multiplying both sides by the positive  $\omega_1(y)$  and rearranging the terms we get,  $\omega(y) := \alpha\omega_1(y) - \omega_2(y) \geq 0$ .

If  $x < \xi$ , then inequality (2.1) gives that

$$\alpha \leq \frac{\omega_2(x)}{\omega_1(x)};$$

multiplying both sides by the negative  $\omega_1(x)$  and rearranging the obtained terms, we arrive at the inequality  $\omega(x) := \alpha\omega_1(x) - \omega_2(x) \geq 0$ .

To complete the proof, it suffices to show that  $\omega$  always differs from zero on the interior of the domain. Assume indirectly that  $\omega(\eta) := \alpha\omega_1(\eta) - \omega_2(\eta) = 0$  for some  $\eta \in I^\circ$ . Clearly,  $\eta \neq \xi$  since  $\omega(\xi) > 0$ . Therefore,  $\omega_1(\eta) \neq 0$  and  $\alpha$  can be expressed explicitly:

$$\alpha = \frac{\omega_2(\eta)}{\omega_1(\eta)}.$$

If  $\xi < \eta$ , choose  $y \in I$  such that  $\eta < y$  hold. By the positivity of  $\omega_1(\eta)\omega_1(y)$  and the Tchebychev property of  $(\omega_1, \omega_2)$ ,

$$\alpha = \frac{\omega_2(\eta)}{\omega_1(\eta)} < \frac{\omega_2(y)}{\omega_1(y)}$$

which contradicts the definition of  $\alpha$ . Similarly, if  $\xi > \eta$ , choose  $x \in I$  such that  $x < \eta$  be valid. Then, the positivity of  $\omega_1(x)\omega_1(\eta)$  and the Tchebychev property of  $(\omega_1, \omega_2)$  imply the inequality

$$\alpha = \frac{\omega_2(\eta)}{\omega_1(\eta)} > \frac{\omega_2(x)}{\omega_1(x)},$$

which contradicts (2.1).  $\square$

As an important consequence of Lemma 2.2, a Tchebychev system can always be replaced equivalently by a “regular” one. In other words, assuming positivity on the first component of a Tchebychev system, as it is required in many further results, is not an essential restriction. Moreover, the next lemma also gives a characterization of pairs of functions to form a Tchebychev system.

LEMMA 2.3. *Let  $(\omega_1, \omega_2)$  be a Tchebychev system on an interval  $I \subset \mathbb{R}$ . Then, there exists a Tchebychev system  $(\omega_1^*, \omega_2^*)$  on  $I$  that possesses the following properties:*

- (i)  $\omega_1^*$  is positive on  $I^\circ$ ;
- (ii)  $\omega_2^*/\omega_1^*$  is strictly monotone increasing on  $I^\circ$ ;
- (iii)  $(\omega_1, \omega_2)$ -convexity is equivalent to  $(\omega_1^*, \omega_2^*)$ -convexity.

Conversely, if  $\omega_1, \omega_2 : I \rightarrow \mathbb{R}$  are continuous functions such that  $\omega_1$  is positive and  $\omega_2/\omega_1$  is strictly monotone increasing, then  $(\omega_1, \omega_2)$  is a Tchebychev system over  $I$ .

PROOF. Lemma 2.2 guarantees the existence of real constants  $\alpha$  and  $\beta$  such that  $\alpha\omega_1 + \beta\omega_2 > 0$  holds for all  $x \in I^\circ$ . Define the functions  $\omega_1^*, \omega_2^* : I \rightarrow \mathbb{R}$  by the formulae

$$\omega_1^* := \alpha\omega_1 + \beta\omega_2 \quad \omega_2^* := -\beta\omega_1 + \alpha\omega_2.$$

Choosing the elements  $x < y$  of  $I$  and applying the product rule of determinants, we get

$$\begin{aligned} \begin{vmatrix} \omega_1^*(x) & \omega_1^*(y) \\ \omega_2^*(x) & \omega_2^*(y) \end{vmatrix} &= \begin{vmatrix} \alpha & \beta \\ -\beta & \alpha \end{vmatrix} \cdot \begin{vmatrix} \omega_1(x) & \omega_1(y) \\ \omega_2(x) & \omega_2(y) \end{vmatrix} \\ &= (\alpha^2 + \beta^2) \begin{vmatrix} \omega_1(x) & \omega_1(y) \\ \omega_2(x) & \omega_2(y) \end{vmatrix} > 0. \end{aligned}$$

Therefore,  $(\omega_1^*, \omega_2^*)$  is also a Tchebychev system over  $I$ . Assuming that  $\omega_1^*$  is positive, as it can easily be checked, the Tchebychev property of  $(\omega_1^*, \omega_2^*)$  yields that the function  $\omega_2^*/\omega_1^*$  is strictly monotone increasing on the interior of  $I$ .

At last, let  $f : I \rightarrow \mathbb{R}$  be an arbitrary function and  $x < y < z$  be arbitrary elements of  $I$ . Then, by the product rule of determinants again,

$$\begin{aligned} \begin{vmatrix} f(x) & f(y) & f(z) \\ \omega_1^*(x) & \omega_1^*(y) & \omega_1^*(z) \\ \omega_2^*(x) & \omega_2^*(y) & \omega_2^*(z) \end{vmatrix} &= \begin{vmatrix} 1 & 0 & 0 \\ 0 & \alpha & \beta \\ 0 & -\beta & \alpha \end{vmatrix} \cdot \begin{vmatrix} f(x) & f(y) & f(z) \\ \omega_1(x) & \omega_1(y) & \omega_1(z) \\ \omega_2(x) & \omega_2(y) & \omega_2(z) \end{vmatrix} \\ &= (\alpha^2 + \beta^2) \cdot \begin{vmatrix} f(x) & f(y) & f(z) \\ \omega_1(x) & \omega_1(y) & \omega_1(z) \\ \omega_2(x) & \omega_2(y) & \omega_2(z) \end{vmatrix}, \end{aligned}$$

which shows that the function  $f$  is generalized convex with respect to the Tchebychev system  $(\omega_1, \omega_2)$  if and only if it is generalized convex with respect to the Tchebychev system  $(\omega_1^*, \omega_2^*)$ .

The proof of the converse assertion is a simple calculation, therefore it is omitted.  $\square$

The following result gives various characterizations of  $(\omega_1, \omega_2)$ -convexity via the monotonicity of the generalized divided difference, the generalized support property and the ‘‘local’’ and the ‘‘global’’ generalized chord properties.

**THEOREM 2.4.** *Let  $(\omega_1, \omega_2)$  be a Tchebychev system over an interval  $I$  such that  $\omega_1$  is positive on  $I^\circ$ . The following statements are equivalent:*

- (i)  $f : I \rightarrow \mathbb{R}$  is  $(\omega_1, \omega_2)$ -convex;
- (ii) for all elements  $x < y < z$  of  $I$  we have that

$$\frac{\begin{vmatrix} f(y) & f(z) \\ \omega_1(y) & \omega_1(z) \end{vmatrix}}{\begin{vmatrix} \omega_1(y) & \omega_1(z) \\ \omega_2(y) & \omega_2(z) \end{vmatrix}} \leq \frac{\begin{vmatrix} f(x) & f(y) \\ \omega_1(x) & \omega_1(y) \end{vmatrix}}{\begin{vmatrix} \omega_1(x) & \omega_1(y) \\ \omega_2(x) & \omega_2(y) \end{vmatrix}};$$

- (iii) for all  $x_0 \in I^\circ$  there exist  $\alpha, \beta \in \mathbb{R}$  such that

$$\begin{aligned} \alpha\omega_1(x_0) + \beta\omega_2(x_0) &= f(x_0), \\ \alpha\omega_1(x) + \beta\omega_2(x) &\leq f(x) \quad (x \in I); \end{aligned}$$

- (iv) for all  $n \in \mathbb{N}$ ,  $x_0, x_1, \dots, x_n \in I$  and  $\lambda_1, \dots, \lambda_n \geq 0$  satisfying the conditions

$$\begin{aligned} \sum_{k=1}^n \lambda_k \omega_1(x_k) &= \omega_1(x_0) \\ \sum_{k=1}^n \lambda_k \omega_2(x_k) &= \omega_2(x_0) \end{aligned}$$

we have that

$$f(x_0) \leq \sum_{k=1}^n \lambda_k f(x_k);$$

(v) for all  $x_0, x_1, x_2 \in I$  and  $\lambda_1, \lambda_2 \geq 0$  satisfying the conditions

$$\lambda_1 \omega_1(x_1) + \lambda_2 \omega_1(x_2) = \omega_1(x_0)$$

$$\lambda_1 \omega_2(x_1) + \lambda_2 \omega_2(x_2) = \omega_2(x_0)$$

we have that

$$f(x_0) \leq \lambda_1 f(x_1) + \lambda_2 f(x_2);$$

(vi) for all elements  $x < p < y$  of  $I$

$$f(p) \leq \alpha \omega_1(p) + \beta \omega_2(p)$$

where the constants  $\alpha, \beta$  are the solutions of the system of linear equations

$$f(x) = \alpha \omega_1(x) + \beta \omega_2(x)$$

$$f(y) = \alpha \omega_1(y) + \beta \omega_2(y).$$

PROOF. (i)  $\Rightarrow$  (ii). Assume indirectly that (ii) is not true. Then, considering the positivity of the denominators, there exist elements  $x < y < z$  of  $I$  such that the inequality

$$\left| \begin{array}{cc} f(y) & f(z) \\ \omega_1(y) & \omega_1(z) \end{array} \right| \cdot \left| \begin{array}{cc} \omega_1(x) & \omega_1(y) \\ \omega_2(x) & \omega_2(y) \end{array} \right| > \left| \begin{array}{cc} f(x) & f(y) \\ \omega_1(x) & \omega_1(y) \end{array} \right| \cdot \left| \begin{array}{cc} \omega_1(y) & \omega_1(z) \\ \omega_2(y) & \omega_2(z) \end{array} \right|$$

holds or equivalently,

$$\begin{aligned} & f(y) \left( \omega_1(x) \left| \begin{array}{cc} \omega_1(y) & \omega_1(z) \\ \omega_2(y) & \omega_2(z) \end{array} \right| + \omega_1(z) \left| \begin{array}{cc} \omega_1(x) & \omega_1(y) \\ \omega_2(x) & \omega_2(y) \end{array} \right| \right) \\ & > \omega_1(y) \left( f(x) \left| \begin{array}{cc} \omega_1(y) & \omega_1(z) \\ \omega_2(y) & \omega_2(z) \end{array} \right| + f(z) \left| \begin{array}{cc} \omega_1(x) & \omega_1(y) \\ \omega_2(x) & \omega_2(y) \end{array} \right| \right). \end{aligned}$$

Subtracting

$$f(y) \omega_1(y) \left| \begin{array}{cc} \omega_1(x) & \omega_1(z) \\ \omega_2(x) & \omega_2(z) \end{array} \right|$$

from both sides and applying the expansion theorem “backwards”, we get

$$f(y) \left| \begin{array}{ccc} \omega_1(x) & \omega_1(y) & \omega_1(z) \\ \omega_1(x) & \omega_1(y) & \omega_1(z) \\ \omega_2(x) & \omega_2(y) & \omega_2(z) \end{array} \right| > \omega_1(y) \left| \begin{array}{ccc} f(x) & f(y) & f(z) \\ \omega_1(x) & \omega_1(y) & \omega_1(z) \\ \omega_2(x) & \omega_2(y) & \omega_2(z) \end{array} \right|.$$

The  $(\omega_1, \omega_2)$ -convexity of  $f$  implies that the right hand side of the inequality is nonnegative, while the left hand side equals zero, which is a contradiction.

(ii)  $\Rightarrow$  (iii). Fix  $x_0 \in I^\circ$ . Then, for all elements  $\xi < x_0 < x$  of  $I$ ,

$$-\frac{\left| \begin{array}{cc} f(\xi) & f(x_0) \\ \omega_1(\xi) & \omega_1(x_0) \end{array} \right|}{\left| \begin{array}{cc} \omega_1(\xi) & \omega_1(x_0) \\ \omega_2(\xi) & \omega_2(x_0) \end{array} \right|} \leq -\frac{\left| \begin{array}{cc} f(x_0) & f(x) \\ \omega_1(x_0) & \omega_1(x) \end{array} \right|}{\left| \begin{array}{cc} \omega_1(x_0) & \omega_1(x) \\ \omega_2(x_0) & \omega_2(x) \end{array} \right|}$$

holds, therefore

$$\beta := \inf_{x > x_0} \left[ \frac{\left| \begin{array}{cc} f(x_0) & f(x) \\ \omega_1(x_0) & \omega_1(x) \end{array} \right|}{\left| \begin{array}{cc} \omega_1(x_0) & \omega_1(x) \\ \omega_2(x_0) & \omega_2(x) \end{array} \right|} \right]$$

is a real number. The positivity assumption on  $\omega_1$  guarantees that the coefficient  $\alpha$  can be chosen such that  $\alpha\omega_1(x_0) + \beta\omega_2(x_0) = f(x_0)$  be satisfied. Rewrite the desired inequality  $\alpha\omega_1(x) + \beta\omega_2(x) \leq f(x)$  into the equivalent form

$$(2.2) \quad \beta \left| \begin{array}{cc} \omega_1(x_0) & \omega_1(x) \\ \omega_2(x_0) & \omega_2(x) \end{array} \right| + \left| \begin{array}{cc} f(x_0) & f(x) \\ \omega_1(x_0) & \omega_1(x) \end{array} \right| \leq 0.$$

The definition of  $\beta$  guarantees that it is valid if  $x_0 < x$ . Assume that  $x < x_0$  and choose  $\xi \in I$  such that  $x < x_0 < \xi$  hold. Then, applying (ii), we have the inequality

$$\frac{\left| \begin{array}{cc} f(x_0) & f(\xi) \\ \omega_1(x_0) & \omega_1(\xi) \end{array} \right|}{\left| \begin{array}{cc} \omega_1(x_0) & \omega_1(\xi) \\ \omega_2(x_0) & \omega_2(\xi) \end{array} \right|} \leq \frac{\left| \begin{array}{cc} f(x) & f(x_0) \\ \omega_1(x) & \omega_1(x_0) \end{array} \right|}{\left| \begin{array}{cc} \omega_1(x) & \omega_1(x_0) \\ \omega_2(x) & \omega_2(x_0) \end{array} \right|}.$$

Observe that the denominator of the right hand side is positive, therefore, after rearranging this inequality, we get

$$-\frac{\left| \begin{array}{cc} f(x_0) & f(\xi) \\ \omega_1(x_0) & \omega_1(\xi) \end{array} \right|}{\left| \begin{array}{cc} \omega_1(x_0) & \omega_1(\xi) \\ \omega_2(x_0) & \omega_2(\xi) \end{array} \right|} \left| \begin{array}{cc} \omega_1(x_0) & \omega_1(x) \\ \omega_2(x_0) & \omega_2(x) \end{array} \right| + \left| \begin{array}{cc} f(x_0) & f(x) \\ \omega_1(x_0) & \omega_1(x) \end{array} \right| \leq 0,$$

which, and the choice of  $\beta$  immediately implies (2.2).

(iii)  $\Rightarrow$  (iv). First assume that  $x_0 = x_1 = \dots = x_n$ . We recall that  $\omega_1(x_0)$  and  $\omega_2(x_0)$  cannot be equal to zero simultaneously due to Lemma 2.1; therefore one of the conditions gives the identity  $\sum_{k=1}^n \lambda_k = 1$ , and the inequality to be proved trivially holds. If  $x_0, x_1, \dots, x_n$  are distinct points of  $I$ , then it necessarily follows  $x_0 \in I^\circ$ . Indeed, if  $\inf(I) \in I$  and indirectly  $x_0 = \inf(I)$ , then we have the inequalities

$$\omega_1(x_0)\omega_2(x_k) - \omega_1(x_k)\omega_2(x_0) \geq 0$$

for all  $k = 1, \dots, n$  since  $(\omega_1, \omega_2)$  is a Tchebychev system on  $I$ ; furthermore, at least one of them is strict. Multiplying the  $k^{\text{th}}$  inequality by the positive  $\lambda_k$  and summing from 1 to  $n$ , we obtain

$$\omega_1(x_0) \sum_{k=1}^n \lambda_k \omega_2(x_k) > \omega_2(x_0) \sum_{k=1}^n \lambda_k \omega_1(x_k).$$

But, due to the conditions, both sides have the common value  $\omega_1(x_0)\omega_2(x_0)$ , which is a contradiction. An analogous argument gives that the case  $x_0 = \sup(I)$  is also impossible, therefore it follows that  $x_0 \in I^\circ$ .

Choose  $\alpha, \beta \in \mathbb{R}$  so that the relations

$$\begin{aligned}\alpha\omega_1(x_0) + \beta\omega_2(x_0) &= f(x_0) \\ \alpha\omega_1(x) + \beta\omega_2(x) &\leq f(x) \quad (x \in I)\end{aligned}$$

be valid. Then, substituting  $x = x_k$  into the last inequality and applying the conditions, we get that

$$\begin{aligned}\sum_{k=1}^n \lambda_k f(x_k) &\geq \sum_{k=1}^n \lambda_k \alpha \omega_1(x_k) + \sum_{k=1}^n \lambda_k \beta \omega_2(x_k) \\ &= \alpha \omega_1(x_0) + \beta \omega_2(x_0) = f(x_0),\end{aligned}$$

which gives the desired implication.

(iv)  $\Rightarrow$  (v). Taking the particular case  $n = 2$  in (iv), we arrive at (v).

(v)  $\Rightarrow$  (vi). According to Cramer's rule, for all elements  $x < p < y$  of  $I$  the system of linear equations

$$\begin{aligned}\lambda_1\omega_1(x) + \lambda_2\omega_1(y) &= \omega_1(p) \\ \lambda_1\omega_2(x) + \lambda_2\omega_2(y) &= \omega_2(p)\end{aligned}$$

has unique nonnegative solutions  $\lambda_1$  and  $\lambda_2$ . Therefore, using the definition of  $\alpha$  and  $\beta$ ,

$$\begin{aligned}f(p) &\leq \lambda_1 f(x) + \lambda_2 f(y) \\ &= \lambda_1(\alpha\omega_1(x) + \beta\omega_2(x)) + \lambda_2(\alpha\omega_1(y) + \beta\omega_2(y)) \\ &= \alpha(\lambda_1\omega_1(x) + \lambda_2\omega_1(y)) + \beta(\lambda_1\omega_2(x) + \lambda_2\omega_2(y)) \\ &= \alpha\omega_1(p) + \beta\omega_2(p).\end{aligned}$$

(vi)  $\Rightarrow$  (i). Expressing the unknowns  $\alpha$  and  $\beta$  with  $\omega_j(x), \omega_j(y)$  and  $\omega_j(p)$ , the inequality  $f(p) \leq \alpha\omega_1(p) + \beta\omega_2(p)$  can be rewritten into the form

$$\begin{vmatrix} \omega_1(x) & \omega_1(y) \\ \omega_2(x) & \omega_2(y) \end{vmatrix} f(p) \leq \begin{vmatrix} f(x) & f(y) \\ \omega_2(x) & \omega_2(y) \end{vmatrix} \omega_1(p) + \begin{vmatrix} f(x) & f(y) \\ \omega_1(x) & \omega_1(y) \end{vmatrix} \omega_2(p)$$

or equivalently

$$0 \leq \begin{vmatrix} f(x) & f(p) & f(y) \\ \omega_1(x) & \omega_1(p) & \omega_1(y) \\ \omega_2(x) & \omega_2(p) & \omega_2(y) \end{vmatrix},$$

which completes the proof.  $\square$

In the particular setting where  $\omega_1(x) := 1$  and  $\omega_2(x) := x$ , this theorem reduces to the well known characterization properties of standard convex functions. Now the last two assertions coincide: both of them state that the function's graph is under the chord joining between the endpoints of the graph. Let us note that in most of the literature the notion of (standard) convexity is defined exactly by this property (see the last assertion of the obtained corollary).

**COROLLARY 2.5.** *Let  $I \subset \mathbb{R}$  be an interval. The following statements are equivalent:*

- (i)  $f : I \rightarrow \mathbb{R}$  is convex (in the standard sense);  
(ii) for all elements  $x < y < z$  of  $I$  we have that

$$\frac{f(y) - f(x)}{y - x} \leq \frac{f(z) - f(y)}{z - y},$$

- (iii) for all  $x_0 \in I^\circ$  there exist  $\alpha, \beta \in \mathbb{R}$  such that

$$\alpha + \beta x_0 = f(x_0), \quad \alpha + \beta x \leq f(x) \quad (x \in I);$$

- (iv) for all  $n \in \mathbb{N}$ ,  $x_0, x_1, \dots, x_n \in I$  and  $\lambda_1, \dots, \lambda_n \geq 0$  satisfying the conditions

$$\sum_{k=1}^n \lambda_k = 1, \quad \sum_{k=1}^n \lambda_k x_k = x_0$$

we have that

$$f(x_0) \leq \sum_{k=1}^n \lambda_k f(x_k);$$

- (v) for all  $x_0, x_1, x_2 \in I$  and  $\lambda_1, \lambda_2 \geq 0$  satisfying the conditions

$$\lambda_1 + \lambda_2 = 1, \quad \lambda_1 x_1 + \lambda_2 x_2 = x_0$$

we have that

$$f(x_0) \leq \lambda_1 f(x_1) + \lambda_2 f(x_2).$$

If the base functions  $\omega_1$  and  $\omega_2$  are twice differentiable with a positive Wronski determinant, then a twice differentiable function  $f : I \rightarrow \mathbb{R}$  is  $(\omega_1, \omega_2)$ -convex if and only if the Wronski determinant of the system  $(f, \omega_1, \omega_2)$  is nonnegative (Bonsall, [Bon50]). This result can also be deduced from Theorem 2.4.

As it is well known, (standard) convex functions are exactly those ones that can be obtained as the pointwise supremum of families of affine functions. As a direct consequence (and also another application) of Theorem 2.4, an analogous statement holds for  $(\omega_1, \omega_2)$ -convex functions.

**COROLLARY 2.6.** *Let  $(\omega_1, \omega_2)$  be a Tchebychev system over an open interval  $I$ . Then, a function  $f : I \rightarrow \mathbb{R}$  is generalized convex with respect to  $(\omega_1, \omega_2)$  if and only if*

$$f(x) = \sup\{\omega(x) \mid \omega \in \mathcal{L}(\omega_1, \omega_2), \omega \leq f\}.$$

**PROOF.** Assertion (iii) of Theorem 2.4 immediately implies the representation above. For the sufficiency, part assertion (v) of Theorem 2.4 is applied. Fix the element  $x_0$  of the open interval  $I$ . Take a generalized line  $\omega = \alpha\omega_1 + \beta\omega_2$  such that  $\omega \leq f$ , furthermore, the elements  $x_1, x_2$  of  $I$  and the nonnegative coefficients  $\lambda_1, \lambda_2$  that fulfill the conditions

$$\begin{aligned} \lambda_1 \omega_1(x_1) + \lambda_2 \omega_1(x_2) &= \omega_1(x_0) \\ \lambda_1 \omega_2(x_1) + \lambda_2 \omega_2(x_2) &= \omega_2(x_0). \end{aligned}$$

Then,

$$\begin{aligned}
\lambda_1 f(x_1) + \lambda_2 f(x_2) &\geq \lambda_1 \omega(x_1) + \lambda_2 \omega(x_2) \\
&= \lambda_1 (\alpha \omega_1(x_1) + \beta \omega_2(x_1)) + \lambda_2 (\alpha \omega_1(x_2) + \beta \omega_2(x_2)) \\
&= \alpha (\lambda_1 \omega_1(x_1) + \lambda_2 \omega_1(x_2)) + \beta (\lambda_1 \omega_2(x_1) + \lambda_2 \omega_2(x_2)) \\
&= \alpha \omega_1(x_0) + \beta \omega_2(x_0) = \omega(x_0).
\end{aligned}$$

That is,  $\lambda_1 f(x_1) + \lambda_2 f(x_2) \geq \omega(x_0)$  for all  $\omega \leq f$ , hence, according to the representation, it follows that  $\lambda_1 f(x_1) + \lambda_2 f(x_2) \geq f(x_0)$ . Therefore  $f$  is convex with respect to  $(\omega_1, \omega_2)$ , indeed.  $\square$

## 2.2. Connection with standard convexity

The convexity notion induced by two dimensional Tchebychev systems turns out to be always reducible to standard convexity with the help of a composite function. This connection enables us to generalize many classical results for the case of  $(\omega_1, \omega_2)$ -convexity directly.

**THEOREM 2.7.** *Let  $(\omega_1, \omega_2)$  be a Tchebychev system on an open interval  $I$  such that  $\omega_1$  is positive. The function  $f : I \rightarrow \mathbb{R}$  is  $(\omega_1, \omega_2)$ -convex if and only if the function  $g : \omega_2/\omega_1(I) \rightarrow \mathbb{R}$  defined by the formula*

$$g := \frac{f}{\omega_1} \circ \left( \frac{\omega_2}{\omega_1} \right)^{-1}$$

is convex in the standard sense.

**PROOF.** In this case the function  $\omega_2/\omega_1$  is continuous and strictly monotone increasing, according to Lemma 2.3. Therefore, the image of the interval  $I$  by the function  $\omega_2/\omega_1$  is a nonempty open interval. Consider the identity

$$\begin{aligned}
&\left| \begin{array}{ccc} f(x) & f(y) & f(z) \\ \omega_1(x) & \omega_1(y) & \omega_1(z) \\ \omega_2(x) & \omega_2(y) & \omega_2(z) \end{array} \right| = \\
&= \omega_1(x)\omega_1(y)\omega_1(z) \left| \begin{array}{ccc} (f/\omega_1)(x) & (f/\omega_1)(y) & (f/\omega_1)(z) \\ 1 & 1 & 1 \\ (\omega_2/\omega_1)(x) & (\omega_2/\omega_1)(y) & (\omega_2/\omega_1)(z) \end{array} \right| \\
&= \omega_1(x)\omega_1(y)\omega_1(z) \left| \begin{array}{ccc} g(u) & g(v) & g(w) \\ 1 & 1 & 1 \\ u & v & w \end{array} \right|
\end{aligned}$$

where

$$u = (\omega_2/\omega_1)(x) \quad v = (\omega_2/\omega_1)(y) \quad w = (\omega_2/\omega_1)(z).$$

The positivity of  $\omega_1$  forces that both sides are simultaneously positive, negative or zero. That is, the function  $f$  is  $(\omega_1, \omega_2)$ -convex if and only if the function  $g$  is convex in the standard sense.  $\square$

Theorem 2.7 yields strong regularity properties for generalized convexity. For example,  $(\omega_1, \omega_2)$ -convex functions defined on compact intervals are integrable which is essential in formulating the main result of the chapter.

**THEOREM 2.8.** *Let  $(\omega_1, \omega_2)$  be a Tchebychev system on an interval  $I$ . If a function  $f : I \rightarrow \mathbb{R}$  is  $(\omega_1, \omega_2)$ -convex, then it is continuous on  $I^\circ$ . Moreover,  $f$  is bounded on each compact subinterval of  $I$ .*

**PROOF.** Without loss of generality we may assume that  $\omega_1$  is positive on  $I^\circ$ . If the function  $f$  is  $(\omega_1, \omega_2)$ -convex on  $I$ , then the composite function  $g$  in the previous theorem is convex in the standard sense on  $J := \omega_2/\omega_1(I)$ . Therefore, by the well known regularity properties of convex functions,  $g$  is continuous on  $J^\circ$ . On the other hand, we have that

$$f = \omega_1 \cdot g \circ \left( \frac{\omega_2}{\omega_1} \right),$$

and the right hand side is continuous on  $I^\circ$  whence the continuity of the function  $f$  follows.

To prove that  $f$  is bounded on the compact subinterval  $[a, b]$  of  $I$ , we shall apply Theorem 2.4. Take a generalized line which supports  $f$  at an arbitrary point  $x_0 \in I^\circ$ . Then, inequality (iii) implies that  $f$  is bounded from below on the whole interval  $I$ . On the other hand, putting  $x := a$  and  $y := b$  into (vi), we get that  $f$  is also bounded by a certain generalized line from above on  $[a, b]$ . Hence  $f$  is bounded, indeed.  $\square$

**DEFINITION.** *Let  $(\omega_1, \omega_2)$  be a Tchebychev system on an interval  $I$  furthermore  $\omega \in \mathcal{L}(\omega_1, \omega_2)$  be a generalized line which is positive on  $I^\circ$ . A function  $f : I \rightarrow \mathbb{R}$  is called generalized  $\omega$ -convex with respect to  $(\omega_1, \omega_2)$  if, for all elements  $x < y < z$  of  $I$ , the following inequality holds:*

$$\begin{vmatrix} f(x) + \omega(x) & f(y) - \omega(y) & f(z) + \omega(z) \\ \omega_1(x) & \omega_1(y) & \omega_1(z) \\ \omega_2(x) & \omega_2(y) & \omega_2(z) \end{vmatrix} \geq 0.$$

Substituting  $\omega_1(x) := 1$ ,  $\omega_2(x) := x$  and  $\omega := \varepsilon/2$ , the definition gives the notion of  $\varepsilon$ -convexity. By well known results,  $\varepsilon$ -convexity is stable: every  $\varepsilon$ -convex function is “close” to a (standard) convex function. As another application of Theorem 2.7, we prove an analogous result for  $(\omega_1, \omega_2)$ -convex functions.

**COROLLARY 2.9.** *Let  $(\omega_1, \omega_2)$  be a Tchebychev system on an interval  $I$  furthermore  $\omega \in \mathcal{L}(\omega_1, \omega_2)$  be a generalized line which is positive on  $I^\circ$ . A function  $f : I \rightarrow \mathbb{R}$  is generalized  $\omega$ -convex with respect to  $(\omega_1, \omega_2)$  if and only if there exist functions  $g, h : I \rightarrow \mathbb{R}$  such that  $g$  is  $(\omega_1, \omega_2)$ -convex,  $\|h\| \leq \|\omega\|$ , and  $f = g + h$ .*

**PROOF.** Assume that  $\omega$  has the representation  $\omega = \alpha\omega_1 + \beta\omega_2$  and define the generalized lines  $\omega_1^*$  and  $\omega_2^*$  by  $\omega_1^* := \alpha\omega_1 + \beta\omega_2$  and  $\omega_2^* := -\beta\omega_1 + \alpha\omega_2$ , respectively. Then, according to Lemma 2.3, the function  $\omega_2^*/\omega_1^*$  is strictly monotone

increasing and the generalized  $\omega$ -convexity of  $f$  is equivalent to the inequality

$$\begin{vmatrix} f(x) + \omega_1^*(x) & f(y) - \omega_1^*(y) & f(z) + \omega_1^*(z) \\ \omega_1^*(x) & \omega_1^*(y) & \omega_1^*(z) \\ \omega_2^*(x) & \omega_2^*(y) & \omega_2^*(z) \end{vmatrix} \geq 0.$$

Dividing both sides by the positive  $\omega_1^*(x)\omega_1^*(y)\omega_1^*(z)$  then substituting the arguments  $u = (\omega_2^*/\omega_1^*)(x)$ ,  $v = (\omega_2^*/\omega_1^*)(y)$  and  $w = (\omega_2^*/\omega_1^*)(z)$ , we get the inequality

$$\begin{vmatrix} F(u) + 1 & F(v) - 1 & F(w) + 1 \\ 1 & 1 & 1 \\ u & v & w \end{vmatrix} \geq 0$$

where

$$F := \frac{f}{\omega_1^*} \circ \left( \frac{\omega_2^*}{\omega_1^*} \right)^{-1}.$$

That is,  $F$  satisfies the inequality of  $\varepsilon$ -convexity with  $\varepsilon = 1$ . Therefore, there exist functions  $G, H : I \rightarrow \mathbb{R}$  such that  $G$  is convex (in the standard sense),  $\|H\| \leq 1$  and  $F = G + H$  or equivalently,

$$f = \omega_1^* \cdot G \circ \left( \frac{\omega_2^*}{\omega_1^*} \right) + \omega_1^* \cdot H \circ \left( \frac{\omega_2^*}{\omega_1^*} \right) =: g + h.$$

Then, Theorem 2.7 and Lemma 2.3 guarantee the  $(\omega_1, \omega_2)$ -convexity of  $g$ , while simple calculations imply  $\|h\| \leq \|\omega\|$ .  $\square$

### 2.3. Hermite–Hadamard-type inequalities

The main result provides Hermite–Hadamard-type inequalities for generalized 2-convex functions.

**THEOREM 2.10.** *Let  $(\omega_1, \omega_2)$  be a Tchebychev system on an interval  $[a, b]$  such that  $\omega_1$  is positive on  $]a, b[$ , furthermore, let  $\rho : [a, b] \rightarrow \mathbb{R}$  be a positive integrable function. Define the point  $\xi$  and the coefficients  $c, c_1, c_2$  by the formulae*

$$\xi = \left( \frac{\omega_2}{\omega_1} \right)^{-1} \left( \frac{\int_a^b \omega_2 \rho}{\int_a^b \omega_1 \rho} \right), \quad c = \frac{\int_a^b \omega_1 \rho}{\omega_1(\xi)}$$

and

$$c_1 = \frac{\begin{vmatrix} \int_a^b \omega_1 \rho & \omega_1(b) \\ \int_a^b \omega_2 \rho & \omega_2(b) \end{vmatrix}}{\begin{vmatrix} \omega_1(a) & \omega_1(b) \\ \omega_2(a) & \omega_2(b) \end{vmatrix}}, \quad c_2 = \frac{\begin{vmatrix} \omega_1(a) & \int_a^b \omega_1 \rho \\ \omega_2(a) & \int_a^b \omega_2 \rho \end{vmatrix}}{\begin{vmatrix} \omega_1(a) & \omega_1(b) \\ \omega_2(a) & \omega_2(b) \end{vmatrix}}.$$

If  $f : [a, b] \rightarrow \mathbb{R}$  is an  $(\omega_1, \omega_2)$ -convex function, then the following Hermite–Hadamard-type inequality holds

$$cf(\xi) \leq \int_a^b f \rho \leq c_1 f(a) + c_2 f(b).$$

PROOF. By the definitions of the point  $\xi$  and the constant  $c$ , we have the formulae

$$\frac{\int_a^b \omega_2 \rho}{\int_a^b \omega_1 \rho} = \frac{\omega_2(\xi)}{\omega_1(\xi)}$$

and

$$\int_a^b \omega_1 \rho = c\omega_1(\xi),$$

which yields the identity

$$\int_a^b \omega_2 \rho = c\omega_2(\xi).$$

That is, the left hand side of the Hermite–Hadamard-type inequality to be proved is exact for  $f = \omega_1$  and  $f = \omega_2$ , respectively. Let  $f : [a, b] \rightarrow \mathbb{R}$  be an arbitrary  $(\omega_1, \omega_2)$ -convex function and choose  $\alpha, \beta \in \mathbb{R}$  such that the relations

$$\begin{aligned} \alpha\omega_1(\xi) + \beta\omega_2(\xi) &= f(\xi) \\ \alpha\omega_1(x) + \beta\omega_2(x) &\leq f(x) \end{aligned}$$

be satisfied for all  $x \in [a, b]$ . By Theorem 2.4 such real numbers exist since  $\xi$  is an interior point of the domain. Multiplying the last inequality by the positive weight function  $\rho$ , we arrive at

$$\int_a^b f \rho \geq \alpha \int_a^b \omega_1 \rho + \beta \int_a^b \omega_2 \rho = \alpha(c\omega_1(\xi)) + \beta(c\omega_2(\xi)) = cf(\xi)$$

which results in the left hand side inequality.

To verify the right hand side one, observe first that the coefficients  $c_1$  and  $c_2$  are the solutions of the following system of linear equations

$$\begin{aligned} \int_a^b \omega_1 \rho &= c_1\omega_1(a) + c_2\omega_1(b) \\ \int_a^b \omega_2 \rho &= c_1\omega_1(a) + c_2\omega_2(b). \end{aligned}$$

In other words, the right hand side of the Hermite–Hadamard-inequality is exact, again, for  $f = \omega_1$  and  $f = \omega_2$ . Let  $f : [a, b] \rightarrow \mathbb{R}$  be an arbitrary  $(\omega_1, \omega_2)$ -convex function. By Theorem 2.4, if the real numbers  $\alpha$  and  $\beta$  are the solutions of the system of linear equations

$$\begin{aligned} f(a) &= \alpha\omega_1(a) + \beta\omega_2(a) \\ f(b) &= \alpha\omega_1(b) + \beta\omega_2(b), \end{aligned}$$

then

$$f(x) \leq \alpha\omega_1(x) + \beta\omega_2(x)$$

for all  $x \in [a, b]$ . Multiplying this inequality by the positive weight function  $\rho$ , we get that

$$\begin{aligned} \int_a^b f\rho &\leq \alpha \int_a^b \omega_1\rho + \beta \int_a^b \omega_2\rho \\ &= \alpha(c_1\omega_1(a) + c_2\omega_1(b)) + \beta(c_1\omega_2(a) + c_2\omega_2(b)) \\ &= c_1(\alpha\omega_1(a) + \beta\omega_2(a)) + c_2(\alpha\omega_1(b) + \beta\omega_2(b)) = c_1f(a) + c_2f(b), \end{aligned}$$

thus the proof is complete.  $\square$

## 2.4. Applications

Simple calculations show that specializing  $\omega_1(x) := 1$ ,  $\omega_2(x) := x$  and  $\rho \equiv 1$ , Theorem 2.10 reduces to the classical Hermite–Hadamard inequality :

COROLLARY 2.11. *If  $f : [a, b] \rightarrow \mathbb{R}$  is a (standard) convex function, then*

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{f(a) + f(b)}{2}.$$

The subsequent corollaries present further Hermite–Hadamard-type inequalities for generalized convex functions where the underlying Tchebychev systems of the induced convexity are the hyperbolic, trigonometric, exponential and power systems (to see that the pairs  $(\omega_1, \omega_2)$  form a Tchebychev system in each case, consult the converse part of Lemma 2.3).

COROLLARY 2.12. *If  $f : [a, b] \rightarrow \mathbb{R}$  is a (cosh, sinh)-convex function, then*

$$2 \sinh\left(\frac{b-a}{2}\right) f\left(\frac{a+b}{2}\right) \leq \int_a^b f(x)dx \leq \tanh\left(\frac{b-a}{2}\right) (f(a) + f(b)).$$

PROOF. If  $\omega_1 := \cosh$  and  $\omega_2 := \sinh$ , then  $\omega_1$  is positive and  $\omega_2/\omega_1 = \tanh$  is strictly monotone increasing; hence, according to Lemma 2.3,  $(\omega_1, \omega_2)$  is a Tchebychev system and  $(\omega_2/\omega_1)^{-1} = \operatorname{artanh}$ . Applying the addition properties of hyperbolic functions for the identities  $b = (b+a)/2 + (b-a)/2$  and  $a = (b+a)/2 - (b-a)/2$ , the integrals of  $\omega_1$  and  $\omega_2$  can be written into product form via the formulae

$$\begin{aligned} \int_a^b \cosh x dx &= \sinh(b) - \sinh(a) = 2 \cosh\left(\frac{b+a}{2}\right) \sinh\left(\frac{b-a}{2}\right) \\ \int_a^b \sinh x dx &= \cosh(b) - \cosh(a) = 2 \sinh\left(\frac{b+a}{2}\right) \sinh\left(\frac{b-a}{2}\right). \end{aligned}$$

Therefore,

$$\xi = \operatorname{artanh}\left(\frac{\int_a^b \sinh x dx}{\int_a^b \cosh x dx}\right) = \frac{b+a}{2}$$

furthermore

$$c = \frac{\int_a^b \cosh x dx}{\cosh \xi} = 2 \sinh \frac{b+a}{2}.$$

To determine the coefficients of the right hand side, first we calculate the numerator of  $c_1$ :

$$\begin{aligned} & \left| \begin{array}{cc} 2 \cosh \left( \frac{b+a}{2} \right) \sinh \left( \frac{b-a}{2} \right) & \cosh b \\ 2 \sinh \left( \frac{b+a}{2} \right) \sinh \left( \frac{b-a}{2} \right) & \sinh b \end{array} \right| = \\ & = 2 \sinh \left( \frac{b-a}{2} \right) \left( \cosh \left( \frac{b+a}{2} \right) \sinh b - \sinh \left( \frac{b+a}{2} \right) \cosh b \right) \\ & = 2 \sinh \left( \frac{b-a}{2} \right) \sinh \left( b - \frac{b+a}{2} \right) = 2 \sinh^2 \left( \frac{b-a}{2} \right). \end{aligned}$$

Similarly, the numerator of the coefficient  $c_2$  can be obtained as follows:

$$\begin{aligned} & \left| \begin{array}{cc} \cosh a & 2 \cosh \left( \frac{b+a}{2} \right) \sinh \left( \frac{b-a}{2} \right) \\ \sinh a & 2 \sinh \left( \frac{b+a}{2} \right) \sinh \left( \frac{b-a}{2} \right) \end{array} \right| = \\ & = 2 \sinh \left( \frac{b-a}{2} \right) \left( \sinh \left( \frac{b+a}{2} \right) \cosh a - \cosh \left( \frac{b+a}{2} \right) \sinh a \right) \\ & = 2 \sinh \left( \frac{b-a}{2} \right) \sinh \left( \frac{b+a}{2} - a \right) = 2 \sinh^2 \left( \frac{b-a}{2} \right). \end{aligned}$$

On the other hand, the denominators in both cases coincide and have the common value

$$\left| \begin{array}{cc} \cosh a & \cosh b \\ \sinh a & \sinh b \end{array} \right| = \sinh(b-a) = 2 \sinh \left( \frac{b-a}{2} \right) \cosh \left( \frac{b-a}{2} \right),$$

therefore

$$c_1 = c_2 = \tanh \left( \frac{b-a}{2} \right).$$

□

Replacing the Tchebychev system  $(\cosh, \sinh)$  with  $(\cos, \sin)$ , the obtained Hermite–Hadamard-type inequality is analogous to the previous one due to the similar additional properties of trigonometric and hyperbolic functions.

**COROLLARY 2.13.** *If  $f : [a, b] \subset ]-\frac{\pi}{2}, \frac{\pi}{2}[ \rightarrow \mathbb{R}$  is a  $(\cos, \sin)$ -convex function, then*

$$2 \sin \left( \frac{b-a}{2} \right) f \left( \frac{a+b}{2} \right) \leq \int_a^b f(x) dx \leq \tan \left( \frac{b-a}{2} \right) (f(a) + f(b)).$$

Observe that both of the previous two Hermite–Hadamard-type inequalities involve the midpoint of the domain; moreover, dividing by  $b-a$  and taking the limit  $a \rightarrow b$ , the coefficient of the left hand sides tends to 1, while the coefficient of the right hand sides tends to  $1/2$ . Therefore these inequalities can be considered as the “local” version of the Hermite–Hadamard inequality.

We say that a function  $f : I \rightarrow \mathbb{R}$  is *log-convex* if the composite function  $f \circ \log : \exp(I) \rightarrow \mathbb{R}$  is convex (in the standard sense). In terms of generalized convexity, log-convex functions are exactly the  $(1, \exp)$ -convex ones (consult Theorem 2.7). The next corollary gives Hermite–Hadamard-type inequality for log-convex functions ([Dra01c], [Fin00]).

**COROLLARY 2.14.** *If  $f : [a, b] \rightarrow \mathbb{R}$  is a  $(1, \exp)$ -convex function, then*

$$\begin{aligned} & (b-a)f\left(\log \frac{\exp(b) - \exp(a)}{b-a}\right) \leq \int_a^b f(x)dx \\ & \leq \left(\frac{(b-a)\exp(b)}{\exp(b) - \exp(a)} - 1\right) f(a) + \left(1 - \frac{(b-a)\exp(a)}{\exp(b) - \exp(a)}\right) f(b). \end{aligned}$$

The last corollary concerning the case of “power convexity” also reduces to the classical Hermite–Hadamard inequality substituting  $p = 0$  and  $q = 1$ :

**COROLLARY 2.15.** *If  $p < q$ ,  $p, q \neq -1$  and  $f : [a, b] \subset ]0, \infty[ \rightarrow \mathbb{R}$  is an  $(x^p, x^q)$ -convex function, then*

$$\begin{aligned} & \left(\frac{b^{p+1} - a^{p+1}}{p+1}\right)^q \left(\frac{q+1}{b^{q+1} - a^{q+1}}\right)^p f\left(\sqrt[q-p]{\frac{(p+1)(b^{q+1} - a^{q+1})}{(q+1)(b^{p+1} - a^{p+1})}}\right) \\ & \leq \int_a^b f(x)dx \\ & \leq \frac{\frac{(b^{p+1} - a^{p+1})b^q}{p+1} - \frac{(b^{q+1} - a^{q+1})b^p}{q+1}}{a^p b^q - a^q b^p} f(a) + \frac{\frac{(b^{q+1} - a^{q+1})a^p}{q+1} - \frac{(b^{p+1} - a^{p+1})a^q}{q+1}}{a^p b^q - a^q b^p} f(b). \end{aligned}$$

The proofs of the last three corollaries need similar calculations as the first one, therefore they are omitted.

## Generalized convexity induced by Tchebychev systems

In this chapter we formulate Hermite–Hadamard-type inequalities for generalized convex functions where the underlying Tchebychev system of the induced convexity is *arbitrary*. The proofs of the main results are based on the Krein–Markov theory of moment spaces induced by Tchebychev systems. According to this theory, the vector integral of a Tchebychev system can uniquely be represented as the linear combination of the values of the system in certain base points of the domain. The number of the points and also the points themselves, depend only on the Tchebychev system and its dimension: it turns out that the cases of odd and even order convexity must be investigated separately. In fact, this is exactly the deeper reason for the analogous phenomenon in the case of polynomial convexity, too. Once the base points of the representations are determined, its coefficients are obtained as the solutions of a system of linear equations. With the help of the representations and the notion of generalized convexity, the Hermite–Hadamard-type inequalities can be verified using integration and pure linear algebraic methods.

In the previous chapters when the basis or the dimension of the studied Tchebychev systems are quite special, the base points of the Hermite–Hadamard-type inequalities can explicitly be given. Unfortunately, under the present general circumstances, we can guarantee only the *existence* (and the uniqueness) of the base points, but *cannot give any explicit formulae for them*.

At last, motivated by Rolle’s mean-value theorem, an alternative and elementary approach is presented for the cases when the Hermite–Hadamard-type inequalities involve at most one interior base point of the domain. Some examples are also presented of these particular cases.

### 3.1. Characterizations and regularity properties

Let  $\boldsymbol{\omega} = (\omega_1, \dots, \omega_n)$  be a Tchebychev system over an interval  $I$  and denote the set of all linear combinations of its members by  $\mathcal{L}(\omega_1, \dots, \omega_n)$ . A function is called *generalized polynomial* (belonging to the system in question) if it is the element of the linear span  $\mathcal{L}(\omega_1, \dots, \omega_n)$ . In terms of generalized polynomials, generalized convexity can be characterized in a very descriptive geometrical manner. Namely, a function is generalized convex if and only if it intersects its generalized polynomial that interpolates the function in any prescribed points alternately. (The number of the points depends on the dimension of the underlying Tchebychev system). More precisely, we have the following

**THEOREM 3.1.** *Let  $\boldsymbol{\omega} = (\omega_1, \dots, \omega_n)$  be a Tchebychev system over an interval  $I$ . Then, for a function  $f : I \rightarrow \mathbb{R}$ , the following statements are equivalent:*

- (i)  *$f$  is generalized convex with respect to  $\boldsymbol{\omega}$ ;*
- (ii) *for all  $y_1 < \dots < y_n$  in  $I$ , the generalized polynomial  $\omega$  of  $\omega_1, \dots, \omega_n$  determined uniquely by the interpolation conditions*

$$f(y_k) = \omega(y_k) \quad (k = 1, \dots, n)$$

*satisfies the inequalities*

$$(-1)^{n+k}(f(y) - \omega(y)) \geq 0 \quad (y_k < y < y_{k+1}, k = 0, \dots, n)$$

*under the conventions  $y_0 := \inf I$  and  $y_{n+1} := \sup I$ ;*

- (iii) *keeping the previous notations and settings, for fixed  $k \in \{0, \dots, n\}$ , the following inequality holds*

$$(-1)^{n+k}(f(y) - \omega(y)) \geq 0 \quad (y_k \leq y \leq y_{k+1}).$$

**PROOF.** First of all, in order to simplify the proof, two useful formulas are derived. Denote the  $n - 1$  tuple obtained by dropping the  $k^{\text{th}}$  component of  $\boldsymbol{\omega}$  by  $\boldsymbol{\omega}_k$ , and define the determinants  $D_0, D_1, \dots, D_n$  furthermore the generalized polynomial  $\omega$  of  $\omega_1, \dots, \omega_n$  by

$$\begin{aligned} D_0 &:= \begin{vmatrix} \boldsymbol{\omega}(y_1) & \cdots & \boldsymbol{\omega}(y_n) \end{vmatrix} \\ D_k &:= \begin{vmatrix} f(y_1) & \cdots & f(y_n) \\ \boldsymbol{\omega}_k(y_1) & \cdots & \boldsymbol{\omega}_k(y_n) \end{vmatrix} \\ \omega &:= \sum_{k=1}^n \frac{(-1)^{k+1} D_k}{D_0} \omega_k. \end{aligned}$$

Due to the Tchebychev property of  $\boldsymbol{\omega}$ , the determinant  $D_0$  is positive hence the definition of  $\omega$  is correct. Fix  $y \in I$ . Applying the expansion theorem to the first column of the following determinant, we get the identity

$$(3.1) \quad \begin{vmatrix} f(y) & f(y_1) & \cdots & f(y_n) \\ \boldsymbol{\omega}(y) & \boldsymbol{\omega}(y_1) & \cdots & \boldsymbol{\omega}(y_n) \end{vmatrix} = D_0(f(y) - \omega(y)).$$

Moreover, if  $y_k \leq y \leq y_{k+1}$  and  $(x_0, x_1, \dots, x_n)$  denotes the increasing rearrangement of  $(y; y_1, \dots, y_n)$ , the previous identity can be written into the form

$$(3.2) \quad \begin{vmatrix} f(x_0) & f(x_1) & \cdots & f(x_n) \\ \boldsymbol{\omega}(x_0) & \boldsymbol{\omega}(x_1) & \cdots & \boldsymbol{\omega}(x_n) \end{vmatrix} = (-1)^k D_0(f(y) - \omega(y)).$$

For the implication (i)  $\implies$  (ii), observe that (3.1) guarantees the required interpolation property of  $\omega$  in the points  $y_1, \dots, y_n$ . Clearly,  $\omega$  is uniquely determined. Suppose that  $f : I \rightarrow \mathbb{R}$  is generalized  $n$ -convex with respect to  $\boldsymbol{\omega}$ . Then, the positivity of  $D_0$  and formula (3.2) yield the inequalities to be proved. The implication (ii)  $\implies$  (iii) is trivial. The proof of (iii)  $\implies$  (i) is completely the same as the proof of the first assertion.  $\square$

In the standard setting and fixing  $k = 1$ , assertion (iii) gives the classical definition of standard convexity: a function is convex (in the standard sense) if and only if it is “under” the chord of the graph. Moreover, substituting  $n = 2$ , we also get a new characterization of generalized 2-convexity that completes Theorem 2.4. However, the most important application of Theorem 3.1 guarantees strong regularity properties for generalized convex functions.

**THEOREM 3.2.** *Let  $\omega = (\omega_1, \dots, \omega_n)$  be a Tchebychev system over an interval  $I$ . If  $f : I \rightarrow \mathbb{R}$  is a generalized  $n$ -convex function with respect to this system and  $n \geq 2$ , then  $f$  is continuous on the interior of  $I$ . Furthermore,  $f$  is bounded on each compact subinterval of  $I$ .*

**PROOF.** Choose  $y_0 \in I^\circ$  and fix  $x_0 < x_1 < \dots < x_n$  in  $I$  so that  $x_1 = y_0$  hold. Denote the generalized polynomials of  $\omega_1, \dots, \omega_n$  that interpolate  $\omega_0$  in the points  $x_0, \dots, x_{n-1}$  and  $x_1, \dots, x_n$  by  $\omega^{(1)}$  and  $\omega^{(2)}$ , respectively. We assume that  $n$  is even (the argument in the odd case is analogous). Then, according to (ii) of Theorem 3.1, we have the inequalities

$$\begin{aligned} \omega^{(1)}(y) &\geq \omega_0(y) \geq \omega^{(2)}(y) & y \in [x_0, x_1] \\ \omega^{(1)}(y) &\leq \omega_0(y) \leq \omega^{(2)}(y) & y \in [x_1, x_2]. \end{aligned}$$

On the other hand,  $\omega^{(1)}(y_0) = \omega_0(y_0)$  and  $\omega^{(2)}(y_0) = \omega_0(y_0)$ . Therefore, due to the continuity of the generalized polynomials  $\omega^{(1)}$  and  $\omega^{(2)}$ , we get that both the left and right hand side limits of  $\omega_0$  exist at the point  $y_0$  and

$$\begin{aligned} \lim_{y \rightarrow y_0 - 0} \omega_0(y) &= \omega_0(y_0) \\ \lim_{y \rightarrow y_0 + 0} \omega_0(y) &= \omega_0(y_0), \end{aligned}$$

which yields the continuity of  $\omega_0$  at the interior point  $y_0$  of  $I$ .

To prove the second assertion, we may assume that  $I = [a, b]$ . It is sufficient to show that  $\omega_0$  is locally bounded at the endpoints of  $I$ . Fix  $x_0 < x_1 < \dots < x_n$  in  $I$  so that  $x_0 = a$  hold, and denote the generalized polynomials of  $\omega_1, \dots, \omega_n$  that interpolate  $\omega_0$  in the points  $x_0, \dots, x_{n-1}$  and  $x_1, \dots, x_n$  by  $\omega^{(1)}$  and  $\omega^{(2)}$ , respectively. We assume that  $n$  is even (the odd case is very similar). Then, by the previous theorem again, we have the inequalities

$$\omega^{(1)}(y) \geq \omega_0(y) \geq \omega^{(2)}(y) \quad y \in [x_0, x_1].$$

On the other hand, the functions  $\omega^{(1)}$  and  $\omega^{(2)}$  are continuous, therefore bounded on  $[a, b]$ . Hence  $\omega_0$  is bounded in a right neighborhood of the endpoint  $a$ . It can be similarly proved that  $\omega_0$  is locally bounded at the left endpoint  $b$ .  $\square$

In particular, generalized convex functions are integrable on any compact subset of the domain. Let us also mention that the special case  $n = 2$  gives the statement of Theorem 2.8 via another approach in the proof.

### 3.2. Moment spaces induced by Tchebychev systems

The geometric study of moment spaces induced by Tchebychev systems was systematically developed by M. G. Krein. Independently and simultaneously, S. Karlin and L. S. Shapley elaborated the geometry of moment spaces induced by the polynomial system. Some of the results of their researches play the key role in the further investigations.

DEFINITION. Let  $\omega := (\omega_1, \dots, \omega_n)$  be a Tchebychev system on  $[a, b]$  and denote the set of all nondecreasing right continuous functions defined on  $[a, b]$  by  $\mathcal{B}([a, b])$ . The set

$$\mathcal{M}_n := \left\{ c \in \mathbb{R}^n \mid c = \int_a^b \omega d\sigma, \sigma \in \mathcal{B}([a, b]) \right\}$$

is called the moment space of  $\omega$ .

It can be shown that  $\mathcal{M}_n$  is a closed convex cone. More precisely, it is the smallest closed convex cone that contains the parameterized curve  $\omega(t)$  where  $t$  traverses the interval  $[a, b]$ . For details, see [KS66, pp. 38-41]. The following notion makes the formulation of many theorems quite convenient.

DEFINITION. The index  $I(c)$  of a point  $c \in \mathcal{M}_n$  is the minimal number of points  $\xi_1, \dots, \xi_{n_0}$  in a representation

$$c = \sum_{k=1}^{n_0} \alpha_k \omega(\xi_k)$$

under the convention that  $\omega(a)$  and  $\omega(b)$  are counted with half multiplicity, while  $\omega(\xi)$  for  $\xi \in ]a, b[$  receives a full count. The points  $\xi_1, \dots, \xi_{n_0}$  are called the roots of the representation.

By the celebrated theorem of Carathéodory (see [Roc70]), each point belonging to the conical hull of a given subset of  $\mathbb{R}^n$  can be represented as a cone combination involving at most  $n$  points of the subset. Due to the Tchebychev property of  $\omega$ , a surprisingly better upper bound can be established than  $n$ : it turns out that the elements of  $\mathcal{M}_n$  are cone combinations of approximately  $n/2$  points of the range of  $\omega$ . More precisely, the boundary and the interior of  $\mathcal{M}_n$ , denoted by  $\text{Bd } \mathcal{M}_n$  and  $\text{Int } \mathcal{M}_n$ , can be characterized via the subsequent two theorems due to Krein and Markov.

THEOREM C. ([KS66, Theorem 2.1. p. 42]) A vector  $c \in \mathcal{M}_n$  is a boundary point of  $\mathcal{M}_n$  if and only if  $I(c) < n/2$ . Moreover, every  $c \in \text{Bd } \mathcal{M}_n$  admits a unique representation

$$c = \sum_{k=1}^{n_0} \alpha_k \omega(\xi_k) \quad (\xi_k \in [a, b], \alpha_k > 0, k = 1, \dots, n_0)$$

where  $n_0 \leq \frac{n+1}{2}$ .

**THEOREM D.** ([KS66, Theorem 3.1. p. 44; Remark 3.1. pp. 45-46; Corollary 3.1. p. 47.]) *For each  $c \in \text{Int } \mathcal{M}_n$  there exist precisely two representations of index  $I(c) = n/2$ . Distinguishing the even and odd cases, the representations in question are the following.*

*Case  $n = 2m$ :*

$$c = \sum_{k=1}^m \alpha_k \boldsymbol{\omega}(\xi_k) \quad (\xi_k \in ]a, b[),$$

$$c = \beta_0 \boldsymbol{\omega}(a) + \sum_{k=1}^{m-1} \beta_k \boldsymbol{\omega}(\eta_k) + \beta_m \boldsymbol{\omega}(b) \quad (\eta_k \in ]a, b[);$$

*Case  $n = 2m + 1$ :*

$$c = \alpha_0 \boldsymbol{\omega}(a) + \sum_{k=1}^m \alpha_k \boldsymbol{\omega}(\xi_k) \quad (\xi_k \in ]a, b[),$$

$$c = \sum_{k=1}^m \beta_k \boldsymbol{\omega}(\eta_k) + \beta_{m+1} \boldsymbol{\omega}(b) \quad (\eta_k \in ]a, b[).$$

*The roots of the representations in both cases strictly interlace.*

Let  $I \subset \mathbb{R}$  be a real interval and  $\boldsymbol{\omega} := (\omega_1, \dots, \omega_n)$  be a Tchebychev system over  $I$ . Then, for pairwise distinct elements  $t_1, \dots, t_n$  of  $I$ , the vectors  $\boldsymbol{\omega}(t_1), \dots, \boldsymbol{\omega}(t_n)$  are linearly independent. This simple observation immediately implies

**THEOREM 3.3.** *The coefficients and the roots of the representations above are uniquely determined.*

Now we present a sufficient condition for a point  $c$  to belong to the interior of the set  $\mathcal{M}_n$ . This condition guarantees that the inequalities of the main results have exactly the required form.

**THEOREM 3.4.** *Let  $\boldsymbol{\omega} = (\omega_1, \dots, \omega_n)$  be a Tchebychev system on  $[a, b]$  and let  $\rho : [a, b] \rightarrow \mathbb{R}$  be a positive integrable function. Then,*

$$c := \int_a^b \boldsymbol{\omega} \rho \in \text{Int } \mathcal{M}_n.$$

**PROOF.** Let us recall that  $\mathcal{M}_n$  is a closed subset of  $\mathbb{R}^n$ . On the other hand, the positivity of  $\rho$  yields  $c \in \mathcal{M}_n$ , therefore it suffices to prove that  $c \notin \text{Bd } \mathcal{M}_n$ . Assume indirectly that  $c \in \text{Bd } \mathcal{M}_n$ . We shall distinguish two cases according to the parity of  $n$ .

*Case  $n = 2m + 1$ .* The indirect assumption and Theorem C implies  $I(c) \leq m$  since  $I(c)$  increases at most  $1/2$ . For simplicity, assume that  $I(c) = m$ . Then there are two further possibilities: the representation of  $c$  involves either  $m$  pairwise distinct interior base points  $\xi_1 < \dots < \xi_m$  or  $m - 1$  pairwise distinct interior base

points  $\xi_1 < \dots < \xi_{m-1}$  plus both the endpoints  $a$  and  $b$ , respectively. In the first case we have the representation

$$c = \sum_{k=1}^m \alpha_k \omega(\xi_k).$$

Due to the Tchebychev property of  $\omega$  and the positivity of  $\rho$ , we arrive at

$$0 < \left| \begin{array}{cccccc} \omega(t_1)\rho(t_1) & \omega(\xi_1) & \cdots & \omega(t_m)\rho(t_m) & \omega(\xi_m) & \omega(t_{m+1})\rho(t_{m+1}) \end{array} \right|$$

for  $t_k \in ]\xi_{k-1}, \xi_k[$  ( $k = 1, \dots, m$ ) where  $\xi_0 := a$  and  $\xi_{m+1} := b$ . After integration with respect to  $(t_1, \dots, t_{m+1})$  and using the above representation of  $c$ , we have

$$\begin{aligned} 0 &< \left| \begin{array}{cccccc} \int_{\xi_0}^{\xi_1} \omega \rho & \omega(\xi_1) & \cdots & \int_{\xi_{m-1}}^{\xi_m} \omega \rho & \omega(\xi_m) & \int_{\xi_m}^{\xi_{m+1}} \omega \rho \end{array} \right| \\ &= \left| \begin{array}{cccccc} \int_{\xi_0}^{\xi_1} \omega \rho & \omega(\xi_1) & \cdots & \int_{\xi_{m-1}}^{\xi_m} \omega \rho & \omega(\xi_m) & \sum_{k=1}^{m+1} \int_{\xi_{k-1}}^{\xi_k} \omega \rho \end{array} \right| \\ &= \left| \begin{array}{cccccc} \int_{\xi_0}^{\xi_1} \omega \rho & \omega(\xi_1) & \cdots & \int_{\xi_{m-1}}^{\xi_m} \omega \rho & \omega(\xi_m) & \int_a^b \omega \rho \end{array} \right| \\ &= \left| \begin{array}{cccccc} \int_{\xi_0}^{\xi_1} \omega \rho & \omega(\xi_1) & \cdots & \int_{\xi_{m-1}}^{\xi_m} \omega \rho & \omega(\xi_m) & \sum_{k=1}^m \alpha_k \omega(\xi_k) \end{array} \right| = 0 \end{aligned}$$

since the last column is the linear combination of the even indexed columns. Thus we get the desired contradiction.

Now consider the other case when  $c$  has the representation

$$c = \alpha_0 \omega(a) + \sum_{k=1}^{m-1} \alpha_k \omega(\xi_k) + \alpha_m \omega(b).$$

Due to the Tchebychev property of  $\omega$  and the positivity of  $\rho$  again, we arrive at

$$0 < \left| \begin{array}{cccccc} \omega(a) & \omega(t_1)\rho(t_1) & \omega(\xi_1) & \cdots & \omega(\xi_{m-1}) & \omega(t_m)\rho(t_m) & \omega(b) \end{array} \right|$$

for  $t_k \in ]\xi_{k-1}, \xi_k[$  ( $k = 1, \dots, m$ ) where  $\xi_0 := a$  and  $\xi_m := b$ . An analogous argument to the previous one leads to contradiction.

*Case  $n = 2m$ .* Similarly to the odd case, now we may assume that  $I(c) = m - 1/2$ . Then there are two possibilities: the representation of  $c$  involves either the endpoint  $a$  and  $m - 1$  pairwise distinct interior base points  $\xi_1 < \dots < \xi_{m-1}$  or the endpoint  $b$  and  $m - 1$  pairwise distinct interior base points  $\xi_1 < \dots < \xi_{m-1}$ . Applying the same method as above, both cases lead to contradiction again.  $\square$

### 3.3. Hermite–Hadamard-type inequalities

The main results concern the cases of even and odd order generalized convexity separately. First we establish Hermite–Hadamard-type inequalities for the odd order one.

**THEOREM 3.5.** *Let  $\boldsymbol{\omega} = (\omega_1, \dots, \omega_{2m+1})$  be a Tchebychev system on  $[a, b]$  and  $\rho : [a, b] \rightarrow \mathbb{R}$  be a positive integrable function. There exist uniquely determined base points  $\xi_1, \dots, \xi_m$  and  $\eta_1, \dots, \eta_m$  of  $]a, b[$  such that*

$$\alpha_0 \boldsymbol{\omega}(a) + \sum_{k=1}^m \alpha_k \boldsymbol{\omega}(\xi_k) = \int_a^b \boldsymbol{\omega} \rho = \sum_{k=1}^m \beta_k \boldsymbol{\omega}(\eta_k) + \beta_{m+1} \boldsymbol{\omega}(b).$$

*The coefficients  $\alpha_0, \dots, \alpha_m$  and  $\beta_1, \dots, \beta_{m+1}$  are positive and uniquely determined, too. Furthermore, for any generalized  $\boldsymbol{\omega}$ -convex function  $f : [a, b] \rightarrow \mathbb{R}$ , the following Hermite–Hadamard-type inequality holds*

$$\alpha_0 f(a) + \sum_{k=1}^m \alpha_k f(\xi_k) \leq \int_a^b f \rho \leq \sum_{k=1}^m \beta_k f(\eta_k) + \beta_{m+1} f(b).$$

**PROOF.** Let us note that  $f\rho$  is integrable on  $[a, b]$  by Theorem 3.2. The proofs of the left and right hand side inequalities need similar methods, therefore, we shall verify only the left hand side one. Theorem 3.4 guarantees that  $\int_a^b \boldsymbol{\omega} \rho$  is an interior point of the moment space  $\mathcal{M}_n$  hence (see Theorem D and Theorem 3.3) it has the representation

$$(3.3) \quad \int_a^b \boldsymbol{\omega} \rho = \alpha_0 \boldsymbol{\omega}(a) + \sum_{k=1}^m \alpha_k \boldsymbol{\omega}(\xi_k)$$

where the coefficients  $\alpha_0, \dots, \alpha_m$  and interior base points  $\xi_1, \dots, \xi_m$  are determined uniquely. Defining  $\xi_0 := a$  and  $\xi_{m+1} := b$ , consider the following system of linear equations

$$\int_{\xi_m}^{\xi_{m+1}} \boldsymbol{\omega} \rho = c_0 \boldsymbol{\omega}(\xi_0) + \sum_{k=1}^m \left( c_k^* \int_{\xi_{k-1}}^{\xi_k} \boldsymbol{\omega} \rho + c_k \boldsymbol{\omega}(\xi_k) \right)$$

where the unknowns are  $c_0, c_1^*, c_1, \dots, c_m^*, c_m$ . Due to the Tchebychev property of  $\boldsymbol{\omega}$  and the positivity of  $\rho$ , its base determinant

$$D := \begin{vmatrix} \boldsymbol{\omega}(\xi_0) & \int_{\xi_0}^{\xi_1} \boldsymbol{\omega} \rho & \boldsymbol{\omega}(\xi_1) & \cdots & \int_{\xi_{m-1}}^{\xi_m} \boldsymbol{\omega} \rho & \boldsymbol{\omega}(\xi_m) \end{vmatrix}$$

is positive. Therefore, the system has a unique solution  $(c_0, c_1^*, c_1, \dots, c_m^*, c_m)$ . On the other hand, representation (3.3) shows that  $(\alpha_0, -1, \alpha_1, \dots, -1, \alpha_m)$  is also a solution. Thus,  $\alpha_0, \alpha_1, \dots, \alpha_n$  can be obtained by Cramer's Rule:

$$\alpha_0 = \frac{1}{D} \begin{vmatrix} \int_{\xi_0}^{\xi_1} \boldsymbol{\omega} \rho & \boldsymbol{\omega}(\xi_1) & \cdots & \int_{\xi_{m-1}}^{\xi_m} \boldsymbol{\omega} \rho & \boldsymbol{\omega}(\xi_m) & \int_{\xi_m}^{\xi_{m+1}} \boldsymbol{\omega} \rho \end{vmatrix}$$

$$\alpha_k = \frac{1}{D} \begin{vmatrix} \boldsymbol{\omega}(\xi_0) & \cdots & \int_{\xi_{k-1}}^{\xi_k} \boldsymbol{\omega} \rho & \int_{\xi_k}^{\xi_{k+1}} \boldsymbol{\omega} \rho & \cdots & \int_{\xi_m}^{\xi_{m+1}} \boldsymbol{\omega} \rho \end{vmatrix}.$$

Suppose now that  $\omega_0 : [a, b] \rightarrow \mathbb{R}$  is generalized  $(2m+1)$ -convex function with respect to  $\boldsymbol{\omega}$ . Then, for all elements  $t_k$  of  $] \xi_k, \xi_{k+1} [$ , the following inequality holds:

$$0 \geq \begin{vmatrix} f(\xi_0) & f(t_0) & \cdots & f(\xi_m) & f(t_m) \\ \boldsymbol{\omega}(\xi_0) & \boldsymbol{\omega}(t_0) & \cdots & \boldsymbol{\omega}(\xi_m) & \boldsymbol{\omega}(t_m) \end{vmatrix}.$$

Multiplying both sides by the positive  $\rho(t_1) \cdots \rho(t_m)$  and integrating on the product  $[\xi_0, \xi_1] \times \cdots \times [\xi_m, \xi_{m+1}]$  with respect to  $(t_0, \dots, t_m)$ , we arrive at the inequality

$$\begin{aligned} 0 &\geq \begin{vmatrix} f(\xi_0) & \int_{\xi_0}^{\xi_1} f\rho & \cdots & f(\xi_m) & \int_{\xi_m}^{\xi_{m+1}} f\rho \\ \omega(\xi_0) & \int_{\xi_0}^{\xi_1} \omega\rho & \cdots & \omega(\xi_m) & \int_{\xi_m}^{\xi_{m+1}} \omega\rho \end{vmatrix} \\ &= \begin{vmatrix} f(\xi_0) & \int_{\xi_0}^{\xi_1} f\rho & \cdots & f(\xi_m) & \int_{\xi_0}^{\xi_1} f\rho + \cdots + \int_{\xi_m}^{\xi_{m+1}} f\rho \\ \omega(\xi_0) & \int_{\xi_0}^{\xi_1} \omega\rho & \cdots & \omega(\xi_m) & \int_{\xi_0}^{\xi_1} \omega\rho + \cdots + \int_{\xi_m}^{\xi_{m+1}} \omega\rho \end{vmatrix} \\ &= \begin{vmatrix} f(\xi_0) & \int_{\xi_0}^{\xi_1} f\rho & \cdots & f(\xi_m) & \int_a^b f\rho \\ \omega(\xi_0) & \int_{\xi_0}^{\xi_1} \omega\rho & \cdots & \omega(\xi_m) & \int_a^b \omega\rho \end{vmatrix}. \end{aligned}$$

Observe that the adjoint determinants of each element  $\int_{\xi_k}^{\xi_{k+1}} f\rho$  in the last expression are equal to zero since their columns are linearly dependent due to (3.3). Therefore, applying the expansion theorem to the first row, it follows that

$$\begin{aligned} 0 &\leq \left| \omega(\xi_0) \int_{\xi_0}^{\xi_1} \omega\rho \quad \omega(\xi_1) \quad \cdots \quad \int_{\xi_{m-1}}^{\xi_m} \omega\rho \quad \omega(\xi_m) \right| \cdot \int_a^b f\rho \\ &\quad - \left| \int_{\xi_0}^{\xi_1} \omega\rho \quad \omega(\xi_1) \quad \cdots \quad \int_{\xi_{m-1}}^{\xi_m} \omega\rho \quad \omega(\xi_m) \quad \int_a^b \omega\rho \right| f(\xi_0) \\ &\quad - \sum_{k=1}^m \left| \omega(\xi_0) \quad \cdots \quad \int_{\xi_{k-1}}^{\xi_k} \omega\rho \quad \int_{\xi_k}^{\xi_{k+1}} \omega\rho \quad \cdots \quad \int_a^b \omega\rho \right| f(\xi_k). \end{aligned}$$

Here the coefficient of  $\int_a^b f\rho$  is the positive determinant  $D$ , while the coefficients of  $f(\xi_0), \dots, f(\xi_m)$  are exactly the numerators of  $\alpha_0, \dots, \alpha_m$  (see above) since the last column  $\int_a^b \omega$  can be replaced by  $\int_{\xi_m}^{\xi_{m+1}} \omega\rho$ . After rearranging, we get the left hand side of the Hermite–Hadamard-type inequality.  $\square$

**THEOREM 3.6.** *Let  $\omega = (\omega_1, \dots, \omega_{2m})$  be a Tchebychev system on  $[a, b]$  and  $\rho : [a, b] \rightarrow \mathbb{R}$  be a positive integrable function. Then, there exist uniquely determined base points  $\xi_1, \dots, \xi_m$  and  $\eta_1, \dots, \eta_{m-1}$  of  $]a, b[$  such that*

$$\sum_{k=1}^m \alpha_k \omega(\xi_k) = \int_a^b \omega\rho = \beta_0 \omega(a) + \sum_{k=1}^{m-1} \beta_k \omega(\eta_k) + \beta_m \omega(b).$$

*The coefficients  $\alpha_1, \dots, \alpha_m$  and  $\beta_0, \dots, \beta_m$  are positive and uniquely determined, too. Furthermore, for any generalized  $\omega$ -convex function  $f : [a, b] \rightarrow \mathbb{R}$ , the following Hermite–Hadamard-type inequality holds*

$$\sum_{k=1}^m \alpha_k f(\xi_k) \leq \int_a^b f\rho \leq \beta_0 f(a) + \sum_{k=1}^{m-1} \beta_k f(\eta_k) + \beta_m f(b).$$

PROOF. To prove the left hand side inequality, take the unique interior base points  $\xi_1, \dots, \xi_m$  and coefficients  $\alpha_1, \dots, \alpha_m$  fulfilling the representation

$$(3.4) \quad \int_a^b \omega \rho = \sum_{k=1}^m \alpha_k \omega(\xi_k)$$

guaranteed by Theorem 3.4. Defining  $\xi_0 := a$  and  $\xi_{m+1} := b$ , consider the following system of linear equations

$$\int_{\xi_m}^{\xi_{m+1}} \omega \rho = \sum_{k=1}^m \left( c_k^* \int_{\xi_{k-1}}^{\xi_k} \omega \rho + c_k \omega(\xi_k) \right)$$

where the unknowns are  $c_1^*, c_1, \dots, c_m^*, c_m$ . Due to the Tchebychev property of  $\omega$  and the positivity of  $\rho$ , its base determinant

$$D_1 := \begin{vmatrix} \int_{\xi_0}^{\xi_1} \omega \rho & \omega(\xi_1) & \cdots & \int_{\xi_{m-1}}^{\xi_m} \omega \rho & \omega(\xi_m) \end{vmatrix}$$

is positive hence the system has a unique solution  $(c_1^*, c_1, \dots, c_m^*, c_m)$ . On the other hand, the representation (3.4) shows that  $(-1, \alpha_1, \dots, -1, \alpha_m)$  is also a solution. Thus, the coefficients can be obtained by Cramer's Rule:

$$\begin{aligned} \alpha_1 &= \frac{1}{D_1} \begin{vmatrix} \int_{\xi_0}^{\xi_1} \omega \rho & \int_{\xi_1}^{\xi_2} \omega \rho & \cdots & \int_{\xi_{m-1}}^{\xi_m} \omega \rho & \omega(\xi_m) & \int_{\xi_m}^{\xi_{m+1}} \omega \rho \end{vmatrix} \\ \alpha_k &= \frac{1}{D_1} \begin{vmatrix} \int_{\xi_0}^{\xi_1} \omega \rho & \cdots & \int_{\xi_{k-1}}^{\xi_k} \omega \rho & \int_{\xi_k}^{\xi_{k+1}} \omega \rho & \cdots & \int_{\xi_m}^{\xi_{m+1}} \omega \rho \end{vmatrix}. \end{aligned}$$

Suppose now that  $f : [a, b] \rightarrow \mathbb{R}$  is a generalized  $(2m)$ -convex function with respect to  $\omega$ . Then, for all elements  $t_k$  of  $] \xi_k, \xi_{k+1} [$ , the following inequality holds:

$$0 \leq \begin{vmatrix} f(t_0) & f(\xi_1) & \cdots & f(\xi_m) & f(t_m) \\ \omega(t_0) & \omega(\xi_1) & \cdots & \omega(\xi_m) & \omega(t_m) \end{vmatrix}.$$

Therefore,

$$\begin{aligned} 0 &\leq \begin{vmatrix} \int_{\xi_0}^{\xi_1} f \rho & f(\xi_1) & \cdots & f(\xi_m) & \int_{\xi_m}^{\xi_{m+1}} f \rho \\ \int_{\xi_0}^{\xi_1} \omega \rho & \omega(\xi_1) & \cdots & \omega(\xi_m) & \int_{\xi_m}^{\xi_{m+1}} \omega \rho \end{vmatrix} \\ &= \begin{vmatrix} \int_{\xi_0}^{\xi_1} f \rho & f(\xi_1) & \cdots & f(\xi_m) & \int_{\xi_0}^{\xi_1} f \rho + \cdots + \int_{\xi_m}^{\xi_{m+1}} f \rho \\ \int_{\xi_0}^{\xi_1} \omega \rho & \omega(\xi_1) & \cdots & \omega(\xi_m) & \int_{\xi_0}^{\xi_1} \omega \rho + \cdots + \int_{\xi_m}^{\xi_{m+1}} \omega \rho \end{vmatrix} \\ &= \begin{vmatrix} \int_{\xi_0}^{\xi_1} f \rho & f(\xi_1) & \cdots & f(\xi_m) & \int_a^b f \rho \\ \int_{\xi_0}^{\xi_1} \omega \rho & \omega(\xi_1) & \cdots & \omega(\xi_m) & \int_a^b \omega \rho \end{vmatrix}. \end{aligned}$$

In the last expression, the adjoint determinant of each element  $\int_{\xi_k}^{\xi_{k+1}} f \rho$  are equal to zero since their columns are linearly dependent due to (3.4). Applying the expansion theorem to the first row, we arrive at the inequality

$$\begin{aligned} 0 \leq & \left| \int_{\xi_0}^{\xi_1} \omega \rho \quad \omega(\xi_1) \quad \cdots \quad \int_{\xi_{m-1}}^{\xi_m} \omega \rho \quad \omega(\xi_m) \right| \cdot \int_a^b f \rho \\ & - \left| \int_{\xi_0}^{\xi_1} \omega \rho \quad \int_{\xi_1}^{\xi_2} \omega \rho \quad \cdots \quad \int_{\xi_{m-1}}^{\xi_m} \omega \rho \quad \omega(\xi_m) \quad \int_a^b \omega \rho \right| f(\xi_1) \\ & - \sum_{k=2}^m \left| \int_{\xi_0}^{\xi_1} \omega \rho \quad \cdots \quad \int_{\xi_{k-1}}^{\xi_k} \omega \rho \quad \int_{\xi_k}^{\xi_{k+1}} \omega \rho \quad \cdots \quad \int_a^b \omega \rho \right| f(\xi_k). \end{aligned}$$

Here the coefficient of  $\int_a^b f \rho$  is the positive  $D_1$ ; moreover, the coefficients of  $f(\xi_1), \dots, f(\xi_m)$  are exactly the numerators of  $\alpha_1, \dots, \alpha_m$  since the last column  $\int_a^b \omega \rho$  can be replaced by  $\int_{\xi_m}^{\xi_{m+1}} \omega \rho$ . After rearranging, we get the left hand side of the Hermite–Hadamard-type inequality.

For the right hand side inequality, take the uniquely determined interior base points  $\eta_1, \dots, \eta_{m-1}$  and coefficients  $\beta_0, \dots, \beta_m$  so that the representation

$$(3.5) \quad \int_a^b \omega \rho = \beta_0 \omega(a) + \sum_{k=1}^{m-1} \beta_k \omega(\eta_k) + \beta_m \omega(b)$$

hold. Defining  $\eta_0 := a$  and  $\eta_m := b$ , consider the following system of linear equations

$$\int_{\eta_{m-1}}^{\eta_m} \omega \rho = c_0 \omega(\eta_0) + \sum_{k=1}^{m-1} \left( c_k^* \int_{\eta_{k-1}}^{\eta_k} \omega \rho + c_k \omega(\eta_k) \right) + c_m \omega(\eta_m)$$

where the unknowns are  $c_0, c_1^*, c_1, \dots, c_{m-1}^*, c_{m-1}, c_m$ . Due to the Tchebychev property of  $\omega$  and the positivity of  $\rho$ , its base determinant

$$D_2 := \left| \omega(\eta_0) \quad \int_{\eta_0}^{\eta_1} \omega \rho \quad \omega(\eta_1) \quad \cdots \quad \int_{\eta_{m-2}}^{\eta_{m-1}} \omega \rho \quad \omega(\eta_{m-1}) \quad \omega(\eta_m) \right|$$

is positive hence the system has a unique solution  $c_0, c_1^*, c_1, \dots, c_{m-1}^*, c_{m-1}, c_m$ . The representation (3.5) shows that  $(\beta_0, -1, \beta_1, \dots, \beta_{m-1}, -1, \beta_m)$  is also a solution, therefore Cramer's Rule can be applied:

$$\begin{aligned} \beta_0 &= \frac{1}{D_2} \left| \int_{\eta_0}^{\eta_1} \omega \rho \quad \omega(\eta_1) \quad \cdots \quad \omega(\eta_{m-1}) \quad \int_{\eta_{m-1}}^{\eta_m} \omega \rho \quad \omega(\eta_m) \right| \\ \beta_k &= \frac{1}{D_2} \left| \omega(\eta_0) \quad \cdots \quad \int_{\eta_{k-1}}^{\eta_k} \omega \rho \quad \int_{\eta_k}^{\eta_{k+1}} \omega \rho \quad \cdots \quad \int_{\eta_{m-1}}^{\eta_m} \omega \rho \quad \omega(\eta_m) \right| \\ \beta_m &= \frac{1}{D_2} \left| \omega(\eta_0) \quad \int_{\eta_0}^{\eta_1} \omega \rho \quad \cdots \quad \int_{\eta_{m-2}}^{\eta_{m-1}} \omega \rho \quad \omega(\eta_{m-1}) \quad \int_{\eta_{m-1}}^{\eta_m} \omega \rho \right|. \end{aligned}$$

These coefficients are positive since even changes are needed to transfer the column  $\int_{\eta_{m-1}}^{\eta_m} \omega \rho$  to the adequate place.

If a function  $f : [a, b] \rightarrow \mathbb{R}$  is a generalized  $(2m)$ -convex with respect to  $\boldsymbol{\omega}$ , then we arrive at the inequality

$$0 \leq \begin{vmatrix} f(\eta_0) & \int_{\eta_0}^{\eta_1} f\rho & f(\eta_1) & \cdots & \int_{\eta_{m-2}}^{\eta_{m-1}} f\rho & f(\eta_m) & \int_a^b f\rho \\ \boldsymbol{\omega}(\eta_0) & \int_{\eta_0}^{\eta_1} \boldsymbol{\omega}\rho & \boldsymbol{\omega}(\eta_1) & \cdots & \int_{\eta_{m-2}}^{\eta_{m-1}} \boldsymbol{\omega}\rho & \boldsymbol{\omega}(\eta_m) & \int_a^b \boldsymbol{\omega}\rho \end{vmatrix},$$

whence an analogous argument to the previous one completes the proof.  $\square$

### 3.4. An alternative approach in a particular case

To prove the main results, the main point is the existence of the representations of Theorem D. These representations can also be considered as systems of nonlinear equations where the unknowns are the coefficients and the base points. The number of the equations and the unknowns coincide in each case. In those cases when only one interior base point is involved, the solubility of the system of equations can directly be verified without applying the Krein–Markov theory of moment spaces.

**THEOREM 3.7.** *Let  $\boldsymbol{\omega} = (\omega_1, \omega_2, \omega_3)$  be a Tchebychev system on  $[a, b]$  and  $\rho : [a, b] \rightarrow \mathbb{R}$  be a positive integrable function. Then, there exist unique elements  $\xi, \eta$  of  $]a, b[$  and uniquely determined positive coefficients  $c_1, c_2$  and  $d_1, d_2$  such that*

$$c_1 \boldsymbol{\omega}(a) + c_2 \boldsymbol{\omega}(\xi) = \int_a^b \boldsymbol{\omega}\rho = d_1 \boldsymbol{\omega}(\eta) + d_2 \boldsymbol{\omega}(b).$$

Furthermore, if a function  $f : [a, b] \rightarrow \mathbb{R}$  is generalized 3-convex with respect to  $\boldsymbol{\omega}$ , then the following Hermite–Hadamard-type inequality holds

$$c_1 f(a) + c_2 f(\xi) \leq \int_a^b f\rho \leq d_1 f(\eta) + d_2 f(b).$$

**PROOF.** We shall restrict the process of the proof only on the existence of the interior point  $\xi$ . To do this, define the function  $F : [a, b] \rightarrow \mathbb{R}$  by the formula

$$F(x) := \begin{vmatrix} \boldsymbol{\omega}(a) & \int_a^x \boldsymbol{\omega}\rho & \int_a^b \boldsymbol{\omega}\rho \end{vmatrix} := \begin{vmatrix} \omega_1(a) & \int_a^x \omega_1\rho & \int_a^b \omega_1\rho \\ \omega_2(a) & \int_a^x \omega_2\rho & \int_a^b \omega_2\rho \\ \omega_3(a) & \int_a^x \omega_3\rho & \int_a^b \omega_3\rho \end{vmatrix}.$$

Then,  $F$  is continuous on  $[a, b]$  and  $F(a) = F(b) = 0$ . Further on,  $F(x) \neq 0$  if  $x \in ]a, b[$  due to the Tchebychev property of  $\boldsymbol{\omega}$  and the positivity of  $\rho$ . For simplicity, we may assume that  $F$  is positive on  $]a, b[$ . Therefore, by Weierstrass' theorem, there exists  $\xi \in ]a, b[$  such that

$$F(\xi) = \max_{[a, b]} F.$$

Assume that  $x \in ]\xi, b]$ . Then, the maximal property of  $\xi$  yields the inequality

$$0 \geq \frac{F(x) - F(\xi)}{\int_\xi^x \rho} = \begin{vmatrix} \boldsymbol{\omega}(a) & \frac{\int_\xi^x \boldsymbol{\omega}\rho}{\int_\xi^x \rho} & \int_a^b \boldsymbol{\omega}\rho \end{vmatrix}.$$

The central column of the determinant tends to  $\omega(\xi)$  as  $x$  tends to  $\xi$  since the following estimations are valid for  $k = 1, 2, 3$ :

$$\min_{[\xi, x]} \omega_k = \frac{\min_{[\xi, x]} \omega_k \int_{\xi}^x \rho}{\int_{\xi}^x \rho} \leq \frac{\int_{\xi}^x \omega_k \rho}{\int_{\xi}^x \rho} \leq \frac{\max_{[\xi, x]} \omega_k \int_{\xi}^x \rho}{\int_{\xi}^x \rho} = \max_{[\xi, x]} \omega_k.$$

Therefore

$$\left| \begin{array}{ccc} \omega(a) & \omega(\xi) & \int_a^b \omega \rho \end{array} \right| \leq 0.$$

Choosing  $x \in [a, \xi[$  and using the maximal property of  $\xi$  again, we get the opposite inequality with the same argument and arrive at the identity

$$\left| \begin{array}{ccc} \omega(a) & \omega(\xi) & \int_a^b \omega \rho \end{array} \right| = 0.$$

Thus, the linear independence of  $\omega(a)$  and  $\omega(\xi)$  yields that there exist coefficients  $c_1$  and  $c_2$  such that

$$c_1 \omega(a) + c_2 \omega(\xi) = \int_a^b \omega \rho.$$

The right hand side inequality can be verified with an analogous argument, therefore the proof is omitted.  $\square$

Let us note, that if the weight function  $\rho$  is continuous, then the function  $F$  is differentiable and Rolle's mean-value theorem can directly be applied.

The representations of Theorem 3.7 are linear with respect to the coefficients. Therefore, in concrete cases, the main difficulty is to determine the interior base points  $\xi$  and  $\eta$ . Without claiming completeness, we list some examples when they can be determined explicitly.

**EXAMPLE 1.** If the Tchebychev system  $(\omega_1, \omega_2, \omega_3)$  is defined on  $[a, b]$  by  $\omega_1(x) = 1$ ,  $\omega_2(x) = \sinh x$ ,  $\omega_3(x) = \cosh x$  and  $\rho \equiv 1$ , then

$$\begin{aligned} \xi &= 2 \operatorname{artanh} \left( \frac{\sinh b - \sinh a - (b-a) \cosh a}{\cosh b - \cosh a - (b-a) \sinh a} \right) - a \\ \eta &= 2 \operatorname{artanh} \left( \frac{\sinh b - \sinh a - (b-a) \cosh b}{\cosh b - \cosh a - (b-a) \sinh b} \right) - b. \end{aligned}$$

**PROOF.** With the above setting, the left hand side representation of Theorem 3.5 reduces to the following system of nonlinear equations

$$\begin{aligned} c_1 + c_2 &= \int_a^b 1 dx = b - a \\ c_1 \sinh a + c_2 \sinh \xi &= \int_a^b \sinh x dx = \cosh b - \cosh a \\ c_1 \cosh a + c_2 \cosh \xi &= \int_a^b \cosh x dx = \sinh b - \sinh a \end{aligned}$$

where the three unknowns are  $c_1$ ,  $c_2$  and  $\xi$ , respectively. Multiplying the first equation by  $\sinh a$  and subtracting it from the second one, then multiplying again the first equation by  $\cosh a$  and subtracting it from the third one, the coefficient  $c_1$  can be eliminated and it follows

$$\begin{aligned} c_2(\sinh \xi - \sinh a) &= \cosh b - \cosh a - (b - a) \sinh a \\ c_2(\cosh \xi - \cosh a) &= \sinh b - \sinh a - (b - a) \cosh a. \end{aligned}$$

Applying the well known additional properties of hyperbolic functions for the identities  $\xi = (\xi + a)/2 + (\xi - a)/2$  and  $a = (\xi + a)/2 - (\xi - a)/2$ , the left hand side of both equations can be written into product form:

$$\begin{aligned} 2c_2 \cosh\left(\frac{\xi + a}{2}\right) \sinh\left(\frac{\xi - a}{2}\right) &= \cosh b - \cosh a - (b - a) \sinh a \\ 2c_2 \sinh\left(\frac{\xi + a}{2}\right) \sinh\left(\frac{\xi - a}{2}\right) &= \sinh b - \sinh a - (b - a) \cosh a. \end{aligned}$$

The left hand side of the first equation differs from zero since  $\xi \neq a$ . Therefore, dividing the second equation by the first one, we get the equation

$$\tanh\left(\frac{\xi + a}{2}\right) = \frac{\sinh b - \sinh a - (b - a) \cosh a}{\cosh b - \cosh a - (b - a) \sinh a},$$

whence the desired expression of  $\xi$  is obtained. For determining  $\eta$ , we shall consider the following system of nonlinear equation:

$$\begin{aligned} d_1 + d_2 &= b - a \\ d_1 \sinh \eta + d_2 \sinh b &= \cosh b - \cosh a \\ d_1 \cosh \eta + d_2 \cosh b &= \sinh b - \sinh a. \end{aligned}$$

In this case, the coefficient  $d_2$  can be eliminated with a similar method to the previous one. The new system of equation, due to the additional formulae again, can be written into the form

$$\begin{aligned} 2d_1 \cosh\left(\frac{b + \eta}{2}\right) \sinh\left(\frac{b - \eta}{2}\right) &= \cosh b - \cosh a - (b - a) \sinh b \\ 2d_1 \sinh\left(\frac{b + \eta}{2}\right) \sinh\left(\frac{b - \eta}{2}\right) &= \sinh b - \sinh a - (b - a) \cosh b. \end{aligned}$$

This system, analogously to the previous case, yields the equation

$$\tanh\left(\frac{b + \eta}{2}\right) = \frac{\sinh b - \sinh a - (b - a) \cosh b}{\cosh b - \cosh a - (b - a) \sinh b},$$

whence the base point  $\eta$  can be expressed easily.  $\square$

The proofs of the subsequent examples are similar to the previous one, therefore they are omitted.

EXAMPLE 2. If the Tchebychev system  $(\omega_1, \omega_2, \omega_3)$  is defined on  $[a, b] \subset ] - \pi, \pi[$  by  $\omega_1(x) = 1$ ,  $\omega_2(x) = \sin x$ ,  $\omega_3(x) = \cos x$  and  $\rho \equiv 1$ , then

$$\begin{aligned}\xi &= 2 \arctan \left( \frac{\sin a - \sin b + (b-a) \cos a}{\cos a - \cos b - (b-a) \sin a} \right) - a \\ \eta &= 2 \arctan \left( \frac{\sin a - \sin b + (b-a) \cos b}{\cos a - \cos b - (b-a) \sin b} \right) - b.\end{aligned}$$

EXAMPLE 3. If the Tchebychev system  $(\omega_1, \omega_2, \omega_3)$  is defined on  $[a, b]$  by  $\omega_1(x) = 1$ ,  $\omega_2(x) = \exp x$ ,  $\omega_3(x) = \exp 2x$  and  $\rho \equiv 1$ , then

$$\begin{aligned}\xi &= \log \left( \frac{\exp 2b - \exp 2a - 2(b-a) \exp 2a}{2(\exp b - \exp a - (b-a) \exp a)} - \exp a \right) \\ \eta &= \log \left( \frac{\exp 2b - \exp 2a - 2(b-a) \exp 2b}{2(\exp b - \exp a - (b-a) \exp b)} - \exp b \right).\end{aligned}$$

EXAMPLE 4. If, for  $p > 0$ , the Tchebychev system  $(\omega_1, \omega_2, \omega_3)$  is defined on  $[a, b] \subset [0, +\infty[$  by  $\omega_1(x) = 1$ ,  $\omega_2(x) = x^p$ ,  $\omega_3(x) = x^{2p}$  and  $\rho \equiv 1$ , then

$$\begin{aligned}\xi &= \left( \frac{p+1}{2p+1} \cdot \frac{b^{2p+1} - a^{2p+1} - (2p+1)(b-a)a^{2p}}{b^{p+1} - a^{p+1} - (p+1)(b-a)a^p} - a^p \right)^{1/p} \\ \eta &= \left( \frac{p+1}{2p+1} \cdot \frac{b^{2p+1} - a^{2p+1} - (2p+1)(b-a)b^{2p}}{b^{p+1} - a^{p+1} - (p+1)(b-a)b^p} - b^p \right)^{1/p}.\end{aligned}$$

The particular case  $p = 1$  of the last example gives a corollary of Theorem 1.10 for polynomially 3-convex functions. For 3 dimensional Tchebychev systems generated by arbitrary power functions, the interior base points in general, cannot be expressed explicitly.

The proof of Theorem 3.7 is applicable for generalized 2-convexity, and gives a different approach followed in Theorem 2.10. We can also state right hand side Hermite–Hadamard-type inequality for generalized 4-convex functions.

THEOREM 3.8. Let  $\omega = (\omega_1, \omega_2, \omega_3, \omega_4)$  be a Tchebychev system on  $[a, b]$  and  $\rho : [a, b] \rightarrow \mathbb{R}$  be a positive integrable function. Then, there exist a unique element  $\xi$  of  $]a, b[$  and uniquely determined positive coefficients  $c_1, c_2, c_3$  such that

$$\int_a^b \omega \rho = c_1 \omega(a) + c_2 \omega(\xi) + c_3 \omega(b).$$

Furthermore, if a function  $f : [a, b] \rightarrow \mathbb{R}$  is generalized 4-convex with respect to  $\omega$ , then the following Hermite–Hadamard-type inequality holds

$$\int_a^b f \rho \leq c_1 f(a) + c_2 f(\xi) + c_3 f(b).$$

HINT. Apply the same argument as in the proof of Theorem 3.7 for the function  $F : [a, b] \rightarrow \mathbb{R}$  defined by the formula

$$F(x) := \left| \begin{array}{cccc} \omega_1(a) & \int_a^x \omega_1 \rho & \omega_1(b) & \int_a^b \omega_1 \rho \\ \omega_2(a) & \int_a^x \omega_2 \rho & \omega_2(b) & \int_a^b \omega_2 \rho \\ \omega_3(a) & \int_a^x \omega_3 \rho & \omega_3(b) & \int_a^b \omega_3 \rho \\ \omega_4(a) & \int_a^x \omega_4 \rho & \omega_4(b) & \int_a^b \omega_4 \rho \end{array} \right|.$$

□

For example, if  $\omega(x) := (\cosh x, \sinh x, \cosh 2x, \sinh 2x)$ , then one can check that the interior base point of the inequality is exactly the midpoint of the domain. Unfortunately, the method fails if someone tries to use it for proving left hand side Hermite–Hadamard-type inequality for a generalized 4-convex function since, by the even case of Theorem D, the existence of two interior base points should be guaranteed. For similar reasons, the “existence” part in the proof of Theorem 3.7 cannot be applied for generalized  $n$ -convex functions if  $n > 4$ .



## Characterizations via Hermite–Hadamard inequalities

Under some weak regularity conditions, the Hermite–Hadamard-inequality characterizes (standard) convexity (see [Kuc85, Excercise 8. p. 205]). The aim of this chapter is to verify analogous results for  $(\omega_1, \omega_2)$ -convexity. To do this, the most important auxiliary tool turns out to be some characterization properties of continuous, *non* generalized 2-convex functions.

### 4.1. Further properties of generalized lines

In the further investigations, two properties of generalized lines are crucial. The first one improves the statement of Lemma 2.2 and states that, on compact intervals, generalized lines are uniformly non bounded.

LEMMA 4.1. *Let  $(\omega_1, \omega_2)$  be a Tchebychev system on an interval  $I$ . Then, for any compact subinterval of  $I$  and positive number  $K$ , there exists  $\omega \in \mathcal{L}(\omega_1, \omega_2)$  such that  $\omega > K$  on the compact subinterval.*

PROOF. According to Lemma 2.2, there exist coefficients  $\alpha, \beta$  such that the generalized line  $\alpha\omega_1 + \beta\omega_2$  is positive on the interior of  $I$ . Therefore, if  $[x, y]$  is a compact subinterval of  $I$ ,  $m := \min\{\alpha\omega_1(t) + \beta\omega_2(t) \mid t \in [x, y]\} > 0$ . Defining the coefficients  $\alpha^*$  and  $\beta^*$  by the formulae

$$\alpha^* := \frac{\alpha K}{m} \quad \beta^* := \frac{\beta K}{m},$$

the generalized line  $\omega := \alpha^*\omega_1 + \beta^*\omega_2$  is strictly greater than  $K$  on  $[x, y]$ .  $\square$

The second important property concerns the convergence of generalized lines. It turns out that pointwise convergence is not only a necessary but a sufficient condition for the uniform convergence of sequences of generalized lines. Let us note that an analogous result remains true for generalized polynomials in the higher-order case.

LEMMA 4.2. *Let  $(\omega_1, \omega_2)$  be a Tchebychev system on an interval  $I$ , furthermore  $\omega = \alpha\omega_1 + \beta\omega_2$  and  $\omega_n = \alpha_n\omega_1 + \beta_n\omega_2$  ( $n \in \mathbb{N}$ ) be generalized lines. Then, the following statements are equivalent:*

- (i) *there exist elements  $x < y$  of  $I$  such that  $\omega_n(x) \rightarrow \omega(x)$  and  $\omega_n(y) \rightarrow \omega(y)$ ;*
- (ii) *the sequences  $\alpha_n$  and  $\beta_n$  are convergent furthermore  $\alpha_n \rightarrow \alpha$  and  $\beta_n \rightarrow \beta$ ;*
- (iii)  *$\omega_n \rightarrow \omega$  uniformly on each compact subset of  $I$ .*

PROOF. (i)  $\Rightarrow$  (ii). Applying Cramer's Rule and the convergence properties of  $\omega_n(x)$  and  $\omega_n(y)$ , one can easily get that

$$\alpha = \frac{\begin{vmatrix} \omega(x) & \omega_2(x) \\ \omega(y) & \omega_2(y) \end{vmatrix}}{\begin{vmatrix} \omega_1(x) & \omega_2(x) \\ \omega_1(y) & \omega_2(y) \end{vmatrix}} = \lim_{n \rightarrow \infty} \frac{\begin{vmatrix} \omega_n(x) & \omega_2(x) \\ \omega_n(y) & \omega_2(y) \end{vmatrix}}{\begin{vmatrix} \omega_1(x) & \omega_2(x) \\ \omega_1(y) & \omega_2(y) \end{vmatrix}} = \lim_{n \rightarrow \infty} \alpha_n.$$

The convergence of  $\beta_n$  can be obtained similarly.

(ii)  $\Rightarrow$  (iii). Let  $[x, y]$  be a compact subinterval of  $I$ , furthermore  $t \in [x, y]$  arbitrary. Due to the continuity of the functions  $\omega_1$  and  $\omega_2$ , there exists  $K > 0$  such that

$$\max \left\{ \sup_{[x,y]} |\omega_1(t)|, \sup_{[x,y]} |\omega_2(t)| \right\} \leq K.$$

Therefore,

$$\begin{aligned} |\omega_n(t) - \omega(t)| &= |\alpha_n \omega_1(t) - \alpha \omega_1(t) + \beta_n \omega_2(t) - \beta \omega_2(t)| \\ &\leq |\alpha_n - \alpha| |\omega_1(t)| + |\beta_n - \beta| |\omega_2(t)| \\ &\leq K (|\alpha_n - \alpha| + |\beta_n - \beta|) \rightarrow 0 \end{aligned}$$

as  $n \rightarrow \infty$ ; hence  $\omega_n \rightarrow \omega$  uniformly on  $[x, y]$ .

(iii)  $\Rightarrow$  (i). Trivial.  $\square$

Under the assumption of continuity, if a function is not convex, then it must be locally strictly concave somewhere. The following theorem generalizes this result for non  $(\omega_1, \omega_2)$ -convexity.

**THEOREM 4.3.** *Let  $(\omega_1, \omega_2)$  be a Tchebychev system on an interval  $I$ , furthermore  $f : I \rightarrow \mathbb{R}$  be a continuous function. Then, the following assertions are equivalent:*

- (i)  $f$  is not  $(\omega_1, \omega_2)$ -convex;
- (ii) there exist elements  $x < y$  of  $I$  such that  $\omega < f$  on  $]x, y[$  where  $\omega$  is the generalized line determined by the properties

$$\omega(x) = f(x) \quad \omega(y) = f(y);$$

- (iii) there exist elements  $x < p < y$  of  $I$  and a generalized line  $\omega$  such that  $\omega \geq f$  on  $[x, y]$ , moreover

$$f(x) < \omega(x) \quad f(p) = \omega(p) \quad f(y) < \omega(y);$$

- (iv) there exists  $p \in I^\circ$  such that  $f$  is locally strictly  $(\omega_1, \omega_2)$ -concave at  $p$ , that is, there exist elements  $x < p < y$  of  $I$  such that, for all  $x < u < p < v < y$ , the following inequality holds:

$$\begin{vmatrix} f(u) & f(p) & f(v) \\ \omega_1(u) & \omega_1(p) & \omega_1(v) \\ \omega_2(u) & \omega_2(p) & \omega_2(v) \end{vmatrix} < 0.$$

PROOF. (i)  $\Rightarrow$  (ii). If  $f$  is not  $(\omega_1, \omega_2)$ -convex, then there exist elements  $x_0 < p < y_0$  of  $I$  such that  $\omega(p) < f(p)$ , where  $\omega$  is the generalized line determined by the properties  $\omega(x_0) = f(x_0)$  and  $\omega(y_0) = f(y_0)$  (see assertion (vi) of Theorem 2.4). Define the function  $F : [x_0, y_0] \rightarrow \mathbb{R}$  by  $F := f - \omega$  furthermore the elements  $x$  and  $y$  by the formulae

$$\begin{aligned} x &:= \sup\{t \mid F(t) = 0, x_0 \leq t < p\} \\ y &:= \sup\{t \mid F(t) = 0, p < t \leq y_0\}. \end{aligned}$$

Clearly,  $x_0 \leq x < p < y \leq y_0$  hold; moreover,  $F(x) = F(y) = 0$  and  $F > 0$  on  $]x, y[$  due to the continuity of  $F$ . That is,  $\omega(x) = f(x)$ ,  $\omega(y) = f(y)$  and  $f(t) > \omega(t)$  for all  $t \in ]x, y[$ .

(ii)  $\Rightarrow$  (iii). Take the elements  $x < y$  of  $I$  and the generalized line  $\omega$  fulfilling the properties  $\omega(x) = f(x)$ ,  $\omega(y) = f(y)$  and  $\omega|_{]x, y[} < f|_{]x, y[}$ . Define, for all  $t \in \mathbb{R}$ , the family of “parallel” generalized lines  $\omega_t$  furthermore the real number  $t_0$  by

$$\omega_t(x) = \omega(x) + t \quad \omega_t(y) = \omega(y) + t$$

Observe first that  $\omega_t|_{[x, y]} > f|_{[x, y]}$  for “sufficiently large”  $t$ . Indeed, take the generalized line  $\omega^*$  satisfying the inequality  $\omega^*|_{[x, y]} > \max f|_{[x, y]}$  and choose  $t > 0$  such that  $\omega_t(x) > \omega^*(x)$  and  $\omega_t(y) > \omega^*(y)$  hold. (The existence of  $\omega^*$  is guaranteed by Lemma 4.1.) Then,  $\omega_t|_{[x, y]} > \omega^*|_{[x, y]}$  due to Lemma 2.1 hence  $\omega_t|_{[x, y]} > f|_{[x, y]}$ . On the other hand, a similar argument to the previous one yields the inequalities  $\omega_t|_{[x, y]} < \omega|_{[x, y]} \leq f|_{[x, y]}$  for all  $t < 0$ . Therefore,

$$t_0 := \inf\{t \in \mathbb{R} \mid \omega_t|_{[x, y]} > f|_{[x, y]}\}.$$

By definition,  $\omega_{t_0} \geq f$  on  $[x, y]$ . Assume indirectly that this inequality is strict. Then, according to the continuity of  $\omega_{t_0}$  and  $f$ , there exists  $\varepsilon > 0$  such that

$$f + \varepsilon < \omega_{t_0}$$

on  $[x, y]$ . Consider the sequence of generalized lines  $\omega_n$  determined by the conditions

$$\omega_n(x) := \omega(x) + t_0 - \frac{1}{n} \quad \omega_n(y) := \omega(y) + t_0 - \frac{1}{n}.$$

Lemma 2.1 implies that  $(\omega_n)$  is strictly monotone increasing; further on, according to Lemma 4.2,  $\omega_n \rightarrow \omega_{t_0}$  uniformly on the compact interval  $[x, y]$  since  $\omega_n(x) \rightarrow \omega_{t_0}(x)$  and  $\omega_n(y) \rightarrow \omega_{t_0}(y)$ . Hence, there exists  $n_0 \in \mathbb{N}$  satisfying the inequalities

$$\omega_{n_0} < \omega_{t_0} < \omega_{n_0} + \frac{\varepsilon}{2}.$$

Comparing this to the previous one, it follows that

$$f + \frac{\varepsilon}{2} < \omega_{n_0} < \omega_{t_0},$$

which contradicts the definition of  $t_0$  since  $\omega_n$  can also be written into the form  $\omega_{t_0 - 1/n}$ . Therefore, the choice  $\omega_{t_0}$  satisfies the requirements.

(iii)  $\Rightarrow$  (iv). Due to the continuity of the functions  $f$  and  $\omega$ , we may assume that  $p$  is the minimal element of  $]x, y[$  fulfilling the properties of the assertion. Then,  $f(u) < \omega(u)$  if  $x < u < p$  and  $f(v) \leq \omega(v)$  if  $p < v < y$ . Therefore,

$$\begin{vmatrix} f(u) & f(p) & f(y) \\ \omega_1(u) & \omega_1(p) & \omega_1(v) \\ \omega_2(u) & \omega_2(p) & \omega_2(v) \end{vmatrix} < \begin{vmatrix} \omega(u) & \omega(p) & \omega(v) \\ \omega_1(u) & \omega_1(p) & \omega_1(v) \\ \omega_2(u) & \omega_2(p) & \omega_2(v) \end{vmatrix}$$

since the adjoint determinants of  $f(u)$  and  $f(v)$  are positive, furthermore,  $f$  and  $\omega$  coincide at  $p$ . However,  $\omega$  is a linear combination of  $\omega_1$  and  $\omega_2$  hence the left hand side of the previous inequality equals zero.

(iv)  $\Rightarrow$  (i). Trivial.  $\square$

The next result shows that  $(\omega_1, \omega_2)$ -convexity, similarly to the standard one, is a pointwise property.

**COROLLARY 4.4.** *Let  $(\omega_1, \omega_2)$  be a Tchebychev system over the open interval  $I$ , furthermore  $f : I \rightarrow \mathbb{R}$  be a given function. Then, the following assertions are equivalent:*

- (i)  $f$  is  $(\omega_1, \omega_2)$ -convex;
- (ii)  $f$  is locally  $(\omega_1, \omega_2)$ -convex, that is, each element of the domain has a neighborhood where it is  $(\omega_1, \omega_2)$ -convex;
- (iii)  $f$  is continuous and, for all  $p \in I$ , there exist elements  $x < p < y$  of  $I$  such that

$$\begin{vmatrix} f(u) & f(p) & f(y) \\ \omega_1(u) & \omega_1(p) & \omega_1(v) \\ \omega_2(u) & \omega_2(p) & \omega_2(v) \end{vmatrix} \geq 0$$

for all  $x < u < p < v < y$  (i. e.,  $f$  is locally convex at each point).

**HINT.** The implications (i)  $\Rightarrow$  (ii) and (ii)  $\Rightarrow$  (iii) are trivial. For the implication (iii)  $\Rightarrow$  (i), the last assertion of Corollary 4.4 can be applied, which, in the case of indirect assumption, immediately leads to contradiction.  $\square$

## 4.2. Hermite–Hadamard-type inequalities and $(\omega_1, \omega_2)$ -convexity

The main results are presented in three theorems. The first and the second ones concern the left and right hand side inequalities of Theorem 2.10 independently, while the third one is analogous to the classical Jensen inequality.

**THEOREM 4.5.** *Let  $(\omega_1, \omega_2)$  be a Tchebychev system on an interval  $[a, b]$  such that  $\omega_1$  is positive on  $]a, b[$ , furthermore  $\rho : [a, b] \rightarrow \mathbb{R}$  be a positive integrable function. Define, for all elements  $x < y$  of  $[a, b]$ , the functions  $\xi(x, y)$  and  $c(x, y)$  by the formulae*

$$\xi(x, y) := \left( \frac{\omega_2}{\omega_1} \right)^{-1} \left( \frac{\int_x^y \omega_2 \rho}{\int_x^y \omega_1 \rho} \right), \quad c(x, y) = \frac{\int_x^y \omega_1 \rho}{\omega_1(\xi(x, y))}.$$

Then, a continuous function  $f : [a, b] \rightarrow \mathbb{R}$  is generalized convex with respect to  $(\omega_1, \omega_2)$  if and only if, for all elements  $x < y$  of  $[a, b]$ , it satisfies the inequality

$$c(x, y)f(\xi(x, y)) \leq \int_x^y f \rho.$$

PROOF. The necessity is due to Theorem 2.10. For the converse assertion, note first that the mapping  $(x, y) \mapsto \xi(x, y)$  is continuous in each variable and takes its value between  $x$  and  $y$  since it is a Lagrange-type mean-value. Further on,  $c(x, y)$  and  $\xi(x, y)$  are so constructed that all generalized lines (i.e., the linear combinations of  $\omega_1$  and  $\omega_2$ ) are the solutions of the functional equation

$$(4.1) \quad c(x, y)\omega(\xi(x, y)) = \int_x^y \omega \rho \quad (x < y).$$

(For the details, see the proof of Theorem 2.10.) Assume that  $f$  satisfies the inequality of the theorem and, indirectly, is not  $(\omega_1, \omega_2)$ -convex. Then, according to assertion (iii) Theorem 4.3, there exist elements  $x < p < y$  of  $I$  and a generalized line  $\omega$  such that  $f \leq \omega$  on  $[x, y]$  and

$$f(x) < \omega(x) \quad f(p) = \omega(p) \quad f(y) < \omega(y).$$

If, for example,  $p \leq \xi(x, y)$ , then there is  $u \in ]p, y]$  such that  $p = \xi(x, u)$  since  $\xi$  is a Lagrange-type mean-value. The inequality  $f(x) < \omega(x)$  and the continuity of  $f$  implies that  $f < \omega$  on a right hand side neighborhood of  $x$  hence, applying (4.1), it follows that

$$c(x, u)f(\xi(x, u)) \leq \int_x^u f \rho < \int_x^u \omega \rho = c(x, u)\omega(\xi(x, u)).$$

On the other hand, both sides have the common value  $c(x, u)f(p)$ , which is a contradiction.  $\square$

THEOREM 4.6. Let  $(\omega_1, \omega_2)$  be a Tchebychev system over an interval  $[a, b]$  such that  $\omega_1$  is positive on  $]a, b[$ , furthermore  $\rho : [a, b] \rightarrow \mathbb{R}$  be a positive integrable function. Define, for all elements  $x < y$  of  $[a, b]$ , the functions  $c_1(x, y)$  and  $c_2(x, y)$  by the formulae

$$c_1(x, y) = \frac{\begin{vmatrix} \int_x^y \omega_1 \rho & \omega_1(y) \\ \int_x^y \omega_2 \rho & \omega_2(y) \end{vmatrix}}{\begin{vmatrix} \omega_1(x) & \omega_1(y) \\ \omega_2(x) & \omega_2(y) \end{vmatrix}}, \quad c_2(x, y) = \frac{\begin{vmatrix} \omega_1(x) & \int_x^y \omega_1 \rho \\ \omega_2(x) & \int_x^y \omega_2 \rho \end{vmatrix}}{\begin{vmatrix} \omega_1(x) & \omega_1(y) \\ \omega_2(x) & \omega_2(y) \end{vmatrix}}.$$

Then, a continuous function  $f : [a, b] \rightarrow \mathbb{R}$  is generalized convex with respect to  $(\omega_1, \omega_2)$  if and only if, for all elements  $x < y$  of  $[a, b]$ , it satisfies the inequality

$$\int_x^y f \rho \leq c_1(x, y)f(x) + c_2(x, y)f(y).$$

PROOF. The necessity is due to Theorem 2.10 again. Conversely, note first that  $c_1(x, y)$  and  $c_2(x, y)$  are constructed such that all generalized lines (i. e., the linear combinations of  $\omega_1$  and  $\omega_2$ ) are the solutions of the functional equation

$$(4.2) \quad \int_x^y \omega \rho = c_1(x, y)\omega(x) + c_2(x, y)\omega(y).$$

(For the details, see the proof of Theorem 2.10.) Assume indirectly that  $f$  is not  $(\omega_1, \omega_2)$ -convex. Then, according to assertion (ii) of Theorem 4.3, there exist elements  $x < y$  of  $I$  and a generalized line  $\omega$  such that  $\omega(x) = f(x)$ ,  $\omega(y) = f(y)$  and  $\omega < f$  on  $]x, y[$ . Therefore,

$$\begin{aligned} \int_x^y \omega \rho < \int_x^y f \rho &\leq c_1(x, y)f(x) + c_2(x, y)f(y) \\ &= c_1(x, y)\omega(x) + c_2(x, y)\omega(y) \end{aligned}$$

which contradicts (4.2).  $\square$

THEOREM 4.7. *Let  $(\omega_1, \omega_2)$  be a Tchebychev system on  $I$  and  $f : I \rightarrow \mathbb{R}$  be a continuous function. Keeping the notions of Theorem 4.6 and Theorem 4.5,  $f$  is  $(\omega_1, \omega_2)$ -convex if and only if, for all elements  $x < y$  of  $I$ , it satisfies the inequality*

$$c(x, y)f(\xi(x, y)) \leq c_1(x, y)f(x) + c_2(x, y)f(y).$$

PROOF. The necessity part has already been proved in Theorem 2.10. For the sufficiency, observe first that the functions  $c$ ,  $c_1$ ,  $c_2$  and  $\xi$  are so constructed that all the generalized lines are the solutions of the functional equation

$$c(x, y)\omega(\xi(x, y)) = c_1(x, y)\omega(x) + c_2(x, y)\omega(y) \quad (x < y)$$

since both sides have the common value  $\int_x^y \omega \rho$ . Assume indirectly that a function  $f : I \rightarrow \mathbb{R}$  satisfies the inequality of the theorem and is not generalized convex with respect to  $(\omega_1, \omega_2)$ . Then, there exist elements  $x < y$  of  $I$  and a generalized line  $\omega$  fulfilling the conditions

$$\omega(x) = f(x) \quad \omega|_{]x, y[} < f|_{]x, y[} \quad \omega(y) = f(y)$$

due to Theorem 4.3. Therefore, taking the above observation into consideration, one can immediately get that

$$\begin{aligned} c(x, y)f(\xi(x, y)) &\leq c_1(x, y)f(x) + c_2(x, y)f(y) \\ &= c_1(x, y)\omega(x) + c_2(x, y)\omega(y) \\ &= c(x, y)\omega(\xi(x, y)) < c(x, y)f(\xi(x, y)) \end{aligned}$$

which is a contradiction.  $\square$

To give a unified view, the previous results are combined in the subsequent corollary. This corollary, Theorem 2.4, Corollary 2.6, Theorem 2.7 and Corollary 4.4 together are a comprehensive characterization of generalized convexity induced by two dimensional Tchebychev systems.

COROLLARY 4.8. *Let  $(\omega_1, \omega_2)$  be a Tchebychev system on  $I$  such that  $\omega_1$  is positive on  $I^\circ$ , furthermore  $\rho : I \rightarrow \mathbb{R}$  be a positive integrable function. Keeping the notions of Theorem 4.5, Theorem 4.6 and Theorem 4.7, the following assertions are equivalent for any function  $f : I \rightarrow \mathbb{R}$ :*

- (i)  *$f$  is generalized convex with respect to  $(\omega_1, \omega_2)$ ;*
- (ii)  *$f$  is continuous and, for all elements  $x < y$  of  $I$ , satisfies the inequality*

$$c(x, y)f(\xi(x, y)) \leq \int_x^y f\rho;$$

- (iii)  *$f$  is continuous and, for all elements  $x < y$  of  $I$ , satisfies the inequality*

$$\int_x^y f\rho \leq c_1(x, y)f(x) + c_2(x, y)f(y);$$

- (iv)  *$f$  is continuous and, for all elements  $x < y$  of  $I$ , satisfies the inequality*

$$c(x, y)f(\xi(x, y)) \leq c_1(x, y)f(x) + c_2(x, y)f(y).$$

The question arises, quite evidently, whether Hermite–Hadamard-type inequalities also characterize generalized convexity in the general case or not. To give an affirmative answer even in the polynomial case remains an open problem and may be the subject of further researches.



## Summary

The notion of convexity can be extended applying *Tchebychev systems* (consult the definitions of the INTRODUCTION). The aim of the dissertation is to generalize the classical Hermite–Hadamard inequality for the extended setting.

In CHAPTER 1 we deal with the case of *polynomial convexity* and apply various methods of numerical analysis, like Gauss-type quadrature formulae (Theorem 1.1, Theorem 1.2, Theorem 1.3, Theorem 1.4) and Hermite-interpolation. Two results of Popoviciu (Theorem A and Theorem B) are also crucial. For technical reasons, further auxiliary tools are developed and applied (see Theorem 1.5, Lemma 1.6, Lemma 1.7). The main results are presented in two theorems (Theorem 1.8 and Theorem 1.9) distinguishing the parity of the order of convexity:

**THEOREM.** *Let  $\rho : [a, b] \rightarrow \mathbb{R}$  be a positive integrable function. Denote the roots of  $P_m$  by  $\xi_1, \dots, \xi_m$  where  $P_m$  is the  $m^{\text{th}}$  degree member of the orthogonal polynomial system on  $[a, b]$  with respect to the weight function  $(x - a)\rho(x)$ , furthermore denote the roots of  $Q_m$  by  $\eta_1, \dots, \eta_m$  where  $Q_m$  is the  $m^{\text{th}}$  degree member of the orthogonal polynomial system on  $[a, b]$  with respect to the weight function  $(b - x)\rho(x)$ . Define the coefficients  $\alpha_0, \dots, \alpha_m$  and  $\beta_1, \dots, \beta_{m+1}$  by the formulae*

$$\begin{aligned}\alpha_0 &:= \frac{1}{P_m^2(a)} \int_a^b P_m^2(x)\rho(x)dx, \\ \alpha_k &:= \frac{1}{\xi_k - a} \int_a^b \frac{(x - a)P_m(x)}{(x - \xi_k)P'_m(\xi_k)}\rho(x)dx\end{aligned}$$

and

$$\begin{aligned}\beta_k &:= \frac{1}{b - \eta_k} \int_a^b \frac{(b - x)Q_m(x)}{(x - \eta_k)Q'_m(\eta_k)}\rho(x)dx, \\ \beta_{m+1} &:= \frac{1}{Q_m^2(b)} \int_a^b Q_m^2(x)\rho(x)dx.\end{aligned}$$

*If a function  $f : [a, b] \rightarrow \mathbb{R}$  is polynomially  $(2m + 1)$ -convex, then it satisfies the following Hermite–Hadamard-type inequality*

$$\alpha_0 f(a) + \sum_{k=1}^m \alpha_k f(\xi_k) \leq \int_a^b f \rho \leq \sum_{k=1}^m \beta_k f(\eta_k) + \beta_{m+1} f(b).$$

**THEOREM.** Let  $\rho : [a, b] \rightarrow \mathbb{R}$  be a positive integrable function. Denote the roots of  $P_m$  by  $\xi_1, \dots, \xi_m$  where  $P_m$  is the  $m^{\text{th}}$  degree member of the orthogonal polynomial system on  $[a, b]$  with respect to the weight function  $\rho(x)$ , and denote the roots of  $Q_{m-1}$  by  $\eta_1, \dots, \eta_{m-1}$  where  $Q_{m-1}$  is the  $(m-1)^{\text{st}}$  degree member of the orthogonal polynomial system on  $[a, b]$  with respect to the weight function  $(b-x)(x-a)\rho(x)$ . Define the coefficients  $\alpha_1, \dots, \alpha_m$  and  $\beta_0, \dots, \beta_{m+1}$  by the formulae

$$\alpha_k := \int_a^b \frac{P_m(x)}{(x - \xi_k)P'_m(\xi_k)} \rho(x) dx$$

and

$$\begin{aligned} \beta_0 &= \frac{1}{(b-a)Q_{m-1}^2(a)} \int_a^b (b-x)Q_{m-1}^2(x)\rho(x)dx, \\ \beta_k &= \frac{1}{(b-\eta_k)(\xi_k-a)} \int_a^b \frac{(b-x)(x-a)Q_{m-1}(x)}{(x-\eta_k)Q'_{m-1}(\eta_k)} \rho(x)dx, \\ \beta_{m+1} &= \frac{1}{(b-a)Q_{m-1}^2(b)} \int_a^b (x-a)Q_{m-1}^2(x)\rho(x)dx. \end{aligned}$$

If a function  $f : [a, b] \rightarrow \mathbb{R}$  is polynomially  $(2m)$ -convex, then it satisfies the following Hermite–Hadamard-type inequality

$$\sum_{k=1}^m \alpha_k f(\xi_k) \leq \int_a^b f \rho \leq \beta_0 f(a) + \sum_{k=1}^{m-1} \beta_k f(\eta_k) + \beta_m f(b).$$

Specializing the weight function  $\rho \equiv 1$ , the roots of the inequalities can be obtained as convex combinations of the endpoints of the domain. The coefficients of the convex combinations are the roots of certain orthogonal polynomials on  $[0, 1]$  in both cases. Observe that interchanging the role of the endpoints in any side of the inequality concerning the odd order case, we obtain the other side of the inequality.

**THEOREM.** Let, for  $m \geq 0$ , the polynomial  $P_m$  be defined by the formula

$$P_m(x) := \begin{vmatrix} 1 & \frac{1}{2} & \cdots & \frac{1}{m+1} \\ x & \frac{1}{3} & \cdots & \frac{1}{m+2} \\ \vdots & \vdots & \ddots & \vdots \\ x^m & \frac{1}{m+2} & \cdots & \frac{1}{2m+1} \end{vmatrix}.$$

Then,  $P_m$  has  $m$  pairwise distinct roots  $\lambda_1, \dots, \lambda_m$  in  $]0, 1[$ . Define the coefficients  $\alpha_0, \dots, \alpha_m$  by

$$\begin{aligned} \alpha_0 &:= \frac{1}{P_m^2(0)} \int_0^1 P_m^2(x) dx, \\ \alpha_k &:= \frac{1}{\lambda_k} \int_0^1 \frac{x P_m(x)}{(x - \lambda_k) P'_m(\lambda_k)} dx. \end{aligned}$$

If a function  $f : [a, b] \rightarrow \mathbb{R}$  is polynomially  $(2m + 1)$ -convex, then it satisfies the following Hermite–Hadamard-type inequality

$$\begin{aligned} \alpha_0 f(a) + \sum_{k=1}^m \alpha_k f((1 - \lambda_k)a + \lambda_k b) &\leq \frac{1}{b - a} \int_a^b f(x) dx \\ &\leq \sum_{k=1}^m \alpha_k f(\lambda_k a + (1 - \lambda_k)b) + \alpha_0 f(b). \end{aligned}$$

THEOREM. Let, for  $m \geq 1$ , the polynomials  $P_m$  and  $Q_{m-1}$  be defined by the formulae

$$\begin{aligned} P_m(x) &:= \begin{vmatrix} 1 & 1 & \cdots & \frac{1}{m} \\ x & \frac{1}{2} & \cdots & \frac{1}{m+1} \\ \vdots & \vdots & \ddots & \vdots \\ x^m & \frac{1}{m+1} & \cdots & \frac{1}{2m} \end{vmatrix}, \\ Q_{m-1}(x) &:= \begin{vmatrix} 1 & \frac{1}{2 \cdot 3} & \cdots & \frac{1}{m(m+1)} \\ x & \frac{1}{3 \cdot 4} & \cdots & \frac{1}{(m+1)(m+2)} \\ \vdots & \vdots & \ddots & \vdots \\ x^{m-1} & \frac{1}{(m+1)(m+2)} & \cdots & \frac{1}{(2m-1)2m} \end{vmatrix}. \end{aligned}$$

Then,  $P_m$  has  $m$  pairwise distinct roots  $\lambda_1, \dots, \lambda_m$  in  $]0, 1[$  and  $Q_{m-1}$  has  $m - 1$  pairwise distinct roots  $\mu_1, \dots, \mu_{m-1}$  in  $]0, 1[$ , respectively. Define the coefficients  $\alpha_1, \dots, \alpha_m$  and  $\beta_0, \dots, \beta_m$  by

$$\alpha_k := \int_0^1 \frac{P_m(x)}{(x - \lambda_k)P'_m(\lambda_k)} dx$$

and

$$\begin{aligned} \beta_0 &:= \frac{1}{Q_{m-1}^2(0)} \int_0^1 (1 - x)Q_{m-1}^2(x) dx, \\ \beta_k &:= \frac{1}{(1 - \mu_k)\mu_k} \int_0^1 \frac{x(1 - x)Q_{m-1}(x)}{(x - \mu_k)Q'_{m-1}(\mu_k)} dx, \\ \beta_m &:= \frac{1}{Q_{m-1}^2(1)} \int_0^1 xQ_{m-1}^2(x) dx. \end{aligned}$$

If a function  $f : [a, b] \rightarrow \mathbb{R}$  is polynomially  $(2m)$ -convex, then it satisfies the following Hermite–Hadamard-type inequality

$$\begin{aligned} \sum_{k=1}^m \alpha_k f((1 - \lambda_k)a + \lambda_k b) &\leq \frac{1}{b - a} \int_a^b f(x) dx \\ &\leq \beta_0 f(a) + \sum_{k=1}^{m-1} \beta_k f((1 - \mu_k)a + \mu_k b) + \beta_m f(b). \end{aligned}$$

In CHAPTER 2 we study the case of *generalized 2-convexity* or, in other terms,  $(\omega_1, \omega_2)$ -convexity. After some technical preambles (such as Lemma 2.1, Lemma 2.2, Lemma 2.3), the first important result of the chapter (Theorem 2.4) gives various characterizations of generalized 2-convex functions:

**THEOREM.** *Let  $(\omega_1, \omega_2)$  be a Tchebychev system over an interval  $I$  such that  $\omega_1$  is positive on  $I^\circ$ . The following statements are equivalent:*

- (i)  $f : I \rightarrow \mathbb{R}$  is  $(\omega_1, \omega_2)$ -convex;  
(ii) for all elements  $x < y < z$  of  $I$  we have that

$$\frac{\begin{vmatrix} f(y) & f(z) \\ \omega_1(y) & \omega_1(z) \end{vmatrix}}{\begin{vmatrix} \omega_1(y) & \omega_1(z) \\ \omega_2(y) & \omega_2(z) \end{vmatrix}} \leq \frac{\begin{vmatrix} f(x) & f(y) \\ \omega_1(x) & \omega_1(y) \end{vmatrix}}{\begin{vmatrix} \omega_1(x) & \omega_1(y) \\ \omega_2(x) & \omega_2(y) \end{vmatrix}};$$

- (iii) for all  $x_0 \in I^\circ$  there exist  $\alpha, \beta \in \mathbb{R}$  such that

$$\begin{aligned} \alpha\omega_1(x_0) + \beta\omega_2(x_0) &= f(x_0), \\ \alpha\omega_1(x) + \beta\omega_2(x) &\leq f(x) \quad (x \in I); \end{aligned}$$

- (iv) for all  $n \in \mathbb{N}$ ,  $x_0, x_1, \dots, x_n \in I$  and  $\lambda_1, \dots, \lambda_n \geq 0$  satisfying the conditions

$$\begin{aligned} \sum_{k=1}^n \lambda_k \omega_1(x_k) &= \omega_1(x_0) \\ \sum_{k=1}^n \lambda_k \omega_2(x_k) &= \omega_2(x_0) \end{aligned}$$

we have that

$$f(x_0) \leq \sum_{k=1}^n \lambda_k f(x_k);$$

- (v) for all  $x_0, x_1, x_2 \in I$  and  $\lambda_1, \lambda_2 \geq 0$  satisfying the conditions

$$\begin{aligned} \lambda_1 \omega_1(x_1) + \lambda_2 \omega_1(x_2) &= \omega_1(x_0) \\ \lambda_1 \omega_2(x_1) + \lambda_2 \omega_2(x_2) &= \omega_2(x_0) \end{aligned}$$

we have that

$$f(x_0) \leq \lambda_1 f(x_1) + \lambda_2 f(x_2);$$

- (vi) for all elements  $x < p < y$  of  $I$

$$f(p) \leq \alpha\omega_1(p) + \beta\omega_2(p)$$

where the constants  $\alpha, \beta$  are the solutions of the system of linear equations

$$\begin{aligned} f(x) &= \alpha\omega_1(x) + \beta\omega_2(x) \\ f(y) &= \alpha\omega_1(y) + \beta\omega_2(y). \end{aligned}$$

In the standard setting this result reduces to the well known properties of convex functions (compare Corollary 2.5) and also gives another characterization via generalized supports (Corollary 2.6). It turns out that generalized convexity is also equivalent to the (standard) convexity of a certain composite function (Theorem 2.7, see below). This connection enables us to state regularity properties for generalized 2-convex functions (Theorem 2.8) and also to generalize the stability result of (standard) convexity (Corollary 2.9).

**THEOREM.** *Let  $(\omega_1, \omega_2)$  be a Tchebychev system on an open interval  $I$  such that  $\omega_1$  is positive. The function  $f : I \rightarrow \mathbb{R}$  is  $(\omega_1, \omega_2)$ -convex if and only if the function  $g : \omega_2/\omega_1(I) \rightarrow \mathbb{R}$  defined by the formula*

$$g := \frac{f}{\omega_1} \circ \left( \frac{\omega_2}{\omega_1} \right)^{-1}$$

*is convex in the standard sense.*

The main result of the chapter states Hermite–Hadamard-type inequality for generalized 2-convex functions (Theorem 2.10).

**THEOREM.** *Let  $(\omega_1, \omega_2)$  be a Tchebychev system on an interval  $[a, b]$  such that  $\omega_1$  is positive on  $]a, b[$ , furthermore, let  $\rho : [a, b] \rightarrow \mathbb{R}$  be a positive integrable function. Define the point  $\xi$  and the coefficients  $c, c_1, c_2$  by the formulae*

$$\xi = \left( \frac{\omega_2}{\omega_1} \right)^{-1} \left( \frac{\int_a^b \omega_2 \rho}{\int_a^b \omega_1 \rho} \right), \quad c = \frac{\int_a^b \omega_1 \rho}{\omega_1(\xi)}$$

*and*

$$c_1 = \frac{\begin{vmatrix} \int_a^b \omega_1 \rho & \omega_1(b) \\ \int_a^b \omega_2 \rho & \omega_2(b) \end{vmatrix}}{\begin{vmatrix} \omega_1(a) & \omega_1(b) \\ \omega_2(a) & \omega_2(b) \end{vmatrix}}, \quad c_2 = \frac{\begin{vmatrix} \omega_1(a) & \int_a^b \omega_1 \rho \\ \omega_2(a) & \int_a^b \omega_2 \rho \end{vmatrix}}{\begin{vmatrix} \omega_1(a) & \omega_1(b) \\ \omega_2(a) & \omega_2(b) \end{vmatrix}}.$$

*If  $f : [a, b] \rightarrow \mathbb{R}$  is an  $(\omega_1, \omega_2)$ -convex function, then the following Hermite–Hadamard-type inequality holds*

$$cf(\xi) \leq \int_a^b f \rho \leq c_1 f(a) + c_2 f(b).$$

The proof is based on the previous two theorems. Theorem 2.8 guarantees the integrability of generalized 2-convex functions defined on compact intervals. The generalized support and chord properties (assertions (iii) and (vi) of Theorem 2.4) give the left and the right hand side inequalities, respectively.

The aim of CHAPTER 3 is to show the existence of Hermite–Hadamard-type inequalities for generalized convexity induced by *arbitrary* Tchebychev systems. To prove regularity properties for generalized convex functions (Theorem 3.2), the following characterization result (Theorem 3.1) plays the key role.

**THEOREM.** Let  $\boldsymbol{\omega} = (\omega_1, \dots, \omega_n)$  be a Tchebychev system over an interval  $I$ . Then, for a function  $f : I \rightarrow \mathbb{R}$ , the following statements are equivalent:

- (i)  $f$  is generalized convex with respect to  $\boldsymbol{\omega}$ ;
- (ii) for all  $y_1 < \dots < y_n$  in  $I$ , the generalized polynomial  $\omega$  of  $\omega_1, \dots, \omega_n$  determined uniquely by the interpolation conditions

$$f(y_k) = \omega(y_k) \quad (k = 1, \dots, n)$$

satisfies the inequalities

$$(-1)^{n+k}(f(y) - \omega(y)) \geq 0 \quad (y_k < y < y_{k+1}, k = 0, \dots, n)$$

under the conventions  $y_0 := \inf I$  and  $y_{n+1} := \sup I$ ;

- (iii) keeping the previous notations and settings, for fixed  $k \in \{0, \dots, n\}$ , the following inequality holds

$$(-1)^{n+k}(f(y) - \omega(y)) \geq 0 \quad (y_k \leq y \leq y_{k+1}).$$

Unfortunately, under such general circumstances the base points of the Hermite–Hadamard-type inequalities cannot be expressed explicitly, we can state only their existence (Theorem 3.4) and uniqueness (Theorem 3.3). Once having this, the inequalities themselves can be verified applying pure linear algebraic methods and the definition of generalized convexity. The main tool of the chapter is the Markov–Krein theory of moment spaces induced by Tchebychev systems (Theorem C and Theorem D). Distinguishing the odd and even order cases, the main results read as follows (Theorem 3.5 and Theorem 3.6).

**THEOREM.** Let  $\boldsymbol{\omega} = (\omega_1, \dots, \omega_{2m+1})$  be a Tchebychev system on  $[a, b]$  and  $\rho : [a, b] \rightarrow \mathbb{R}$  be a positive integrable function. There exist uniquely determined base points  $\xi_1, \dots, \xi_m$  and  $\eta_1, \dots, \eta_m$  of  $]a, b[$  such that

$$\alpha_0 \boldsymbol{\omega}(a) + \sum_{k=1}^m \alpha_k \boldsymbol{\omega}(\xi_k) = \int_a^b \boldsymbol{\omega} \rho = \sum_{k=1}^m \beta_k \boldsymbol{\omega}(\eta_k) + \beta_{m+1} \boldsymbol{\omega}(b).$$

The coefficients  $\alpha_0, \dots, \alpha_m$  and  $\beta_1, \dots, \beta_{m+1}$  are positive and uniquely determined, too. Furthermore, for any generalized  $\boldsymbol{\omega}$ -convex function  $f : [a, b] \rightarrow \mathbb{R}$ , the following Hermite–Hadamard-type inequality holds

$$\alpha_0 f(a) + \sum_{k=1}^m \alpha_k f(\xi_k) \leq \int_a^b f \rho \leq \sum_{k=1}^m \beta_k f(\eta_k) + \beta_{m+1} f(b).$$

**THEOREM.** Let  $\boldsymbol{\omega} = (\omega_1, \dots, \omega_{2m})$  be a Tchebychev system on  $[a, b]$  and let  $\rho : [a, b] \rightarrow \mathbb{R}$  be a positive integrable function. Then, there exist uniquely determined base points  $\xi_1, \dots, \xi_m$  and  $\eta_1, \dots, \eta_{m-1}$  of  $]a, b[$  such that

$$\sum_{k=1}^m \alpha_k \boldsymbol{\omega}(\xi_k) = \int_a^b \boldsymbol{\omega} \rho = \beta_0 \boldsymbol{\omega}(a) + \sum_{k=1}^{m-1} \beta_k \boldsymbol{\omega}(\eta_k) + \beta_m \boldsymbol{\omega}(b).$$

The coefficients  $\alpha_1, \dots, \alpha_m$  and  $\beta_0, \dots, \beta_m$  are positive and uniquely determined, too. Furthermore, for any generalized  $\omega$ -convex function  $f : [a, b] \rightarrow \mathbb{R}$ , the following Hermite–Hadamard-type inequality holds

$$\sum_{k=1}^m \alpha_k f(\xi_k) \leq \int_a^b f \rho \leq \beta_0 f(a) + \sum_{k=1}^{m-1} \beta_k f(\eta_k) + \beta_m f(b).$$

Motivated by Rolle’s mean-value theorem, an elementary approach can also be followed (see Theorem 3.7 and Theorem 3.8) in some particular cases (that is, when the dimension of the underlying Tchebychev system is “small”).

The classical Hermite–Hadamard inequality immediately follows from any of the main results of the first three chapters. Without claiming completeness, at the end of these chapters several applications and examples are presented.

CHAPTER 4 is devoted to proving that the Hermite–Hadamard-type inequality (Theorem 2.10) obtained for generalized 2-convex functions characterize generalized 2-convexity. The most important tool is the following characterization of continuous, non generalized 2-convex functions (Theorem 4.3):

**THEOREM.** *Let  $(\omega_1, \omega_2)$  be a Tchebychev system on an interval  $I$ , furthermore  $f : I \rightarrow \mathbb{R}$  be a continuous function. Then, the following assertions are equivalent:*

- (i)  *$f$  is not  $(\omega_1, \omega_2)$ -convex;*
- (ii) *there exist elements  $x < y$  of  $I$  such that  $\omega < f$  on  $]x, y[$  where  $\omega$  is the generalized line determined by the properties*

$$\omega(x) = f(x) \quad \omega(y) = f(y);$$

- (iii) *there exist elements  $x < p < y$  of  $I$  and a generalized line  $\omega$  such that  $\omega \geq f$  on  $[x, y]$ , moreover*

$$f(x) < \omega(x) \quad f(p) = \omega(p) \quad f(y) < \omega(y);$$

- (iv) *there exists  $p \in I^\circ$  such that  $f$  is locally strictly  $(\omega_1, \omega_2)$ -concave at  $p$ , that is, there exist elements  $x < p < y$  of  $I$  such that, for all  $x < u < p < v < y$ , the following inequality holds:*

$$\begin{vmatrix} f(u) & f(p) & f(v) \\ \omega_1(u) & \omega_1(p) & \omega_1(v) \\ \omega_2(u) & \omega_2(p) & \omega_2(v) \end{vmatrix} < 0.$$

The main results are presented in the subsequent three theorems (see Theorem 4.5, Theorem 4.6 and Theorem 4.7). The first and the second one can be considered as the left and right hand side of the Hermite–Hadamard-type inequality for generalized 2-convex functions (Theorem 2.10); the third one corresponds to the classical Jensen inequality.

**THEOREM.** Let  $(\omega_1, \omega_2)$  be a Tchebychev system over an interval  $[a, b]$  such that  $\omega_1$  is positive on  $]a, b[$ , furthermore  $\rho : [a, b] \rightarrow \mathbb{R}$  be a positive integrable function. Define, for all elements  $x < y$  of  $[a, b]$ , the functions  $\xi(x, y)$  and  $c(x, y)$  by the formulae

$$\xi(x, y) := \left( \frac{\omega_2}{\omega_1} \right)^{-1} \left( \frac{\int_x^y \omega_2 \rho}{\int_x^y \omega_1 \rho} \right), \quad c(x, y) = \frac{\int_x^y \omega_1 \rho}{\omega_1(\xi(x, y))}.$$

Then, a continuous function  $f : [a, b] \rightarrow \mathbb{R}$  is generalized convex with respect to  $(\omega_1, \omega_2)$  if and only if, for all elements  $x < y$  of  $[a, b]$ , it satisfies the inequality

$$c(x, y)f(\xi(x, y)) \leq \int_x^y f \rho.$$

**THEOREM.** Let  $(\omega_1, \omega_2)$  be a Tchebychev system over an interval  $[a, b]$  such that  $\omega_1$  is positive on  $]a, b[$ , furthermore  $\rho : [a, b] \rightarrow \mathbb{R}$  be a positive integrable function. Define, for all elements  $x < y$  of  $[a, b]$ , the functions  $c_1(x, y)$  and  $c_2(x, y)$  by the formulae

$$c_1(x, y) = \frac{\begin{vmatrix} \int_x^y \omega_1 \rho & \omega_1(y) \\ \int_x^y \omega_2 \rho & \omega_2(y) \end{vmatrix}}{\begin{vmatrix} \omega_1(x) & \omega_1(y) \\ \omega_2(x) & \omega_2(y) \end{vmatrix}}, \quad c_2(x, y) = \frac{\begin{vmatrix} \omega_1(x) & \int_x^y \omega_1 \rho \\ \omega_2(x) & \int_x^y \omega_2 \rho \end{vmatrix}}{\begin{vmatrix} \omega_1(x) & \omega_1(y) \\ \omega_2(x) & \omega_2(y) \end{vmatrix}}.$$

Then, a continuous function  $f : [a, b] \rightarrow \mathbb{R}$  is generalized convex with respect to  $(\omega_1, \omega_2)$  if and only if, for all elements  $x < y$  of  $[a, b]$ , it satisfies the inequality

$$\int_x^y f \rho \leq c_1(x, y)f(x) + c_2(x, y)f(y).$$

**THEOREM.** Let  $(\omega_1, \omega_2)$  be a Tchebychev system on  $I$ , furthermore  $f : I \rightarrow \mathbb{R}$  be a continuous function. Keeping the notions of the previous two theorems,  $f$  is  $(\omega_1, \omega_2)$ -convex if and only if, for all elements  $x < y$  of  $I$ , it satisfies the inequality

$$c(x, y)f(\xi(x, y)) \leq c_1(x, y)f(x) + c_2(x, y)f(y).$$

These theorems together give an additional characterization of generalized 2-convexity (Corollary 4.8). Verifying analogous results in the general setting (or even in the polynomial case) remains an open problem and may be the subject of further researches.

## Összefoglaló

A konvexitás fogalma *Csebisev rendszerek* alkalmazásával természetes módon kiterjeszhető (lásd a bevezetés definícióit). A disszertáció célja a klasszikus analízisből ismert Hermite–Hadamard egyenlőtlenség általánosítása erre a kiterjesztett esetre.

Az első fejezetben a *polinomiális konvexitás* esetével foglalkozunk. A vizsgálatok során a numerikus analízis módszereit használjuk, mégpedig Gauss-típusú kvadraturákat (Theorem 1.1, Theorem 1.2, Theorem 1.3, Theorem 1.4) és Hermite-interpolációt. Fontos szerepet játszik Popoviciu két eredménye is (Theorem A és Theorem B). A különféle regularitási feltételek kiküszöbölése miatt további segéderedmények szükségesek (például Theorem 1.5, Lemma 1.6, Lemma 1.7). A fő eredményt a konvexitás rendjének paritása szerint két tételben igazoljuk:

**TÉTEL.** Legyen  $\rho : [a, b] \rightarrow \mathbb{R}$  pozitív integrálható függvény. Jelölje  $P_m$  gyökeit  $\xi_1, \dots, \xi_m$ , ahol  $P_m$  az  $[a, b]$  intervallumon az  $(x - a)\rho(x)$  súlyfüggvényre nézve ortogonális polinomrendszer  $m$ -ed fokú tagja, továbbá jelölje  $Q_m$  gyökeit  $\eta_1, \dots, \eta_m$ , ahol  $Q_m$  az  $[a, b]$  intervallumon a  $(b - x)\rho(x)$  súlyfüggvényre nézve ortogonális polinomrendszer  $m$ -ed fokú tagja. Definiáljuk az  $\alpha_0, \dots, \alpha_m$  illetve a  $\beta_1, \dots, \beta_{m+1}$  együtthatókat az

$$\alpha_0 := \frac{1}{P_m^2(a)} \int_a^b P_m^2(x) \rho(x) dx,$$

$$\alpha_k := \frac{1}{\xi_k - a} \int_a^b \frac{(x - a)P_m(x)}{(x - \xi_k)P'_m(\xi_k)} \rho(x) dx,$$

illetve

$$\beta_k := \frac{1}{b - \eta_k} \int_a^b \frac{(b - x)Q_m(x)}{(x - \eta_k)Q'_m(\eta_k)} \rho(x) dx,$$

$$\beta_{m+1} := \frac{1}{Q_m^2(b)} \int_a^b Q_m^2(x) \rho(x) dx.$$

formulákkal. Ha az  $f : [a, b] \rightarrow \mathbb{R}$  függvény polinomiálisan  $(2m + 1)$ -konvex, akkor teljesíti az alábbi Hermite–Hadamard típusú egyenlőtlenséget:

$$\alpha_0 f(a) + \sum_{k=1}^m \alpha_k f(\xi_k) \leq \int_a^b f \rho \leq \sum_{k=1}^m \beta_k f(\eta_k) + \beta_{m+1} f(b).$$

TÉTEL. Legyen  $\rho : [a, b] \rightarrow \mathbb{R}$  pozitív integrálható függvény. Jelölje  $P_m$  gyökeit  $\xi_1, \dots, \xi_m$ , ahol  $P_m$  az  $[a, b]$  intervallumon a  $\rho(x)$  súlyfüggvényre nézve ortogonális polinomrendszer  $m$ -ed fokú tagja, továbbá jelölje  $Q_{m-1}$  gyökeit  $\eta_1, \dots, \eta_{m-1}$ , ahol  $Q_{m-1}$  az  $[a, b]$  intervallumon a  $(b-x)(x-a)\rho(x)$  súlyfüggvényre nézve ortogonális polinomrendszer  $(m-1)$ -ed fokú tagja. Defináljuk az  $\alpha_1, \dots, \alpha_m$  illetve a  $\beta_0, \dots, \beta_{m+1}$  együtthatókat az

$$\alpha_k := \int_a^b \frac{P_m(x)}{(x - \xi_k)P'_m(\xi_k)} \rho(x) dx$$

illetve

$$\begin{aligned} \beta_0 &= \frac{1}{(b-a)Q_{m-1}^2(a)} \int_a^b (b-x)Q_{m-1}^2(x)\rho(x) dx, \\ \beta_k &= \frac{1}{(b-\eta_k)(\xi_k-a)} \int_a^b \frac{(b-x)(x-a)Q_{m-1}(x)}{(x-\eta_k)Q'_{m-1}(\eta_k)} \rho(x) dx, \\ \beta_{m+1} &= \frac{1}{(b-a)Q_{m-1}^2(b)} \int_a^b (x-a)Q_{m-1}^2(x)\rho(x) dx. \end{aligned}$$

formulákkal. Ha az  $f : [a, b] \rightarrow \mathbb{R}$  függvény polinomiálisan  $(2m)$ -konvex, akkor teljesíti az alábbi Hermite–Hadamard típusú egyenlőtlenséget:

$$\sum_{k=1}^m \alpha_k f(\xi_k) \leq \int_a^b f \rho \leq \beta_0 f(a) + \sum_{k=1}^{m-1} \beta_k f(\eta_k) + \beta_m f(b).$$

Speciálisan, a  $\rho \equiv 1$  választás mellett az egyenlőtlenségek alappontjai az értelmezési tartomány végpontjainak konvex kombinációjaként állnak elő. A konvex kombinációs együtthatók bizonyos ortogonális polinomok gyökei. Figyeljük meg, hogy a páratlan rendű esetre vonatkozó egyenlőtlenség egyik oldalán fölcserélve a végpontok szerepét, épp az egyenlőtlenség másik oldalát kapjuk.

TÉTEL. Defináljuk  $m \geq 0$  esetén a  $P_m$  polinomot a következő formulával:

$$P_m(x) := \begin{vmatrix} 1 & \frac{1}{2} & \cdots & \frac{1}{m+1} \\ x & \frac{1}{3} & \cdots & \frac{1}{m+2} \\ \vdots & \vdots & \ddots & \vdots \\ x^m & \frac{1}{m+2} & \cdots & \frac{1}{2m+1} \end{vmatrix}.$$

Ekkor  $P_m$ -nek  $m$  páronként különböző gyöke van a  $]0, 1[$  intervallumban. Jelölje ezeket a gyököket  $\lambda_1, \dots, \lambda_m$ , és definiáljuk az  $\alpha_0, \dots, \alpha_m$  együtthatókat az alábbi módon:

$$\begin{aligned} \alpha_0 &:= \frac{1}{P_m^2(0)} \int_0^1 P_m^2(x) dx, \\ \alpha_k &:= \frac{1}{\lambda_k} \int_0^1 \frac{x P_m(x)}{(x - \lambda_k) P'_m(\lambda_k)} dx. \end{aligned}$$

Ha az  $f : [a, b] \rightarrow \mathbb{R}$  függvény polinomiálisan  $(2m + 1)$ -konvex, akkor teljesíti a következő Hermite–Hadamard típusú egyenlőtlenséget:

$$\begin{aligned} \alpha_0 f(a) + \sum_{k=1}^m \alpha_k f((1 - \lambda_k)a + \lambda_k b) &\leq \frac{1}{b-a} \int_a^b f(x) dx \\ &\leq \sum_{k=1}^m \alpha_k f(\lambda_k a + (1 - \lambda_k)b) + \alpha_0 f(b). \end{aligned}$$

TÉTEL. Definiáljuk  $m \geq 1$  esetén a  $P_m$  és  $Q_{m-1}$  polinomokat a következő formulákkal:

$$\begin{aligned} P_m(x) &:= \begin{vmatrix} 1 & 1 & \cdots & \frac{1}{m+1} \\ x & \frac{1}{2} & \cdots & \frac{1}{m+1} \\ \vdots & \vdots & \ddots & \vdots \\ x^m & \frac{1}{m+1} & \cdots & \frac{1}{2m} \end{vmatrix}, \\ Q_{m-1}(x) &:= \begin{vmatrix} 1 & \frac{1}{2 \cdot 3} & \cdots & \frac{1}{m(m+1)} \\ x & \frac{1}{3 \cdot 4} & \cdots & \frac{1}{(m+1)(m+2)} \\ \vdots & \vdots & \ddots & \vdots \\ x^{m-1} & \frac{1}{(m+1)(m+2)} & \cdots & \frac{1}{(2m-1)2m} \end{vmatrix}. \end{aligned}$$

Ekkor  $P_m$ -nek  $m$ ,  $Q_{m-1}$ -nek  $m - 1$  páronként különböző gyöke van a  $]0, 1[$  intervallumban. Jelölje a gyököket rendre  $\lambda_1, \dots, \lambda_m$  illetve  $\mu_1, \dots, \mu_{m-1}$ , és definiáljuk az  $\alpha_1, \dots, \alpha_m$  és  $\beta_0, \dots, \beta_m$  együttthatókat az alábbi módon:

$$\begin{aligned} \alpha_k &:= \int_0^1 \frac{P_m(x)}{(x - \lambda_k)P'_m(\lambda_k)} dx; \\ \beta_0 &:= \frac{1}{Q_{m-1}^2(0)} \int_0^1 (1-x)Q_{m-1}^2(x) dx, \\ \beta_k &:= \frac{1}{(1 - \mu_k)\mu_k} \int_0^1 \frac{x(1-x)Q_{m-1}(x)}{(x - \mu_k)Q'_{m-1}(\mu_k)} dx, \\ \beta_m &:= \frac{1}{Q_{m-1}^2(1)} \int_0^1 xQ_{m-1}^2(x) dx. \end{aligned}$$

Ha az  $f : [a, b] \rightarrow \mathbb{R}$  függvény polinomiálisan  $(2m)$ -konvex, akkor teljesíti a következő Hermite–Hadamard típusú egyenlőtlenséget:

$$\begin{aligned} \sum_{k=1}^m \alpha_k f((1 - \lambda_k)a + \lambda_k b) &\leq \frac{1}{b-a} \int_a^b f(x) dx \\ &\leq \beta_0 f(a) + \sum_{k=1}^{m-1} \beta_k f((1 - \mu_k)a + \mu_k b) + \beta_m f(b). \end{aligned}$$

A második fejezetben az *általánosított 2-konvexitás*, más néven  $(\omega_1, \omega_2)$ -konvexitás esetével foglalkozunk. Néhány technikai jellegű előzményt követően (úgy-mint Lemma 2.1, Lemma 2.2, Lemma 2.3) az első fontos eredmény (Theorem 2.4) az általánosított 2-konvex függvények különféle jellemzéseit adja:

**TÉTEL.** Legyen  $(\omega_1, \omega_2)$  Csebisev rendszer a nemüres  $I$  intervallumon úgy, hogy  $\omega_1$  pozitív. Ekkor az alábbi állítások ekvivalensek:

- (i)  $f : I \rightarrow \mathbb{R}$   $(\omega_1, \omega_2)$ -konvex;  
(ii)  $I$ -nek bármely  $x < y < z$  eleme esetén

$$\frac{\begin{vmatrix} f(y) & f(z) \\ \omega_1(y) & \omega_1(z) \end{vmatrix}}{\begin{vmatrix} \omega_1(y) & \omega_1(z) \\ \omega_2(y) & \omega_2(z) \end{vmatrix}} \leq \frac{\begin{vmatrix} f(x) & f(y) \\ \omega_1(x) & \omega_1(y) \end{vmatrix}}{\begin{vmatrix} \omega_1(x) & \omega_1(y) \\ \omega_2(x) & \omega_2(y) \end{vmatrix}};$$

- (iii) bármely  $x_0 \in I^\circ$  esetén létezik  $\alpha, \beta \in \mathbb{R}$  úgy, hogy

$$\begin{aligned} \alpha\omega_1(x_0) + \beta\omega_2(x_0) &= f(x_0), \\ \alpha\omega_1(x) + \beta\omega_2(x) &\leq f(x) \quad (x \in I); \end{aligned}$$

- (iv) bármely  $n \in \mathbb{N}$ ,  $x_0, x_1, \dots, x_n \in I$  és  $\lambda_1, \dots, \lambda_n \geq 0$  esetén, amelyek eleget tesznek a

$$\begin{aligned} \sum_{k=1}^n \lambda_k \omega_1(x_k) &= \omega_1(x_0) \\ \sum_{k=1}^n \lambda_k \omega_2(x_k) &= \omega_2(x_0) \end{aligned}$$

egyenleteknek, teljesül, hogy

$$f(x_0) \leq \sum_{k=1}^n \lambda_k f(x_k);$$

- (v) bármely  $x_0, x_1, x_2 \in I$  és  $\lambda_1, \lambda_2 \geq 0$  esetén, amelyek eleget tesznek a

$$\begin{aligned} \lambda_1 \omega_1(x_1) + \lambda_2 \omega_1(x_2) &= \omega_1(x_0) \\ \lambda_1 \omega_2(x_1) + \lambda_2 \omega_2(x_2) &= \omega_2(x_0) \end{aligned}$$

egyenleteknek, teljesül, hogy

$$f(x_0) \leq \lambda_1 f(x_1) + \lambda_2 f(x_2);$$

- (vi)  $I$ -nek minden  $x < p < y$  elemére fönnáll, hogy

$$f(p) \leq \alpha\omega_1(p) + \beta\omega_2(p),$$

ahol az  $\alpha, \beta$  együtthatók az alábbi lineáris egyenletrendszer megoldásai:

$$\begin{aligned} f(x) &= \alpha\omega_1(x) + \beta\omega_2(x) \\ f(y) &= \alpha\omega_1(y) + \beta\omega_2(y). \end{aligned}$$

A standard esetben ez az eredmény a konvexitás jól ismert jellemzési tulajdonságaira vezet (lásd Corollary 2.5), továbbá egy újabb karakterizációt is lehetővé tesz általánosított tartók segítségével (Corollary 2.6). Kiderül, hogy az általánosított konvexitás ekvivalens egy bizonyos összetett függvény standard értelemben vett konvexitásával (Theorem 2.7, lásd alább). Ez a kapcsolat biztosítja, hogy regularitási tulajdonságokat igazoljunk általánosított 2-konvex függvényekre (Theorem 2.8), és hogy a (standard) konvex függvényekre vonatkozó stabilitási tételt általánosítsuk (Corollary 2.9).

**TÉTEL.** *Legyen  $(\omega_1, \omega_2)$  Csebisev rendszer a nyílt  $I$  intervallumon úgy, hogy  $\omega_1$  pozitív. Az  $f : I \rightarrow \mathbb{R}$  függvény akkor és csak akkor  $(\omega_1, \omega_2)$ -konvex, ha a*

$$g : \omega_2/\omega_1(I) \rightarrow \mathbb{R}, \quad g := \frac{f}{\omega_1} \circ \left( \frac{\omega_2}{\omega_1} \right)^{-1}$$

*módon definiált függvény konvex (a standard értelemben).*

A fejezet fő eredménye Hermite–Hadamard típusú egyenlőtlenséget állít általánosított 2-konvex függvényekre (Theorem 2.10).

**TÉTEL.** *Legyen  $(\omega_1, \omega_2)$  Csebisev rendszer az  $[a, b]$  intervallumon úgy, hogy  $\omega_1$  pozitív az  $]a, b[$  intervallumon, és legyen  $\rho : [a, b] \rightarrow \mathbb{R}$  pozitív integrálható függvény. Definiáljuk a  $\xi$  alappontot és a  $c, c_1, c_2$  együtthatókat a következő formulákkal:*

$$\xi = \left( \frac{\omega_2}{\omega_1} \right)^{-1} \left( \frac{\int_a^b \omega_2 \rho}{\int_a^b \omega_1 \rho} \right), \quad c = \frac{\int_a^b \omega_1 \rho}{\omega_1(\xi)}$$

*továbbá*

$$c_1 = \frac{\begin{vmatrix} \int_a^b \omega_1 \rho & \omega_1(b) \\ \int_a^b \omega_2 \rho & \omega_2(b) \end{vmatrix}}{\begin{vmatrix} \omega_1(a) & \omega_1(b) \\ \omega_2(a) & \omega_2(b) \end{vmatrix}}, \quad c_2 = \frac{\begin{vmatrix} \omega_1(a) & \int_a^b \omega_1 \rho \\ \omega_2(a) & \int_a^b \omega_2 \rho \end{vmatrix}}{\begin{vmatrix} \omega_1(a) & \omega_1(b) \\ \omega_2(a) & \omega_2(b) \end{vmatrix}}.$$

*Ha az  $f : [a, b] \rightarrow \mathbb{R}$  függvény  $(\omega_1, \omega_2)$ -konvex, akkor teljesül rá az alábbi Hermite–Hadamard típusú egyenlőtlenség:*

$$cf(\xi) \leq \int_a^b f \rho \leq c_1 f(a) + c_2 f(b).$$

A bizonyítás a korábbi eredményekből adódik. A regularitási tétel (Theorem 2.8) biztosítja a kompakt intervallumon értelmezett általánosított 2-konvex függvények integrálhatóságát. Az általánosított tartó és húr tulajdonságok (Theorem 2.4 (iii) és (vi) állításai) adják az egyenlőtlenség bal illetve jobb oldalát.

A harmadik fejezetben Hermite–Hadamard típusú egyenlőtlenségek létezését igazoljuk *tetszőleges Csebisev rendszer által indukált általánosított konvexitás* esetére. Az általánosított konvex függvények regularitási tulajdonságainak igazolásában (Theorem 3.2) a következő jellemzési tétel játszik kulcsszerepet.

TÉTEL. Legyen  $\omega = (\omega_1, \dots, \omega_n)$  Csebisev rendszer az  $I$  intervallumon, továbbá  $f : I \rightarrow \mathbb{R}$  adott függvény. Ekkor az alábbi állítások ekvivalensek:

- (i)  $f$  általánosított  $\omega$ -konvex;  
(ii)  $I$ -nek bármely  $y_1 < \dots < y_n$  eleme esetén

$$(-1)^{n+k}(f(y) - \omega(y)) \geq 0 \quad (y_k < y < y_{k+1}, k = 0, \dots, n),$$

ahol  $y_0 := \inf I$ ,  $y_{n+1} := \sup I$  és  $\omega$  az az egyértelműen meghatározott általánosított polinom, amelyre

$$f(y_k) = \omega(y_k) \quad (k = 1, \dots, n);$$

- (iii) megtartva (ii) jelöléseit, bármely rögzített  $k \in \{0, \dots, n\}$  esetén fennáll a következő egyenlőtlenség:

$$(-1)^{n+k}(f(y) - \omega(y)) \geq 0 \quad (y_k \leq y \leq y_{k+1}).$$

Sajnos, ebben az általánosságban az Hermite–Hadamard típusú egyenlőtlenségek alappontjai nem adhatók meg explicit formában, csupán azok létezését (Theorem 3.4) és egyértelműségét (Theorem 3.3) állíthatjuk. Ezek birtokában maguk az egyenlőtlenségek egyszerű lineáris algebrai módszerekkel illetve az általánosított konvexitás definíciója segítségével igazolhatók. A fejezet legfontosabb eszköze a Csebisev rendszerek által indukált momentumterek Krein–Markov-féle elmélete (Theorem C és Theorem D). A fő eredmények (Theorem 3.5 és Theorem 3.6) külön tárgyalják a páros és páratlan rendű eseteket:

TÉTEL. Legyen  $\omega = (\omega_1, \dots, \omega_{2m+1})$  Csebisev rendszer az  $[a, b]$  intervallumon, és legyen  $\rho : [a, b] \rightarrow \mathbb{R}$  pozitív integrálható függvény. Egyértelműen léteznek az  $]a, b[$  nyílt intervallumban  $\xi_1, \dots, \xi_m$  és  $\eta_1, \dots, \eta_m$  alappontok úgy, hogy

$$\alpha_0 \omega(a) + \sum_{k=1}^m \alpha_k \omega(\xi_k) = \int_a^b \omega \rho = \sum_{k=1}^m \beta_k \omega(\eta_k) + \beta_{m+1} \omega(b).$$

Az  $\alpha_0, \dots, \alpha_m$  és  $\beta_1, \dots, \beta_{m+1}$  együtthatók pozitívak és szintén egyértelműen meghatározottak. Továbbá, bármely  $f : [a, b] \rightarrow \mathbb{R}$  általánosított  $\omega$ -konvex függvény esetén fennáll az alábbi Hermite–Hadamard típusú egyenlőtlenség:

$$\alpha_0 f(a) + \sum_{k=1}^m \alpha_k f(\xi_k) \leq \int_a^b f \rho \leq \sum_{k=1}^m \beta_k f(\eta_k) + \beta_{m+1} f(b).$$

TÉTEL. Legyen  $\omega = (\omega_1, \dots, \omega_{2m})$  Csebisev rendszer az  $[a, b]$  intervallumon, és legyen  $\rho : [a, b] \rightarrow \mathbb{R}$  pozitív integrálható függvény. Egyértelműen léteznek az  $]a, b[$  nyílt intervallumban  $\xi_1, \dots, \xi_m$  és  $\eta_1, \dots, \eta_{m-1}$  alappontok úgy, hogy

$$\sum_{k=1}^m \alpha_k \omega(\xi_k) = \int_a^b \omega \rho = \beta_0 \omega(a) + \sum_{k=1}^{m-1} \beta_k \omega(\eta_k) + \beta_m \omega(b).$$

Az  $\alpha_1, \dots, \alpha_m$  és  $\beta_0, \dots, \beta_m$  együtthatók pozitívak és szintén egyértelműen meghatározottak. Továbbá, bármely  $f : [a, b] \rightarrow \mathbb{R}$  általánosított  $\omega$ -konvex függvény esetén fennáll az alábbi Hermite–Hadamard típusú egyenlőtlenség:

$$\sum_{k=1}^m \alpha_k f(\xi_k) \leq \int_a^b f \rho \leq \beta_0 f(a) + \sum_{k=1}^{m-1} \beta_k f(\eta_k) + \beta_m f(b).$$

A Rolle-féle középértéktétel alap gondolatából kiindulva egy elemi megközelítés is követhető bizonyos speciális esetekben, nevezetesen amikor a konvexitást indukáló Csebisev rendszer dimenziója “kicsi” (lásd Theorem 3.7 valamint Theorem 3.8).

A klasszikus Hermite–Hadamard egyenlőtlenség természetesen az első három fejezet fő eredményeinek bármelyikéből egyszerűn következik. E három fejezet végén, a teljesség igénye nélkül, számos további példa és alkalmazás is található.

A negyedik fejezetben megmutatjuk, hogy az általánosított 2-konvex függvényekre vonatkozó Hermite–Hadamard típusú egyenlőtlenség (Theorem 2.10) egyben jellemzése is az általánosított 2-konvexitásnak. A vizsgálatok során a legfontosabb segédeszköz a folytonos, általánosított *nem* 2-konvex függvények jellemzése általánosított egyenesekkel, vagyis a konvexitást indukáló Csebisev rendszer tagjainak lineáris kombinációival (Theorem 4.3):

**TÉTEL.** Legyen  $(\omega_1, \omega_2)$  Csebisev rendszer az  $I$  intervallumon, továbbá legyen  $f : I \rightarrow \mathbb{R}$  folytonos függvény. Az alábbi állítások ekvivalensek:

- (i)  $f$  nem  $(\omega_1, \omega_2)$ -konvex;
- (ii) léteznek  $x < y$  elemei  $I$ -nek úgy, hogy  $\omega < f$  az  $]x, y[$  nyílt intervallumon, ahol  $\omega$  az az egyértelműen meghatározott általánosított egyenes, amelyre

$$\omega(x) = f(x) \quad \omega(y) = f(y);$$

- (iii) léteznek  $x < p < y$  elemei  $I$ -nek és létezik  $\omega$  általánosított egyenes úgy, hogy  $\omega \geq f$  az  $[x, y]$  intervallumon, továbbá

$$f(x) < \omega(x) \quad f(p) = \omega(p) \quad f(y) < \omega(y);$$

- (iv) létezik  $p \in I^\circ$  úgy, hogy  $f$  lokálisan szigorúan  $(\omega_1, \omega_2)$ -konkáv  $p$ -ben, azaz, léteznek  $I$ -nek  $x < p < y$  elemei úgy, hogy bármely  $x < u < p < v < y$  esetén fennáll az alábbi egyenlőtlenség:

$$\begin{vmatrix} f(u) & f(p) & f(v) \\ \omega_1(u) & \omega_1(p) & \omega_1(v) \\ \omega_2(u) & \omega_2(p) & \omega_2(v) \end{vmatrix} < 0.$$

A fejezet fő eredményeit az alábbi három tétel foglalja össze (lásd Theorem 4.5, Theorem 4.6 és Theorem 4.7). Az első és a második az általánosított 2-konvexitás esetére vonatkozó Hermite–Hadamard típusú egyenlőtlenség (Theorem 2.10) bal illetve jobb oldalának a megfelelője, míg a harmadik a klasszikus Jensen-egyenlőtlenség általánosítása.

**TÉTEL.** Legyen  $(\omega_1, \omega_2)$  Csebisev rendszer az  $[a, b]$  intervallumon úgy, hogy  $\omega_1$  pozitív  $]a, b[$ -n, és legyen  $\rho : [a, b] \rightarrow \mathbb{R}$  pozitív integrálható függvény. Defináljuk az  $[a, b]$  bármely  $x < y$  elempárja esetén a  $\xi(x, y)$  és  $c(x, y)$  függvényeket a következő formulákkal:

$$\xi(x, y) := \left( \frac{\omega_2}{\omega_1} \right)^{-1} \left( \frac{\int_x^y \omega_2 \rho}{\int_x^y \omega_1 \rho} \right), \quad c(x, y) = \frac{\int_x^y \omega_1 \rho}{\omega_1(\xi(x, y))}.$$

Ekkor az  $f : [a, b] \rightarrow \mathbb{R}$  folytonos függvény pontosan akkor általánosított konvex  $(\omega_1, \omega_2)$ -re nézve, ha  $I$ -nek bármely  $x < y$  eleme esetén teljesül az alábbi egyenlőtlenség:

$$c(x, y)f(\xi(x, y)) \leq \int_x^y f \rho.$$

**TÉTEL.** Legyen  $(\omega_1, \omega_2)$  Csebisev rendszer az  $[a, b]$  intervallumon úgy, hogy  $\omega_1$  pozitív  $]a, b[$ -n, és legyen  $\rho : [a, b] \rightarrow \mathbb{R}$  pozitív integrálható függvény. Defináljuk az  $[a, b]$  bármely  $x < y$  elempárja esetén a  $c_1(x, y)$  és  $c_2(x, y)$  függvényeket a következő formulákkal:

$$c_1(x, y) = \frac{\begin{vmatrix} \int_x^y \omega_1 \rho & \omega_1(y) \\ \int_x^y \omega_2 \rho & \omega_2(y) \end{vmatrix}}{\begin{vmatrix} \omega_1(x) & \omega_1(y) \\ \omega_2(x) & \omega_2(y) \end{vmatrix}}, \quad c_2(x, y) = \frac{\begin{vmatrix} \omega_1(x) & \int_x^y \omega_1 \rho \\ \omega_2(x) & \int_x^y \omega_2 \rho \end{vmatrix}}{\begin{vmatrix} \omega_1(x) & \omega_1(y) \\ \omega_2(x) & \omega_2(y) \end{vmatrix}}.$$

Ekkor az  $f : [a, b] \rightarrow \mathbb{R}$  folytonos függvény pontosan akkor általánosított konvex  $(\omega_1, \omega_2)$ -re nézve, ha  $I$ -nek bármely  $x < y$  eleme esetén teljesül az alábbi egyenlőtlenség:

$$\int_x^y f \rho \leq c_1(x, y)f(x) + c_2(x, y)f(y).$$

**TÉTEL.** Legyen  $(\omega_1, \omega_2)$  Csebisev rendszer az  $I$  intervallumon, továbbá legyen  $f : I \rightarrow \mathbb{R}$  folytonos függvény. Megtartva az előző két tétel jelöléseit,  $f$  pontosan akkor  $(\omega_1, \omega_2)$ -konvex ha  $I$ -nek bármely  $x < y$  eleme esetén teljesül a következő egyenlőtlenség:

$$c(x, y)f(\xi(x, y)) \leq c_1(x, y)f(x) + c_2(x, y)f(y).$$

E három tétel együttesen az általánosított 2-konvexitás egy újabb jellemzését adja (Corollary 4.8). Hasonló típusú eredmények igazolása az általános esetben, vagy akár a polinomiális konvexitás esetében is, nyílt probléma és további kutatások témáját képezheti.

## Bibliography

- [AD96] S. Ambikumar and S. W. Drury, *A reverse Hadamard inequality*, Linear Algebra Appl. **247** (1996), 83–95.
- [AGP99] G. Allasia, C. Giordano, and J. E. Pečarić, *Hadamard-type inequalities for  $(2r)$ -convex functions with applications*, Atti Accad. Sci. Torino Cl. Sci. Fis. Mat. Natur. **133** (1999), 187–200.
- [AK92] J. Aczél and M. Kuczma, *Solutions of a functional equation, convex of higher order*, General Inequalities, 6 (Oberwolfach, 1990) (W. Walter, ed.), International Series of Numerical Mathematics, Birkhäuser, Basel, 1992, pp. 209–213.
- [Alz89] H. Alzer, *A note on Hadamard's inequalities*, C. R. Math. Rep. Acad. Sci. Canada **11** (1989), no. 6, 255–258.
- [Amb67] J. Ambrosiewicz, *A note on Hadamard's inequality*, Wiadom. Mat. (2) **10** (1967), 7–8 (1967).
- [AP88] St. A. Andersson and M. D. Perlman, *Group-invariant analogues of Hadamard's inequality*, Linear Algebra Appl. **110** (1988), 91–116.
- [Azp94] A. G. Azpeitia, *Convex functions and the Hadamard inequality*, Rev. Colombiana Mat. **28** (1994), no. 1, 7–12.
- [BDB96] C. Buşe, S. S. Dragomir, and D. Barbu, *The convergence of some sequences connected to Hadamard's inequality*, Demonstratio Math. **29** (1996), no. 1, 53–59.
- [Bel43] R. Bellman, *A note on determinants and Hadamard's inequality*, Amer. Math. Monthly **50** (1943), 550–551.
- [Bes04] M. Bessenyei, *Hermite–Hadamard-type inequalities for generalized 3-convex functions*, Publ. Math. Debrecen **65** (2004), no. 1-2, 223–232.
- [Bje95] M. Bjelica, *Hadamard's inequality and fixed-point method*, Filomat (1995), no. 9, part 3, 599–602, Algebra, Logic & Discrete Mathematics (Niš, 1995).
- [Bon50] F. F. Bonsall, *The characterization of generalized convex functions*, Quart. J. Math., Oxford Ser. (2) **1** (1950), 100–111.
- [BP02] M. Bessenyei and Zs. Páles, *Higher-order generalizations of Hadamard's inequality*, Publ. Math. Debrecen **61** (2002), no. 3-4, 623–643.
- [BP03] M. Bessenyei and Zs. Páles, *Hadamard-type inequalities for generalized convex functions*, Math. Inequal. Appl. **6** (2003), no. 3, 379–392.
- [BP04] M. Bessenyei and Zs. Páles, *On generalized higher-order convexity and Hermite–Hadamard-type inequalities*, Acta Sci. Math. (Szeged) **70** (2004), 13–24.
- [BP05] M. Bessenyei and Zs. Páles, *Hermite–Hadamard inequalities for generalized convex functions*, Aequationes Math., (2005), to appear.
- [BP] M. Bessenyei and Zs. Páles, *Characterization of convexity via Hadamard's inequality*, manuscript.
- [CF99] M. Cardin and P. Ferretti, *On convex functions of higher order*, Generalized convexity and optimization for economic and financial decisions (Verona, 1998), Pitagora, Bologna, 1999, pp. 95–109.

- [Cho92] B. S. Choi, *A maximum likelihood proof of the Hadamard inequality*, Math. Sci. **17** (1992), no. 1, 68.
- [CP04] P. Czinder and Zs. Páles, *An extension of the Hermite-Hadamard inequality and an application for Gini and Stolarsky means*, J. Inequal. Pure Appl. Math. **5** (2004), no. 2, Article 42, pp. 8 (electronic).
- [Cra94] R. Craigen, *A direct approach to Hadamard's inequality*, Bull. Inst. Combin. Appl. **12** (1994), 28–32.
- [CS98] F. Constantinescu and G. Scharf, *Generalized Gram-Hadamard inequality*, J. Inequal. Appl. **2** (1998), no. 4, 381–386.
- [DA98] S. S. Dragomir and R. P. Agarwal, *Two new mappings associated with Hadamard's inequalities for convex functions*, Appl. Math. Lett. **11** (1998), no. 3, 33–38.
- [Dan85] S. Sh. Dang, *A note on the extended Hadamard inequalities*, J. Math. Res. Exposition **5** (1985), no. 1, 16.
- [DB95] S. S. Dragomir and C. Buşe, *Refinements of Hadamard's inequality for multiple integrals*, Utilitas Math. **47** (1995), 193–198.
- [DF98] S. S. Dragomir and S. Fitzpatrick, *Hadamard's inequality for  $s$ -convex functions in the first sense and applications*, Demonstratio Math. **31** (1998), no. 3, 633–642.
- [DF99] S. S. Dragomir and S. Fitzpatrick, *The Hadamard inequalities for  $s$ -convex functions in the second sense*, Demonstratio Math. **32** (1999), no. 4, 687–696.
- [dG90] A. M. de Guzman, *An information-theoretic proof of Hadamard's inequality*, Matimyás Mat. **13** (1990), no. 1, 1–10.
- [DM97] S. S. Dragomir and B. Mond, *On Hadamard's inequality for a class of functions of Godunova and Levin*, Indian J. Math. **39** (1997), no. 1, 1–9.
- [DM98] S. S. Dragomir and B. Mond, *Integral inequalities of Hadamard type for log-convex functions*, Demonstratio Math. **31** (1998), no. 2, 355–364.
- [DM99] S. S. Dragomir and D. M. Milošević, *Some refinements of Hadamard's inequalities*, J. Indian Math. Soc. (N.S.) **66** (1999), no. 1–4, 97–104.
- [DMS93] S. S. Dragomir, D. M. Milošević, and J. Sándor, *On some refinements of Hadamard's inequalities and applications*, Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. **4** (1993), 3–10.
- [DP98] S. S. Dragomir and C. E. M. Pearce, *Quasi-convex functions and Hadamard's inequality*, Bull. Austral. Math. Soc. **57** (1998), no. 3, 377–385.
- [DP00] S. S. Dragomir and C. E. M. Pearce, *Selected Topics on Hermite-Hadamard Inequalities*, RGMIA Monographs ([http://rgmia.vu.edu.au/monographs/hermite\\_hadamard.html](http://rgmia.vu.edu.au/monographs/hermite_hadamard.html)), Victoria University, 2000.
- [DPP95] S. S. Dragomir, J. E. Pečarić, and L.-E. Persson, *Some inequalities of Hadamard type*, Soochow J. Math. **21** (1995), no. 3, 335–341.
- [DPS90] S. S. Dragomir, J. E. Pečarić, and J. Sándor, *A note on the Jensen-Hadamard inequality*, Anal. Numér. Théor. Approx. **19** (1990), no. 1, 29–34.
- [Dra90] S. S. Dragomir, *Two refinements of Hadamard's inequalities*, Zb. Rad. (Kragujevac) (1990), no. 11, 23–26.
- [Dra91] S. S. Dragomir, *A mapping in connection to Hadamard's inequalities*, Anz. Österreich. Akad. Wiss. Math.-Natur. Kl. **128** (1991), 17–20 (1992).
- [Dra92a] S. S. Dragomir, *On Hadamard's inequalities for convex functions*, Math. Balkanica (N.S.) **6** (1992), no. 3, 215–222.
- [Dra92b] S. S. Dragomir, *Two mappings in connection to Hadamard's inequalities*, J. Math. Anal. Appl. **167** (1992), no. 1, 49–56.
- [Dra93a] S. S. Dragomir, *A note on Hadamard's inequalities*, Mathematica (Cluj) **35(58)** (1993), no. 1, 21–24.

- [Dra93b] S. S. Dragomir, *A refinement of Hadamard's inequality for isotonic linear functionals*, Tamkang J. Math. **24** (1993), no. 1, 101–106.
- [Dra94] S. S. Dragomir, *Some remarks on Hadamard's inequalities for convex functions*, Extracta Math. **9** (1994), no. 2, 88–94.
- [Dra00a] S. S. Dragomir, *On Hadamard's inequality for the convex mappings defined on a ball in the space and applications*, Math. Inequal. Appl. **3** (2000), no. 2, 177–187.
- [Dra00b] S. S. Dragomir, *On Hadamard's inequality on a disk*, JIPAM. J. Inequal. Pure Appl. Math. **1** (2000), no. 1, Article 2, 11 pp. (electronic).
- [Dra01a] S. S. Dragomir, *On the Hadamard's inequality for convex functions on the co-ordinates in a rectangle from the plane*, Taiwanese J. Math. **5** (2001), no. 4, 775–788.
- [Dra01b] S. S. Dragomir, *Refinements of the Hermite-Hadamard inequality for convex functions*, Tamsui Oxf. J. Math. Sci. **17** (2001), no. 2, 131–137.
- [Dra01c] S. S. Dragomir, *Refinements of the Hermite-Hadamard integral inequality for log-convex functions*, Austral. Math. Soc. Gaz. **28** (2001), no. 3, 129–134.
- [Dra02a] S. S. Dragomir, *An inequality improving the first Hermite-Hadamard inequality for convex functions defined on linear spaces and applications for semi-inner products*, JIPAM. J. Inequal. Pure Appl. Math. **3** (2002), no. 2, Article 31, 8 pp. (electronic).
- [Dra02b] S. S. Dragomir, *An inequality improving the second Hermite-Hadamard inequality for convex functions defined on linear spaces and applications for semi-inner products*, JIPAM. J. Inequal. Pure Appl. Math. **3** (2002), no. 3, Article 35, 8 pp. (electronic).
- [Dra02c] S. S. Dragomir, *On some new inequalities of Hermite-Hadamard type for  $m$ -convex functions*, Tamkang J. Math. **33** (2002), no. 1, 55–65.
- [Fen85] C. H. Feng, *On Hadamard inequalities for convex functions*, Chinese Ann. Math. Ser. A **6** (1985), no. 4, 443–446, An English summary appears in Chinese Ann. Math. Ser. B **6** (1985), no. 3, 380.
- [Fin98] A. M. Fink, *A best possible Hadamard inequality*, Math. Inequal. Appl. **1** (1998), no. 2, 223–230.
- [Fin00] A. M. Fink, *Hadamard's inequality for log-concave functions*, Math. Comput. Modelling **32** (2000), no. 5-6, 625–629, Boundary value problems and related topics.
- [Fri75] Sh. Friedland, *Generalized Hadamard inequality and its applications*, Linear and Multilinear Algebra **2** (1975), 327–334.
- [Gaj86] Z. Gajda, *Christensen measurability of polynomial functions and convex functions of higher orders*, Ann. Polon. Math. **47** (1986), no. 1, 25–40.
- [Gal91] S. Gh. Gal, *Calculus of higher order modulus of smoothness for convex functions of higher order and applications*, Calcolo **28** (1991), no. 3-4, 275–282 (1992).
- [Gal92] S. Gh. Gal, *Calculus of higher order averaged modulus of smoothness in  $L^p$ -norm for convex functions of higher order*, Serdica **18** (1992), no. 3-4, 232–239.
- [Gav00] B. Gavrea, *On Hadamard's inequality for the convex mappings defined on a convex domain in the space*, JIPAM. J. Inequal. Pure Appl. Math. **1** (2000), no. 1, Article 9, 6 pp. (electronic).
- [Geř65] S. P. Geřsberg, *A generalization of Hadamard's inequality*, Leningrad. Meh. Inst. Sb. Naučn. Trudov No. **50** (1965), 42–54.
- [Ger90] J. Ger, *Convex functions of higher orders in some metric spaces*, Rad. Mat. **6** (1990), no. 1, 55–69.
- [GG87] I. Gavrea and M. Gurzău, *Inequalities of the Levinson-Popoviciu type for convex functions of higher order*, Itinerant Seminar on Functional Equations, Approximation and Convexity (Cluj-Napoca, 1987), Preprint, vol. 87, Univ. “Babeş-Bolyai”, Cluj, 1987, pp. 163–166.
- [GJMW84] B. Grone, Ch. Johnson, E. Marques de Sa, and H. Wolkowicz, *Improving Hadamard's inequality*, Linear and Multilinear Algebra **16** (1984), no. 1-4, 305–322.

- [GPP97] P. M. Gill, C. E. M. Pearce, and J. E. Pečarić, *Hadamard's inequality for  $r$ -convex functions*, J. Math. Anal. Appl. **215** (1997), no. 2, 461–470.
- [GS01] S. Guillaume and A. Seeger, *A higher-order smoothing technique for polyhedral convex functions: geometric and probabilistic considerations*, J. Convex Anal. **8** (2001), no. 1, 109–126.
- [Had93] J. Hadamard, *Étude sur les propriétés des fonctions entières et en particulier d'une fonction considérée par Riemann*, J. Math. Pures Appl. **58** (1893), 171–215.
- [HB83] D. Hershkowitz and A. Berman, *Necessary conditions and a sufficient condition for the Fischer-Hadamard inequalities*, Linear and Multilinear Algebra **13** (1983), no. 1, 67–72.
- [Hey92] P. Heyfron, *A generalization of Hadamard's inequality*, Linear and Multilinear Algebra **32** (1992), no. 1, 75–84.
- [HH94] G. Hämmerlin and K. H. Hoffmann, *Numerische Mathematik*, Springer, Berlin, 1994.
- [Joh66] F. John, *Lectures on Advanced Numerical Analysis*, Gordon and Breach, London, 1966.
- [Joh75] Ch. R. Johnson, *Hadamard's inequality for matrices with positive-definite Hermitian component*, Michigan Math. J. **22** (1975), no. 3, 225–228 (1976).
- [Kar68] S. Karlin, *Total positivity. Vol. I*, Stanford University Press, Stanford, California, 1968.
- [KK92] E. Klafszky and P. Kas, *A proof of the generalized Hadamard inequality via information theory*, Ann. Univ. Sci. Budapest. Sect. Comput. **13** (1992), 21–24.
- [KS66] S. Karlin and W. J. Studden, *Tchebycheff systems: With applications in analysis and statistics*, Pure and Applied Mathematics, Vol. XV, Interscience Publishers John Wiley & Sons, New York-London-Sydney, 1966.
- [Kuc85] M. Kuczma, *An Introduction to the Theory of Functional Equations and Inequalities*, Prace Naukowe Uniwersytetu Śląskiego w Katowicach, vol. 489, Państwowe Wydawnictwo Naukowe — Uniwersytet Śląski, Warszawa-Kraków-Katowice, 1985.
- [Lan00] E. G. Lanina, *On the generalized Hadamard inequality for multiple integrals*, Vestnik Moskov. Univ. Ser. I Mat. Mekh. (2000), no. 1, 55–58.
- [Liu94] J. Zh. Liu, *A minimal value of the determinant of a positive definite self-conjugate matrix, and a further improvement of the Hadamard inequality*, J. Math. Res. Exposition **14** (1994), no. 1, 73–78.
- [LT00] K.-Ch. Lee and K.-L. Tseng, *On weighted generalization of Hadamard's inequality for  $g$ -convex functions*, Tamsui Oxf. J. Math. Sci. **16** (2000), no. 1, 91–104.
- [Lup76] A. Lupaş, *A generalization of Hadamard inequalities for convex functions*, Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz. (1976), no. 544–576, 115–121.
- [Lup97] A. Lupaş, *The Jensen-Hadamard inequality for convex functions of higher order*, Octogon Math. Mag. **5** (1997), no. 2, 8–9.
- [LV79] I. B. Lacković and P. M. Vasić, *Notes on convex functions. V. Positive linear operators and functions convex of order  $n$* , Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz. (1979), no. 634–677, 239–243.
- [Mak79] D. M. Maksimović, *A short proof of generalized Hadamard's inequalities*, Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz. (1979), no. 634–677, 126–128.
- [Mer99] M. Merkle, *Remarks on Ostrowski's and Hadamard's inequality*, Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. **10** (1999), 113–117.
- [ML85] D. S. Mitrinović and I. B. Lacković, *Hermite and convexity*, Aequationes Math. **28** (1985), 229–232.
- [MO82] A. W. Marshall and I. Olkin, *A convexity proof of Hadamard's inequality*, Amer. Math. Monthly **89** (1982), no. 9, 687–688.
- [Nic02] C. P. Niculescu, *The Hermite-Hadamard inequality for convex functions of a vector variable*, Math. Inequal. Appl. **5** (2002), no. 4, 619–623.

- [Nis79] A. Nishi, *Generalized Hadamard's inequalities and their applications to statistics*, Bull. Math. Statist. **18** (1978/79), no. 3-4, 45–49.
- [NP03] E. Neuman and Zs. Páles, *On comparison of Stolarsky and Gini means*, J. Math. Anal. Appl. **278** (2003), no. 2, 274–284.
- [NP04] C. Niculescu and L.-E. Persson, *Old and new on the Hermite–Hadamard inequality*, Real Anal. Exchange (2004).
- [Pál86] Zs. Páles, *Hölder-type inequalities for quasarithmetic means*, Acta Math. Hungar. **47** (1986), no. 3-4, 395–399.
- [Pál99] Zs. Páles, *Strong Hölder and Minkowski inequalities for quasarithmetic means*, Acta Sci. Math. (Szeged) **65** (1999), no. 3-4, 493–503.
- [Pál00a] Zs. Páles, *Bernstein–Doetsch-type results for general functional inequalities*, Rocznik Nauk.-Dydakt. Prace Mt. **17** (2000), 197–206, Dedicated to Professor Zenon Moszner on the occasion of his seventieth birthday.
- [Pál00b] Zs. Páles, *Nonconvex functions and separation by power means*, Math. Inequal. Appl. **3** (2000), no. 2, 169–176.
- [PČ02] J. E. Pečarić and V. Čuljak, *Interpolation polynomials and inequalities for convex functions of higher order*, Math. Inequal. Appl. **5** (2002), no. 3, 369–386, Inequalities, 2001 (Timișoara).
- [PČF99] J. E. Pečarić, V. Čuljak, and A. M. Fink, *On some inequalities for convex functions of higher order*, Nonlinear Stud. **6** (1999), no. 2, 131–140.
- [PČR01] J. Pečarić, V. Čuljak, and M. Rogina, *On some inequalities for convex functions of higher order. II*, Nonlinear Anal. **45** (2001), no. 3, Ser. A: Theory Methods, 281–294.
- [PD91] J. E. Pečarić and S. S. Dragomir, *A generalization of Hadamard's inequality for isotonic linear functionals*, Rad. Mat. **7** (1991), no. 1, 103–107.
- [Peč89] J. E. Pečarić, *A note on Hadamard's inequality*, Makedon. Akad. Nauk. Umet. Oddel. Mat.-Tehn. Nauk. Prilozi **10** (1989), no. 1-2, 21–23 (1991).
- [Peč92a] J. E. Pečarić, *Notes on convex functions*, General inequalities, 6 (Oberwolfach, 1990), Internat. Ser. Numer. Math., vol. 103, Birkhäuser, Basel, 1992, pp. 449–454.
- [Peč92b] J. E. Pečarić, *Remarks on two interpolations of Hadamard's inequalities*, Makedon. Akad. Nauk. Umet. Oddel. Mat.-Tehn. Nauk. Prilozi **13** (1992), no. 1, 9–12.
- [Pop36] T. Popoviciu, *Notes sur les fonctions convexes d'ordre supérieur (I)*, Mathematica (Cluj) **12** (1936), 81–92.
- [Pop38a] T. Popoviciu, *Deux remarques sur les fonctions convexes*, Bull. Sect. Sci. Acad. Roum. **20** (1938), 187–191 (or 45–49) (1939).
- [Pop38b] T. Popoviciu, *Sur l'approximation des fonctions convexes d'ordre supérieur*, Bull. Sect. Sci. Acad. Roum. **20** (1938), 192–195 (or 50–53) (1939).
- [Pop39a] T. Popoviciu, *Notes sur les fonctions convexes d'ordre supérieur (VI)*, Rev. Math. Union Interbalkan. **2** (1939), 31–40.
- [Pop39b] T. Popoviciu, *Notes sur les fonctions convexes d'ordre supérieur (VII)*, Acad. Roum. Bull. Sect. Sci. **22** (1939), 29–33.
- [Pop40a] T. Popoviciu, *Introduction à la théorie des différences divisées*, Bull. Math. Soc. Roumaine Sci. **42** (1940), no. 1, 65–78.
- [Pop40b] T. Popoviciu, *Notes sur les fonctions convexes d'ordre supérieur. III*, Mathematica, Cluj **16** (1940), 74–86.
- [Pop40c] T. Popoviciu, *Notes sur les fonctions convexes d'ordre supérieur. IV*, Disquisit. Math. Phys. **1** (1940), 163–171.
- [Pop40d] T. Popoviciu, *Notes sur les fonctions convexes d'ordre supérieur (V)*, Acad. Roum. Bull. Sect. Sci. **22** (1940), 351–356.
- [Pop40e] T. Popoviciu, *Notes sur les généralisations des fonctions convexes d'ordre supérieur. I*, Disquisit. Math. Phys. **1** (1940), 35–42.

- [Pop40f] T. Popoviciu, *Notes sur les généralisations des fonctions convexes d'ordre supérieur (II)*, Acad. Roum. Bull. Sect. Sci. **22** (1940), 473–477.
- [Pop41] T. Popoviciu, *Notes sur les fonctions convexes d'ordre supérieur. IX. Inégalités linéaires et bilinéaires entre les fonctions convexes. Quelques généralisations d'une inégalité de Tchebycheff*, Bull. Math. Soc. Roumaine Sci. **43** (1941), 85–141.
- [Pop42a] T. Popoviciu, *Notes sur les fonctions convexes d'ordre supérieur. X. Sur quelques propriétés des différences divisées et des polynomes de Lagrange*, Ann. Sci. Univ. Jassy. Sect. I. **28** (1942), 161–207.
- [Pop42b] T. Popoviciu, *Notes sur les généralisations des fonctions convexes d'ordre supérieur. IV*, Disquisit. Math. Phys. **2** (1942), 127–148.
- [Pop42c] T. Popoviciu, *Sur l'approximation des fonctions continues d'une variable réelle par des polynomes*, Ann. Sci. Univ. Jassy. Sect. I. **28** (1942), 208.
- [Pop43] T. Popoviciu, *Notes sur les généralisations des fonctions convexes d'ordre supérieur. III*, Acad. Roum. Bull. Sect. Sci. **24** (1943), 409–416.
- [Pop44] T. Popoviciu, *Les fonctions convexes*, Hermann et Cie, Paris, 1944.
- [PPŠ98] C. E. M. Pearce, J. E. Pečarić, and V. Šimić, *Stolarsky means and Hadamard's inequality*, J. Math. Anal. Appl. **220** (1998), no. 1, 99–109.
- [Pré01] A. Prékopa, *Discrete higher order convex functions and their applications*, Generalized convexity and generalized monotonicity (Karlovassi, 1999), Lecture Notes in Econom. and Math. Systems, vol. 502, Springer, Berlin, 2001, pp. 21–47.
- [Roc70] R. T. Rockafellar, *Convex Analysis*, Princeton University Press, Princeton, 1970.
- [RV73] A. W. Roberts and D. E. Varberg, *Convex Functions*, Pure and Applied Mathematics, vol. 57, Academic Press, New York–London, 1973.
- [Sad97] E. Sadowska, *Hadamard inequality and a refinement of Jensen inequality for set-valued functions*, Results Math. **32** (1997), no. 3-4, 332–337.
- [Saf92] B. Saffari, *On the Landau-Hadamard inequality*, General Inequalities, 6 (Oberwolfach, 1990) (E. F. Beckenbach, ed.), International Series of Numerical Mathematics, Birkhäuser, Basel, 1992, pp. 459–462.
- [Sán90] J. Sándor, *An application of the Jensen-Hadamard inequality*, Nieuw Arch. Wisk. (4) **8** (1990), no. 1, 63–66.
- [Sán91] J. Sándor, *On the Jensen-Hadamard inequality*, Studia Univ. Babeş-Bolyai Math. **36** (1991), no. 1, 9–15.
- [SJ90] Y. X. Sun and Zh. Sh. Jiang, *A further improvement of Hadamard's inequality*, Math. Appl. **3** (1990), no. 2, 79–82.
- [Sun93] M. B. Sun, *A strengthening of Hadamard's inequality*, Hunan Jiaoyu Xueyuan Xuebao (Ziran Kexue) **11** (1993), no. 5, 156–157.
- [Szeg39] G. Szegő, *Orthogonal Polynomials*, American Math. Society, Providence, Rhode Island, 1939.
- [Toa94] Gh. Toader, *Superadditivity and Hermite-Hadamard's inequalities*, Studia Univ. Babeş-Bolyai Math. **39** (1994), no. 2, 27–32.
- [Vag74] A. I. Vagabov, *A generalization of Hadamard's inequality*, Akad. Nauk Azerbaïdžan. SSR Dokl. **30** (1974), no. 2, 7–8.
- [VL77] P. M. Vasić and I. B. Lacković, *Notes on convex functions. I. A new proof of Hadamard's inequalities*, Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz. (1977), no. 577-598, 21–24.
- [VLM80] P. M. Vasić, I. B. Lacković, and D. M. Maksimović, *Notes on convex functions. IV. On Hadamard's inequalities for weighted arithmetic means*, Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz. (1980), no. 678-715, 199–205 (1981).
- [Wac94] A. Wach, *A note on Hadamard's inequality*, Univ. Iagel. Acta Math. (1994), no. 31, 87–92.

- [Wal92] R. Waldi, *Über ein Analogon zu Hadamards Ungleichung*, Arch. Math. (Basel) **59** (1992), no. 1, 15–20.
- [Woj64] M. Wojtas, *On Hadamard's inequality for the determinants of order non-divisible by 4*, Colloq. Math. **12** (1964), 73–83.
- [WW82] Zh. L. Wang and X. H. Wang, *On an extension of Hadamard inequalities for convex functions*, Chinese Ann. Math. **3** (1982), no. 5, 567–570.
- [YH97] G.-Sh. Yang and M.-Ch. Hong, *A note on Hadamard's inequality*, Tamkang J. Math. **28** (1997), no. 1, 33–37.
- [YT99] G.-Sh. Yang and K.-L. Tseng, *On certain integral inequalities related to Hermite-Hadamard inequalities*, J. Math. Anal. Appl. **239** (1999), no. 1, 180–187.
- [YT03] G.-Sh. Yang and K.-L. Tseng, *Inequalities of Hermite-Hadamard-Fejér type for convex functions and Lipschitzian functions*, Taiwanese J. Math. **7** (2003), no. 3, 433–440.
- [YW96] G.-Sh. Yang and H.-L. Wu, *A refinement of Hadamard's inequality for isotonic linear functionals*, Tamkang J. Math. **27** (1996), no. 4, 327–336.
- [YW97a] G.-Sh. Yang and Ch.-Sh. Wang, *On refinements of Hadamard's inequalities*, Tamkang J. Math. **28** (1997), no. 3, 183–191.
- [YW97b] G.-Sh. Yang and Ch.-Sh. Wang, *Some refinements of Hadamard's inequalities*, Tamkang J. Math. **28** (1997), no. 2, 87–92.
- [Zei86] E. Zeidler, *Nonlinear Functional Analysis and its Applications I.*, Springer, New York—Berlin—Heidelberg, 1986.
- [Zha87] Y. Zhang, *Hadamard inequalities for convex functions*, J. Math. (Wuhan) **7** (1987), no. 4, 385–386.
- [ZY97] X. D. Zhang and Sh. J. Yang, *A note on Hadamard's inequality*, Acta Math. Appl. Sinica **20** (1997), no. 2, 269–274.



**Hermite–Hadamard-type inequalities  
for generalized convex functions**

Értekezés a doktori (Ph.D.) fokozat megszerzése érdekében  
a matematika tudományágban.

Írta: Bessenyei Mihály okleveles matematikus.

Készült a Debreceni Egyetem Matematika doktori programja Analízis  
alprogramja keretében.

Témavezető: Dr. Páles Zsolt

A doktori szigorlati bizottság:

elnök: Dr. ....

tagok: Dr. ....

Dr. ....

A doktori szigorlat időpontja: 200... ..

Az értekezés bírálói:

Dr. ....

Dr. ....

Dr. ....

A bírálóbizottság:

elnök: Dr. ....

tagok: Dr. ....

Dr. ....

Dr. ....

Dr. ....

Az értekezés védésének időpontja: 200... ..