

2009

REVIEW OF FACULTY OF ENGINEERING

Analecta Technica Szegedinensia



UNIVERSITY OF SZEGED
UNIVERSITAS SCIENTIARUM SZEGEDIENSIS
FACULTY OF ENGINEERING



PUBLISHER:

Dr. Antal Véha
Dean, Head of Department
UNIVERSITY OF SZEGED FACULTY OF ENGINEERING

EDITED BY:

Dr. Cecília Hodúr
Vice Dean

Dr. Elisabeth T. Kovács
Professor

Dr. József Gál
Assoc. Professor

Mónika Szilágyi
administrator

PUBLISHER'S-READERS

Dr. József Csanádi, PhD
Dr. Tamás Endődy, PhD
Dr. József Gál, PhD
Dr. László Gulyás, PhD
Dr. Ernő Gyimes, PhD
Dr. János Gyeviki, PhD
Dr. Ottília Bara- Herczegh, PhD
Dr. Cecília Hodúr, PhD
Dr. Katalin Horváth-Almássy, CSc
Dr. Zsuzsanna H. Horváth, PhD
Dr. Gabriella Keczer, PhD
Dr. Elisabeth T. Kovács, CSc
Dr. Judit Krisch, PhD
Dr. Erzsébet Markovics, PhD
Dr. Tibor Marosi, PhD
Dr. György Mészáros, PhD
Dr. Zsuzsanna László, PhD
Dr. Robert Rajkó, PhD
Dr. József Soós, PhD
Dr. István Tibor Tóth, CSc
Dr. Antal Véha, CSc
Dr. Edina Vincze-Lendvai, PhD

NUMBER OF COPIES PRINTED: 100

Norma Nyomdász Kft. Kiadó és Nyomda
6800 Hódmezővásárhely, Rárósi u. 10.

ISSN 1788-6392
UNIVERSITY OF SZEGED FACULTY OF ENGINEERING
H-6724 Szeged, Mars tér 7.
Phone: +36 (62) 546 000

Tünde Baló
*FIAT PANIS! LET THERE BE
WORLD'S FOOD SUPPLY I.*

Ágnes Pongráczné Barancsi, Zoltán
*RESEARCH ON ALVEOLAR
AESTIVUM) VARIETIES*

Sándor Csató, Mária Fekete
*EXAMINATION OF PHISIC
FRUIT PRESSCAKES*

Francis Vasile Dulf
*THE AUTHENTICITY CON-
PROFILE*

János Gyeviki, József Sárosi, Tamás
*SLIDING MODE CONTROL
APPLICATION*

László Hegedűs
HACCP SYSTEM IN FOOD I

Gabriella Keczer
REGIONAL BIASES OF RESI

Richárd Kicsiny
*DEVELOPMENT OF AN EN
HEATING SYSTEMS*

Ágnes Szerenesi, Erika Lakatos, Attila
*NON-THERMAL EFFECT OF
PART I: WATER ELECTROLY-
SIS*

Erika Lakatos, Attila J. Kovács, Ágnes
*NON-THERMAL EFFECT OF
PART II: CELLULOSE ENZY-
ME*

Edina Vincze-Lendvai
*PRIMER KVANTITATÍVUS SU-
MMA*

Ágota Panyor
*THE IMPORTANCE OF NI
"SPECIAL FOODS"*

Péter Ozsváth, Csaba Németh, László
Kinga Horváth, Csilla Vén, Csaba Ba
*RETAIL STORAGE OF PEELI-
NG*

Alfredas Rimkus
*INVESTIGATION OF CAR EN-
ZYME*

Ferenc Szabó
*SYNERGIC DISORDERS IN
WASTE MANAGEMENT PRO-
BLEMS*

Balázs P. Szabó, András Fekete, Anta
*PHYSICO-MECHANICAL IN
VARIETIES*

Orsolya Szigeti, Viktória Szente, Zsolt
Zoltán Szalkály
*CONSUMERS' PREFERENC-
ES FOR TRADITIONAL HUNGARIAN
FOODS*

Dénes Szilágyi
*DYNAMIC MODEL OF ROTO-
RY*

J. Tarek-Tilistyák, J. Agócs, M. D-Tóth
*EVALUATION OF OIL SEED I-
NDICES*

Mohamed Tarek, Zoltán Györi
*ESTIMATING SHELF-LIFE OF
FOODS*

RESEARCH ON ALVEOGRAPHICAL PARAMETERS OF WINTER WHEAT (*T. AESTIVUM*) VARIETIES

Ágnes Pongráczné Barancsi^a, Zoltán Mezei^b, Zoltán Győri^c, Péter Sipos^c

^aTechnological and Agriculture Faculty of Szolnok Collage, 1 Petőfi Square, Mezőtúr H-5400

^bHajdúsági Gabonaiipari Inc., 13 Széchenyi Street, Debrecen H-4025

^cUniversity of Debrecen Centre for Agricultural Sciences and Engineering Faculty of Agronomy, Institute of Food Science, Quality Assurance and Microbiology, 138 Böszörményi Street, Debrecen H-4032
e-mail: postmaster@turagro.t-online.hu

ABSTRACT

Nowadays, many methods have been worked up in the interests of objective expression of winter wheat rheological quality. These define wide spectrum in view of their complexity and apparatus claim.

We have to know the new, untraditional quality parameters because of satisfy both costumers of the Europe Union and the export market. The ambition of importers is shown by the fact that needs of certain suppliers and costumers for the alveographical parameter has increased.

Hungary could come in the uniform EU wheat market with special Hungarian wheat and stabilised good quality if we innovate the rheological characteristics. We have to find extra information about the baking value of winter wheat varieties and the qualification of export rate and we have to select special quality types for wheat growing.

In our research we examined the alveographical parameters of 10 winter wheat varieties (GK Kalász, GK Csillag, GK Élet, GK Garaboly, GK Verecke, GK Csongrád, GK Petur, GK Piacos, GK Kapos, GK Hattyú) in Szeged (Hungary) on the basis of the results in three years (2005-2007).

We established with correlation analysis that is no statistical relation between the W and G; W and L parameters.

1. INTRODUCTION

The definition of quality has widened with meeting the market needs and costumer-oriented approach. More and more claim appear for special rheological characteristics of dough on Hungarian and international wheat market.

The quality of wheat is a complex concept (Lásztity 1980, Véha and Gyimesi 1999, Matuz et al., 2007, Pepó 2007, Tanács 2007).

The Alveograph is suitable for the examination of rheological characteristics which characterises the extensibility of dough (Rakszegi et al., 2005).

The Alveograph qualification of wheat samples is a current method in the French professional word, but it is preferred in West-Europe and Europe Union, too. This method gives extra information for backing tests (Zsikla, 2005). The Alveograph is a modern model of Hankóczy Farinometer. The specialisation of the Alveograph is that: this machine deforms the dough not only in one, but more directions. In the meantime it originates a dough bubble from the dough; in other words it imitates the enlargements in the rising dough. The indexes of alveogram are the followings: P (mm) as pressure, L (mm) as break abscissa, G (ml) as swelling index, W (10^{-4} J) as deform work and P/L as a configure or relative of graph. The interpretation of the alveographical characteristics are equal with the extensographical characteristics (Markovics 2004). The attitude of graph means the resistance to extension, the length of graph means the extensibility, the subarea of graph is the energy (Faridi and Rasper 1987).

Vida et al. (1996) analysed the relation between the alveographical and other baking industry quality characteristics of 19 winter wheat varieties and they established the close positive correlation between the alveographical G, W and gluten index with statistical method. The alveographical G and W are in satisfactory significant relation with the wet gluten content.

Matuz et al. (1999) established the values and the value relation of 13 parameters (among others alveographical P, L, P/L, W, G wet gluten content, spreading of wet gluten) of 29 winter wheat varieties produced in 1995, 1996 and 1997. They used double correlation, factor analysis, polyvariable regression analysis and stepwise regression. The aim of their analyses was to define the parameter that has the closest correlation with the alveographical W. They stated that the majority of correlation is different annually.

2. MATERIALS AND METHODS

The 10 representative winter wheat samples (GK Kalász, GK Csillag, GK Élet, GK Garaboly, GK Verecke, GK Csongrád, GK Petur, GK Piacos, GK Kapos, GK Hattyú), which were produced in 2005-2007, originate from the Keckés breeding station of the Cereal Research Non Profit Company (Hungary). The plot size was 20 m²/variety, and the wheat types were produced without repetition.

The flour was made in Metefém FQC 109 Labor mill. The quality complied with the requirements of the MSZ 6367/9-1989 standard. The specification of the moisture complied with the requirements of the MSZ 6369/4-87 standard. We made Alveograph examinations with Alveograph by Chopin (AACC 1983.54.30.). The Alveograph examinations were performed in the Laboratory of University of Debrecen Centre for Agricultural Sciences and Engineering Faculty of Agronomy, Institute of Food Science, Quality Assurance and Microbiology. We used SPSS 12.0 for Windows, which runs on Microsoft Windows 2000 operation system.

3. RESULTS AND DISCUSSION

The changes of alveographical parameters

The four types in baking industry are grouped according to alveographical W (deform work) in some member states of Europe Union. In table I the GK Kalász variety presents the highest parameter in tree years (in 2005 469*10⁻⁴J/g, in 2006 459*10⁻⁴J/g, in 2007 421*10⁻⁴J/g) and GK Garaboly shows the lowest value in 2005 (83*10⁻⁴J/g), in 2006 (89*10⁻⁴J/g) and GK Hattyú in 2007 (144*10⁻⁴J/g).

We can see that similar situation in the case of average P value, too. GK Kalász shows P=95 mm in 2005, P=93 mm in 2006, P=90 mm in 2007. We analysed the lowest values of GK Garaboly. It was 23 mm in 2005, 29 mm in 2006 (GK Csongrád, too) and 30 mm in 2007.

We can see high L parameters in view of GK Petur in 2005 (253 mm), GK Csongrád in 2006 (203 mm) and in 2007 (209 mm). We analysed low L parameter in view of GK Kapos in 2005 (113 mm), GK Piacos in 2006 (121 mm) and GK Élet in 2007 (103 mm).

We count the G parameter from L, so the G values present similar tendency in tree years. In 2005 and in 2007 the L and G values in view of Petur were the highest parameters and GK Kapos presented the lowest parameters.

Table 1. The alveographical parameter (2007)

year	variety	W (10 ⁻⁴ J)
2005	GK Kalász	469
	GK Csillag	262
	GK Élet	162
	GK Garaboly	83
	GK Verecke	316
	GK Csongrád	116
	GK Petur	199
	GK Piacos	220
	GK Kapos	179
	GK Hattyú	127
2006	GK Kalász	459
	GK Csillag	247
	GK Élet	188
	GK Garaboly	89
	GK Verecke	30
	GK Csongrád	12
	GK Petur	21
	GK Piacos	25
	GK Kapos	21
	GK Hattyú	14
2007	GK Kalász	421
	GK Csillag	247
	GK Élet	21
	GK Garaboly	16
	GK Verecke	27
	GK Csongrád	17
	GK Petur	29
	GK Piacos	29
	GK Kapos	17
	GK Hattyú	144

Correlation among the analyses

In table 2 are close linear pos W and P (0,886), W and P/L (0,886), P and L (-0,750), P and G (-0,750), so there is no sense of the correlation between W and G and W and L.

Table 1. The alveographical parameters of analysed winter wheat varieties (Debrecen, Hungary 2005-2007)

year	variety	W (10 ⁻⁴ J/g)	P (mm)	L (mm)	P/L	G (cm ³)
2005	GK Kalász	469	95	150	0,63	27,19
	GK Csillag	262	82	121	0,68	24,42
	GK Élet	162	47	159	0,30	27,99
	GK Garaboly	83	23	225	0,10	33,30
	GK Verecke	316	74	133	0,56	25,60
	GK Csongrád	116	40	160	0,25	28,08
	GK Petur	194	33	253	0,13	35,31
	GK Piacos	220	53	152	0,35	27,37
	GK Kapos	179	64	113	0,57	23,60
GK Hattyú	122	27	197	0,14	31,16	
2006	GK Kalász	459	93	146	0,63	28,24
	GK Csillag	247	70	130	0,53	25,61
	GK Élet	183	43	163	0,26	25,00
	GK Garaboly	89	29	181	0,16	23,15
	GK Verecke	302	76	152	0,50	28,06
	GK Csongrád	127	29	203	0,14	29,56
	GK Petur	212	53	174	0,30	30,12
	GK Piacos	257	72	121	0,59	24,70
	GK Kapos	214	55	192	0,28	29,56
GK Hattyú	146	52	186	0,27	30,98	
2007	GK Kalász	421	90	147	0,61	27,00
	GK Csillag	248	74	134	0,55	25,80
	GK Élet	213	70	103	0,68	22,60
	GK Garaboly	169	39	167	0,24	28,80
	GK Verecke	271	58	176	0,33	29,50
	GK Csongrád	171	40	209	0,19	32,20
	GK Petur	249	51	201	0,25	31,50
	GK Piacos	243	68	135	0,51	25,80
	GK Kapos	192	72	105	0,69	22,80
GK Hattyú	144	30	200	0,15	31,50	

Correlation among the analysed parameters

In table 2 are close linear positive correlation between the following parameters in 2005: W and P (0,886), W and P/L (0,747). Negative linear correlation is between the followings: P and L (-0,750), P and G (-0,757). We count the P/L value from P and L, the G from L, so there is no sense of the correlation in this situation. There is no statistical relation between W and G and W and L.

Table 2. Correlation among the analysed parameter in 2005

	W (10 ⁻⁴ J/g)	P (mm)	L (mm)	P/L	G (cm ³)
W (10 ⁻⁴ J/g)	1				
P (mm)	**0,886	1			
L (mm)	-0,412	*-0,750	1		
P/L	*0,747	**0,965	**0,856	1	
G (cm ³)	-0,413	*-0,757	**0,998	**0,869	1

** Correlation is significant at the 0.01 level
* Correlation is significant at the 0.05 level

We can see in table 3 similar situation as in table 2. There are close linear positive correlation between W and P (0,936), W and P/L (0,868). There are negative linear correlation between W and L (-0,606), P and L (-0,735). There is no statistical relation between W and G, W and L, P and G.

Table 3. Correlation among the analysed parameter in 2006

	W (10 ⁻⁴ J/g)	P (mm)	L (mm)	P/L	G (cm ³)
W (10 ⁻⁴ J/g)	1				
P (mm)	**0,936	1			
L (mm)	-0,606	*-0,735	1		
P/L	**0,868	**0,957	**0,887	1	
G (cm ³)	0,098	0,076	0,508	-0,151	1

** Correlation is significant at the 0.01 level
* Correlation is significant at the 0.05 level

We present the date in 2007 in table 4. There are close linear positive correlation between W and P (0,769) too. There are negative linear correlation between P and L (-0,743), P and G (-0,730). There is no statistical relation between W and G, W and L, W and P/L.

Table 4. Correlation among the analysed parameter in 2007

	W (10 ⁻⁴ J/g)	P (mm)	L (mm)
W (10 ⁻⁴ J/g)	1		
P (mm)	**0,769	1	
L (mm)	-0,200	*-0,743	1
P/L	0,445	**0,902	**0,902
G (cm ³)	-0,175	*-0,730	**0,902

** Correlation is significant at the 0.01 level
* Correlation is significant at the 0.05 level

The results of correlation analysis positive and negative connection are to each other between W and L parameters.

4. SUMMARY

We have to know the new, untraditional customers of the Europe Union and the gluten examination test is dominant in Hungary could come in the uniform stabilised good quality if we innovate about the conditions of EU and Hungary at them.

The pizza-, strudel-, puff- and creck Polish, German, Dutch and Italian countries have the followings: extensographical parameters. More and more laboratories have claimed parameters, the market and important to give the correct, fast speed extra information about the baking volume export rate and we have to select speed. In our research we examined the alveographical parameters of winter wheat (GK Kalász, GK Csillag, GK Élet, GK Piacos, GK Kapos, GK Hattyú) in years (2005-2007), which were produced at the breeding station of the Cereal Research Institute. The alveographical W value plays an important role in the selection of winter wheat in Europe Union. The GK Kalász variety has the lowest value in 2007. We established with correlation analysis between the W and G; W and L parameters.

Table 4. Correlation among the analysed parameter in 2007

	W (10 ⁻⁴ J/g)	P (mm)	L (mm)	P/L	G (cm ³)
W (10 ⁻⁴ J/g)	1				
P (mm)	**0,769	1			
L (mm)	-0,200	*-0,743	1		
P/L	0,445	**0,902	**0,944	1	
G (cm ³)	-0,175	*-0,730	**0,999	**0,943	1

** Correlation is significant at the 0.01 level
* Correlation is significant at the 0.05 level

The results of correlation analysis present in three years, that W and P positive, P and L negative connection are to each other and there is no statistical relation between W and G, W and L parameters.

4. SUMMARY

We have to know the new, untraditional quality parameters in Hungary, which satisfy the costumers of the Europe Union and the export market, too.

The gluten examination test is dominant in Hungary. It has its own reasons and traditions. Hungary could come in the uniform EU wheat market with special Hungarian wheat and stabilised good quality if we innovate the rheological characteristics. The costumers decide about the conditions of EU and Hungarian market and the research innovations often aim at them.

The pizza-, strudel-, puff- and creeker flour types are curiosity flour types in the French, Polish, German, Dutch and Italian dough flour market. These flour types are required to have the followings: extensographical extensibility and alveographical "P" parameter.

More and more laboratories have Alveograph in Hungary. If we can standardize the claimed parameters, the market and the costumers will be satisfied with Hungary. It is important to give the correct, fast specification of the claimed parameters. We have to find extra information about the baking value of winter wheat varieties and the qualification of export rate and we have to select special quality types for wheat growing.

In our research we examined the alveographical parameters of 10 winter wheat varieties (GK Kalász, GK Csillag, GK Élet, GK Garaboly, GK Verecke, GK Csongrád, GK Petur, GK Piacos, GK Kapos, GK Hattyú) in Szeged (Hungary) on the basis of the results in three years (2005-2007), which were produced in 2005-2007, originate from the Kecskés breeding station of the Cereal Research Non Profit Company.

The alveographical W value plays impotent roll in baking industry in some member states of Europe Union. The GK Kalász variety presents the highest parameter in tree years and GK Garaboly the lowest value in 2005, 2006 and GK Hattyú in 2007.

We established with correlation analysis in three years that is no statistical relation between the W and G; W and L parameters.

This extra information can give us help to select special quality types for wheat growing and qualify the different export rate.

REFERENCES

1. American Association of Cereal Chemists. (1983): Alveograph Method for Soft and Hard Wheat Flour. 54.30A. The Association: St. Paul, MN.
2. Györi és Györiné Mile.I. (1998): A búza minősége és minősítése. Mezőgazda Szaktudás Kiadó, Budapest. 37-44p.
3. Faridi, H., Rasper, V.F. (1987): The Alveograph Handbook. AACC. St.Paul, Minnesota, USA.
4. Lásztity, R. (1980): Correlation between chemical structure and rheological properties of gluten. Ann.Technol. Agric. 29.339-361 p.
5. Lásztity R.-Molnár P. (2001): Búzatermelés és búza minőség az EU országokban. Molnárak Lapja. 106. évf.2001. évi 3. szám.5-7.
6. Markovics E. (2004): Őszi búza lisztek sütési tulajdonságainak összefüggésvizsgálata. PhD Értekezés. DE-ATC, Debrecen.
7. Matuz, J., Markovics E., Ács, E., Véha, A. (1999): Őszi búza fajták lisztjének tulajdonságai közötti összefüggések vizsgálata. Növénytermelés. 3. 243-254 p.
8. Matuz, J., Kirsch J., Véha A., Petróczy, I. M., Tanács L. (2007): Effect of the fertilization and the fungicide treatment on the alveographic quality of winter wheat. Cereal Research Communications, 35: 2. 1193-1196 p.
9. MSZ 6367/9-1989. Élelmezési, takarmányozási, ipari magvak és hántolt termények vizsgálata. A búzaliszt laboratóriumi előállítása.
10. Pepó, P. (2007): The role of fertilization and genotype in sustainable winter wheat (Triticum aestivum L.) production. Cereal Research Communications, 35: 2. 917-920 p.
11. Rakszegi, M., Láng, L., Bedő, Z. (2005): Tészta nyújthatóság vizsgálatok a búzanevelésben. Martonvásár, 2005/1. 12-13 p.
12. Tanács, L. (2007): Seasonal and genotype effect on the alveographical value of winter wheats. Cereal Research Communications, 35: 2. 1197-1200 p.
13. Véha, A., Gyimes, E. (1999): Investigation of kernel hardness in winter wheat varieties with hammermill. Cereal Research Communications, 27. 4. 463-470 p.
14. Vida, Gy., Láng, L., Bedő, Z. (1996): Őszi búzák alveográfus és más sütőipari minőségi tulajdonságai közötti összefüggések elemzése főkomponensanalízissel. Növénytermelés. MTA Mezőgazdasági Kutatóintézet, Martonvásár. Tom.45.No.56.435-445 p.
15. Zsikla, A. (2005): Az alveográfus tézstaviselkedés és a sütési teljesítmény kapcsolatának vizsgálata. <http://www.food.kel.hu/tdk/2004/szekcio3.pdf>. 2005.12.02.

EXAMINATION OF PECTIN EXTRACTED

Sán
University

In our study we aimed to investigate extracted from presscakes of berries, fruits such as black-currant, red-currant examinations. Two commercial mate rheological examinations showed that best gellificator properties. The analysis pectin used as control is very clear a pectin gels of black-currant, blackberry colourants, in this way they are very d

1. INTRODUCTION

The presscake obtained by the quantity of technological was compost. However in case of the remain in the presscake after pectin colorants for the berries. Pectin is well as stabilizer. The presscake quantities of pectin, which is well further examination.

In our study we aimed to examine gels extracted from berries, and commercial apple and citrus pectin

2. MATERIALS AND METHODS

The pectins extracted from the pectin examinations, namely: black-currant commercial materials were applied

Rheological analysis

For the examinations a gel was prepared from 150 ml water+ 5g pectin+ 60g sucrose at the boiling point applying continuous stirring. The analysis was carried out using a rheometer at 20°C. S1 plate was used for red-currant and black-currant pectin