


2-Phenylethyl 1-thio- β -D-galactopyranoside hemihydrate

Iván Brito, László Szilágyi, Ambati Ashok Kumar, Joselyn Albanez and Michael Bolte

Acta Cryst. (2011). E67, o2308–o2309

This open-access article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.



Acta Crystallographica Section E: Structure Reports Online is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. The journal publishes over 4000 structures per year. The average publication time is less than one month.

Crystallography Journals **Online** is available from journals.iucr.org

2-Phenylethyl 1-thio- β -D-galactopyranoside hemihydrate

Iván Brito,^{a*} László Szilágyi,^b Ambati Ashok Kumar,^b Joselyn Albanez^a and Michael Bolte^c

^aDepartamento de Química, Facultad de Ciencias Básicas, Universidad de Antofagasta, Casilla 170, Antofagasta, Chile, ^bDepartment of Organic Chemistry, University of Debrecen, H-4010 Debrecen Pf 20., Hungary, and ^cInstitut für Anorganische Chemie der Goethe-Universität Frankfurt, Max-von-Laue-Strasse 7, D-60438 Frankfurt am Main, Germany
Correspondence e-mail: ivanbritob@yahoo.com


Received 25 July 2011; accepted 5 August 2011

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.010$ Å; R factor = 0.058; wR factor = 0.117; data-to-parameter ratio = 13.8.

The title compound, $\text{C}_{14}\text{H}_{20}\text{O}_5\text{S} \cdot 0.5\text{H}_2\text{O}$, crystallizes with two organic molecules and a solvent water molecule in the asymmetric unit. In both molecules, the hexapyranosyl rings adopt a slightly distorted chair conformation ($^5\text{C}_2$) with four substituents in equatorial positions and one substituent in an axial position. The main difference between the organic molecules is the dihedral angle between the phenyl ring and the best plane defined by the $\text{O}-\text{C}_1-\text{C}_2-\text{C}_3$ atoms (r.m.s deviations = 0.003 and 0.043 Å) of the hexapyranosyl rings [47.4 (4) and 86.5 (4)°]. In the asymmetric unit, molecules are linked by two strong $\text{O}-\text{H} \cdots \text{O}$ hydrogen bonds. In the crystal, the components are linked by a total of 10 distinct $\text{O}-\text{H} \cdots \text{O}$ hydrogen bonds, resulting in the formation of a two-dimensional network parallel to the ab plane.

Related literature

For synthetic methods see: Helferich & Türk (1956). For pharmacological properties of the title compound, see: De Bruyne *et al.* (1977); Choi *et al.* (2003). Gutiérrez *et al.* (2011). For puckering parameters see: Cremer & Pople (1975).



Experimental

Crystal data

$\text{C}_{14}\text{H}_{20}\text{O}_5\text{S} \cdot 0.5\text{H}_2\text{O}$
 $M_r = 309.37$

Orthorhombic, $P2_12_12_1$
 $a = 4.8358$ (4) Å

$b = 14.8218$ (16) Å
 $c = 41.390$ (3) Å
 $V = 2966.6$ (5) Å³
 $Z = 8$

Mo $K\alpha$ radiation
 $\mu = 0.24$ mm⁻¹
 $T = 173$ K
 $0.20 \times 0.09 \times 0.08$ mm

Data collection

Stoe IPDS II two-circle diffractometer
Absorption correction: multi-scan (*MULABS*; Spek, 200; Blessing, 1995)
 $T_{\min} = 0.954$, $T_{\max} = 0.981$

14206 measured reflections
5217 independent reflections
2395 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.169$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.058$
 $wR(F^2) = 0.117$
 $S = 0.68$
5217 reflections
377 parameters
H-atom parameters constrained

$\Delta\rho_{\max} = 0.28$ e Å⁻³
 $\Delta\rho_{\min} = -0.30$ e Å⁻³
Absolute structure: Flack (1983),
2269 Friedel pairs
Flack parameter: -0.25 (16)

Table 1
Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{O31}-\text{H31} \cdots \text{O1W}$	0.84	1.92	2.759 (7)	179
$\text{O41}-\text{H41} \cdots \text{O41A}$	0.84	1.95	2.779 (7)	169
$\text{O51}-\text{H51} \cdots \text{O1W}^{\text{i}}$	0.84	1.98	2.773 (6)	156
$\text{O61}-\text{H61} \cdots \text{O31A}^{\text{ii}}$	0.84	1.86	2.697 (7)	172
$\text{O31A}-\text{H31A} \cdots \text{O61}^{\text{iii}}$	0.84	1.92	2.650 (7)	145
$\text{O41A}-\text{H41A} \cdots \text{O41}^{\text{iv}}$	0.84	2.05	2.788 (7)	147
$\text{O51A}-\text{H51A} \cdots \text{O61A}^{\text{v}}$	0.84	2.10	2.785 (7)	138
$\text{O61A}-\text{H61A} \cdots \text{O31}^{\text{ii}}$	0.84	2.02	2.744 (6)	145
$\text{O1W}-\text{H1WA} \cdots \text{O31}^{\text{v}}$	0.84	2.23	2.989 (8)	150
$\text{O1W}-\text{H1WB} \cdots \text{O41A}^{\text{v}}$	0.84	2.43	3.270 (6)	180

Symmetry codes: (i) $-x, y - \frac{1}{2}, -z + \frac{3}{2}$; (ii) $-x + 1, y - \frac{1}{2}, -z + \frac{3}{2}$; (iii) $-x, y + \frac{1}{2}, -z + \frac{3}{2}$; (iv) $x + 1, y, z$; (v) $x - 1, y, z$.

Data collection: *X-Area* (Stoe & Cie, 2001); cell refinement: *X-Area*; data reduction: *X-Area*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* in *SHELXTL-Plus* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2009).

We thank OTKA, the Hungarian Scientific Research Fund (grant Nos. IN-79731 and NK-68578) for financial support. IB thanks the Spanish Research Council (CSIC) for the provision of a free-of-charge license to the Cambridge Structural Database.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: OM2454).

References

- Blessing, R. H. (1995). *Acta Cryst.* **A51**, 33–38.
- Choi, J. H., Choe, Y. S., Lee, K.-H., Choi, Y., Kim, S. E. & Kim, B.-T. (2003). *Carbohydr. Res.* **338**, 29–34.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- De Bruyne, C. K. & Yde, M. (1977). *Carbohydr. Res.* **56**, 153–164.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.

- Gutiérrez, B., Muñoz, C., Osorio, L., Ambati, A. K., Kövér, K. E., Sagua, H., Araya, J. E., Morales, P., Szilágyi, L. & González, J. (2011). *Acta Trop.* Submitted.
- Helferich, B. & Türk, D. (1956). *Chem. Ber.* **89**, 2215–2219.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Stoe & Cie (2001). *X-AREA*. Stoe & Cie, Darmstadt, Germany.

supplementary materials

Acta Cryst. (2011). E67, o2308–o2309 [doi:10.1107/S1600536811031667]

2-Phenylethyl 1-thio- β -D-galactopyranoside hemihydrate

I. Brito, L. Szilágyi, A. A. Kumar, J. Albanez and M. Bolte

Comment

2-Phenylethyl-1-thio- β -D-galactopyranoside is one of the most potent inhibitors of β -galactosidase (EC 3.2.1.23) (De Bruyne, *et al.*, 1977) and a radiologically labeled derivative has also been used for imaging of *LacZ* gene expression. (Choi *et al.*, 2003). It was recently found to be moderately active in tests against *Trypanosoma cruzi*, the causal agent of Chagas disease (Gutiérrez *et al.*, 2011). In the title compound, it crystallizes with two organic molecules and a solvent water molecule in the asymmetric unit, Fig. 1. In both molecules the hexapyranosyl rings adopts a slightly distorted chair conformation (5C_2) ($Q_T = 0.574$ (6) Å, $\theta = 3.4$ (6)°, $\varphi_2 = 8$ (9)°; $Q_T = 0.587$ (6) Å, $\theta = 9.2$ (7)°, $\varphi_2 = 299$ (4)° for both molecules respectively), (Cremer & Pople, 1975) with four substituents in equatorial positions and one substituent in an axial position. The main difference between the organic molecules is the dihedral angle between the phenyl ring and the best plane defined by the atoms O1/C2/C3/C4 and O1A/C2A/C3A/C4A (r.m.s deviation 0.003 Å; 0.043 Å respectively), of the hexapyranosyl rings [47.4 (4) and 86.5 (4)°]. The max. deviation for the best planes of the hexapyranosyl rings are: 0.290 (7) Å and 0.050 (6) Å for molecules A and B respectively. The mean bond distances are: C—O 1.425 (7) Å, Csp3—Csp3 1.524 (9) Å and aromatic C—C 1.386 (11) Å. In the asymmetric unit the three molecules are linked by two strong O—H...O hydrogen bonds and the crystal packing is stabilized by eight O—H...O hydrogen bonding leading to the formation of a two-dimensional network parallel to the *ab* plane, Fig. 2, Table 1.

Experimental

1-thio-2,3,4,6-tetra-*O*-acetyl- β -D-galactopyranose (0.364 g, 1 mmol) was dissolved in acetonitrile (2 ml) and 1-bromo-2-phenylethane (0.185 g, 1 mmol) and triethyl amine (242 μ l, 2 mmol) added. The reaction mixture was stirred at RT until disappearance of the starting materials (TLC, 60 min). The solvent was removed under reduced pressure, and the residue purified by column chromatography (EtOAc: hexane - 8: 2) to give 2-phenylethyl 2,3,4,6-tetra-*O*-acetyl-1-thio- β -D-galactopyranoside (**1**). Syrup, 402 mg (86%). $[\alpha]_D -22.9$ (CHCl₃, c 0.15), Lit. (Helferich & Türk, 1956). $[\alpha]_D -19.2$ (CHCl₃). HR—MS: *m/z* calcd. for C₂₂H₂₈O₉S[M+Na]⁺: 491.135. Found: 491.138. ¹H-NMR (CDCl₃, 500 MHz): δ 7.20–7.35 (m, 5H, Phenyl-H); 5.44 (br.s, 1H, H-4); 5.26 (t, 1H, H-2, $J_{2,3}$ 9.9 Hz); 5.03 (br.d, 1H, H-3); 4.44 (d, 1H, H-1, $J_{1,2}$ 10.1 Hz); 4.10–4.20 (m, 2H, H-6a,b); 3.90 (m, 1H, H-5); 2.92 (m, 1H, S-CH_{2a}); 2.94 (m, 2H, Ph-CH₂); 3.00 (m, 1H, S-CH_{2b}); 2.16, 2.06, 2.04, 1.99 (s, 4x3H, 4x COCH₃); ¹³C-NMR (CDCl₃, 125 MHz): δ 130.5 (Phenyl-C); 85.6 (C-1); 76.5 (C-5); 73.3 (C-3); 69.0 (C-4); 68.7 (C-2); 62.2 (C-6); 38.2 (Ph-CH₂); 33.1 (S-CH₂); 22.3 (4xCOCH₃). The product (0.300 g, 0.64 mmol) was deacetylated by treatment with catalytic amount of NaOMe in methanol. The reaction mixture was stirred at room temperature until completion (TLC 20 min). After neutralization with a cation exchanger (Amberlyst 15) the solvent was removed under reduced pressure and the title molecule, 2-phenylethyl-1-thio- β -D-galactopyranoside, was isolated as a white solid (MeOH: EtOAc - 2:8), 185 mg (96.3%). $[\alpha]_D -22.4$ (MeOH, c 0.11). Lit. (Helferich & Türk, 1956) $[\alpha]_D -32.2$ (MeOH). HR—MS: *m/z* calcd. for C₁₄H₂₀O₅S[M+Na]⁺: 323.094. Found: 323.094.

$^1\text{H-NMR}$ (CD_3OD , 500 MHz): δ 7.15–7.30 (m, 5H, Phenyl-H); 4.34 (d, 1H, H-1, $J_{1,2}$ 9.6 Hz); 3.90 (dd, 1H, H-4, $J_{4,5} \sim 1$ Hz); 3.77(dd, 1H, H-6a, $J_{6a,6b}$ 11.5 Hz, $J_{5,6a}$ 6.6 Hz); 3.71(dd, 1H, H-6 b, $J_{5,6b}$ 5.3 Hz); 3.57 (t, 1H, H-2, $J_{2,3}$ 9.6 Hz); 3.53(m, 1H, H-5); 3.46 (1H, dd, H-3, $J_{3,4}$ 3.4 Hz); 2.92 (m, 1H, $S\text{-CH}_2\text{a}$); 2.95 (m, 2H, Ph-CH_2); 3.02(m, 1H, $S\text{-CH}_2\text{b}$); $^{13}\text{C-NMR}$ (CDCl_3 , 125 MHz): δ 131.3, 131.0, 128.9 (Phenyl-C); 88.4 (C-1); 81.5 (C-5); 77.1 (C-3); 72.3 (C-2); 71.3 (C-4); 63.5 (C-6); 38.6 (Ph-CH_2); 33.2 ($S\text{-CH}_2$). Colourless single crystals suitable for X-ray analysis were obtained by slow evaporation of an aqueous solution.

Refinement

All H atoms could be located by difference Fourier synthesis but were ultimately placed in calculated positions using a riding model with C—H = 0.95 - 1.00 Å and O—H = 0.84 Å with fixed individual displacement parameters [$U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$ or $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{O})$].

Figures




Fig. 1. Perspective view of the asymmetric unit of the title compound, with the atom numbering. Displacement ellipsoids are shown at the 50% probability level.




Fig. 2. A Packing diagram, viewed down the c axis.

2-Phenylethyl 1-thio- β -D-galactopyranoside monohydrate

Crystal data

$\text{C}_{14}\text{H}_{20}\text{O}_5\text{S} \cdot 0.5\text{H}_2\text{O}$

$M_r = 309.37$

Orthorhombic, $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a = 4.8358$ (4) Å

$b = 14.8218$ (16) Å

$F(000) = 1320$

$D_x = 1.385$ Mg m $^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3425 reflections

$\theta = 2.8\text{--}25.6^\circ$

$\mu = 0.24$ mm $^{-1}$

$c = 41.390 (3) \text{ \AA}$
 $V = 2966.6 (5) \text{ \AA}^3$
 $Z = 8$

$T = 173 \text{ K}$
 Needle, colourless
 $0.20 \times 0.09 \times 0.08 \text{ mm}$

Data collection

Stoe IPDS II two-circle
 diffractometer
 Radiation source: fine-focus sealed tube
 graphite
 ω scans
 Absorption correction: multi-scan
 (MULABS; Spek, 200; Blessing, 1995)
 $T_{\min} = 0.954$, $T_{\max} = 0.981$
 14206 measured reflections

5217 independent reflections
 2395 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.169$
 $\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 2.8^\circ$
 $h = -4 \rightarrow 5$
 $k = -17 \rightarrow 17$
 $l = -49 \rightarrow 47$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.058$
 $wR(F^2) = 0.117$
 $S = 0.68$
 5217 reflections
 377 parameters
 0 restraints
 Primary atom site location: structure-invariant direct
 methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring
 sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0007P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.28 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.30 \text{ e \AA}^{-3}$
 Absolute structure: Flack (1983), 2269 Friedel pairs
 Flack parameter: $-0.25 (16)$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.3816 (4)	0.92470 (10)	0.85163 (4)	0.0217 (4)
O1	0.2305 (9)	0.7771 (2)	0.82113 (11)	0.0186 (11)

supplementary materials

C2	0.1463 (14)	0.8684 (3)	0.82448 (14)	0.0171 (14)
H2	−0.0432	0.8697	0.8341	0.020*
C3	0.1355 (15)	0.9128 (3)	0.79134 (14)	0.0192 (15)
H3	0.3253	0.9114	0.7818	0.023*
C4	−0.0578 (14)	0.8611 (4)	0.76924 (14)	0.0150 (14)
H4	−0.2469	0.8656	0.7789	0.018*
C5	0.0165 (15)	0.7603 (4)	0.76836 (16)	0.0206 (17)
H5	−0.1313	0.7263	0.7566	0.025*
C6	0.0370 (14)	0.7259 (4)	0.80286 (15)	0.0176 (15)
H6	−0.1488	0.7321	0.8132	0.021*
C7	0.1644 (18)	0.9413 (4)	0.88681 (16)	0.0298 (18)
H7A	0.2706	0.9759	0.9031	0.036*
H7B	0.0036	0.9785	0.8804	0.036*
C8	0.0570 (17)	0.8546 (4)	0.90277 (18)	0.0335 (19)
H8A	−0.0909	0.8706	0.9183	0.040*
H8B	−0.0258	0.8156	0.8859	0.040*
C11	0.2787 (17)	0.8018 (4)	0.92039 (17)	0.0302 (19)
C12	0.3759 (19)	0.8276 (4)	0.95047 (17)	0.036 (2)
H12	0.3035	0.8805	0.9603	0.043*
C13	0.5745 (19)	0.7786 (5)	0.96658 (19)	0.040 (2)
H13	0.6369	0.7982	0.9872	0.049*
C14	0.684 (2)	0.7011 (4)	0.9531 (2)	0.044 (2)
H14	0.8212	0.6671	0.9641	0.053*
C15	0.588 (2)	0.6744 (5)	0.9230 (2)	0.050 (3)
H15	0.6588	0.6205	0.9137	0.060*
C16	0.395 (2)	0.7234 (4)	0.90633 (19)	0.042 (2)
H16	0.3392	0.7048	0.8854	0.050*
O31	0.0486 (12)	1.0048 (3)	0.79387 (11)	0.0352 (14)
H31	−0.0953	1.0262	0.7855	0.053*
O41	−0.0739 (10)	0.8992 (3)	0.73781 (10)	0.0234 (11)
H41	0.0863	0.9052	0.7303	0.035*
O51	0.2726 (10)	0.7497 (2)	0.75186 (12)	0.0239 (12)
H51	0.3191	0.6951	0.7521	0.036*
C61	0.1245 (16)	0.6275 (4)	0.80496 (16)	0.0254 (16)
H61B	0.0055	0.5902	0.7908	0.030*
H61C	0.3182	0.6210	0.7976	0.030*
O61	0.0999 (11)	0.5973 (3)	0.83768 (11)	0.0276 (11)
H61	0.2498	0.5739	0.8436	0.041*
S1A	0.1800 (4)	0.92289 (11)	0.58806 (4)	0.0207 (4)
O1A	0.4361 (10)	0.7963 (2)	0.62109 (10)	0.0185 (11)
C2A	0.4278 (15)	0.8930 (4)	0.61866 (14)	0.0174 (15)
H2A	0.6142	0.9150	0.6118	0.021*
C3A	0.3507 (14)	0.9383 (3)	0.65050 (15)	0.0162 (14)
H3A	0.1519	0.9264	0.6556	0.019*
C4A	0.5327 (14)	0.9033 (4)	0.67780 (14)	0.0166 (15)
H4A	0.7246	0.9268	0.6747	0.020*
C5A	0.5420 (14)	0.8010 (4)	0.67805 (16)	0.0170 (15)
H5A	0.6784	0.7805	0.6947	0.020*
C6A	0.6377 (15)	0.7696 (3)	0.64495 (15)	0.0171 (14)

H6A	0.8174	0.7998	0.6398	0.021*
C7A	0.3763 (16)	0.8986 (4)	0.55177 (15)	0.0246 (16)
H7A1	0.5560	0.9305	0.5526	0.030*
H7A2	0.4135	0.8330	0.5504	0.030*
C8A	0.2146 (16)	0.9289 (5)	0.52203 (15)	0.0298 (18)
H8A1	0.0333	0.8979	0.5216	0.036*
H8A2	0.1805	0.9946	0.5233	0.036*
C11A	0.3715 (16)	0.9077 (4)	0.49135 (15)	0.0265 (16)
C12A	0.5655 (18)	0.9652 (5)	0.47895 (18)	0.0323 (19)
H12A	0.6040	1.0207	0.4896	0.039*
C13A	0.7059 (17)	0.9425 (5)	0.45085 (19)	0.042 (2)
H13A	0.8432	0.9822	0.4426	0.050*
C14A	0.650 (2)	0.8640 (6)	0.43483 (19)	0.046 (2)
H14A	0.7484	0.8492	0.4157	0.055*
C15A	0.4537 (19)	0.8077 (5)	0.44622 (19)	0.039 (2)
H15A	0.4146	0.7536	0.4348	0.047*
C16A	0.3062 (18)	0.8271 (4)	0.47450 (17)	0.034 (2)
H16A	0.1666	0.7874	0.4822	0.041*
O31A	0.3948 (11)	1.0327 (2)	0.64734 (11)	0.0213 (10)
H31A	0.2432	1.0585	0.6437	0.032*
O41A	0.4250 (10)	0.9385 (3)	0.70761 (10)	0.0241 (11)
H41A	0.5551	0.9456	0.7208	0.036*
O51A	0.2770 (10)	0.7663 (3)	0.68599 (11)	0.0261 (12)
H51A	0.2409	0.7220	0.6740	0.039*
C61A	0.6753 (15)	0.6677 (4)	0.64172 (16)	0.0196 (15)
H61D	0.5043	0.6362	0.6484	0.023*
H61E	0.7153	0.6517	0.6190	0.023*
O61A	0.8991 (13)	0.6415 (3)	0.66183 (13)	0.0420 (15)
H61A	0.8688	0.5898	0.6693	0.063*
O1W	−0.4231 (11)	1.0767 (3)	0.76658 (13)	0.0445 (14)
H1WA	−0.5761	1.0770	0.7762	0.067*
H1WB	−0.4623	1.0412	0.7514	0.067*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0215 (10)	0.0253 (7)	0.0184 (9)	0.0002 (8)	−0.0020 (8)	−0.0018 (7)
O1	0.016 (3)	0.018 (2)	0.022 (3)	0.0012 (18)	−0.004 (2)	0.0002 (19)
C2	0.016 (4)	0.020 (3)	0.015 (4)	0.003 (3)	−0.001 (3)	0.002 (3)
C3	0.029 (4)	0.015 (3)	0.014 (3)	0.003 (3)	−0.005 (3)	−0.004 (2)
C4	0.012 (4)	0.024 (3)	0.009 (4)	0.004 (3)	0.003 (3)	0.006 (2)
C5	0.020 (4)	0.025 (3)	0.017 (4)	−0.003 (3)	0.003 (3)	−0.003 (3)
C6	0.019 (4)	0.017 (3)	0.017 (4)	−0.002 (3)	−0.001 (3)	−0.003 (3)
C7	0.038 (5)	0.035 (4)	0.017 (4)	0.004 (4)	0.000 (4)	0.000 (3)
C8	0.033 (5)	0.039 (4)	0.028 (5)	0.007 (4)	0.003 (4)	0.003 (3)
C11	0.041 (5)	0.024 (3)	0.025 (5)	−0.010 (3)	0.002 (4)	0.007 (3)
C12	0.044 (6)	0.038 (4)	0.024 (4)	0.011 (4)	0.011 (4)	0.002 (3)
C13	0.051 (6)	0.040 (4)	0.030 (5)	−0.010 (4)	0.004 (5)	0.008 (4)

supplementary materials

C14	0.063 (7)	0.028 (4)	0.041 (5)	0.004 (4)	−0.013 (5)	0.007 (3)
C15	0.066 (7)	0.033 (4)	0.053 (6)	0.011 (5)	−0.001 (6)	−0.004 (4)
C16	0.062 (6)	0.034 (4)	0.029 (5)	−0.005 (4)	−0.013 (5)	−0.004 (3)
O31	0.055 (4)	0.021 (2)	0.029 (3)	0.020 (2)	−0.018 (3)	0.001 (2)
O41	0.021 (3)	0.029 (2)	0.021 (3)	0.000 (2)	−0.005 (2)	0.0035 (19)
O51	0.034 (3)	0.013 (2)	0.024 (3)	0.0057 (19)	0.002 (3)	0.0008 (19)
C61	0.019 (4)	0.024 (3)	0.033 (4)	−0.009 (3)	0.002 (4)	0.001 (3)
O61	0.023 (3)	0.028 (2)	0.032 (3)	0.001 (2)	0.002 (2)	0.0133 (19)
S1A	0.0180 (9)	0.0286 (8)	0.0154 (9)	0.0032 (7)	−0.0022 (7)	−0.0005 (7)
O1A	0.020 (3)	0.0161 (19)	0.020 (3)	0.0050 (19)	−0.006 (2)	0.0004 (17)
C2A	0.017 (4)	0.021 (3)	0.014 (4)	0.002 (3)	−0.003 (3)	0.004 (2)
C3A	0.015 (4)	0.015 (3)	0.019 (3)	−0.003 (3)	0.000 (3)	−0.002 (3)
C4A	0.020 (4)	0.021 (3)	0.008 (3)	−0.005 (3)	−0.007 (3)	−0.002 (3)
C5A	0.014 (4)	0.019 (3)	0.018 (4)	−0.002 (3)	−0.002 (3)	0.002 (3)
C6A	0.015 (4)	0.018 (3)	0.019 (4)	0.003 (3)	−0.003 (3)	0.002 (3)
C7A	0.029 (4)	0.033 (3)	0.012 (4)	0.004 (3)	0.003 (4)	0.007 (3)
C8A	0.043 (5)	0.031 (3)	0.016 (4)	0.014 (4)	−0.003 (3)	−0.003 (3)
C11A	0.026 (4)	0.035 (4)	0.018 (4)	0.007 (4)	−0.006 (3)	0.001 (3)
C12A	0.034 (5)	0.035 (4)	0.028 (4)	0.002 (4)	−0.004 (4)	0.002 (3)
C13A	0.025 (5)	0.067 (5)	0.032 (5)	−0.001 (4)	0.004 (4)	0.016 (4)
C14A	0.042 (6)	0.074 (5)	0.022 (5)	0.023 (5)	0.000 (4)	0.000 (4)
C15A	0.044 (6)	0.042 (4)	0.032 (5)	0.013 (4)	0.005 (5)	−0.011 (4)
C16A	0.051 (6)	0.031 (4)	0.022 (4)	0.007 (4)	0.002 (4)	0.005 (3)
O31A	0.022 (3)	0.0197 (19)	0.023 (3)	−0.002 (2)	−0.008 (3)	0.002 (2)
O41A	0.032 (3)	0.027 (2)	0.014 (2)	−0.005 (2)	−0.003 (2)	−0.0025 (19)
O51A	0.029 (3)	0.028 (2)	0.021 (3)	−0.012 (2)	0.003 (2)	−0.001 (2)
C61A	0.020 (4)	0.021 (3)	0.018 (4)	0.001 (3)	0.003 (3)	0.000 (3)
O61A	0.050 (4)	0.019 (2)	0.057 (4)	0.005 (3)	−0.021 (3)	0.007 (2)
O1W	0.026 (3)	0.049 (3)	0.058 (4)	−0.015 (3)	0.005 (3)	−0.002 (3)

Geometric parameters (\AA , $^\circ$)

S1—C2	1.804 (7)	S1A—C7A	1.813 (7)
S1—C7	1.812 (7)	O1A—C2A	1.437 (6)
O1—C2	1.420 (7)	O1A—C6A	1.443 (8)
O1—C6	1.422 (7)	C2A—C3A	1.526 (8)
C2—C3	1.522 (8)	C2A—H2A	1.0000
C2—H2	1.0000	C3A—O31A	1.421 (6)
C3—O31	1.432 (7)	C3A—C4A	1.523 (8)
C3—C4	1.516 (9)	C3A—H3A	1.0000
C3—H3	1.0000	C4A—O41A	1.437 (7)
C4—O41	1.420 (7)	C4A—C5A	1.517 (8)
C4—C5	1.537 (8)	C4A—H4A	1.0000
C4—H4	1.0000	C5A—O51A	1.419 (8)
C5—O51	1.423 (8)	C5A—C6A	1.519 (9)
C5—C6	1.519 (9)	C5A—H5A	1.0000
C5—H5	1.0000	C6A—C61A	1.527 (7)
C6—C61	1.521 (8)	C6A—H6A	1.0000
C6—H6	1.0000	C7A—C8A	1.525 (9)

C7—C8	1.535 (9)	C7A—H7A1	0.9900
C7—H7A	0.9900	C7A—H7A2	0.9900
C7—H7B	0.9900	C8A—C11A	1.512 (9)
C8—C11	1.515 (10)	C8A—H8A1	0.9900
C8—H8A	0.9900	C8A—H8A2	0.9900
C8—H8B	0.9900	C11A—C12A	1.368 (11)
C11—C12	1.385 (10)	C11A—C16A	1.419 (9)
C11—C16	1.416 (10)	C12A—C13A	1.388 (10)
C12—C13	1.376 (11)	C12A—H12A	0.9500
C12—H12	0.9500	C13A—C14A	1.366 (11)
C13—C14	1.383 (10)	C13A—H13A	0.9500
C13—H13	0.9500	C14A—C15A	1.348 (12)
C14—C15	1.386 (11)	C14A—H14A	0.9500
C14—H14	0.9500	C15A—C16A	1.400 (10)
C15—C16	1.371 (12)	C15A—H15A	0.9500
C15—H15	0.9500	C16A—H16A	0.9500
C16—H16	0.9500	O31A—H31A	0.8400
O31—H31	0.8395	O41A—H41A	0.8400
O41—H41	0.8400	O51A—H51A	0.8400
O51—H51	0.8400	C61A—O61A	1.420 (8)
C61—O61	1.431 (8)	C61A—H61D	0.9900
C61—H61B	0.9900	C61A—H61E	0.9900
C61—H61C	0.9900	O61A—H61A	0.8400
O61—H61	0.8400	O1W—H1WA	0.8394
S1A—C2A	1.799 (7)	O1W—H1WB	0.8399
C2—S1—C7	101.4 (4)	C2A—O1A—C6A	109.8 (4)
C2—O1—C6	111.8 (5)	O1A—C2A—C3A	112.7 (5)
O1—C2—C3	109.5 (5)	O1A—C2A—S1A	108.3 (4)
O1—C2—S1	108.7 (4)	C3A—C2A—S1A	109.7 (4)
C3—C2—S1	112.5 (4)	O1A—C2A—H2A	108.7
O1—C2—H2	108.7	C3A—C2A—H2A	108.7
C3—C2—H2	108.7	S1A—C2A—H2A	108.7
S1—C2—H2	108.7	O31A—C3A—C4A	108.5 (5)
O31—C3—C4	110.2 (5)	O31A—C3A—C2A	108.5 (5)
O31—C3—C2	110.9 (5)	C4A—C3A—C2A	110.4 (5)
C4—C3—C2	110.2 (5)	O31A—C3A—H3A	109.8
O31—C3—H3	108.5	C4A—C3A—H3A	109.8
C4—C3—H3	108.5	C2A—C3A—H3A	109.8
C2—C3—H3	108.5	O41A—C4A—C5A	111.6 (5)
O41—C4—C3	112.7 (5)	O41A—C4A—C3A	107.7 (5)
O41—C4—C5	112.2 (5)	C5A—C4A—C3A	111.3 (5)
C3—C4—C5	111.2 (5)	O41A—C4A—H4A	108.7
O41—C4—H4	106.8	C5A—C4A—H4A	108.7
C3—C4—H4	106.8	C3A—C4A—H4A	108.7
C5—C4—H4	106.8	O51A—C5A—C4A	109.7 (5)
O51—C5—C6	110.9 (6)	O51A—C5A—C6A	111.9 (5)
O51—C5—C4	108.8 (5)	C4A—C5A—C6A	108.0 (5)
C6—C5—C4	108.6 (5)	O51A—C5A—H5A	109.1
O51—C5—H5	109.5	C4A—C5A—H5A	109.1

supplementary materials

C6—C5—H5	109.5	C6A—C5A—H5A	109.1
C4—C5—H5	109.5	O1A—C6A—C5A	109.1 (5)
O1—C6—C5	111.3 (5)	O1A—C6A—C61A	106.9 (5)
O1—C6—C61	107.4 (5)	C5A—C6A—C61A	114.7 (5)
C5—C6—C61	113.2 (5)	O1A—C6A—H6A	108.6
O1—C6—H6	108.3	C5A—C6A—H6A	108.6
C5—C6—H6	108.3	C61A—C6A—H6A	108.6
C61—C6—H6	108.3	C8A—C7A—S1A	110.0 (5)
C8—C7—S1	115.4 (5)	C8A—C7A—H7A1	109.7
C8—C7—H7A	108.4	S1A—C7A—H7A1	109.7
S1—C7—H7A	108.4	C8A—C7A—H7A2	109.7
C8—C7—H7B	108.4	S1A—C7A—H7A2	109.7
S1—C7—H7B	108.4	H7A1—C7A—H7A2	108.2
H7A—C7—H7B	107.5	C11A—C8A—C7A	111.1 (6)
C11—C8—C7	113.6 (7)	C11A—C8A—H8A1	109.4
C11—C8—H8A	108.8	C7A—C8A—H8A1	109.4
C7—C8—H8A	108.8	C11A—C8A—H8A2	109.4
C11—C8—H8B	108.8	C7A—C8A—H8A2	109.4
C7—C8—H8B	108.8	H8A1—C8A—H8A2	108.0
H8A—C8—H8B	107.7	C12A—C11A—C16A	119.6 (7)
C12—C11—C16	117.5 (7)	C12A—C11A—C8A	122.0 (6)
C12—C11—C8	122.0 (6)	C16A—C11A—C8A	118.4 (7)
C16—C11—C8	120.5 (7)	C11A—C12A—C13A	119.9 (7)
C13—C12—C11	121.8 (7)	C11A—C12A—H12A	120.1
C13—C12—H12	119.1	C13A—C12A—H12A	120.1
C11—C12—H12	119.1	C14A—C13A—C12A	121.1 (8)
C12—C13—C14	120.7 (8)	C14A—C13A—H13A	119.5
C12—C13—H13	119.7	C12A—C13A—H13A	119.5
C14—C13—H13	119.7	C15A—C14A—C13A	119.8 (8)
C13—C14—C15	118.2 (8)	C15A—C14A—H14A	120.1
C13—C14—H14	120.9	C13A—C14A—H14A	120.1
C15—C14—H14	120.9	C14A—C15A—C16A	121.6 (7)
C16—C15—C14	121.9 (8)	C14A—C15A—H15A	119.2
C16—C15—H15	119.0	C16A—C15A—H15A	119.2
C14—C15—H15	119.0	C15A—C16A—C11A	118.0 (8)
C15—C16—C11	119.9 (7)	C15A—C16A—H16A	121.0
C15—C16—H16	120.0	C11A—C16A—H16A	121.0
C11—C16—H16	120.0	C3A—O31A—H31A	109.5
C3—O31—H31	125.0	C4A—O41A—H41A	109.5
C4—O41—H41	109.5	C5A—O51A—H51A	109.5
C5—O51—H51	109.5	O61A—C61A—C6A	108.1 (5)
O61—C61—C6	109.3 (5)	O61A—C61A—H61D	110.1
O61—C61—H61B	109.8	C6A—C61A—H61D	110.1
C6—C61—H61B	109.8	O61A—C61A—H61E	110.1
O61—C61—H61C	109.8	C6A—C61A—H61E	110.1
C6—C61—H61C	109.8	H61D—C61A—H61E	108.4
H61B—C61—H61C	108.3	C61A—O61A—H61A	109.5
C61—O61—H61	109.5	H1WA—O1W—H1WB	99.1
C2A—S1A—C7A	100.7 (3)		

C6—O1—C2—C3	−63.2 (7)	C6A—O1A—C2A—C3A	−59.8 (7)
C6—O1—C2—S1	173.6 (4)	C6A—O1A—C2A—S1A	178.7 (4)
C7—S1—C2—O1	−109.5 (4)	C7A—S1A—C2A—O1A	−76.8 (5)
C7—S1—C2—C3	129.0 (5)	C7A—S1A—C2A—C3A	159.8 (4)
O1—C2—C3—O31	179.5 (5)	O1A—C2A—C3A—O31A	169.9 (5)
S1—C2—C3—O31	−59.5 (6)	S1A—C2A—C3A—O31A	−69.4 (6)
O1—C2—C3—C4	57.2 (7)	O1A—C2A—C3A—C4A	51.1 (7)
S1—C2—C3—C4	178.2 (4)	S1A—C2A—C3A—C4A	171.8 (4)
O31—C3—C4—O41	57.8 (7)	O31A—C3A—C4A—O41A	69.2 (6)
C2—C3—C4—O41	−179.5 (5)	C2A—C3A—C4A—O41A	−171.9 (5)
O31—C3—C4—C5	−175.2 (5)	O31A—C3A—C4A—C5A	−168.2 (5)
C2—C3—C4—C5	−52.5 (7)	C2A—C3A—C4A—C5A	−49.3 (7)
O41—C4—C5—O51	57.6 (7)	O41A—C4A—C5A—O51A	53.7 (7)
C3—C4—C5—O51	−69.6 (6)	C3A—C4A—C5A—O51A	−66.6 (7)
O41—C4—C5—C6	178.4 (6)	O41A—C4A—C5A—C6A	175.9 (5)
C3—C4—C5—C6	51.3 (7)	C3A—C4A—C5A—C6A	55.6 (7)
C2—O1—C6—C5	63.8 (7)	C2A—O1A—C6A—C5A	66.0 (6)
C2—O1—C6—C61	−171.8 (5)	C2A—O1A—C6A—C61A	−169.4 (5)
O51—C5—C6—O1	63.5 (6)	O51A—C5A—C6A—O1A	57.2 (6)
C4—C5—C6—O1	−56.0 (7)	C4A—C5A—C6A—O1A	−63.6 (6)
O51—C5—C6—C61	−57.5 (7)	O51A—C5A—C6A—C61A	−62.7 (7)
C4—C5—C6—C61	−177.1 (5)	C4A—C5A—C6A—C61A	176.5 (6)
C2—S1—C7—C8	61.9 (6)	C2A—S1A—C7A—C8A	−174.0 (5)
S1—C7—C8—C11	71.3 (7)	S1A—C7A—C8A—C11A	−178.9 (5)
C7—C8—C11—C12	77.3 (9)	C7A—C8A—C11A—C12A	−86.0 (8)
C7—C8—C11—C16	−102.6 (8)	C7A—C8A—C11A—C16A	96.4 (8)
C16—C11—C12—C13	−1.1 (12)	C16A—C11A—C12A—C13A	−3.0 (11)
C8—C11—C12—C13	179.0 (8)	C8A—C11A—C12A—C13A	179.5 (7)
C11—C12—C13—C14	−0.1 (13)	C11A—C12A—C13A—C14A	1.4 (12)
C12—C13—C14—C15	0.0 (13)	C12A—C13A—C14A—C15A	0.5 (12)
C13—C14—C15—C16	1.4 (14)	C13A—C14A—C15A—C16A	−0.7 (13)
C14—C15—C16—C11	−2.7 (14)	C14A—C15A—C16A—C11A	−0.9 (12)
C12—C11—C16—C15	2.5 (12)	C12A—C11A—C16A—C15A	2.7 (11)
C8—C11—C16—C15	−177.7 (8)	C8A—C11A—C16A—C15A	−179.6 (7)
O1—C6—C61—O61	65.0 (7)	O1A—C6A—C61A—O61A	172.2 (5)
C5—C6—C61—O61	−171.8 (6)	C5A—C6A—C61A—O61A	−66.6 (7)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O31—H31 \cdots O1W	0.84	1.92	2.759 (7)	179.
O41—H41 \cdots O41A	0.84	1.95	2.779 (7)	169.
O51—H51 \cdots O1W ⁱ	0.84	1.98	2.773 (6)	156.
O61—H61 \cdots O31A ⁱⁱ	0.84	1.86	2.697 (7)	172.
O31A—H31A \cdots O61 ⁱⁱⁱ	0.84	1.92	2.650 (7)	145.
O41A—H41A \cdots O41 ^{iv}	0.84	2.05	2.788 (7)	147.
O51A—H51A \cdots O61A ^v	0.84	2.10	2.785 (7)	138.
O61A—H61A \cdots O31 ⁱⁱ	0.84	2.02	2.744 (6)	145.

supplementary materials

O1W—H1WA...O31 ^v	0.84	2.23	2.989 (8)	150.
O1W—H1WB...O41A ^v	0.84	2.43	3.270 (6)	180.

Symmetry codes: (i) $-x, y-1/2, -z+3/2$; (ii) $-x+1, y-1/2, -z+3/2$; (iii) $-x, y+1/2, -z+3/2$; (iv) $x+1, y, z$; (v) $x-1, y, z$.

Fig. 1




Fig. 2

