

Geophysical Research Letters[®]



RESEARCH LETTER

10.1029/2022GL100698

Key Points:

- Sulfur ions were implanted into CO₂ ices at 20 and 70 K to simulate Jovian magnetospheric radiation chemistry at the surface of Europa
- SO₂ was observed to be among the radiolytic products at 20 K, but not at the more Europa-relevant temperature of 70 K
- Alternative explanations for the formation of SO₂ on the surface of Europa should be considered

Correspondence to:

D. V. Mifsud and Z. Kaňuchová,
dm618@kent.ac.uk;
zkanuch@ta3.sk

Citation:

Mifsud, D. V., Kaňuchová, Z., Herczku, P., Juhász, Z., Kovács, S. T. S., Lakatos, G., et al. (2022). Sulfur ion implantations into condensed CO₂: Implications for Europa. *Geophysical Research Letters*, 49, e2022GL100698. <https://doi.org/10.1029/2022GL100698>

Received 5 AUG 2022

Accepted 7 DEC 2022

Author Contributions:

Conceptualization: D. V. Mifsud, Z. Kaňuchová

Data curation: Z. Kaňuchová

Formal analysis: Z. Kaňuchová, Z.

Juhász, B. Sulik

Funding acquisition: D. V. Mifsud, Z. Kaňuchová, Z. Juhász, B. Sulik, S. Ioppolo, N. J. Mason

Investigation: D. V. Mifsud, P. Herczku, Z. Juhász, S. T. S. Kovács, G. Lakatos, K. K. Rahul, B. Sulik

Methodology: D. V. Mifsud, P. Herczku, G. Lakatos, K. K. Rahul, S. Ioppolo, R. W. McCullough

Project Administration: D. V. Mifsud, I. Rajta, I. Vajda
















Resources: R. Rácz, S. Biri, I. Rajta, I. Vajda

Supervision: N. J. Mason

© 2022 The Authors.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial License](https://creativecommons.org/licenses/by/4.0/), which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

Sulfur Ion Implantations Into Condensed CO₂: Implications for Europa

D. V. Mifsud^{1,2} , Z. Kaňuchová³ , P. Herczku² , Z. Juhász² , S. T. S. Kovács² , G. Lakatos^{2,4} , K. K. Rahul² , R. Rácz² , B. Sulik² , S. Biri² , I. Rajta² , I. Vajda² , S. Ioppolo⁵ , R. W. McCullough⁶ , and N. J. Mason^{1,2} 

¹Centre for Astrophysics and Planetary Science, School of Physical Sciences, University of Kent, Canterbury, UK, ²Institute for Nuclear Research (Atomki), Debrecen, Hungary, ³Astronomical Institute, Slovak Academy of Sciences, Tatranská Lomnica, Slovakia, ⁴Institute of Chemistry, University of Debrecen, Debrecen, Hungary, ⁵School of Electronic Engineering and Computer Science, Queen Mary University of London, London, UK, ⁶Department of Physics and Astronomy, School of Mathematics and Physics, Queen's University Belfast, Belfast, UK

Abstract The ubiquity of sulfur ions within the Jovian magnetosphere has led to suggestions that the implantation of these ions into the surface of Europa may lead to the formation of SO₂. However, previous studies on the implantation of sulfur ions into H₂O ice (the dominant species on the European surface) have failed to detect SO₂ formation. Other studies concerned with similar implantations into CO₂ ice, which is also known to exist on Europa, have offered seemingly conflicting results. In this letter, we describe the results of a study on the implantation of 290 keV S⁺ ions into condensed CO₂ at 20 and 70 K. Our results demonstrate that SO₂ is observed after implantation at 20 K, but not at the Europa-relevant temperature of 70 K. We conclude that this process is likely not a reasonable mechanism for SO₂ formation on Europa, and that other mechanisms should be explored instead.

Plain Language Summary SO₂ ice is known to exist at the surface of one of Jupiter's moons; Europa. However, the method by which this ice forms is still uncertain. Due to the orbit of Europa being within the giant magnetosphere of Jupiter, it has been proposed that sulfur ions within the magnetosphere could implant into the cold surface ices on Europa and subsequently react to form SO₂. However, laboratory experiments looking into the implantation of such ions into H₂O ice (the dominant ice on Europa's surface) and CO₂ ice have either failed to yield SO₂ or have provided inconclusive results. We have therefore performed an experiment in which we have implanted high-energy sulfur ions into CO₂ ice at two temperatures. Our results indicate that such implantations are unlikely to be the mechanism by which the SO₂ on Europa is formed, and that other chemical processes should be considered instead.

1. Introduction

Sulfur is a ubiquitous element in the cosmos and is an important participant in many biochemical, atmospheric, and geochemical processes (Mifsud, Kaňuchová, et al., 2021). Within our own Solar System, sulfur astrochemistry is perhaps best associated with the Galilean moon system of Jupiter. The innermost of the Galilean moons, Io, is the most volcanically active celestial body in the Solar System; emitting approximately one tonne of sulfurous molecules per second (Thomas et al., 2004). Some of these molecules are subsequently stripped from the Ionian exosphere and dissociated and ionized within the Jovian magnetosphere, whereupon the resultant sulfur ions may interact with the surfaces of the other, icy Galilean moons (Cooper et al., 2001). The detection of SO₂ ice on the trailing hemisphere of Europa by the *International Ultraviolet Explorer* spacecraft (Lane et al., 1981) led many researchers to suggest that an influx of magnetospheric sulfur ions could contribute to its formation there as part of a wider radiolytic sulfur cycle (Carlson et al., 1999), in which sulfur is chemically transformed to various molecular forms as a result of its interaction with ionising radiation.

Indeed, the distribution of SO₂ on the European trailing hemisphere is such that it adopts a so-called bulls-eye pattern (Becker et al., 2022; Carlson et al., 2005; Hendrix et al., 2011). This is suggestive of an exogenic sulfur source in which magnetospheric sulfur ions implanting into the surface ices yield SO₂ as a primary product. However, laboratory experiments considering the implantation of such ions into H₂O ices, which represent the dominant icy species at the surface of Europa, have thus far failed to detect any SO₂ among the radiolytic products. Instead, H₂SO₄ and its hydrates are efficiently formed (Ding et al., 2013; Strazzulla et al., 2007, 2009).

Validation: S. Ioppolo, R. W. McCullough, N. J. Mason
Writing – original draft: D. V. Mifsud
Writing – review & editing: D. V. Mifsud, Z. Kaňuchová, P. Herczku, Z. Juhász, S. T. S. Kovács, G. Lakatos, K. K. Rahul, R. Rácz, B. Sulik, S. Biri, I. Rajta, I. Vajda, S. Ioppolo, R. W. McCullough, N. J. Mason

Computational simulations have also been unsuccessful in demonstrating a reasonable reaction pathway by which sulfur ion implantations yield SO₂ as a primary product (Anders & Urbassek, 2019a, 2019b).

Sulfur ion implantations into other oxygen-bearing ices known to be present on the surface of Europa have also been considered more recently. Lv et al. (2014) performed experiments that demonstrated that such implantations into pure CO and CO₂ ices at 15 K did indeed yield SO₂, and proposed that a geologically reasonable time-scale of 2 × 10⁴ years is sufficient to produce the abundance of SO₂ observed on Europa. However, this estimation relied on the assumption that sulfur ion implantation experiments carried out at 15 K are representative of analogous processes occurring on the surface of Europa, which is characterized by significantly higher temperatures (Ashkenazy, 2019). Follow-up studies by Boduch et al. (2016) did not detect SO₂ after the implantation of sulfur ions into pure CO₂ ice at 16 K, clearly contrasting with the previous results of Lv et al. (2014). It should be noted, however, that Boduch et al. (2016) made use of ultraviolet absorption spectroscopy as their product detection method, and it is therefore possible that the formation of any SO₂ was masked by the stronger absorptions of sulfur oxyanions which were detected after implantation. Nonetheless, it is apparent that the question of a possible exogenic sulfur source for SO₂ on the surface of Europa remains an open one.

We have therefore made an experimental attempt to address the possibility of SO₂ formation as a result of sulfur ion implantation into a carbon oxide ice. In this letter, we present the results of high-fluence (>10¹⁶ ions cm⁻²) implantations of 290 keV S⁺ ions into pure CO₂ ices at 20 and 70 K. By performing our experiment at two different temperatures (one that is similar to previous experimental work conducted in this field, and one that is relevant to some of the colder European surface ices at the mid- to high-latitudes), we have been able to determine the influence of temperature on the radiation chemistry taking place within our ices and thus extend the results described by previous studies. Our results and their implications are discussed in light of the radiolytic chemistry present on the European surface.

2. Materials and Methods

Our experiments were carried out using the Ice Chamber for Astrophysics–Astrochemistry (ICA) located at the Institute for Nuclear Research (Atomki) in Debrecen, Hungary. The technical details of this set-up have been extensively described in previous publications (Herczku et al., 2021; Mifsud, Juhász, et al., 2021), and so only a short overview is provided here. The ICA is an ultra-high vacuum compatible steel chamber containing a gold-coated oxygen-free copper sample holder at its center. This sample holder is able to host a maximum of four ZnSe substrates on which astrophysical ice analogs may be prepared. The substrates may be cooled to 20 K using a closed-cycle helium cryostat, although an operational temperature range of 20–300 K is available. The base pressure of the chamber is maintained in the region of a few 10⁻⁹ mbar by the combined use of a dry rough vacuum pump and a turbomolecular pump. The chamber is also connected to a 2 MV Tandatron accelerator (Biri et al., 2021; Rajta et al., 2018) which allows ion beams to be delivered to the prepared astrophysical ice analogs at a nominal incidence angle of 36° to the normal. A simplified schematic diagram of the ICA is given in Figure 1.

CO₂ ices were prepared on the ZnSe substrates at 20 and 70 K via background deposition of the gas (Linde, 99.995% purity), which was dosed into the main chamber through an all-metal needle valve at a chamber pressure of a few 10⁻⁶ mbar. Ice growth and processing were monitored in situ using Fourier-transform mid-infrared (FTIR) transmission absorption spectroscopy over a spectral range of 4,000–650 cm⁻¹ using a nominal resolution of 1 cm⁻¹. Acquired FTIR spectra of the CO₂ ices allowed for quantitative measurements of their molecular column densities *N* (molecules cm⁻²) to be made by using Equation 1:

$$N = \frac{\int \tau(\nu) d\nu}{A_\nu} = \frac{P \ln(10)}{A_\nu} \quad (1)$$

where $\tau(\nu)$ is the optical depth of the molecular ice at a given wavenumber (cm⁻¹), A_ν is the integrated band strength constant for the infrared absorption band over which Equation 1 is integrated (cm molecule⁻¹), and P is the measured area of this absorption band (cm⁻¹). The molecular column density of a deposited ice may be related to its thickness d (μm) through Equation 2:

$$d = 10^4 \frac{NZ}{N_{Ap}} \quad (2)$$




Figure 1. Simplified top-view schematic diagram of the Ice Chamber for Astrophysics-Astrochemistry set-up. Implantations of S^+ ions were performed using the arrangement shown here, with ions impacting the ice targets at angles of 36° to the normal. The Fourier-transform mid-infrared spectroscopic beam, detected by an external mercury-cadmium-telluride detector, remained orthogonal to the substrates and ices. Other attachments to the side of the chamber (e.g., Faraday cups, pressure gauges, etc.) are omitted for clarity.

where Z and ρ are respectively the molar mass (44 g mol^{-1}) and density ($\rho_{20 \text{ K}} = 0.98 \text{ g cm}^{-3}$; $\rho_{70 \text{ K}} = 1.48 \text{ g cm}^{-3}$) of the deposited CO_2 ice (Satorre et al., 2008), and N_A is the Avogadro number ($6.02 \times 10^{23} \text{ molecules mol}^{-1}$).

The most prominent infrared absorption feature of condensed CO_2 is the asymmetric stretching mode at around $2,343 \text{ cm}^{-1}$ (Isokoski et al., 2013; Mifsud et al., 2022a). However, this band saturates fairly quickly during ice deposition; indeed, saturation occurs before the ice reaches a suitable thickness for ion implantation to be performed. Therefore, we have made use of the less intense absorption band related to the asymmetric stretching mode of the $^{13}\text{CO}_2$ isotopologue at around $2,283 \text{ cm}^{-1}$ to quantify the column densities and thicknesses of the deposited ices. To do this, a FTIR spectrum of the depositing ice was acquired just prior to the saturation of the $^{12}\text{CO}_2$ asymmetric stretching mode and the column density ratio of the $^{12}\text{CO}_2$ and $^{13}\text{CO}_2$ isotopologues was measured using Equation 1 assuming A_ν for the asymmetric stretching modes of these isotopologues to be 7.6×10^{-17} and $7.8 \times 10^{-17} \text{ cm molecule}^{-1}$, respectively (Gerakines et al., 1995). Continued deposition then resulted in the saturation of the $^{12}\text{CO}_2$ asymmetric stretching mode, after which the measured $^{13}\text{CO}_2$ column densities were used in combination with the experimental isotopologue abundance ratio to determine the total molecular column densities of the deposited ices as well as their thicknesses (Equation 2). A similar approach had been previously adopted by Lv et al. (2014).

Pure CO_2 ices were deposited to thicknesses of about $3 \mu\text{m}$. This thickness is greater than the penetration depths of the projectile 290 keV S^+ ions supplied by the Tandatron accelerator as calculated using the *Stopping and Range of Ions in Matter* (SRIM) program ($0.8\text{--}1.1 \mu\text{m}$; Ziegler et al., 2010), thus ensuring implantation of the ions into the ices. Once an ice was deposited to this desired thickness, a pre-irradiation FTIR spectrum was acquired after which the ice was exposed to the ion beam with further spectra acquired at pre-defined fluence steps until a total fluence of $>10^{16} \text{ ions cm}^{-2}$ was implanted. During implantation, however, it was noted that sputtering resulted in a gradual thinning of the ice. To compensate for this sputtering, a simultaneous deposition-irradiation method was used. In this method, after initial deposition of the ice to a thickness of $3 \mu\text{m}$, the ices were irradiated by the S^+ ion beam with concurrent background deposition of more CO_2 occurring at a chamber pressure of about 10^{-5} mbar . Both irradiation and deposition were halted during FTIR spectral acquisition. We do not anticipate the use of this simultaneous deposition-irradiation method to impact the chemical evolution of our ices any differently to what would be expected under standard ion implantation conditions, and have provided evidence supporting this claim in the Appendix A section of this letter.

3. Results and Discussion

In this study, the implantation of 290 keV S^+ ions into CO_2 ices at 20 and 70 K was investigated with the aim of determining whether such a mechanism could account for the formation of SO_2 on the cold surface of Europa. CO_2 ice is a minor component of the European surface (Hansen & McCord, 2008), having an estimated abundance of 0.036% by number relative to H_2O (Hand et al., 2007). As such, it is likely to interact to at least some extent with incident magnetospheric sulfur ions. The fluxes of keV-MeV S^{n+} ions vary between 2×10^6 and $10^8 \text{ ions cm}^{-2} \text{ s}^{-1}$, depending on the location on the surface of Europa (Dalton et al., 2013). Therefore, a fluence of $10^{16} \text{ ions cm}^{-2}$ as was supplied in this study could be expected to be delivered to the European surface within 160 years.

The implantation of 290 keV S^+ ions into CO_2 ice at 20 K resulted in the appearance of several new absorption features in the FTIR spectrum (Figure 2). In particular, two new bands were observed at $1,336$ and $1,150 \text{ cm}^{-1}$ which we have respectively attributed to the asymmetric and symmetric stretching modes of SO_2 (Gomis & Strazzulla, 2008; Sandford et al., 1991). These absorption bands are, however, fairly small and are located in a region of the spectrum where several other, more intense absorption bands are also present. In light of this, we have performed two control experiments so as to be confident in our detection of SO_2 . First, we compared the wavenumber peak positions of these bands with those of SO_2 in an unirradiated $\text{CO}_2:\text{SO}_2$ (6:1) ice mixture (Figure 2), and found a very good agreement (within 1 cm^{-1}) with the suspected SO_2 band positions in the




Figure 2. Fourier-transform mid-infrared (FTIR) spectra of condensed CO_2 before (black trace) and after (red trace) the implantation of 290 keV S^+ ions at 20 K. Also shown are the FTIR spectra acquired during control experiments, including an unirradiated $\text{CO}_2:\text{SO}_2$ (6:1) ice mixture at 20 K (blue trace) and a CO_2 ice after the implantation of 300 keV He^+ ions at 20 K (green trace). The stopping powers of the S^+ and He^+ ions in the CO_2 ice are respectively 4.33 and 9.74 eV \AA^{-1} , while the radiation doses supplied after delivering 10^{16} ions cm^{-2} are 325 and 730 eV per CO_2 molecule. Note that the spectra of the irradiated ices are difference spectra yielded after the subtraction of the “as deposited” spectra.

irradiated ice. Second, we implanted a similar fluence of 300 keV He^+ ions into pure CO_2 ice at 20 K under similar experimental conditions. After such implantations, absorption bands due to the sulfur chemistry initiated by the implantation of S^+ ions would be absent while all other bands produced as a result of the energetic processing of the ice would still be present. This outcome was indeed observed, with the bands at 1,336 and 1,150 cm^{-1} not being observed at any point during the implantation of the He^+ ions.

The mechanistic chemistry leading to the formation of SO_2 is presumed to proceed after the neutralization of the implanted sulfur ion. As the ion traverses the ice it dissipates energy into its surroundings resulting in the dissociation of CO_2 to CO and an oxygen atom (Pilling et al., 2022). The neutralized sulfur ion may then react with one such oxygen atom to produce SO (Tevault & Smardzewski, 1978); alternatively, SO may result from the abstraction of an oxygen atom from a surviving CO_2 molecule by the neutralized sulfur ion (Froese & Goddard, 1993). SO is a very reactive species, and so will rapidly oxidize to yield SO_2 (Herron & Huie, 1980; Rolfes et al., 1965).

Interestingly, when the implantation of 290 keV S^+ ions was performed at 70 K, no SO_2 was detected in the FTIR spectrum (Figure 3). Indeed, this spectrum shows significantly fewer absorption bands than does its 20 K counterpart. One striking difference is the size of the absorption band due to the asymmetric stretching mode of O_3 located at about 1,041 cm^{-1} (Chaabouni et al., 2000). In the 20 K spectrum (Figure 2), this band is very intense and is indicative of the efficient formation of O_3 in the irradiated CO_2 ice (Mifsud et al., 2022a, 2022b). In the 70 K spectrum (Figure 3), however, this band is significantly smaller, suggesting an inefficient O_3 formation pathway. It is this observation that gives us an insight into why the formation of SO_2 as a result of S^+ ion implantation into condensed CO_2 is favorable at 20 K, but not at 70 K.

The energetic processing of CO_2 ice by ions, electrons, and ultraviolet photons is known to yield several oxygen-bearing products, including O_3 (Martín-Doménech et al., 2015; Mifsud et al., 2022a; Sivaraman et al., 2013). The formation of this radiolytic product is, however, dependent upon the prior synthesis of O_2 within the ice which then furnishes O_3 upon the barrierless addition of a supra-thermal oxygen atom. In our 20 K experiment, the O_2 is stable within the ice as the experimental temperature is lower than its sublimation




Figure 3. Fourier-transform mid-infrared (FTIR) spectra of condensed CO₂ before (black trace) and after (red trace) the implantation of 290 keV S⁺ ions at 70 K. Also shown is the FTIR spectrum of an unirradiated CO₂:SO₂ (6:1) ice mixture at 20 K (blue trace). Note that the spectrum of the irradiated ice is a difference spectrum yielded after the subtraction of the “as deposited” spectrum.

temperature, and so O₃ formation may occur efficiently. Indeed, O₃ is one of the major products of this implantation process (Figure 2). In the 70 K experiment, however, it is evident that O₂ sublimation from the ice is fairly efficient (Jones et al., 2014), thus effectively depleting the ice of its oxygen content. The result of this is that there are fewer oxygen atoms available within the bulk ice that may react with the implanted sulfur to yield SO₂, thus explaining the absence of this latter species at 70 K. Previous studies have suggested that otherwise volatile molecules such as O₂ may be stabilized at temperatures beyond their sublimation point via their encapsulation within clathrate-like structures (Hand et al., 2006). We speculate that, in our experiments, any such CO₂-based clathrate-like structures may not have been sufficiently stable to retain O₂ due to the fact that CO₂ is itself a volatile species. The presence of a radiolytically-derived tenuous O₂ exosphere on Europa (Milillo et al., 2016) is consistent with this interpretation.

Our results build upon and extend the previous findings of Lv et al. (2014) and Boduch et al. (2016). In their study, Lv et al. (2014) implanted multiply charged sulfur ions into condensed CO and CO₂ at 15 K, and recorded the formation of SO₂ in each ice. Based on their reported SO₂ formation efficiency of 0.38 molecules ion⁻¹ for the implantation of 90 keV S⁹⁺ ions into CO₂ ice, Lv et al. (2014) suggested that the observed abundances of SO₂ on Europa could be formed within a geologically reasonable time-scale of 2 × 10⁴ years. This result was not reproduced in the later study by Boduch et al. (2016), who did not observe any SO₂ in the ultraviolet absorption spectra of CO₂ ice into which 144 keV S⁹⁺ ions had been implanted at 16 K. This non-detection was ascribed to one of two reasons: either the accumulated column density of SO₂ formed as a result of ion implantation was below the spectroscopic detection limits of their instrument, or else the absorption bands attributable to SO₂ were masked by the intense absorptions of sulfur oxyanions such as SO₃⁻.

The results of this present study suggest that, although the formation of SO₂ as a result of sulfur ion implantation into CO₂ ice is indeed possible at low temperatures (15–20 K), this is not true for higher temperatures more representative of the surface of Europa, such as the 70 K temperature considered here. This is actually a somewhat unexpected result, as other ion implantation processes in which the implanted ion is incorporated into an oxygen-bearing product molecule have been shown to be unaffected by changes in the experimental temperature (Ding et al., 2013; Lv et al., 2012). We conclude, therefore, that sulfur ion implantation into CO₂ ices at the surface

of Europa is not an efficient mechanism by which the SO₂ observed on the surface may form. This is consistent with the lack of association in the spatial distributions of SO₂ and CO₂ ices on Europa (Hansen & McCord, 2008). Other formation mechanisms must therefore be considered instead. The correlation in the distribution of SO₂ and H₂SO₄ hydrates on the surface of Europa, as well as their bulls-eye patterns, hints at a related synthetic chemistry. Indeed, an early study by Hochanadel et al. (1955) demonstrated that the irradiation of concentrated H₂SO₄ solution yielded SO₂, and a similar result was more recently observed by Loeffler et al. (2011) for condensed H₂SO₄ and its hydrates at low temperatures. Therefore, it is reasonable to suggest that the implantation of sulfur ions into the European H₂O surface ices yields H₂SO₄ hydrates (Ding et al., 2013; Strazzulla et al., 2007, 2009), which are then dissociated to yield SO₂ upon further irradiation. It is also possible that alternative formation mechanisms, such as the radiolytic decomposition of sulfate minerals (McCord et al., 2001) and the implantation of cold (sub-keV) magnetospheric sulfur ions into the European surface ice (Becker et al., 2022), may further contribute to the presence of SO₂.

4. Conclusions

In this letter, we present the results of a study considering the implantation of 290 keV S⁺ ions into condensed CO₂ at 20 and 70 K as a possible mechanism by which the SO₂ ice on the surface of Europa may form. We have found that, although SO₂ is observed amongst the radiolytic products after implantation of the ions into CO₂ ice at 20 K, it is not observed after implantation at 70 K; a temperature more relevant to the European surface. We have attributed this to the fact that, at this higher temperature, any O₂ formed as a result of the combination of radiolytically-derived oxygen atoms will efficiently sublime from the bulk ice, thus depleting it of the oxygen necessary for SO₂ synthesis. Based on the results of previous studies that have demonstrated the efficient synthesis of H₂SO₄ and its hydrates after sulfur ion implantation into H₂O ice, as well as the observed spatial correlation between these hydrates and SO₂ on the surface of Europa, we suggest that a possible major source of SO₂ may be the radiolytic dissociation of H₂SO₄ hydrates, themselves formed as a result of the implantation of sulfur ions into the European surface.

Appendix A: Justification of the Experimental Methodology

The above experiment made use of a simultaneous deposition-irradiation system in which pure CO₂ ices were first deposited to a thickness of about 3 μm, after which the ices were irradiated with a 290 keV S⁺ ion beam with concurrent background deposition of further CO₂ ice at a chamber pressure of 10⁻⁵ mbar. It is possible that the interaction of the ion beam with gas-phase CO₂ molecules may result in the formation of fragment species that may then deposit onto the growing ice and potentially influence any solid-phase radiation chemistry occurring there. However, under the experimental conditions described above, we consider the incorporation of such fragments into the depositing ice to be a negligible process and we provide calculations to support this statement herein.

The expected number n_e of CO₂ molecules undergoing fragmentation as a result of their collision with S⁺ ions (molecules ion⁻¹) is given as:

$$n_e = \sigma L \rho_{\text{mol}} \quad (\text{A1})$$

where σ is the fragmentation cross-section of the molecule (cm² ion⁻¹), L is the pathlength of the ion in the chamber before it collides with the ice layer (cm), and ρ_{mol} is the molecular gas density (molecules cm⁻³). A value for this latter term may be calculated by first considering the Ideal Gas Law, which gives the ratio of the amount of gas present n (mol) to its volume V (cm³) to be:

$$\frac{n}{V} = \frac{P}{RT} = 4 \times 10^{-7} \text{ mol m}^{-3} \quad (\text{A2})$$

where P is the gas pressure (Pa; 10⁻⁵ mbar = 10⁻³ Pa), R is the molar gas constant (8.314 J K⁻¹ mol⁻¹) and T is the temperature of the gas which we have taken to be 298 K (i.e., room temperature). Converting this value

to the molecular gas density ρ_{mol} (molecules cm^{-3}) may be achieved by multiplying by the Avogadro number (6.02×10^{23} molecules mol^{-1}):

$$\rho_{\text{mol}} = N_A \frac{n}{V} = 2.4 \times 10^{17} \text{ molecules m}^{-3} = 2.4 \times 10^{11} \text{ molecules cm}^{-3} \quad (\text{A3})$$

Assuming that $\sigma = 10^{-15} \text{ cm}^2 \text{ ion}^{-1}$ (Mejía et al., 2015) and taking $L = 20 \text{ cm}$ (known from the geometry of the chamber) and substituting into Equation A1, a value of $4.8 \times 10^{-3} \text{ molecules ion}^{-1}$ may be calculated for n_e . Consider now the rate of ion delivery to the ICA chamber, r_{ion} (ions s^{-1}): this may be determined by taking the ratio of the nominal S^+ beam current ($I = 100 \text{ nA}$) to the Coulombic charge of the projectile S^+ ions ($q = 1.602 \times 10^{-19} \text{ C}$). This yields a value of $6.2 \times 10^{11} \text{ ions s}^{-1}$ for r_{ion} . The rate of molecular dissociation to yield fragments, r_{diss} (molecules s^{-1}) is therefore given as:

$$r_{\text{diss}} = n_e r_{\text{ion}} = 3 \times 10^9 \text{ molecules s}^{-1} \quad (\text{A4})$$

The pumping speed of the turbomolecular pump maintaining the vacuum in the ICA is 400 L s^{-1} (equivalent to $0.4 \text{ m}^3 \text{ s}^{-1}$). Such a value is valid for N_2 ; however, the pumping speed of CO_2 gas S is estimated to be 90% of this value and is thus $0.36 \text{ m}^3 \text{ s}^{-1}$. By once again making use of the Ideal Gas Law, the partial pressure of those molecules undergoing dissociation due to interaction with the S^+ beam, P_p , may be given as:

$$P_p = \frac{r_{\text{diss}} RT}{S N_A} = 3.4 \times 10^{-11} \text{ Pa} = 3.4 \times 10^{-13} \text{ mbar} \quad (\text{A5})$$

Following Dalton's Law of Partial Pressures, the ratio of the total number of gas-phase CO_2 molecules in the chamber at any one time to the number of dissociated CO_2 molecules is in excess of 3×10^7 . As such, it is highly unlikely that the fragments formed by molecular interaction with the S^+ ion beam could contribute in any significant way to the radiation chemistry and physics occurring within the irradiated bulk ice (for which the column density is in excess of $4 \times 10^{18} \text{ molecules cm}^{-2}$), even if all the fragments formed were to deposit and be incorporated into the ice (a scenario which in of itself is unlikely, since at least some of the fragments should be pumped out of the chamber by the turbomolecular and dry rough vacuum pumps before they can condense). Therefore, we conclude that the simultaneous deposition-irradiation method used in this study was appropriate.

Conflict of Interest

The authors declare no conflicts of interest relevant to this study.

Data Availability Statement

The raw spectroscopic data collected during experimentation is available on the Mendeley Data online repository: <https://doi.org/10.17632/zr2mdbk5dt.1> (Mifsud et al., 2022c). Data analysis was performed using the OriginPro 2018 software: <https://www.originlab.com/>.

References

- Anders, C., & Urbassek, H. M. (2019a). Energetic sulphur ion impacts into cometary ice surfaces: A molecular dynamics study. *Monthly Notices of the Royal Astronomical Society*, 482(2), 2374–2388. <https://doi.org/10.1093/mnras/sty2770>
- Anders, C., & Urbassek, H. M. (2019b). High-energy ion impacts into the sulphur-bearing ice surface of Europa: An atomistic study of chemical transformations. *Astronomy and Astrophysics*, 625, 140. <https://doi.org/10.1051/0004-6361/201935367>
- Ashkenazy, Y. (2019). The surface temperature of Europa. *Heliyon*, 5(6), e01908. <https://doi.org/10.1016/j.heliyon.2019.e01908>
- Becker, T. M., Trumbo, S. K., Molyneux, P. M., Retherford, K. D., Hendrix, A. R., Roth, L., et al. (2022). Mid-ultraviolet *hubble* observations of Europa and the global surface distribution of SO_2 . *Planetary Science Journal*, 3(6), 129. <https://doi.org/10.3847/PSJ/ac69eb>
- Biri, S., Vajda, I. K., Hajdu, P., Rácz, R., Csík, A., Kormány, Z., et al. (2021). The atomik accelerator centre. *European Physical Journal Plus*, 136(2), 247. <https://doi.org/10.1140/epjp/s13360-021-01219-z>
- Boduch, P., Brunetto, R., Ding, J. J., Domaracka, A., Kaňuchová, Z., Palumbo, M. E., et al. (2016). Ion processing of ices and the origin of SO_2 and O_3 on the icy surfaces of the icy Jovian satellites. *Icarus*, 277, 424–432. <https://doi.org/10.1016/j.icarus.2016.05.026>
- Carlson, R. W., Anderson, M. S., Mehlman, R., & Johnson, R. E. (2005). Distribution of hydrate on Europa: Further evidence for sulphuric acid hydrate. *Icarus*, 177(2), 461–471. <https://doi.org/10.1016/j.icarus.2005.03.026>
- Carlson, R. W., Johnson, R. E., & Anderson, M. S. (1999). Sulphuric acid on Europa and the radiolytic sulphur cycle. *Science*, 286(5437), 97–99. <https://doi.org/10.1126/science.286.5437.97>
- Chaabouni, H., Schriver-Mazzuoli, L., & Schriver, A. (2000). Infrared spectroscopy of neat solid ozone and that of ozone in interaction with amorphous and crystalline water ice. *Journal of Physical Chemistry A*, 104(30), 6962–6969. <https://doi.org/10.1021/jp0008290>

Acknowledgments

The authors acknowledge support from the Europlanet 2024 RI which has received funding from the European Union's Horizon 2020 Research Innovation Programme under grant agreement No. 871149. The main components of the ICA set-up were purchased using funds from the Royal Society obtained through Grants UF130409, RGF/EA/180306, and URF/R/191018. Further developments of the installation were supported in part by the Eötvös Loránd Research Network through Grants ELKH IF-2/2019 and ELKH IF-5/2020. This work has also received support from the European Union and the State of Hungary; co-financed by the European

Regional Development Fund through Grant GINOP-2.3.3-15-2016-00005. Support has also been received from the Research, Development, and Innovation Fund of Hungary through Grant K128621. DVM is the recipient of a University of Kent Vice-Chancellor's Research Scholarship. The research of ZK is supported by the Slovak Grant Agency for Science (Grant 2/0059/22) and the Slovak Research and Development Agency (contract No. APVV-19-0072). ZJ is grateful for the support of the Hungarian Academy of Sciences through the János Bolyai Research Scholarship. SI acknowledges the Royal Society for financial support.

- Cooper, J. F., Johnson, R. E., Mauk, B. H., Garrett, H. B., & Gehrels, N. (2001). Energetic ion and electron irradiation of the icy galilean satellites. *Icarus*, 149(1), 133–159. <https://doi.org/10.1006/icar.2000.6498>
- Dalton, J. B., Cassidy, T., Paranicas, C., Shirley, J. H., Prockter, L. M., & Kamp, L. W. (2013). Exogenic controls on sulphuric acid hydrate production at the surface of Europa. *Planetary and Space Science*, 77, 45–63. <https://doi.org/10.1016/j.pss.2012.05.013>
- Ding, J. J., Boduch, P., Domaracka, A., Guillous, S., Langlinay, T., Lv, X. Y., et al. (2013). Implantation of multiply charged sulphur ions in water ice. *Icarus*, 226(1), 860–864. <https://doi.org/10.1016/j.icarus.2013.07.002>
- Froese, R. D. J., & Goddard, J. D. (1993). *Ab initio* studies of the lowest singlet and triplet potential energy surfaces of CO₂S. *Molecular Physics*, 79(4), 685–697. <https://doi.org/10.1080/00268979300101541>
- Gerakines, P. A., Schutte, W. A., Greenberg, J. M., & van Dishoeck, E. F. (1995). The infrared band strengths of H₂O, CO, and CO₂ in laboratory simulations of astrophysical ice mixtures. *Astronomy and Astrophysics*, 296, 810.
- Gomis, O., & Strazzulla, G. (2008). Ion irradiation of H₂O ice on top of sulphurous solid residues and its relevance to the galilean satellites. *Icarus*, 194(1), 146–152. <https://doi.org/10.1016/j.icarus.2007.09.015>
- Hand, K. P., Carlson, R. W., & Chyba, C. F. (2007). Energy, chemical disequilibrium, and geological constraints on Europa. *Astrobiology*, 7(6), 1006–1022. <https://doi.org/10.1089/ast.2007.0156>
- Hand, K. P., Chyba, C. F., Carlson, R. W., & Cooper, J. F. (2006). Clathrate hydrates of oxidants in the ice shell of Europa. *Astrobiology*, 6(3), 463–482. <https://doi.org/10.1089/ast.2006.6.463>
- Hansen, G. B., & McCord, T. B. (2008). Widespread CO₂ and other non-ice compounds on the anti-Jovian and trailing sides of Europa from galileo/NIMS observations. *Geophysical Research Letters*, 35(1), L01202. <https://doi.org/10.1029/2007GL031748>
- Hendrix, A. R., Cassidy, T. A., Johnson, R. E., Paranicas, C., & Carlson, R. W. (2011). Europa's disc-resolved ultraviolet spectra: Relationships with plasma flux and surface terrains. *Icarus*, 212(2), 736–743. <https://doi.org/10.1016/j.icarus.2011.01.023>
- Herczku, P., Mifsud, D. V., Ioppolo, S., Juhász, Z., Kaňuchová, Z., Kovács, S. T. S., et al. (2021). The ice chamber for astrophysics-astrochemistry (ICA): A new experimental facility for ion impact studies of astrophysical ice analogues. *Review of Scientific Instruments*, 92(8), 084501. <https://doi.org/10.1063/5.0050930>
- Herron, J. T., & Huie, R. E. (1980). Rate constants at 298 K for the reactions SO + SO + M → (SO)₂ + M and SO + (SO)₂ → SO₂ + S₂O. *Chemical Physics Letters*, 76(2), 322–324. [https://doi.org/10.1016/0009-2614\(80\)87032-1](https://doi.org/10.1016/0009-2614(80)87032-1)
- Hochanadel, C. J., Ghormley, J. A., & Sworski, T. J. (1955). The decomposition of sulphuric acid by cobalt γ -rays. *Journal of the American Chemical Society*, 77(12), 3215. <https://doi.org/10.1021/ja01617a017>
- Isokoski, K., Poteet, C. A., & Linnartz, H. (2013). Highly resolved infrared spectra of pure CO₂ ice (15–75 K). *Astronomy and Astrophysics*, 555, 85. <https://doi.org/10.1051/0004-6361/201321517>
- Jones, B. M., Kaiser, R. I., & Strazzulla, G. (2014). UV-vis, infrared, and mass spectroscopy of electron irradiated frozen oxygen and carbon dioxide mixtures with water. *The Astrophysical Journal*, 781(2), 85. <https://doi.org/10.1088/0004-637X/781/2/85>
- Lane, A. L., Nelson, R. M., & Matson, D. L. (1981). Evidence for sulphur implantation in Europa's UV absorption band. *Nature*, 292(5818), 38–39. <https://doi.org/10.1038/292038a0>
- Loeffler, M. J., Hudson, R. L., Moore, M. H., & Carlson, R. W. (2011). Radiolysis of sulphuric acid, sulphuric acid monohydrate, and sulphuric acid tetrahydrate and its relevance to Europa. *Icarus*, 215(1), 370–380. <https://doi.org/10.1016/j.icarus.2011.06.008>
- Lv, X. Y., Boduch, P., Ding, J. J., Domaracka, A., Langlinay, T., Palumbo, M. E., et al. (2014). Sulphur implantation in CO and CO₂ ices. *Monthly Notices of the Royal Astronomical Society*, 438(2), 922–929. <https://doi.org/10.1093/mnras/stt2004>
- Lv, X. Y., de Barros, A. L. F., Boduch, P., Bordalo, V., da Silveira, E. F., Domaracka, A., et al. (2012). Implantation of multiply charged carbon ions in water ice. *Astronomy and Astrophysics*, 546, 81. <https://doi.org/10.1051/0004-6361/201219886>
- Martín-Doménech, R., Manzano-Santamaría, J., Muñoz-Caro, G. M., Cruz-Díaz, G. A., Chen, Y.-J., Herrero, V. J., & Tanarro, I. (2015). UV photoprocessing of CO₂ ice: A complete quantification of photochemistry and photon-induced desorption processes. *Astronomy and Astrophysics*, 584, 14. <https://doi.org/10.1051/0004-6361/201526003>
- McCord, T. B., Orlando, T. M., Teeter, G., Hansen, G. B., Sieger, M. T., Petrik, N. G., & van Keulen, L. (2001). Thermal and radiation stability of the hydrated salt minerals epsomite, mirabilite, and natron under Europa environmental conditions. *Journal of Geophysical Research*, 106(E2), 3311–3319. <https://doi.org/10.1029/2000JE001282>
- Mejía, C., Bender, M., Severin, D., Trautmann, C., Boduch, P., Bordalo, V., et al. (2015). Radiolysis and sputtering of carbon dioxide ice induced by swift Ti, Ni, and Xe ions. *Nuclear Instruments and Methods in Physics Research B: Beam Interactions with Materials and Atoms*, 365, 477–481. <https://doi.org/10.1016/j.nimb.2015.09.039>
- Mifsud, D. V., Juhász, Z., Herczku, P., Kovács, S. T. S., Ioppolo, S., Kaňuchová, Z., et al. (2021). Electron-irradiation and thermal chemistry studies of interstellar and planetary ice analogues at the ICA astrochemistry facility. *European Physical Journal D: Atomic, Molecular, Optical, and Plasma Physics*, 75(6), 182. <https://doi.org/10.1140/epjd/s10053-021-00192-7>
- Mifsud, D. V., Kaňuchová, Z., Herczku, P., Ioppolo, S., Juhász, Z., Kovács, S. T. S., et al. (2021). Sulphur ice astrochemistry: A review of laboratory studies. *Space Science Reviews*, 217(1), 14. <https://doi.org/10.1007/s11214-021-00792-0>
- Mifsud, D. V., Kaňuchová, Z., Herczku, P., Juhász, Z., Kovács, S. T. S., Lakatos, G., et al. (2022c). Raw data for “sulphur ion implantations into condensed CO₂: Implications for Europa”. *Mendeley Data*, V1. <https://doi.org/10.17632/zr2mdbk5dt.1>
- Mifsud, D. V., Kaňuchová, Z., Ioppolo, S., Herczku, P., Traspas Muiña, A., Field, T. A., et al. (2022a). Mid-IR and VUV spectroscopy characterisation of thermally processed and electron irradiated CO₂ astrophysical ice analogues. *Journal of Molecular Spectroscopy*, 385, 111599. <https://doi.org/10.1016/j.jms.2022.111599>
- Mifsud, D. V., Kaňuchová, Z., Ioppolo, S., Herczku, P., Traspas Muiña, A., Sulik, B., et al. (2022b). Ozone production in CO₂:O₂ ices. *Physical Chemistry Chemical Physics*, 24(30), 18169–18178. <https://doi.org/10.1039/D2CP01535H>
- Milillo, A., Plainaki, C., De Angelis, E., Mangano, V., Massetti, S., Mura, A., et al. (2016). Analytical model of Europa's O₂ exosphere. *Planetary and Space Science*, 130, 3–13. <https://doi.org/10.1016/j.pss.2015.10.011>
- Pilling, S., Carvalho, G. A., & Rocha, W. R. M. (2022). Chemical evolution of CO₂ ices under processing by ionising radiation: Characterisation of nonobserved species and chemical equilibrium phase with the employment of PROCODA code. *The Astrophysical Journal*, 925(2), 147. <https://doi.org/10.3847/1538-4357/ac3d8a>
- Rajta, I., Vajda, I., Gyürky, G., Csedreki, L., Kiss, Á. Z., Biri, S., et al. (2018). Accelerator characterisation of the new ion beam facility at MTA atomki in debrecen, Hungary. *Nuclear Instruments and Methods in Physics Research: Accelerators, Spectrometers, Detectors, and Associated Equipment*, 880, 125–130. <https://doi.org/10.1016/j.nima.2017.10.073>
- Rolfes, T. R., Reeves, R. R., & Harteck, P. (1965). The chemiluminescent reaction of oxygen atoms with sulphur monoxide at low pressures. *Journal of Physical Chemistry*, 69(3), 849–853. <https://doi.org/10.1021/j100887a024>

- Sandford, S. A., Salama, F., Allamandola, L. J., Trafton, L. M., Lester, D. F., & Ramseyer, T. F. (1991). Laboratory studies of the newly discovered infrared band at 4705.2 cm^{-1} ($2.1253\text{ }\mu\text{m}$) in the spectrum of Io: The tentative identification of CO_2 . *Icarus*, *91*(1), 125–144. [https://doi.org/10.1016/0019-1035\(91\)90132-D](https://doi.org/10.1016/0019-1035(91)90132-D)
- Satorre, M. Á., Domingo, M., Millán, C., Luna, R., Vilaplana, R., & Santonja, C. (2008). Density of CH_4 , N_2 , and CO_2 ices at different temperatures of deposition. *Planetary and Space Science*, *56*(13), 1748–1752. <https://doi.org/10.1016/j.pss.2008.07.015>
- Sivaraman, B., Rajasekhar, B. N., Fulvio, D., Hunniford, A., McCullough, R. W., Palumbo, M. E., & Mason, N. J. (2013). Ozone and carbon trioxide synthesis by low energy ion implantation onto solid carbon dioxide and implications to astrochemistry. *Journal of Chemical Physics*, *139*(7), 074706. <https://doi.org/10.1063/1.4818166>
- Strazzulla, G., Baratta, G. A., Leto, G., & Gomis, O. (2007). Hydrate sulphuric acid after sulphur implantation in water ice. *Icarus*, *192*(2), 623–628. <https://doi.org/10.1016/j.icarus.2007.08.004>
- Strazzulla, G., Garozzo, M., & Gomis, O. (2009). The origin of sulphur-bearing species on the surfaces of icy satellites. *Advances in Space Research*, *43*(9), 1442–1445. <https://doi.org/10.1016/j.asr.2009.01.007>
- Tevault, D. E., & Smardzewski, R. R. (1978). Chemiluminescent reactions of sulphur atoms and oxygen atoms in solid argon matrices. SO chemiluminescence. *Journal of Chemical Physics*, *69*(7), 3182–3189. <https://doi.org/10.1063/1.436967>
- Thomas, N., Bagenal, F., Hill, T. W., & Wilson, J. K. (2004). The Io neutral clouds and plasma torus. In F. Bagenal, T. Dowling, & W. McKinnon. (Eds.), *Jupiter: The planet, satellites and magnetosphere*, (pp. 561–592). Cambridge University Press.
- Ziegler, J. F., Ziegler, M., & Biersack, J. (2010). SRIM – The stopping and range of ions in matter. *Nuclear Instruments and Methods in Physics Research B*, *268*(11–12), 1818–1823. <https://doi.org/10.1016/j.nimb.2010.02.091>