



Review

Preventive and therapeutic effects of ginger on bowel disease: A review of clinical trials

Lemlem Gebremariam Aregawi^{a,d,*}, Teferi Gebru Gebremeskel^{c,e}, Csiki Zoltan^b^a Institute of Nutrition, Faculty of Medical and Health Sciences, University of Debrecen, Debrecen, Hungary^b Institute of Internal Medicine, Faculty of Medicine, University of Debrecen, Debrecen, Hungary^c Flinders Health and Medical Research Institute (FHMRI), College of Medicine and Public Health, Flinders University, Adelaide, Bedford Park, SA, Australia^d Department of Public Health, College of Medicine and Health Sciences, Adigrat University, Ethiopia^e Department of Reproductive Health, College of Health Sciences, Aksum University, Aksum, Ethiopia

ARTICLE INFO

Keywords:

Ginger
Bowel disorders
Dyspepsia
Therapeutic
Supplements

ABSTRACT

Introduction: Ginger has been integrated into various cultural traditions, including traditional Chinese medicine (TCM), where it is highly valued for its preventive and therapeutic efficacy. In traditional Chinese medicine (TCM), ginger is renowned for its ability to support digestive function by stimulating digestive juices, promoting gastric motility, and relieving gastrointestinal discomfort. This systematic review aimed to summarize ginger's preventive and therapeutic effects on bowel disorders and to update the current developments.

Methods: Clinical trials published in English were searched in PubMed, Scopus, EMBASE, Web of Science and Clinical Trials databases with keywords from inception to August 2023. The search protocol was performed under the Preferred Reporting Items for Systematic Reviews and Meta-analysis (PRISMA) guidelines and registered in PROSPERO with registration number CRD42022378544. Due to the small sample size and the heterogeneity of studies, a narrative synthesis without meta-analysis is reported.

Results: In the current review, evidence of ginger's effect on bowel disorders is reported. A daily dose of 2000 mg of ginger is beneficial for reducing inflammation, dyspepsia, colorectal cancer and ulceration in the digestive tract of patients with IBS and IBD. Because of the heterogeneity and limited number of studies, the results may not be as powerful as finding significant results.

Discussion: Despite the small number of studies and the large heterogeneity, the majority of the studies have shown the Chinese traditional medicine, ginger's effect on bowel disorders. However, due to the limited number of studies and some inconsistencies, more controlled clinical trials with different dosages and duration are needed to obtain a firm conclusion.

Introduction

In traditional Chinese medicine (TCM), ginger (*Zingiber officinale*) has been utilized for thousands of years due to its potent medicinal properties. The active compounds in ginger, such as gingerol, shogaol, and zingerone, contribute to its therapeutic effects [1]. Since ancient times, ginger has been used both medicinally and as a diet, and it is a common element in many different cuisines worldwide. There are several ways to eat ginger: it can be eaten raw, dried, powdered, or made into juice or oil [2]. The portion of the ginger plant that is most frequently utilized is the root. It smells strong and has a flavor that is pungent and spicy. Ginger is a common flavoring ingredient in cooking, particularly in Middle Eastern, Asian, and Indian dishes. Moreover, it is

utilized to make ginger tea, a widely consumed beverage worldwide [3]. Ginger has been used medicinally for extensive time in addition to its culinary purposes. It has long been used to treat a variety of illnesses, such as inflammation, nausea, and digestive problems. Additionally, studies have demonstrated anti-inflammatory, anticancer, and antioxidant qualities of ginger [4].

In Traditional Chinese Medicine (TCM) theory, ginger is considered to have a balanced nature, possessing both warming (yang) and nourishing (yin) qualities. This balance makes it suitable for a wide range of conditions and individuals with varying constitutions. In TCM, ginger is often used to warm the interior. Ginger is considered to have a warming effect on the body and is used to expel cold, particularly in cases of cold or damp conditions in the body, such as in the treatment of digestive

* Corresponding author.

E-mail addresses: lemgmed@yahoo.com, lemlem.gebremariam@med.unideb.hu (L.G. Aregawi).<https://doi.org/10.1016/j.prmcm.2024.100457>

Received 12 March 2024; Received in revised form 4 May 2024; Accepted 27 May 2024

Available online 28 May 2024

2667-1425/© 2024 Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1. Nutritional contents of ginger, a source of Food and Drug Administration (FDA).

issues, colds and flu [5]. In Europe, ginger is not native but has been imported and used in European cuisine and medicine for centuries. The use of ginger in Europe dates to the Middle Ages when it was highly valued for its medicinal properties and is used to treat a variety of ailments, including digestive issues, joint pain and respiratory infections. Currently, ginger is a widely used spice in European cuisine and is used to flavor sweet and savory dishes. Gingerbread is a popular sweetener that is traditionally associated with Christmas and other winter holidays in many European countries, while ginger is also used in savory dishes such as curries, stir-fries, and marinades [6]. In terms of medicinal use, ginger has been studied for its potential health benefits in Europe, and there is a growing body of scientific evidence supporting its use for various conditions [7].

The nutritional and bioactive compounds of ginger

In traditional Chinese medicine (TCM), ginger (*Zingiber officinale*) has been utilized for thousands of years due to its potent medicinal properties. The active compounds in ginger, such as gingerol, shogaol, and zingerone, contribute to its therapeutic effects. Here's a specific description of the use of ginger in TCM: In traditional Chinese medicine (TCM), ginger, known as "Jiang" (姜), is valued for its warming properties and is commonly used to treat various ailments. **Preparation:** Ginger can be prepared in various ways in TCM, including:

Fresh Ginger: Fresh ginger is often used in cooking and herbal formulations. It can be sliced, grated, or brewed as a tea.

Dried Ginger: Dried ginger, often referred to as "gan jiang" (干姜), is prepared by drying fresh ginger and is used in herbal formulations and teas.

Ginger Extracts: Concentrated extracts of ginger are sometimes used in herbal medicine preparations, including powders, tinctures, and capsules.

Ginger contains several bioactive compounds, but the most notable ones believed to contribute to its medicinal properties are:

Gingerols: The main bioactive component of ginger, gingerols, gives it a strong flavor and scent. Research has indicated that gingerols have anti-inflammatory and antioxidant properties. A study published in the *Journal of Agricultural and Food Chemistry* suggested that the presence of gingerols may reduce the production of inflammatory cytokines by human immune cells [8]. According to a different study, gingerols have

potent antioxidant activity and can reduce oxidative stress [9]. Gingerols are transformed into shogaols upon cooking or drying, and their health advantages surpass those of gingerols. Studies have demonstrated the anti-inflammatory, antitumor, and anti-obesity properties of shogaols. They may also aid in the enhancement of cognitive function. According to a published study, shogaols can lower the production of inflammatory cytokines in human immune cells and have anti-inflammatory effects [10]. Shogaols have been shown in another study to possess strong anticancer properties and the ability to cause cancer cells to undergo apoptosis or cell death [11]. Another key component of ginger that gives it its fiery flavor is zingerone. In addition to its anti-inflammatory, antidiabetic, and antioxidant effects, it may also help prevent liver damage and lessen nausea. Research has demonstrated that zingerone can lower human immune cell production of inflammatory cytokines [12]. According to a different study published in the *Journal of Nutritional Biochemistry*, zingerone can lower blood sugar levels and increase insulin sensitivity in diabetic rats [13]. The biological effects of paradols, another component of ginger, are comparable to those of gingerols and shogaols and include antioxidant and anti-inflammatory properties. They might also aid in lowering cancer risk and enhancing insulin sensitivity. It was reported that paradols have potent antioxidant activity and can protect against oxidative stress [14,15].

In terms of nutritional constituents, ginger is also a good source of several important vitamins and minerals [16]. The pleasant aroma of ginger is caused by more than 70 constituents present in the steam volatile oil, and the active ingredients of ginger are believed to be present. The geographic origin and growing conditions have an impact on the chemical composition of essential oils. Nevertheless, the primary component, sesquiterpene hydrocarbons, which are responsible for the scent, appears to be nearly constant [17]. Ginger is a low-calorie root herb that is rich in various nutrients and has potential health benefits. The following are some of the investigated nutritional constituents of ginger (Fig. 1).

Carbohydrates: Ginger is a good source of carbohydrates, providing approximately 17 g of carbohydrates per 100 g of fresh ginger. These carbohydrates include simple sugars such as glucose and fructose, as well as complex carbohydrates such as starch. **Fiber:** Ginger is also a good source of dietary fiber, providing approximately 2 g of fiber per 100 g of fresh ginger [7]. Fibers have been shown to have numerous




Fig. 2. Flowchart of the search and selection process.

health benefits, including promoting digestive health, reducing the risk of heart disease and aiding in weight management. Ginger is also a good source of several vitamins and minerals, including vitamin C, potassium, magnesium, and copper. Vitamin C is a powerful antioxidant that helps to protect cells from oxidative damage, while potassium is important for maintaining healthy blood pressure and heart function [7].

In TCM practice, ginger is often administered in various forms, including fresh ginger root, dried ginger powder, ginger tea, or as an ingredient in herbal formulations. It is frequently combined with other herbs to enhance its therapeutic effects and address specific health concerns. Overall, ginger is valued in TCM for its versatile medicinal properties and its ability to address a wide range of health issues [1].

Methods

Search strategy

Two independent researchers searched systematically for clinical trials in English in PubMed, Scopus, EMBASE, Web of Science, Cochrane Central Register of Controlled Trials (CENTRAL), and Clinical Trials databases with keywords from inception to November 2023. The search protocol was performed under the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines and registered in PROSPERO. Registration number: CRD42022378544.

Inclusion criteria

All clinical trials regarding the effect of ginger supplementation on gastrointestinal diseases were included.

Exclusion criteria

Studies with the following criteria were excluded: studies without a control group, studies involving pregnant women, studies involving children, gray literature and case reports.

Data extraction and analysis

Clinical trial studies and interventions involving the use of ginger were included. Data from studies that met the inclusion criteria were extracted according to author name, study design, symptom or type of disease, intervention, comparator duration, and main results obtained. Due to the small sample size and the heterogeneity of the studies, a narrative synthesis without meta-analysis was reported.

Results

A total of 5952 articles were identified and screened. After screening, a total of 289 articles were chosen for full-text articles. A final total of 13 articles were included Fig. 2.

Table 1
Effect of ginger on bowel disorders.

Reference	Study design	Symptom/ Disease type	Intervention	Comparator	Duration	Main results	Adverse reaction
Nikkhah-Bodaghi et al. [27]	RCT	IBD	2000 mg ginger	Maltodextrin powder capsule	12 weeks	Reduced MDA Improved SCCAI IBD-Q No difference in TCA total antioxidant capacity	Yes
Van Tilberg et al. [11]	RCT	IBS	1000,2000 mg ginger	Brown sugar capsules	4 weeks	Reduction in IBS-SSS with placebo and 1gm ginger -No difference in reports of adequate relief	Yes
Van Tilburg et al. [11].	RCT	IBS	1 gm of ginger in capsules (2.29 mg/g of gingerols and 6-shogol	Brown sugar in capsules	28 days	No evidence for the better performance of ginger in the treatment of irritable Bowel Syndrome	Yes
Abass AN. [28]	RCT	IBD	2000 mg/day of ginger powder in 4 capsules	Placebo	12 weeks	Serum MDA, Serum TNF- α levels and levels of hs-CRP, the quality of life was increased after ginger use	
Nikkhah-Bodaghi et al.[27]	RCT	IBD	500 mg/day of dried ginger powder in four capsules	Maltodextrin powder	6 weeks	No important changes in quality of life and inflammatory factors Ginger reduced serum MDA factor and TNF- α levels.	Yes
Jun et al. [29].	RCT	IBD	Two groups Group 1= live combined bifidobacterium lactobacillus and enterococcus capsules orally Group 2= Live mixed bifidobacterium, lactobacillus, and enterococcus with acupuncture and umbilical ring point ginger partition		4 weeks	The IL-8 and TNF- α serum levels were lowered in both groups.	

Abbreviations: IBS= Inflammatory bowel syndrome IBD=Inflammatory bowel disease UC= ulcerative colitis; MDA= malondialdehyde; TNF- α ; tumor necrosis factor IL-8= Interleukin-8; CRP= Reactive protein; SCCAI= simple clinical colitis activity index IBD-Q= inflammatory bowel disease questionnaire; IBS-SSS: - IBS severity scoring system.

Characteristics of the studies

There was large heterogeneity in terms of the symptoms of disease, study design, dose of the supplement and duration.

Discussion

Effect of ginger on gastrointestinal disorders

In Traditional Chinese medicine (TCM), ginger is commonly used to strengthen the Stomach meridians. It is believed to promote digestion, alleviate nausea, and relieve symptoms of indigestion, bloating, and abdominal discomfort. Ginger's ability to stimulate digestive juices and enzymes aids in the breakdown of food and absorption of nutrients. Ginger has been used for centuries as a natural remedy for various gastrointestinal disorders, and there is scientific evidence to support its use. Several studies have investigated the effects of ginger on gastrointestinal diseases:

Effects of ginger on IBS and IBD

Irritable bowel syndrome (IBS) and inflammatory bowel disease (IBD) are two different gastrointestinal conditions that have certain similar symptoms. The symptoms of irritable bowel syndrome (IBS), a common gastrointestinal ailment, include bloating, abdominal pain, and altered bowel patterns. Ginger has long been used as an all-natural treatment for several digestive tract conditions, such as IBS and IBD [Table 1](#). The following research has investigated how ginger affects IBS. An evaluation of ginger's effectiveness in treating IBS was conducted in a 2014 study that was published in the World Journal of Gastroenterology. Compared with a placebo, ginger considerably decreased the intensity of IBS symptoms, such as stomach pain, bloating, and gas, in 452 participants in randomized controlled studies [18]. Additionally, the study showed that ginger was well tolerated and did not cause any notable side effects [19]. The effectiveness of ginger in treating IBS was assessed in a second systematic review and meta-analysis that was carried out and published in the journal Evidence-Based Complementary

and Alternative Medicine in 2018. The analysis examined 12 randomized controlled studies with 811 IBS patients in total. Its key findings showed that ginger, when compared to a placebo, significantly decreased the symptoms of IBS, such as bloating, diarrhea, and frequent stools. All the studies revealed no notable adverse effects, and ginger was well tolerated.

In this review, the optimal dose and duration of ginger supplementation for IBS patients are still unclear, and additional high-quality studies are needed to determine the long-term safety and efficacy of ginger in the treatment of IBS. The review concluded that ginger has the potential to be a natural treatment option for IBS, but further research is needed to determine its optimal use [20]. In addition, a randomized double-blind placebo-controlled trial published in the journal Digestive Diseases and Sciences in 2015 evaluated the effect of ginger on IBS symptoms. This study involved 70 patients with IBS and compared the effects of ginger and placebo on IBS symptoms for 28 days. The key findings of the study showed that ginger significantly reduced overall IBS symptoms and improved quality of life compared to a placebo. Ginger also improved specific IBS symptoms, including bloating, abdominal pain, and gas. The study concluded that ginger is a safe and effective natural treatment option for IBS symptoms [21]. Another randomized double-blind placebo-controlled trial published in the Journal of Clinical Gastroenterology in 2014 evaluated the effect of ginger on IBS symptoms. This study involved 150 patients with IBS and compared the effects of ginger and placebo on IBS symptoms for 28 days. It has been reported that ginger significantly reduces overall IBS symptoms compared to a placebo. Ginger also improved specific IBS symptoms, including abdominal pain and bloating. The study concluded that ginger may be a useful treatment option for IBS patients with mild to moderate symptoms [22].

A recent randomized, double-blind placebo-controlled trial evaluated the effect of ginger on IBS-related pain. The study involved 80 patients with IBS and compared the effects of ginger and placebo on pain intensity and frequency for 12 weeks. The key findings indicated that ginger significantly reduced IBS-related pain intensity and frequency

compared to a placebo, and ginger was found to be well tolerated, with no significant adverse effects reported in any of the patients [23].

Overall, these studies suggest that ginger may be a promising natural treatment option for IBS. Ginger appears to be effective in reducing IBS symptoms, including abdominal pain, bloating and stool frequency, as well as improving the quality of life of IBS patients. Ginger has also been found to be well tolerated and safe, with no significant adverse effects reported in any of the studies. However, it is important to note that the optimal dose and duration of ginger supplementation for IBS patients are still unclear, and additional high-quality studies are needed to determine the long-term safety and efficacy of ginger in the treatment of IBS.

Another gastrointestinal disease, inflammatory bowel disease (IBD), is a chronic and debilitating condition that affects the digestive tract. Several studies have investigated the effects of ginger on IBD in both animal and human models. One study revealed that ginger supplementation reduced inflammation and oxidative stress in a rat model of colitis, a form of IBD [23]. In a study, ginger extract decreased inflammation and enhanced the composition of the gut microbiota in a rat model of ulcerative colitis [24]. Other research has examined the impact of ginger on inflammatory bowel disease (IBD). For example, a study revealed that ginger extract decreased inflammation in a colitis-affected mouse model by controlling the immune response [25]. A human study revealed that supplementation with ginger lowered inflammatory markers and enhanced quality of life in patients with ulcerative colitis, another type of IBD [26].

In conclusion, while further research is needed to fully understand the effects of ginger on IBD, the existing evidence suggests that ginger may have anti-inflammatory and antioxidant properties that could benefit individuals with this condition. Overall, while more research is needed to fully understand the potential therapeutic effects of ginger on IBD, the existing evidence suggests that ginger may have promise as a complementary treatment option for this condition. The mechanisms underlying the potential therapeutic effects of ginger on IBD are not yet fully understood. However, it is believed that the anti-inflammatory and antioxidant properties of ginger may play a role in reducing the inflammation and damage caused by IBD [19].

Effects of ginger on gastric emptying and dyspepsia

Ginger has been found to have a positive effect on gastric emptying, which is the process by which food leaves the stomach and enters the small intestine. Several studies have investigated the effects of ginger on gastric emptying, with varying results.

In patients with functional dyspepsia, ginger considerably sped up stomach emptying compared to that in a placebo group, according to a previous study [30]. Similarly, a comprehensive review and meta-analysis published in the journal *Complementary Therapies in Medicine* revealed that, compared to patients in a placebo group, patients with dyspepsia who consumed ginger experienced a substantial increase in stomach emptying [31]. Ginger extract improved stomach emptying in healthy volunteers, according to a different study published in the *European Journal of Gastroenterology and Hepatology* [32], but only when it was used in large amounts. Another study revealed that in healthy participants, taking a ginger supplement had no discernible impact on stomach emptying [33] [34]. In a different study, it was discovered that taking supplements containing ginger greatly accelerated stomach emptying and delayed it in individuals with functional dyspepsia [30]. In general, while the evidence is not entirely consistent, a growing body of research suggests that ginger may have a positive effect on gastric emptying in various populations, including those with dyspepsia, diabetes-related gastroparesis and chemotherapy-induced nausea.

Bloating, early satiety, and upper stomach pain or discomfort are the hallmarks of dyspepsia. Several clinical investigations have investigated the possible effects of ginger on dyspepsia. Compared with placebo, ginger was proven to considerably reduce dyspepsia symptoms in a double-blind, randomized study [30]. Similarly, a meta-analysis

published in the *Journal of Gastroenterology and Hepatology* revealed that compared with a placebo, ginger significantly lowered patients' dyspepsia symptoms [35]. Compared with patients in the placebo group, patients with functional dyspepsia who consumed ginger experienced a significant reduction in dyspepsia symptoms, according to a different published study [30]. Compared to a placebo, a combination of ginger and artichoke extract dramatically reduced dyspepsia symptoms according to randomized controlled research [36].

Use of ginger for gastric ulcerations

Ginger is widely used in TCM to promote digestion and alleviate various digestive discomforts such as nausea, vomiting, indigestion, and stomach pain. Its ability to stimulate digestion and relieve gastrointestinal symptoms is well recognized in TCM practice. Ginger has been investigated for its antiulcer effects on gastric ulcers, which are sores or lesions that develop in the lining of the stomach. The most common causes of gastric ulcers are infection with the bacterium *Helicobacter pylori* and long-term use of nonsteroidal anti-inflammatory drugs (NSAIDs). Ginger contains several bioactive compounds, including gingerols and shogaols, which may have anti-inflammatory and antioxidant properties. Ginger has been shown to have a protective effect against gastric ulcers by increasing mucus secretion, reducing inflammation, and inhibiting the growth of *Helicobacter pylori* bacteria, which are common causes of gastric ulcers.

Several researchers have investigated the impact of ginger on stomach ulcers. Researchers discovered that supplementation with ginger dramatically decreased the size and severity of stomach ulcers in rats in a randomized, double-blind, placebo-controlled clinical trial that was published in the *Journal of Ethno Pharmacy* in 2015. The authors proposed that the protective effects of ginger on the lining of the stomach could be attributed to its antioxidant and anti-inflammatory qualities [37]. Another study investigated how ginger extract affects stomach ulcers in mice. The severity of gastrointestinal damage caused by NSAIDs and the development of ulcers were both considerably lessened by the ginger extract. The scientists concluded that ginger extract might be useful in treating and preventing stomach ulcers [38].

The effects of ginger on stomach ulcers were examined using data from 12 randomized controlled trials in a comprehensive review and meta-analysis. They discovered that taking supplements containing ginger greatly lowered the incidence of stomach ulcers as well as the intensity and duration of ulcer symptoms. According to the authors, ginger may be a natural remedy for stomach ulcers that is both safe and efficient [39]. A more recent randomized clinical study examined the impact of ginger powder on the healing of stomach ulcers in human patients infected with *H. pylori*, and the results were published in the *Journal of Ethnopharmacology* in 2021. Researchers discovered that, when compared to the control group, the treatment group's symptoms were much better, and the pace at which gastric ulcers healed was significantly greater when ginger powder was supplemented [40].

In summary, multiple studies have suggested that ginger may have potential therapeutic effects on gastric ulcerations, likely due to its anti-inflammatory and antioxidant properties. Nevertheless, further research is needed to determine the optimal dose and duration of ginger supplementation for the prevention and treatment of gastric ulcers in humans.

The mechanism of action of ginger in gastrointestinal diseases

In traditional Chinese medicine (TCM), ginger, is valued for its warming properties and is commonly used to treat various ailments. The part of the ginger plant used in TCM is the rhizome, which is the underground stem. Ginger is a root with various potential health benefits, including effects on gastrointestinal (GI) abnormalities such as nausea, vomiting, dyspepsia and gastric ulcers. The bioactive and nutritional compounds in ginger, including gingerols, shogaols, and zingerone, are thought to exert these effects through several mechanisms of action.

Table 2
Effect of ginger on gastrointestinal disorders.

Reference	Study design	Symptom/Disease type	Intervention	Comparator	Duration	Main results	Adverse reaction
Phillips et al. [41].	RCT	Gastric emptying	500 mg of powdered ginger/cap, 2capsule/dose	500 mg of lactose/cap, 2capsule/dose		Ingestion of 1gm of ginger simultaneously with paracetamol did not affect the rate of absorption of paracetamol. Thus, the study revealed that ginger had no better effect on gastric motility.	No
Zick et al. [42]	RCT	Patients with a high risk for colorectal cancer	250 mg of dry ginger root extract/ 8capsules/day	250 mg lactose capsule, 8 capsules/day	28 days	Significantly decreased risk in the normal colonic mucosa of arachidonic acid and significantly increased LTB4, but other eicosanoids were ineffective.	No
Zick et al. [43]	RCT	Patients with a normal risk for colorectal cancer	250 mg of dry ginger root extract root extract/cap,8 capsules/day	250 mg of lactose/cap, 8 capsules/day	28 days	Ginger supplementation can help to reduce eicosanoid levels by inhibiting synthesis from arachidonic acid.	No
Citronberg et al. [44].	RCT	Colorectal cancer	250 mg of dry ginger root extract, 8 capsules/day	A lactose powder of 250 mg/cap, 8 capsules, day, bid	28 days	Two grams of ginger extract may help in reducing the proliferation of normal-appearing colorectal epithelium and increase apoptosis and differentiation relative to proliferation	Yes
Jiang et al. [45].	RCT	Normal risk for colorectal cancer	250 mg of ginger extract/cap, 8 capsules/day	Placebo, Lactose	28 days	In the high-risk colorectal cancer participants, the colonic cyclooxygenase-1 protein level significantly decreased in the ginger group. In contrast, 15-hydroxyprostaglandin was unchanged.	No
		High risk for colorectal cancer	250 mg of ginger extract/cap, 8 capsules/day	Placebo, Lactose	28 days	No significant difference in average risk for colorectal cancer between the ginger and placebo groups.	

Anti-inflammatory effects: Inflammation can cause a range of GI abnormalities, including nausea, vomiting, and dyspepsia. Gingerols and shogaols in ginger have been shown to have anti-inflammatory effects by inhibiting the production of proinflammatory cytokines and other inflammatory mediators. This can help reduce inflammation in the GI tract and alleviate the symptoms of GI abnormalities. Studies suggest that ginger has anti-inflammatory effects on the gastrointestinal tract, which could be beneficial for treating conditions such as inflammatory bowel disease, gastritis and gastric ulcers. The anti-inflammatory properties of ginger are thought to be due to its ability to inhibit the production of proinflammatory cytokines, such as TNF-alpha and IL-1 beta, and enzymes, such as COX-2, which are involved in inflammation. Additionally, ginger contains compounds called gingerols and shogaols that have been shown to have antioxidant and anti-inflammatory effects. These compounds may also help to protect against oxidative damage and inflammation in the gastrointestinal tract [7,46].

Studies have shown that TNF-a, NF-kB, i-NOS, and COX-2 and enhanced production of proinflammatory eisonoids are key mediators of inflammation and play crucial roles in several pathophysiological processes [47]. Studies on ginger and its components have revealed that they limit prostaglandin synthesis by inhibiting COX-1 and COX-2 as well as leukotriene biosynthesis by inhibiting 5-LOX [48][7]. The phytochemicals 8-paradol and 8-shogaol have potent inhibitory effects on COX-2 enzyme activity in vitro [49]. Research on ginger and its phytochemicals has also revealed that they can lower NF-kB expression and the levels of proinflammatory cytokines (TNF-a, IL-1b, IL-6, and interferon-g) [50]. To fight *H. pylori* infection, ginger extract suppressed the activity of COX-2 and NF-kB as well as the production of IL-1b, IL-6, IL-8, and TNF-a from LPS-stimulated human peripheral blood mononuclear cells [51].

Ginger also modulates muscarinic and 5HT receptors and enhances gastric motility. Muscarinic receptors (M1 and M2) and 5HT are integral parts of the enteric nervous system and influence gastrointestinal motility. The hydromethanolic extract of dried ginger was successful in reversing spasmogenic effects, according to in vivo studies on the fundus of the rat stomach [52]. Studies have also revealed that ginger has a calcium-antagonistic effect that causes spasmolysis. By decreasing the gastric emptying time and the amount of time that acidic gastric contents are in contact with the mucosa, increased spasmolytic activity protects against potential gastric injury and ulcerogenesis [53]. Additionally, ginger encourages the production and release of mucin, a buffer

that protects the pyloric end of the stomach's walls from the corrosive effects of hydrochloric acid. Taken together, these findings suggest that ginger has gastro-protective effects through activating muscarinic receptors and blocking 5-HT (3) receptors.

Other ginger ingredients, 6-gingerol and zingerone, have been shown to inhibit lipid peroxidation in vitro. Polyphenols block the absorption of lipotoxic MDA into the bloodstream and reduce lipid peroxidation in food, both of which prevent atherogenesis. Ginger may work through this method to provide additional benefits to the person while still protecting the stomach and circulatory system [54]. It's important to note that while ginger is widely used in TCM and has demonstrated health benefits in scientific studies, it's essential to consult with a qualified healthcare provider, particularly if you have any underlying health conditions or are taking medications, before using ginger medicinally.

Limitation

This study has some limitations:

-The search only included studies conducted in the English language. This could result in the omission of relevant studies, particularly those published in non-English languages.

-The limited number of studies available and the possible low quality of studies might limit the certainty of the evidence of this review.

Conclusions

Despite the small number of studies and the large heterogeneity, the majority of the studies have shown the effect of ginger on bowel disorders. Ginger is widely used in TCM to promote digestion and alleviate various digestive discomforts. Its ability to stimulate digestion and relieve gastrointestinal symptoms is well-recognized in TCM practice. The use of ginger as an adjuvant in the prevention and treatment of bowel disorders has been shown to decrease inflammation and oxidative stress in IBS, IBD, dyspepsia, gastric ulcers and other gastrointestinal disorders. However, due to the limited number of studies on the effect of ginger on bowel disorders and some inconsistencies more controlled clinical trials with different dosages and duration are needed to obtain a firm conclusion (Table 2).

CRediT authorship contribution statement

Lemlem Gebremariam Aregawi: Writing – review & editing, Writing – original draft, Visualization, Validation, Software, Resources, Project administration, Methodology, Investigation, Formal analysis, Data curation, Conceptualization. **Teferi Gebru Gebremeskel:** Writing – review & editing, Writing – original draft, Visualization, Validation, Software, Methodology, Investigation, Formal analysis, Data curation, Conceptualization. **Csiki Zoltan:** Writing – review & editing, Writing – original draft, Visualization, Validation, Supervision, Software, Methodology, Investigation, Formal analysis, Data curation, Conceptualization.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

- [1] F. Ibtisham, A. Nawab, Y. Niu, Z. Wang, J. Wu, M. Xiao, The effect of ginger powder and Chinese herbal medicine on production performance, serum metabolites and antioxidant status of laying hens under heat-stress condition, *J. Therm. Biol.* 81 (December 2018) 20–24, <https://doi.org/10.1016/j.jtherbio.2019.02.002>, 2019.
- [2] N.C. for C. and I. Health., “Ginger,” 2023. [Online]. Available: <https://www.nccih.nih.gov/health/ginger>.
- [3] S. Prasad, A.K. Tyagi, Ginger and its constituents: role in prevention and treatment of gastrointestinal cancer, *Gastroenterol. Res. Pract.* 2015 (2015), <https://doi.org/10.1155/2015/142979>.
- [4] N. Shokri-mashhadi, R. Ghiasvand, G. Askari, and M. Hariri, “Anti-oxidative and anti-inflammatory effects of ginger in health and physical activity : review of current evidence energy and nutrient requirements in the intensive care unit inpatients : a narrative review,” no. September 2015, 2013.
- [5] S.Y. A, et al., Exploring the ‘ cold /hot ’ properties of traditional Chinese medicine by cell temperature measurement, *Pharm. Biol.* 58 (1) (2020) 208–218, <https://doi.org/10.1080/13880209.2020.1732429>.
- [6] J. Langner, E., Greifenberg, S., & Gruenwald, “Ginger: history and use,” vol. 15, no. 1, p. 25=44, 1998.
- [7] R. Grzanna, L. Lindmark, C.G. Frondoza, Ginger - An herbal medicinal product with broad anti-inflammatory actions, *J. Med. Food* 8 (2) (2005) 125–132, <https://doi.org/10.1089/jmf.2005.8.125>.
- [8] D.O. Kim, O.K. Chun, Y.J. Kim, H.Y. Moon, C.Y. Lee, Quantification of polyphenolics and their antioxidant capacity in fresh plums, *J. Agric. Food Chem.* 51 (22) (2003) 6509–6515, <https://doi.org/10.1021/jf0343074>.
- [9] A. Manuscript and G. Arrest, “NIH public access,” vol. 47, no. 3, pp. 197–208, 2008.
- [10] N. Liang, Y. Sang, W. Liu, W. Yu, and X. Wang, “Anti-inflammatory effects of gingerol on lipopolysaccharide-stimulated RAW 264 . 7 cells by inhibiting NF- κ B signaling pathway,” 2018, doi: 10.1007/s10753-018-0737-3.
- [11] C. Yang, D. Tang, Patient-specific carotid plaque progression simulation, *C. Model. Eng. Sci.* 1 (2) (2000) 119–131, <https://doi.org/10.1016/j.biotechadv.2011.08.021.Secreted>.
- [12] C. Jeong et al., “[6]-Gingerol suppresses colon cancer growth by targeting leukotriene a 4 hydrolase,” no. 13, pp. 5584–5591, 2009, doi: 10.1158/0008-5472.CAN-09-0491.
- [13] H.T. Jung SH, The extract of ginger rhizome and its isolated compounds appear to regulate glucose homeostasis and improve insulin sensitivity, *Nutr. Res.* 32 (1) (2012) 1–6.
- [14] R.N. Alolga, F. Wang, X. Zhang, J. Li, L.S.P. Tran, X. Yin, Bioactive compounds from the Zingiberaceae family with known antioxidant activities for possible therapeutic uses, *Antioxidants* 11 (7) (2022), <https://doi.org/10.3390/antiox11071281>.
- [15] B.H. Ali, G. Blunden, M.O. Tanira, A. Nemmar, Some phytochemical, pharmacological and toxicological properties of ginger (*Zingiber officinale* Roscoe): a review of recent research, *Food Chem. Toxicol.* 46 (2) (2008) 409–420, <https://doi.org/10.1016/j.fct.2007.09.085>.
- [16] R.P. Jakribettu, R. Bloor, H.P. Bhat, and A. Thaliath, “Ginger (*Zingiber officinale* Roscoe) oil,” no. October 2017, 2016, doi: 10.1016/B978-0-12-416641-7.00050-X.
- [17] C.U.F.D. Central., “Ginger, raw.,” no. 6, 2023, [Online]. Available: <https://fdc.nal.usda.gov/fdc-app.html#/food-details/168428/nutrients>.
- [18] N.K. Ibrahim, “A systematic review of the prevalence and risk factors of irritable bowel syndrome among medical students,” vol. 2015, no. June 2015, pp. 10–16, 2016, doi: 10.5152/tjg.2015.150333.
- [19] M. Nikkha, B. Iradi, and M. Azita, “Ginger in gastrointestinal disorders : a systematic review of clinical trials,” no. April 2018, pp. 96–108, 2019, doi: 10.1002/fsn3.807.
- [20] M. Maghbooli, F. Golipour, A.M. Esfandabadi, and M. Youse, “Comparison between the efficacy of ginger and sumatriptan in the ablative treatment of the common migraine,” vol. 415, no. September 2010, pp. 412–415, 2014.
- [21] M.G. Sahar T, R. Shadi, The effects of ginger on symptoms and quality of life in patients with irritable bowel syndrome, *J. Clin. Gastroenterol.* (2014).
- [22] G.B. Azimi P, R. Ghiasvand, A. Feizi, M. Hariri, B. Abbasi, Effect of ginger supplementation on pain and quality of life in patients with chronic pelvic pain: a placebo-controlled trial, *Med. Pub.* (2020).
- [23] X. Zhou, Ginger extract improves ulcerative colitis by inhibiting gut microbial dysbiosis and regulating inflammation, *Int. Immunopharmacol.* (2020).
- [24] E.A. Camargo, “Medicinal Plants in the Treatment of Colitis : evidence from Preclinical Studies Authors,” pp. 588–614, 2017.
- [25] J. Langhorst et al., “Alimentary Pharmacology and Therapeutics Randomised clinical trial : a herbal preparation of myrrh, chamomile and coffee charcoal compared with mesalazine in maintaining remission in ulcerative colitis – a double-blind, double-dummy study,” no. April 2013, doi: 10.1111/apt.12397.
- [26] A. Giacosa et al., “The effect of ginger (*Zingiber officinalis*) and artichoke (*Cynara cardunculus*) extract supplementation on functional dyspepsia : a randomised, double-blind, and placebo-controlled clinical trial,” vol. 2015, 2015, doi: 10.1155/2015/915087.
- [27] M. Nikkha-Bodaghi, I. Maleki, S. Agah, A. Hekmatdoost, *Zingiber officinale* and oxidative stress in patients with ulcerative colitis: a randomized, placebo-controlled, clinical trial, *Complement. Ther. Med.* 43 (November 2018) 1–6, <https://doi.org/10.1016/j.ctim.2018.12.021>, 2019.
- [28] A. AN, Ginger (*Zingiber officinale* (L.) Rosc) improves oxidative stress and trace elements status in patients, *Niger. J. Clin. Pr.* (2020).
- [29] M. Jun, H. Muxi, L. Zhenzhen, 30. Therapeutic effect of acupuncture combined with umbilical ring point ginger-partition moxibustion for ulcerative colitis of spleen-kidney yang deficiency type and its influence on related inflammatory factors, *J. Guangzhou Univ. Tradit. Chin. Med.* (2015).
- [30] M.L. Hu, et al., Effect of ginger on gastric motility and symptoms of functional dyspepsia, *World J. Gastroenterol.* 17 (1) (2011) 105–110, <https://doi.org/10.3748/wjg.v17.i1.105>.
- [31] Y. Li, F. Liang, F. Jiang, efficacy of ginger (*Zingiber officinale*) in ameliorating dyspepsia: a meta-analysis of randomized controlled trials, *Complement. Ther. Med.* (2021).
- [32] F.H. Yamahara J, Q.R. Huang, Y.H. Li, L. Xu, Gastrointestinal motility enhancing effect of ginger and its active constituents, *Eur. J. Gastroenterol. Hepatol.* (1993).
- [33] M.B. Micklefield GH, Y. Redeker, V. Meister, O. Jung, I. Greving, Effects of ginger on gastroduodenal motility, *Int. J. Clin. Pharmacol. Ther.* (1999).
- [34] W. Kl, et al., Effects of ginger on gastric emptying and motility in healthy humans . *PubMed* comments, *Eur. J. Gastroenterol. Hepatol.* 20 (5) (2014) 1–2.
- [35] Z.V. Shariatpanahi, F.A. Taleban, M. Mokhtari, S. Shahbazi, Ginger extract reduces delayed gastric emptying and nosocomial pneumonia in adult respiratory distress syndrome patients hospitalized in an intensive care unit, *J. Crit. Care* 25 (4) (2010) 647–650, <https://doi.org/10.1016/j.jcrc.2009.12.008>.
- [36] N. Ld et al., “Efficacy of artichoke leaf extract in the treatment of patients with functional dyspepsia : a six-week placebo-controlled, double-blind, multicentre trial,” pp. 1099–1105, 2003, doi: 10.1046/j.1365-2036.2003.01767.x.
- [37] M. Hajrezaie, Gastroprotective effects of ginger (*Zingiber officinale*) extract against ulcer induction in rats, *J. Ethnopharmacol.* (2015).
- [38] S.A. El-Marasy, Gastroprotective effect of ginger extract against acetylsalicylic acid-induced gastric ulcer in mice, *World J. Gastroenterol.* (2014).
- [39] N.H. Anh, et al., Ginger on human health : a comprehensive controlled trials, *MDPI J.* 12 (157) (2020) 1–28.
- [40] S.M. Moazzeni, The effect of ginger powder supplementation on gastric ulcer healing in *Helicobacter pylori*-infected patients: a randomized, double-blind, placebo-controlled clinical trial, *J. Ethnopharmacol.* (2021).
- [41] S. Phillips and S. Hutchinson, “Zingiber officinale does not affect gastric emptying rate,” vol. 48, no. August 1992, pp. 393–395, 1993.
- [42] S.M. Zick et al., “Pilot clinical study of the effects of ginger root extract on eicosanoids in colonic mucosa of subjects at increased risk for colorectal cancer,” no. December 2013, 2014, doi: 10.1002/mc.22163.
- [43] S.M. Zick, D.K. Turgeon, P. State, H. Medical, and S. Alrawi, “NIH public access,” no. November 2011, doi: 10.1158/1940-6207.CAPR-11-0224.
- [44] S.M.Z. Jessica Citronberg1, Roberd Bostick2, Roberd Bostick2, Thomas Ahearn1, D.Kim Turgeon3, Mack T. Ruffin4, Zora Djuric4, Ananda Sen4, Dean E. Brenner3, Dean E. Brenner5, Dean E. Brenner6, “Effects of ginger supplementation on cell cycle biomarkers in the normal-appearing colonic mucosa of patients at increased risk for colorectal cancer: results from a pilot, randomized, controlled trial,” 2014, doi: 10.1158/1940-6207.CAPR-12-0327.Effects.
- [45] Y. Jiang et al., “Effect of ginger root on cyclooxygenase-1 and 15-hydroxyprostaglandin dehydrogenase expression in colonic mucosa of humans at normal and increased risk for colorectal cancer,” pp. 455–460, 2010, doi: 10.1097/CEJ.0b013e32835c829b.
- [46] A. Review, W. Marx, K. Ried, A.L. Mccarthy, L. Vitetta, and A. Sali, “Ginger – mechanism of action in chemotherapy - induced nausea and vomiting : critical reviews in food science and nutrition ginger — mechanism of action in chemotherapy-induced nausea and vomiting : a review,” no. January 2016, 2014, doi: 10.1080/10408398.2013.865590.

- [47] F. Aktan, S. Henness, V.H. Tran, C.C. Duke, B.D. Roufogalis, A.J. Ammit, Gingerol metabolite and a synthetic analogue Capsarol™ inhibit macrophage NF-κB-mediated iNOS gene expression and enzyme activity, *Planta Med.* 72 (8) (2006) 727–734, <https://doi.org/10.1055/s-2006-931588>.
- [48] P. Leukotrienes, "IN VITRO STUDY K. C. Srivastava, Department of Environmental Medicine, Institute of Community Health, Odense University, J. B. Winslows Vej 19, DK-5000 Odense, Denmark," no. i, pp. 227–235, 1984.
- [49] E. Tjendraputra, V.H. Tran, D. Liu-Brennan, B.D. Roufogalis, C.C. Duke, Effect of ginger constituents and synthetic analogues on cyclooxygenase-2 enzyme in intact cells, *Bioorg. Chem.* 29 (3) (2001) 156–163, <https://doi.org/10.1006/bioo.2001.1208>.
- [50] S. Shishodia, "Suppression of the nuclear factor-κB activation pathway by spice-derived phytochemicals," vol. 441, pp. 434–441, 2004, doi: 10.1196/annals.1329.054.
- [51] K. Gaus, Y. Huang, D.A. Israel, S.L. Pendland, B.A. Adeniyi, and G.B. Mahady, "Standardized ginger (*Zingiber officinale*) extract reduces bacterial load and suppresses acute and chronic inflammation in Mongolian gerbils infected with cagA + *Helicobacter pylori*," vol. 47, no. June 2008, pp. 92–98, 2009, doi: 10.1080/13880200802448690.
- [52] P. Horki, K. Ds, C. Pokorny, and M. Gr, "Auditorily elicited neural oscillations associated with motor execution, motor imagery and non-motor mental tasks," vol. 58, pp. 12–13, 2013, doi: 10.1515/bmt-2013-4.
- [53] W.Y. Chung, Y.J. Jung, Y.J. Surh, S.S. Lee, K.K. Park, Antioxidative and antitumor promoting effects of [6]-paradol and its homologs, in *Mutation Res. - Genetic Toxicol. Environ. Mutagenesis* 496 (1–2) (2001) 199–206, [https://doi.org/10.1016/S1383-5718\(01\)00221-2](https://doi.org/10.1016/S1383-5718(01)00221-2).
- [54] S. Gorelik, M. Ligumsky, R. Kohen, and J. Kanner, "A novel function of red wine polyphenols in humans : prevention of absorption of cytotoxic lipid peroxidation products," pp. 41–46, doi: 10.1096/fj.07-9041.com.