

The Effects of drought stress on Soybean (*Glycine max* (L.) Merr.) growth, physiology and quality: A Review

Oqba Basal

PhD student, Department of Crop Production and Applied Ecology

Institute of Crop Sciences, University of Debrecen

oqba@agr.unideb.hu

Summary

Abiotic stresses are one of the most limiting factors inhibit plant's growth, leading to a serious production loss. Drought stress is one of the most destructive abiotic stresses and is still increasing year after year resulting in serious yield losses in many regions of the world, consequently, affecting world's food security for the increasing world population. Soybean is an important grain legume. It is one of the five major crops in the world, an essential source of oil, protein, macronutrients and minerals, and it is known as the main source of plant oil and protein. Harvested area of soybean is increasing globally year after year. However, soybean is the highest drought stress sensitive crop, the water deficit influences the physiology, production and seed composition of this crop. We introduce a review for literatures concerning the changes of the above traits of soybean exposed to drought stress, with past explanations for these changes.

1. Drought stress

Biotic and abiotic stresses are one of the most limiting factors inhibit plant's growth, leading to a serious production loss (Kang et al., 2002; Mahajan and Tuteja, 2005). To adapt with such stresses, plants get morphological, physiological and molecular changes (Bray, 1993; Seki et al., 2003; Yamaguchi-Shinozaki and Shinozaki, 2006). Water scarcity is a major factor limiting agricultural production all over the world, consequently, many efforts were and should be made to adapt edible crops preventing drought stress from making critical harms to them.

2. Soybean

Soybean is an important grain legume. It is one of the five major crops in the world, an essential source of oil (18–22%), protein (35 – 40%), macronutrients and minerals, and it is known as the main source of plant oil and protein (Lei et al., 2006; Maleki et al., 2013). Among crops, soybean is the highest drought stress sensitive (Maleki et al., 2013). Its water requirement is relatively high (Yang et al., 2003), so it is quite important to use the genotypes which best exploit available water and use it more effectively.

3. Effects of drought stress on soybean

3.1 Effects of drought stress on soybean morphological and physiological traits

Drought stress was demonstrated to affect soybean morphology and physiology. Navari-Izzo et al., (1990) found that drought-stressed soybean resulted in a 25% reduction in leaf water potential (LWP). Five days of drought stress during pre-flowering stage resulted in a drop of the LWP to -2.2 MPa, compared to -1.4 MPa for control plants (Lei et al., 2006), which was consistent with Liu et al., (2003) results by which a decrease of WLP was recorded for drought-stressed soybean compared to watered control. Many studies reported similar results (Pennypacker et al., 1990; Siddique et al., 2000; Fu and Huang, 2001; Shaw et al., 2002; Liu et al., 2004a; Liu et al., 2004b; Liu et al., 2005; Hao et al., 2013). Moreover, it was concluded that drought stress affected not only soybean leaf water potential, but also pod water potential causing both to fell (Liu et al., 2004c). Makbul et al., (2011) recorded a significant decline of soybean leaf water potential from -0.88 MPa in unstressed leaves to -1.18 MPa in drought stressed ones, moreover, they suggested that this decline may have resulted in the significant decrease chlorophyll content by 28% in drought-stressed soybean. Hao et al., (2013) found that the chlorophyll content of the drought-stressed soybean decreased by 31% compared to control plants. Similar results were provided by Atti et al., (2004).

Schulze, (1993) reported that drought stress decreased the leaf water potential, resulting in a reduction of the swelling pressure and consequently causing the stomatal closure. Giorio et al., (1999) also reported good positive relationships between leaf water potential and stomatal conductance. Stomatal control was considered as a main physiological factor for optimizing water use under drought conditions (Makbul et al., 2011), preventing excessive water loss under extended drought conditions (Ku et al., 2013). Stomatal conductance, as compared to control, decreased 60% under drought stress (Hao et al., 2013), this result was consistent with previous studies (Liu et al., 2005; Makbul et al., 2011), and was confirmed later (Mak et al., 2014; Mutava et al., 2015). Lei et al., (2006) noticed that the stomatal conductance decreased as the drought progressed, confirming the conclusion that the impact of moisture deficit was greater at the severe stress than at the medium stress (Atti et al., 2004), who demonstrated a reduction in stomatal conductance, transpiration rate and photosynthetic rate by 92%, 85.4% and 78.4%, respectively, under water stress. They showed that stomatal conductance was correlated with transpiration rate more than with photosynthetic rate, which was reported earlier (De Souza et al., 1997). Transpiration rate decreased 53% (Hao et al., 2013), and 57% (Mak et al., 2014) under drought stress, this decrease was due mainly to a decrease of stomatal conductance which was controlled by root-originated ABA, as a 50-fold xylem ABA was measured under drought stress (Liu et al., 2005), that, however, was reported earlier (Samet et al., 1984; Liu et al., 2003) with a significant increase as the stress became severe (Liu et al., 2004_b). Moreover, Mutava et al., (2015) reported a significant ABA increase in a drought tolerant soybean genotype more than in a drought susceptible one under drought stress conditions.

Photosynthetic rates, after 9 days of drought stress, were significantly lower than in the control plants (Liu et al., 2004_a). Similar results were obtained for soybean seedlings (Hao et al., 2013). These results were consistent with other reports at different soybean growing stages (Lei et al., 2006; Mak et al., 2014; Mutava et al., 2015).

Soybean seedling height decreased 4.3% under drought stress (Navari-Izzo et al., 1990), which was reported later at different stages (Atti et al., 2004; Hao et al., 2013; Mak et al., 2014). Particularly, drought stress at V4 stage reduced the plant height more than at R2 (full flowering), R4 (full pod formation) and R6 (full seed filling) stages (Maleki et al., 2013; Li et al., 2013). Although Sionit and Kramer, (1977) reported no significant differences on plant height under drought stress, most researchers reported the opposite (Brady et al., 1974; Kadhem et al., 1985; Demirtas et al., 2010; Maleki et al., 2013). This reduction might be caused by a drought tolerance mechanism, as cell swelling, cell wall and synthesis enzymes are reduced, consequently, growth and plant height are decreased (Levitt, 1980; Austin, 1989).

It was concluded that drought stress increased the root length as the roots searched for water, in addition to the formation of fine roots able to penetrate smaller soil pores (Turner, 1986; Komatsu and Hossain, 2013; Maleki et al., 2013; Khan and Setsuko Komatsu, 2016). However, the root weight of the drought-stressed soybean plants was reduced as compared to the controls (Kausar et al., 2012; Mohammadi et al., 2012; Hao et al., 2013; Oh and Komatsu, 2015; Khan and Setsuko Komatsu, 2016).

Moreover, root to shoot ratio increased under water deficit conditions, because roots are less sensitive than shoots to growth inhibition by drought stress (Wu and Cosgrove, 2000; Franco et al., 2011). Soybean shoot dry matter was reduced under drought stress (Liu et al., 2005; Hao et al., 2013; Mak et al., 2014). The reduction was higher when stress was applied during podding than during seed filling stage, which in part was higher compared to drought stress application during flowering stage (Li et al., 2013).

Relative water content decreased 33% under drought stress (Hao et al., 2013), and it was decreased more for a drought susceptible soybean genotype compared to drought tolerant one (Mutava et al., 2015).

Karam et al. (2005) reported a reduced soybean leaf area index under drought stress conditions by an average of 30% compared to controls. Similar results were obtained later (Atti et al., 2004; Demirtas et al., 2010; Li et al., 2013).

Both establishment and activity of the legume–Rhizobium symbiosis were reported to be extremely sensitive to drought stress (Zablotowicz et al., 1981; Kirda et al., 1989). Consequently, legume productivity can be greatly reduced both by moderate and severe drought (Saxena et al., 1993; Subbarao et al., 1995). Smith et al. (1988) reported substantial decreases in nodule mass in drought-stressed soybean. However, Sinclair et al. (1988) noticed a decrease in nodule number and dry weight only after a severe drought.

Samarah et al., (2009) and Heatherly, (1993) reported the germination rate to be reduced for soybean under drought stress during seed filling period. Flowering period, Podding period, flower number and

leaf number were reported to be reduced under drought stress (Atti et al., 2004; Mak et al., 2014; He et al., 2016).

3.2 Effects of drought stress on soybean production traits

Frederick et al. (2001) noticed a large effect of drought stress on the final number of soybean branches formed. Later, Atti et al., (2004) reported that water reduced the mean number of branches by 28% compared to the controls. The number of pods per plant was reported to be reduced under drought stress by many researchers (Kadhem et al., 1985; Desclaux and Roumet, 2000; De Costa and Shanmugathan, 2002; Demirtas et al., 2010; Sadeghipour and Abbasi, 2012; Li et al., 2013; Mak et al., 2014). At the beginning of pod development, drought stress reduced pod number by 92.7%, while during pod lengthening, the reduction was 81.6% compared to the controls, due to the cumulative effects of reductions in pod induction, young pods (due to abortion, Kokubun, 2011) and pod enlargement (Atti et al., 2004) and to the reductions in flower number (He et al., 2016).

Reduced seed number per plant under drought stress was recorded (Demirtas et al., 2010; Li et al., 2013; He et al., 2016), this results, however, was reported earlier (Kadhem et al., 1985; Karam et al., 2005). Highest reduction in seed numbers per plant occurred in flowering stage (Sionit and Kramer, 1977; Smiciklas et al., 1992; Maleki et al., 2013). Bord and Hartville, (1998) suggested that drought stress during flower formation led to a shorter flowering period and produced fewer flowers, fewer pods, and consequently, significantly smaller number of seeds per plant. However, it was concluded earlier that drought stress during the seed set period reduces seed number (Meckel et al., 1984; Rotundo and Westgate, 2010).

Weight of soybean seeds was affected by drought stress (Sionit and Kramer, 1977; Rose, 1988; Liu et al., 2003; Karam et al., 2005; Demirtas et al., 2010; Sadeghipour and Abbasi, 2012; Li et al., 2013). 15.2% reduction in the weight of 100 seeds grown under drought stress during R5 (beginning of seed filling) stage was noticed compared with non-stressed plants (Gutierrez-Gonzalez et al., 2010), which was confirmed later by He et al., (2016). It was suggested that the weight of grain is determined in the late reproductive stage, and therefore, more affected by drought stress at R5 (beginning of seed filling) stage, this decrease could be due to loss of assimilate to seeds (Yordanov et al., 2003), or to a shortened seed filling period (Demirtas et al., 2010).

Soybean seed yield, when exposed to drought stress, was reduced (Rose, 1988; Kokubun et al., 2001; Sadeghipour and Abbasi, 2012; Li et al., 2013). More specifically, severe drought stress reduced the seed yield of soybean more than moderate drought stress (Dornbos and Mullen, 1992).

The soybean genotype was found to have a role on yield loss (Bellaloui and Mengistu, 2008; He et al., 2016), which is consistent with the conclusions of Brown et al., (1985) who found the reduction to be significant (Maleki et al., 2013).

Doss et al., (1974) and Sionet and Kramer (1977) found that the drought stress during R3 (beginning of pod formation) and R4 (full pod formation) stages resulted in greater yield reduction than that occurred

during R1 (beginning of flowering) and R2 (full flowering) stages. Moreover, Demirtas et al., (2010) and Maleki et al., (2013) concluded that drought stress during R5 (beginning of seed filling) stage caused the most reduction, which was previously reported (Ashley and Ethridge 1978; Huck et al., 1983; Eck et al., 1987; Foroud et al., 1993; Karam et al., 2005;). It was suggested that drought stress during R5 (beginning of seed filling) stage shortened the seed-filling period and reduced seed yield (Meckel et al., 1984; Frederick et al., 1991; Smiciklas et al., 1992). Others suggested it to be due to the reduction of pod numbers per plant (Atti et al., 2004), seeds number (Dornbos et al., 1989), and seeds weight (Samarah et al., 2006; Demirtas et al., 2010; Maleki et al., 2013). However, seed yield was found to have a significantly positive correlation with plant height, number of pods and seeds per plant, seed weight, harvest index and days to maturity (Georgiev, 2004; Maleki et al., 2013). A reduction in soybean biomass due to drought stress was early reported (Read and Bartlett, 1972), and confirmed lately (Khan and Setsuko Komatsu, 2016). Particularly, the biomass was significantly decreased, when drought stress was applied in the pod filling period (Demirtas et al., 2010) more than V4 stage (Maleki et al., 2013). The effect of drought stress on harvest index was significant, as drought occurrence at R5 (beginning of seed filling) stage reduced it by 27.9% compared to control (Maleki et al., 2013). Ashley and Ethridge, (1978) suggested that the harvest index was reduced due to the loss of flowers and the decrease in seed numbers per plant.

$WUE_{\text{yield, biomass}}$ (Water Use Efficiency) for soybean was improved under a certain range of drought stress (Liu et al., 2005; Lei et al., 2006; Demirtas et al., 2010; Li et al., 2013). Moreover, Karam et al. (2005) found that soybean WUE at R2 (full flowering), R6 (full seed filling) stage was 13 and 4% higher than the control, respectively, whereas at R5 (beginning of seed filling) was 17% lower than the control. That was recently demonstrated (He et al., 2016).

3.3 Effects of drought stress on soybean quality traits

The protein and oil contents of soybean seed are the major parameters determining the nutritional value (Chung et al., 2003). The relationship between drought stress and soybean seed composition, however, still remains controversial (Medic and Atkinson 2014). Dornbos and Mullen (1992), Kumar et al. (2006) and Rotundo and Westgate (2009; 2010) reported 2–23 % increased protein contents under drought stress compared to controls. Environmental factors affect protein concentrations increasing them in the harvested parts (Rose, 1988; Dornbos and Mullen, 1992; Bellaloui and Mengistu, 2008; Wang and Frei, 2011). However, few studies showed no effect (Sionit and Kramer, 1977) or lower protein concentration (Carrera et al., 2009; Specht et al., 2001; Rose, 1988; Boydak et al., 2002). Differences among the reported conclusions were suggested to be due to the timing and intensity of the drought stress during the different stages (Carrera et al., 2009). Bellaloui and Mengistu (2008) suggested that the plant's response to drought stress might be cultivar-dependent.

In general, soybean seed protein content is negatively correlated with the amount of seed oil (Chung et al., 2003). Results of most studies indicated that the drought stress reduced oil content in the seed (Rose,

1988; Specht et al., 2001; Bellaloui and Mengistu, 2008; Rotundo and Westgate, 2009), whereas few other reports showed increased oil content with the water deficit (Specht et al., 2001; Boydak et al., 2002). However, Gao et al. (2009) reported that drought stress had little effect on the oil content. The effect of drought stress at different stages was significant on oil content, and the lowest oil percentage was obtained when drought stress was applied at V5 stage (Dornbos and Mullen, 1992; Smiciklas et al., 1992; Maleki et al., 2013). The oil content had a significantly positive correlation with seed weight (Maleki et al., 2013).

The fatty acid composition differed when soybean was subjected to drought stress; a decrease in Palmitic, Linoleic Linolenic acid and an increase in Stearic and Oleic acid were noticed (Bellaloui and Mengistu, 2008; Gao et al., 2009). The lipid content also increased 11.4% under drought stress conditions (Navari-Izzo et al., 1990), particularly, an increase in glycolipids, diacylglycerols, triacylglycerols and free sterols was recorded while free fatty acids and phospholipids decreased by drought stress (Navari-Izzo et al., 1990). Dornbos and Mullen, (1992) suggested that the increase in triacylglycerols and diacylglycerols of stressed soybean increased the proportion of stearic acid, while oleic acid content decreased.

Bennet et al. (2004) found that isoflavones were reduced 2.5-fold under drought stress applied on soybeans. These results were confirmed as unstressed soybean seeds were found to have increased total isoflavone, daidzein, and genistein contents by 1.2-folds but without differences in the content of glycitein (Lozovaya et al., 2005). Britz and Kremer, (2002) concluded that the tocopherol content of soybeans was insignificantly affected by the drought stress. Drought stress decreased the non-structural carbohydrate quantity in soybean pods (Liu et al., 2004c).

Free proline increased in drought stressed soybean plants (De Ronde et al., 2004) because of the increase in the of enzymes involved in proline biosynthesis and the decrease of those involved in proline degradation (Kiyosue et al., 1996).

Significant decrease in seed isoflavones in soybean seeds under drought stress compared to control was noticed (Bennett et al., 2004, Al-Tawaha et al., 2007); the reduction increased as the stress increased (Gutierrez Gonzalez et al., 2010).

Drought stress increased alfa-tocopherol concentration (Britz and Kremer, 2002).

4. Conclusion

Drought stress is an important abiotic stress affecting soybean traits, resulting in considerable yield losses by different mechanisms. More studies should be generated in order to better understand the soybean seed quality under water deficit conditions, as the past studies demonstrated different results. In addition, different soybean genotypes need to be compared in order to find tolerant genotypes.

5. References

- Al-Tawaha, A. M., Seguin, P., Smith, D. L., Bonnell, R. B., 2007. Irrigation level affects isoflavone concentrations of early maturing soya bean cultivars. *J. Agronom. Crop Sci.* 193, 238 – 246.
- Ashley, D. A., Ethridge, W. J., 1978. Irrigation effects on vegetative and reproductive development of three soybeans cultivars. *Agron. J.* 70, 467-471.
- Atti, S., Bonnell, R., Smith, D., Prasher, S., 2004. Response of an Indeterminate Soybean {*Glycine Max* (L.) Merr} to Chronic Water Deficit During Reproductive Development Under Greenhouse Conditions. *Canadian Water Resources Journal / Revue canadienne des ressources hydriques.* 29(4), 209-222.
- Austin, R. B., 1989. Maximizing crop production in water limited environments. 13-25. In: F. W. G. Baker. *Drought resistance in cereals.* CAB International, Wallingford, England 2220.
- Bellaloui, N., Mengistu, A. 2008. Seed composition is influenced by irrigation regimes and cultivar differences in soybean. *Irrig Sci.* 26, 261–268.
- Bennett, J. O., Yu, O., Heatherly, L. G., Krishnan, H. B. 2004. Accumulation of genistein and daidzein, soybean isoflavones implicated in promoting human health, is significantly elevated by irrigation. *J. Agric. Food Chem.* 52, 7574 – 7579.
- Bord, J. E., Harville, B. G. 1998. Late -planted soybean yield response to reproductive source/sink stress. *Crop Science* 38, 763-771.
- Boydak, E., Alpaslan, M., Hayta, M., Gercek, S., Simsek, M. 2002. Seed composition of soybeans grown in the Harran region of Turkey as affected by row spacing and irrigation. *J Agric Food Chem.* 50, 4718–4720.
- Brady, R. A., Stone, L. R., Nickell, C. D., Powers, W. L. 1974. Water conservation through proper timing of soybean irrigation. *J. Soil Water Conserv.* 29, 266-268.
- Bray, E. A. 1993. Molecular responses to water deficit. *Plant Physiol.* 103, 1035–1040.
- Britz, S. J., Kremer, D. F., 2002. Warm temperatures or drought during seed maturation increase free alpha-tocopherol in seeds of soybean (*Glycine max* L, Merr.). *J. Agric. Food Chem.* 50, 6058–6063.
- Brown, E., Brown, D., Caviness, C. 1985. Response of selected soybean cultivars to soil moisture deficit. *Agronomy Journal* 77(2), 274-278.
- Carrera, C., Martinez, M.J., Dardanelli, J., Balzarini, M. 2009. Water deficit effect on the relationship between temperature during the seed fill period and soybean seed oil and protein concentrations. *Crop Sci.* 49, 990–998.
- Chung, J., Babka, H. L., Graef, G. L., Staswick, P. E., Lee, D. J., Cregan, P. B., Shoemaker, R. C., Specht, J. E. 2003. The seed protein, oil, and yield QTL on soybean linkage group I. *Crop Science*, 43(3), 1053-1067.
- De Costa, W. A. J. M., Shanmugathan, K. N. 2002. Physiology of yield determination of soybean (*Glycine max* (L.) Merr.) under different irrigation regimes in the sub-humid zone of Sri Lanka, *Field Crops Res.* 75, 23-35.
- Demirtas, Ç. D., Yazgan, S., Candogan, B. C., Sincik, M., Büyükçangaz, H., Göksoy, A. T. 2010. Quality and yield response of soybean (*Glycine max* (L.) Merrill) to drought stress in sub-humid environment. *African Journal of Biotechnology* 9(41), 6873-6881, 11.
- Desclaux, D. H. T., Roumet, P. 2000. Identification of soybean plant characteristics that indicate the timing of drought stress. *Crop Sci. Madison*, 40(3), 716-722.
- De Ronde, J. A., Cress, W. A., Kruger, G. H. J., Strasser, R. J., Van Staden, J. 2004. Photosynthetic response of transgenic soybean plants, containing an Arabidopsis P5CR gene, during heat and drought stress. *Journal of Plant Physiology* 161, 1211–1224.

- De Souza, P. I., Egli, D. B., Bruening, W. P. 1997. Water Stress During Seed Filling and Leaf Senescence in Soybean. *Agronomy Journal* 89, 807-812.
- Dornbos, D. L., Mullen, R. E., Shibles, R. E. 1989. Drought stress effects during seed fill on soybean seed germination and vigor. *Crop Science* 29(2), 476-480.
- Dornbos, D. L., Mullen, R. E. 1992. Soybean seed protein and oil contents and fatty acid composition adjustments by drought and temperature. *Journal of the American Oil Chemists' Society* 69(3), 228-231.
- Doss, B. D., Pearson R. W., Rogers H. T. 1974. Effect of soil water stress at various growth stages on soybean yield. *Agron. J.* 66, 297-299.
- Eck, H. V., Mathers, A. C., Musick, J. T. 1987. Plant water stress at various growth stages and growth and yield of soybeans. *Field Crops Research* 17(1), 1-16.
- Foroud, N., Mundel, H. H., Saindon, G., Entz, T. 1993. Effect of level and timing of moisture stress on soybean yield components. *Irrigat. Sci.* 13, 149-155.
- Franco, J. A., Banon, S., Vicente, M. J., Miralles, J., Martinez-Sanchez, J. J. 2011. Root development in horticultural plants grown under abiotic stress conditions - a review, *J. Hortic. Sci. Biotechnol.* 86, 543-556.
- Frederick, J.R., Camp, C. R., Bauer, P. J. 2001. Drought-Stress effects on branch and main stem seed yield and yield components of determinate soybean. *Crop Sci.* 41, 759-763.
- Frederick, J. R., Woolley, J. T., Hesketh, J. D., Peters, D. B. 1991. Seed yield and agronomic traits of old and modern soybean cultivars under irrigation and soil water-deficit. *Field Crops Res.* 27, 71-82.
- Fu, J., Huang, B. 2001. Involvement of antioxidants and lipid peroxidation in the adaptation of two cool-season grasses to localized drought stress. *Environ Exp. Bot.* 45, 105-114.
- Gao, J., Hao, X., Thelen, K. D., Robertson, G. P. 2009. Agronomic management system and precipitation effects on soybean oil and fatty acid profiles. *Crop Sci.* 49, 1049-1057.
- Georgiev, G. 2004. Influence of moisture conditions on the yield of soybean varieties. *Plant Sci.* 5, 406-410.
- Giorio, P., Sorrentino, G., d'Andria, R. 1999. Stomatal behavior, leaf water status and photosynthetic response in field-grown olive trees under water deficit. *Environ Exp. Bot.* 42, 95-104.
- Gutierrez-Gonzalez, J. J., Guttikonda, S. K., Tran, L. S., Aldrich, D. L., Zhong, R., Yu, O., Nguyen, H. T., Sleper, D. A. 2010. Differential Expression of Isoflavone Biosynthetic Genes in Soybean During Water Deficits. *Plant Cell Physiol.* 51(6), 936-948.
- Hao, L., Wang, Y., Zhang, J., Xie, Y., Zhang, M., Duan, L., Li, Z. 2013. Coronatine enhances drought tolerance via improving antioxidative capacity to maintaining higher photosynthetic performance in soybean. *Plant Science* 210, 1-9.
- He, J., Du, Y-L., Wang, T., Turner, N. C., Yang, R-P., Jin, Y., Xi, Y., Zhang, C., Cui, T., Fang, X-W., Li, F-M. 2016. Conserved water use improves the yield performance of soybean (*Glycine max* (L.) Merr.) under drought. *Agric. Water Manage.* <http://dx.doi.org/10.1016/j.agwat.2016.07.008>.
- Heatherly, L. G. 1993. Drought stress and irrigation effects on germination of harvested soybean seed. *Crop Science* 33(4), 777-781.
- Huck, M. G., Ishihara, K., Peterson, C. M., Ushijima, T. 1983. Soybean adaptation to water stress at selected stages of growth. *Plant Physiol.* 73, 422-427.
- Karam, F., Masaad, R., Sfeir, T., Mounzer, O., Roupheal, Y. 2005. Evapotranspiration and seed yield of field grown soybean under deficit irrigation conditions. *Agric. Water Manage.* 75, 226-244.

- Kadhem, F. A., Specht, J. E., Williams, J. H. 1985. Soybean irrigation serially timed during stages R1 to R6. II. Yield component responses. *Agron. J.* 77, 299-304.
- Kang, J. Y., Choi, H. I., Im, M. Y., Kim, S. Y. 2002. Arabidopsis basic leucine zipper proteins that mediate stress-responsive abscisic acid signaling. *Plant Cell.* 14, 343–357.
- Kausar, R., Hossain, Z., Makino, T., Komatsu, S. 2012. Characterization of ascorbate peroxidase in soybean under flooding and drought stresses. *Mol. Biol. Rep.* 39, 10573–10579.
- Khan, M. N., Komatsu, S. 2016. Proteomic analysis of soybean root including hypocotyl during recovery from drought stress. *Journal of Proteomics* 144, 39–50.
- Kirda, C., Danso, S. K. A., Zapata, F. 1989. Temporal water stress effects on nodulation, nitrogen accumulation and growth of soybean. *Plant and Soil* 120, 49–55.
- Kiyosue, T., Yoshiba, Y., Yamaguchi-Shinozaki, K., Shinozaki, K. 1996. A nuclear gene encoding mitochondrial proline dehydrogenase, an enzyme involved in proline metabolism, is up-regulated by proline but down-regulated by dehydration in Arabidopsis. *Plant Cell* 8, 23–35.
- Kokubun, M., Shimada, S., Takahashi, M. 2001. Flower Abortion Caused by Preanthesis Water Deficit is Not Attributed to Impairment of Pollen in Soybean. *Crop Science* 41, 1517-1521.
- Kokubun, M. 2011. Physiological Mechanisms Regulating Flower Abortion in Soybean, *Soybean - Biochemistry, Chemistry and Physiology*, Prof. Tzi-Bun Ng (Ed.), ISBN: 978-953-307-219-7, InTech, Available from: <http://www.intechopen.com/books/soybean-biochemistry-chemistry-and-physiology/physiologicalmechanisms-regulating-flower-abortion-in-soybean>.
- Kumar, V., Rani, A., Solanki, S., Hussain, S. M. 2006. Influence of growing environment on the biochemical composition and physical characteristics of soybean seed. *J. Food Compos. Anal.* 19, 188–195.
- Komatsu, S., Hossain, Z. 2013. Organ-specific proteome analysis for identification of abiotic stress response mechanism in crop. *Front. Plant Sci.* 4, 71.
- Ku, Y-S, Au-Yeung, W-K., Yung, Y-L., Li, M-W., Wen, C-Q., Liu, X., Lam, H-M. 2013. Drought Stress and Tolerance in Soybean, *A Comprehensive Survey of International Soybean Research - Genetics, Physiology, Agronomy and Nitrogen Relationships*, Prof. James Board (Ed.), InTech, DOI: 10.5772/52945. Available from: <http://www.intechopen.com/books/a-comprehensive-survey-of-international-soybean-research-genetics-physiology-agronomy-and-nitrogen-relationships/drought-stress-and-tolerance-in-soybean>.
- Lei, W., Tong, Z., Shengyan, D. 2006. Effect of drought and rewatering on photosynthetic physioecological characteristics of soybean. *Acta Ecologica Sinica* 26(7), 2073–2078.
- Levitt, J. 1980. Responses of plants to environmental stresses. Academic Press. New York and London. 697 pp.
- Li, D., Liu, H., Qiao, Y., Wang, Y., Cai, Z., Dong, B., Shi, Ch., Liu, Y., Li, X., Liu, M. 2013. Effects of elevated CO₂ on the growth, seed yield, and water use efficiency of soybean (*Glycine max* (L.) Merr.) under drought stress. *Agricultural Water Management* 129, 105–112.
- Liu, F., Andersen, M. N., Jensen, Ch. R. 2003. Loss of Pod Set Caused by Drought Stress Is Associated with Water Status and ABA Content of Reproductive Structures in Soybean. *Functional Plant Biology* 30, 271-280.
- Liu, F. Andersen, M. N., Jensen, Ch. R. 2004_b. Root signal controls pod growth in drought-stressed soybean during the critical, abortion-sensitive phase of pod development. *Field Crops Research* 85, 159–166.

- Liu, F., Andersen, M. N., Jacobsen, S-E., Jensen, Ch. R. 2005. Stomatal control and water use efficiency of soybean (*Glycine max* (L.) Merr.) during progressive soil drying. *Environmental and Experimental Botany* 54, 33–40.
- Liu, F., Jensen, Ch. R., Andersen, M. N. 2004_a. Pod Set Related to Photosynthetic Rate and Endogenous ABA in Soybeans Subjected to Different Water Regimes and Exogenous ABA and BA at Early Reproductive Stages. *Annals of Botany* 94, 405–411.
- Liu, F., Jensen, Ch. R., Andersen, M. N. 2004_c. Drought stress effect on carbohydrate concentration in soybean leaves and pods during early reproductive development: its implication in altering pod set. *Field Crops Research* 86, 1–13.
- Lozovaya, V. V., Lygin, A. V., Ulanov, A. V., Nelson, R. L., Dayde, J., Widholm, J. M. 2005. Effect of temperature and soil moisture status during seed development on soybean seed isoflavone concentration and composition. *Crop Sci.* 45, 1934–1940.
- Mahajan, S., Tuteja, N. 2005. Cold, salinity and drought stresses: An overview. *Arch. Biochem. Biophys.* 444, 139–158.
- Mak, M., Babla, M., Xu, S. C., O’Carrigan, A., Liu, X. H., Gong, Y. M., Holford, P., Chen, Z. H. 2014. Leaf mesophyll K⁺, H⁺ and Ca²⁺ fluxes are involved in drought-induced decrease in photosynthesis and stomatal closure in soybean. *Environmental and Experimental Botany* 98, 1– 12.
- Makbul, S., Saruhan Guler, N., Durmus, N., Guven, S. 2011. Changes in anatomical and physiological parameters of soybean under drought stress. *Turk. J. Bot.* 35, 369-377.
- Maleki, A., Naderi, A., Naseri, R., Fathi, A., Bahamin, S. Maleki, R. 2013. Physiological Performance of Soybean Cultivars under Drought Stress. *Bull. Env. Pharmacol. Life Sci.* 2(6), 38-44.
- Meckel, L., Egli, D. B., Philips, R. E., Radcliffe, D., Leggett, J. E. 1984. Effect of moisture stress on seed growth in soybeans. *Agron. J.* 76, 647–650.
- Medic, J., Atkinson, Hurburgh Jr, Ch. R. 2014. Current Knowledge in Soybean Composition. *J. Am. Oil Chem. Soc.* 91, 363–384.
- Mohammadi, P. P., Moieni, A., Hiraga, S., Komatsu, S. 2012. Organ-specific proteomic analysis of drought-stressed soybean seedlings. *J. Proteome.* 75, 1906–1923.
- Mutava, R. N., Prince, S. J. K., Syed, N. H., Song, L., Valliyodan, B., Chen, W., Nguyen, H., T. 2015. Understanding abiotic stress tolerance mechanisms in soybean: A comparative evaluation of soybean response to drought and flooding stress. *Plant Physiology and Biochemistry* 86, 109–120.
- Navari-Izzo, F., Vangioni, N., Quartacci, M. F. 1990. Lipids of soybean and sunflower seedlings grown under drought conditions. *Phytochemistry* 29(7), 2119-2123.
- Oh, M., Komatsu, S. 2015. Characterization of proteins in soybean roots under flooding and drought stresses. *Journal Of Proteomics* 114, 161–181.
- Pennypacker, B. W., Leath, K. L., Stout, W. L. Hill, R. R. 1990. Technique for simulating field drought stress in greenhouse. *Agron. J.* 82, 951-957.
- Read, D. J., Bartlett, E. M. 1972. The physiology of drought resistance in soybean plant (*Glycine max* (L.) Merr.), The relationship between drought resistance and growth. *J. Appl. Ecol.* 9, 487–489.
- Rose, I. 1988. Effects of moisture stress on the oil and protein components of soybean seeds. *Crop Pasture Sci.* 39, 163–170.

- Rotundo, J. L., Westgate, M. E., 2009. Meta-analysis of environmental effects on soybean seed composition. *Field Crop Res.* 110, 147–156.
- Rotundo, J. L., Westgate, M. E. 2010. Rate and duration of seed component accumulation in water-stressed soybean. *Crop Sci.* 50, 676–684.
- Sadeghipour, O., Abbasi, S. 2012. Soybean Response to Drought and Seed Inoculation. *World Applied Sciences Journal* 17(1), 55-60.
- Samarah, N. H., Mullen, R. E., Anderson, I. 2009. Soluble sugar contents, germination, and vigor of soybean seeds in response to drought stress. *Journal of New Seeds* 10(2), 63-73.
- Samarah, N. H., Mullen, R. E., Cianzio, S. R., Scott, P. 2006. Dehydrin-like proteins in soybean seeds in response to drought stress during seed filling. *Crop Sci.* 46, 2141-2150.
- Samet, J. S., Cortes, P. M., Sinclair, T. R. 1984. Diurnal and Seasonal Changes in Abscisic Acid of Field grown Soybeans Subjected to Drought Stress. *Field Crops Research* 8, 49-59.
- Saxena, N. P., Johansen, C., Saxena, M. C., Silim, S.N. 1993. Selection for drought and salinity tolerance in cool season food legumes. In: Singh KB, MC Saxena, eds. *Breeding for stress tolerance in cool season food legumes*. UK: Wiley, 245–270.
- Schulze, E. D. 1993. Soil water deficit and atmospheric humidity as environmental signals. In: Smith J A C, Griffiths H Eds. *Water Deficits*. Oxford: Bios Publisher, 129–145.
- Seki, M., Kamei, A., Yamaguchi-Shinozaki, K., Shinozaki, K. 2003. Molecular responses to drought, salinity and frost: Common and different paths for plant protection. *Curr. Opin. Biotech.* 14, 194–199.
- Shaw, B., Thomas, T. H., Cooke, D. T. 2002. Responses of sugar beet (*Beta vulgaris* L.) to drought and nutrient deficiency stress. *Plant Growth Regul.* 37, 77-83.
- Siddique, M. R. B., Hamid, A., Islam, M. S. 2000. Drought stress effects on water relations of wheat. *Bot. Bull. Acad. Sin.* 41, 35-39.
- Sinclair, T. R., Zimet, A. R., Muchow, R. C. 1988. Changes in soybean nodule number and dry weight in response to drought. *Field Crops Research* 18(2–3), 197-202.
- Sionit, N., Kramer, P. J. 1977. Effect of water stress during different stages of growth of soybeans, *Agronomy Journal* 69, 274-278.
- Smiciklas, K. D., Mullen, R. E., Carlson, R. E., Knapp, A. D. 1992. Soybean seed quality response to drought stress and pod position. *Agron. J.* 84, 166-170.
- Smith, D. L., Dijak, M., Hume, D. J. 1988. The effect of water deficit on N₂ (C₂H₂) fixation by white bean and soybean. *Canadian Journal of Plant Science* 68, 957–967.
- Specht, J., Chase, K., Markwell, J., Germann, M., Lark, K., Graef, G., Macrander, M., Orf, J., Chung, J. 2001. Soybean response to water. *Crop Sci.* 41, 493–509.
- Subbarao, G. V., Johansen, C., Slinkard, A. E., Nageswara, R. R. C., Saxena, N. P., Chauhan, Y. S. 1995. Strategies for improving drought resistance in grain legumes. *Critical Review of Plant Science* 14, 469–523.
- Turner, N.C. 1986. Adaptation to water deficits: a changing perspective. *Aust. J. Plant Physiol.* 13, 175–190.
- Wang, Y., Frei, M. 2011. Stressed food – The impact of abiotic environmental stresses on crop quality. *Agriculture, Ecosystems and Environment* 141, 271– 286.
- Wu, Y., Cosgrove, D. J. 2000. Adaptation of roots to low water potentials by changes in cell wall extensibility and cell wall proteins. *Journal of Experimental Botany* 51(350), 1543-1553.

- Yamaguchi-Shinozaki, K., Shinozaki, K. 2006. Transcriptional regulatory networks in cellular responses and tolerance to dehydration and cold stresses. *Annu. Rev. Plant Biol.*57, 781-803.
- Yang, P. H., Li, G. Q., Guo, L., Wu, S. J. 2003. Effect of drought stress on plasma membrane permeability of soybean varieties during flowering-podding stage. *Agricultural Research in the Arid Areas* 21(3), 127–130.
- Yordanov, I., Velikova, V., Tsonev, T. 2003. Plant response to drought and stress tolerance. *Bulg. J. Plant Physiol.* 187-206.
- Zablutowicz, R. M., Focht, D.D., Cannell, G. H. 1981. Nodulation and N fixation of field-grown California cowpeas as influenced by well-irrigated and droughted conditions. *Agronomy Journal* 73, 9–12.