

**Mesenchymal stromal-like cells set the balance of stimulatory and inhibitory signals in monocyte-derived dendritic cells**

Ildikó Bacskai<sup>a</sup>, Anett Mázló<sup>a</sup>, Katalin Kis-Tóth<sup>b</sup>, Attila Szabó<sup>a</sup>, György Panyi<sup>c</sup>, Balázs Sarkadi<sup>d</sup>, Ágota Apáti<sup>d#</sup>, Éva Rajnavölgyi<sup>a\*</sup>#

<sup>a</sup>Department of Immunology, University of Debrecen, Medical and Health Science Center, Debrecen, Hungary

<sup>b</sup>Department of Rheumatology, Beth Israel Deaconess Medical Center, Boston, MA, USA

<sup>c</sup>Department of Biophysics and Cell Biology, University of Debrecen, Faculty of Medicine, 98 Nagyerdei krt., Debrecen 4032, Hungary

<sup>d</sup>Institute of Enzymology, Research Centre for Natural Sciences, Hungarian Academy of Sciences, Budapest, Hungary

**Keywords:** mesenchymal stromal cell-like cell, dendritic cell, RIG-like receptors, immunosuppression, matrix metalloproteinases

# Ágota Apáti and Éva Rajnavölgyi contributed equally to this work

\*Corresponding Author: Éva Rajnavölgyi

Department of Immunology

Medical and Health Science Center

University of Debrecen

1 Egyetem sq Debrecen, H-4032 Hungary

Tel/Fax: +36 52 417 159

E-mail: evaraj@med.unideb.hu

## **ABSTRACT**

The major reservoir of human multipotent mesenchymal stem/stromal cells (MSC) is the bone marrow (BM) with the capability to control hematopoietic stem cell (HSC) development. The regenerative potential of MSC is associated with enhanced endogenous repair and healing mechanisms that modulate inflammatory responses. Our previous results revealed that MSC-like (MSCl) cells derived from pluripotent human embryonic stem cells resemble BM-derived MSC in morphology, phenotype and differentiating potential. Here we investigated the effects of MSCl cells on the phenotype and functions of dendritic cells (DC). To assess how anti-viral immune responses could be regulated by intracellular pattern recognition receptors (PRR) of DC in the presence of MSCl cells we activated DC with the specific ligands of retinoic acid-inducible gene I (RIG-I) helicases and found that activated DC co-cultured with MSCl cells exhibited reduced expression of CD1a and CD83 cell surface molecules serving as phenotypic indicators of DC differentiation and activation, respectively. However, RIG-I-mediated stimulation of DC via specific ligands in the presence of MSCl cells resulted in significantly higher expression of the co-stimulatory molecules CD80 and CD86 than in the presence of BM-MSC. In line with these results the concentration of IL-6, IL-10 and CXCL8 was increased in the supernatant of the DC-MSCl co-cultures, while the secretion of TNF- $\alpha$ , CXCL10, IL-12 and IFN $\gamma$  was reduced. Furthermore, the concerted action of mechanisms involved in the regulation of DC migration resulted in the blockade of cell migration indicating altered DC functionality mediated by MSCl cell-derived signals and mechanisms resulting in a suppressive microenvironment.

## INTRODUCTION

Mesenchymal stromal cells (MSC) developing from multipotent hematopoietic stem cells (HSC) are preferentially localized to the bone marrow [1], but can also be obtained from umbilical cord, adipose tissue, placenta, muscle, vein wall, peripheral blood [2] and corneal stroma[3]. The main functions of MSC involve the maintenance of the bone marrow niche via inhibiting the differentiation of HSC through direct cell-to-cell contacts [4] and the release of galectin-1, angiopoietin-1, osteopontin, thrombospondin-1 and 2 together with other factors to support tissue repair through regulating cell differentiation and immunomodulation [5]. Based on their wide spread physiological functions MSC were also considered in tissue engineering and the first successful therapy based on the intravenous infusion of haplo-identical MSC for treating graft vs host disease was published in 2004 [6-8]. The inhibitory effects of MSC on DC functional activities and antigen-specific T- and B-lymphocyte expansion through blocking G0/G1 transition to the S phase were also reported [9-12]. Recently, different approaches have been developed for using MSC in clinical trials [13] ([www.clinicaltrials.gov](http://www.clinicaltrials.gov)). However, the establishment and the production of reproducible, stable and well characterized human MSC has remained a challenge hampered by the heterogeneity of tissue-derived MSC [14]. Other studies focused to the phenotypic and functional characterization of MSC-like cell lines derived from different human tissues [15] and to confirm pluripotency of human ES cell-derived MSC by generating un-limited numbers of early passage MSC of consistent quality and immune suppressive properties [16]. In the present study we used a previously characterized human MSC1 cell line generated from pluripotent HUES9 embryonic stem cells. The phenotype, morphology and functional attributes of these cells have previously been described [17,18] and its stability up to 15 passages has been verified. They expressed the typical cell surface markers of MSC including

CD44, CD73, CD90, CD105, but were negative for the hematopoietic markers CD34, CD14, CD133, CD45 and the embryonic markers Oct4, Nanog, ABCG2, PODXL and SSEA4. They were able to differentiate to osteogenic, chondrogenic and adipogenic cell lineages with similar potency as BM-derived MSC [17]. MSC1 cells also had the capability to support the growth of undifferentiated human ESC via acting as feeder layer considered as an important MSC-associated property. Cells with potent self renewing capacity were also used as an *in vitro* model of human MSC to obtain high numbers of well characterized cells and for analyzing the immune suppressive potential and the mode of action in human primary monocyte-derived DC with both inflammatory and tolerogenic potential [19].

MSC are known to modulate immune responses via acting on multiple cell types such as DC, natural killer cells (NK), T- and B-lymphocytes [8,20] and were shown to suppress the differentiation, activation, migration and antigen presenting functions of conventional DC [20], a cell population essential for linking innate and adaptive immune mechanisms through presenting antigenic peptides to naive T-lymphocytes. This series of events can induce cell expansion and polarization of effector T-lymphocytes to Th1, Th2, Th17 and Treg directions [21]. It has also been demonstrated that BM-derived MSC can shift the differentiation of naive CD4<sup>+</sup> T-cells to an anti-inflammatory Th2 direction [22] underpinning the relevance of this type of regulation for utilization in clinical settings, exemplified by bone marrow transplantation, contact allergy and autoimmune disorders [23]. Conventional DC continuously binding and internalizing antigens use a wide spectrum of pattern recognition receptors (PRR). Upon sensing endogenous and exogenous danger signals DC become activated and migrate into the draining lymph nodes where they present their antigenic cargo to naïve T-lymphocytes. Retinoic acid-inducible gene I (RIG-I) and melanoma differentiation-associated protein 5 (MDA5) are cytoplasmic PRR expressed by a wide array of cell types and play essential roles in recognizing and eliminating viruses. DC activated by the specific

ligands (ssRNA, dsRNA or DNA) of the RIG-I receptors results in the secretion of pro-inflammatory cytokines and chemokines [24]. RIG-I is also expressed and is functional in MSC and has an important role in supporting cell survival indicating its possible contribution to the regulation of MSC functional activities [25]. Human MSC express several TLR, which are involved in the regulation of cell migration and the production of immune modulatory factors. The functional properties of human MSC was suggested to depend on the type of TLR as TLR4-primed MSC gave rise to inflammatory MSC1, while TLR3-induced MSC to suppressive MSC2 cell populations [26].

Encounter of conventional DC with naïve T-lymphocytes depends on the timely migration of activated DC to the draining lymph nodes [27,28] driven by the CCR7 chemokine receptor expressed by stimulated DC, as well as by the expression of the CCR7 chemokine receptor ligands CCL19 and/or CCL21 [28]. This receptor-ligand interaction up-regulates the expression of matrix metalloproteinases (MMP), responsible for the degradation and remodelling of the extracellular matrix upon cell migration [29], while the proteolytic activity of MMP is regulated by the tissue inhibitors of MMP enzymes (TIMP). Maintaining the balance between these opposing activities MMP and TIMP are crucial for preventing uncontrolled enzymatic degradation of the extracellular matrix known to contribute to the initiation of inflammation, autoimmune disorders and cancer metastasis [30].

In a previous work we have characterized the expression patterns of MMP enzymes and TIMP inhibitors in monocytes and monocyte-derived DC subpopulations [31] and discovered that the chemokine driven migration of activated DC is regulated by the expression of the voltage gated sodium channel Nav1.7 [32]. Based on these findings in this study we aimed to investigate the effects of MSC1 cells on DC activation triggered by ligand-specific RIG-I stimulation. Our results demonstrated for the first time that MSC1 cells are able to modify the activation status of DC, the secretion level of cytokines and chemokines as well as the

outcome of DC-mediated T-cell polarization suggesting its profound impact on the outcome of DC functionality. We also propose that MSC1 cells could be used as a feasible model for human *in vitro* studies by offering means for developing novel MSC-based strategies to design directed immune suppressive cellular therapies.

## **MATERIALS AND METHODS**

### *Generation of monocyte-derived dendritic cells*

Monocytes were separated from peripheral blood mononuclear cells (PBMC) of healthy blood donors drawn at the Regional Blood Center of the Hungarian National Blood Transfusion Service in accordance with the written approval of the Director of the National Blood Transfusion Service according to the directives of the European Union. PBMC were separated by Ficoll Pacque Plus (GE Healthcare Bio-Sciences AB, Uppsala, Sweden) gradient centrifugation followed by positive selection of CD14<sup>+</sup> monocytes by using anti-CD14 coated magnetic beads (Miltenyi Biotec, Bergisch Gladbach, Germany). Monocytes were plated at  $2 \times 10^6$  cell/ml concentration in RPMI (Hyclone, South Logan, Utah) supplemented by 10% FCS (Gibco, Paisley, Scotland) and 1% antibiotic/antimycotic solution (Hyclone, South Logan, Utah) in the presence of 100 ng/ml IL-4 and 75 ng/ml GM-CSF (Peprotech EC, London, UK) added on days 0 and 2.

### *Generation of mesenchymal stromal cell-like cells*

Mesenchymal stromal cell-like cells derived from the human embryonic stem cell lines HUES9 and HUES1 were kindly provided by Douglas Melton, HHMI. MSC1 cells were used according to the ethical permission 6681/2012/EHR. The cells were cultured on mitotically inactivated mouse embryonic fibroblast (MEF) to form embryonic bodies (EB) followed by trypsinization to obtain single cell cultures, which were further cultured on gelatin covered 10

cm plates in DMEM containing 10% FBS [18]. Cells of the confluent cultures exhibited fibroblast like morphology and were further characterized in a collaborative study with the laboratory of Balázs Sarkadi, Membrane Research Group of the Hungarian Academy of Sciences, Semmelweis University and National Blood Service, Budapest, Hungary [17]. MSC1 cell passages in opto-mechanical-treated polystyrene flasks (TPP, Trasadingen, Switzerland) provided coherent cell layers in the presence of L-glutamine, 10% FCS and 1% anti-mycoticum/anti-bioticum solution (Hyclone, South Logan, Utah) in low glucose DMEM. (Hyclone, South Logan, Utah). The cultured MSC1 cells were used after 10 passages.

#### *Co-cultures of DC and MSC1 cells*

MSC1 cells were collected by using 0.05 –0.02% Trypsin/EDTA solution in Dulbecco's PBS (DPBS), washed and cultured at a cell density of  $4 \times 10^5$  cells in opto-mechanical treated six well plates for optimal cell growth on a  $8.9 \text{ cm}^2/\text{well}$  area (TPP, Trasadingen, Switzerland). To achieve confluent cell layers the MSC1 cells were cultured in 2.5 ml RPMI supplemented with 10% FCS containing 1% antimycotic/antibiotic solution (Hyclone, South Logan, Utah) for minimum 6 hours.  $2 \times 10^6$  freshly isolated monocytes were placed directly on the top of the adherent MSC1 cells and the co-cultured DC were differentiated in the presence of GM-CSF and IL-4 at the same concentration as used for DC differentiation.

#### *Activation of DC by inflammatory stimuli*

On day 5 of *in vitro* DC differentiation the cells were activated by different inflammatory stimuli in the presence or absence of MSC1 cells. These involved inflammatory cocktail containing 10 ng/ml TNF- $\alpha$ , 5 ng/ml IL-1 $\beta$ , 75 ng/ml GM-CSF, 20 ng/ml IL-6 and 1  $\mu\text{g}/\text{ml}$  PGE<sub>2</sub> (Peprtech EC, London, UK), the RIG-I ligand poly(I:C) used at 25  $\mu\text{g}/\text{ml}$ , or with 1  $\mu\text{g}/\text{ml}$  5'ppp-dsRNA introduced by the Lyovect transfection reagent (Invivogen, San Diego,

CA, USA). 24h after activation DC expressing CD209/DC-SIGN on the cell surface were separated by magnetic beads (Miltenyi Biotec).

#### *Phenotypic characterization of DC differentiated in the presence or absence of MSC1 cells*

The phenotypic characterization of resting and activated DC was performed by flow cytometry on day 6 of *in vitro* cell differentiation and was compared to cells co-cultured with MSC1 cells (BD Biosciences, Franklin Lakes, NJ, USA), and the data were analyzed by the FlowJo software (Tree Star, Ashland, OR, USA). MSC1 cells were identified by the expression of CD105, whereas DC were captured by CD209/DC-SIGN (BioLegend, San Diego, CA, USA). To measure changes in the expression levels of the CD80, CD83, CD86 and CD1a cell surface markers fluorochrome-labeled antibodies were used along with the respective isotype matched control antibodies (BD Pharmingen, San Diego, CA, USA).

#### *RNA isolation, cDNA synthesis and quantitative RT-PCR*

Total RNA was isolated by using TRIzol reagent (MRC, Cincinnati, OH USA). 2 µg of RNA was reverse-transcribed at 37°C for 120 minutes using the High Capacity cDNA Archive Kit (Applied Biosystems, Foster City, CA) and oligo-(dT) primers (Promega, Madison, WI, USA). Quantitative real-time PCR was performed by using gene-specific TaqMan assays (Applied Biosystems, Foster City, CA, USA), DreamTaq DNA polymerase (Fermentas St. Leon-Rot, Germany) in a final volume of 12.5 µl and in ABI StepOnePlus real-time PCR instrument (Applied Biosystems). The housekeeping gene h36B4 was used for data normalization. Cycle thresholds were determined by using the StepOne Software v2.1 (Applied Biosystems).

#### *Cytokine measurements*



Concentration of the pro-inflammatory cytokines IL-1 $\beta$ , TNF- $\alpha$ , IL-6, the chemokines CXCL8 and CXCL10, and that of the T-cell polarizing cytokines IL-12, IL-10 and IFN $\gamma$  was measured in the supernatants of cell cultures by ELISA kits (BD Biosciences) following the manufacturer's instructions. Optical densities were determined by microplate reader (Biotek, Winooski, VT, USA).

#### *ELISPOT assay*

Activated DC previously co-cultured or not with MSC1 cells were cultured with allogeneic T-cells in serum-free RPMI medium for 3 days at 37°C. The number of IFN $\gamma$  secreting T-cells was detected by the avidin-HRP system (NatuTec, GmbH, Germany) and the results were analyzed by the ImmunoScan plate Reader (CTL, Shaker Heights OH, USA).

#### *Migration assays*

Migration of the differentiated DC co-cultured or not with MSC1 cells was tested in a Transwell system of 6.5 mm diameter and 5 $\mu$ m pore size (Corning Inc., Glendale, Arizona, USA). DCs were co-cultured with MSC1 cells for 6 days and were activated by PolyI:C or 5'ppp-RNA for 24h followed by the separation of DCs from the MSC1 cells by using CD209/DC-SIGN magnetic beads. 10<sup>6</sup> isolated DC were added to the upper chamber of the transwell plate and the migration of DC was measured in the presence or absence of 200ng/ml CCL19 chemokine placed to the lower chamber. After 24h incubation at 37° the migrated cells from the lower chamber were collected and counted by flow cytometry using poly-styrene beads as controls (Fluka St. Gallen, Switzerland).

#### *Statistical analysis*

Statistical analyses were performed by the one-way ANOVA with Bonferroni post-hoc test using the GraphPad Prism v.6 software (GraphPad Software Inc. La Jolla, CA, USA). Differences were considered to be statistically significant at \* $P < 0.05$ ; \*\* $P < 0.01$ ; \*\*\* $P < 0.005$ .

## RESULTS

### MSCI cells inhibit the differentiation and activation of monocyte-derived dendritic cells

Freshly isolated CD14<sup>+</sup> monocytes were cultured in the presence of GM-CSF and IL-4 for 5 days to generate competent DC used as control. Another fraction of DC was differentiated at similar conditions together with MSCI cells added at 5:1 monocyte to MSCI ratio present throughout the differentiation process. On day 5 both the control DC and the DC co-cultured with MSCI cells were activated with poly(I:C), 5'pppRNA or with an inflammatory cytokine cocktail containing GM-CSF, IL-1 $\beta$ , TNF- $\alpha$ , IL-6 and PGE2. On day 6 the activated DC were separated from the MSCI cells by the positive selection of CD209/DC-SIGN expressing cells and their proportion was measured in both the control and the MSCI-DC co-cultures. This procedure resulted in cell populations with  $>90 \pm 5\%$  DC in both the *in vitro* differentiated and the MSCI co-cultured cells indicating complete DC differentiation in both cases (data not shown). Cell surface expression of the CD1a (**Figure 1A**) and CD83 (**Figure 1B**) membrane proteins, used as indicators of DC activation, showed significant decrease when the DC were co-cultured with MSCI cells irrespective of the mode of activation induced by poly(I:C), 5'pppRNA or the inflammatory cocktail. Interestingly, stimulation of cells in the DC-MSCI co-cultures by the specific RIG-I ligand 5'pppRNA resulted in significant up-regulation of the co-stimulatory molecules CD80 (**Figure 1C**) and CD86 (**Figure 1D**) on the DC surface as compared to DC cultured without MSCI cells, whereas the presence of MSCI cells did not affect the expression levels of these markers in DC stimulated by the inflammatory cocktail or

by poly(I:C) (**Figure 1 C, D**). These results indicated the potential of MSC1 cells to modulate DC activation and consequently its other functional activities.

### **MSC1 cells modulate the cytokine secretion of activated DC**

To further analyse the functional cross-talk of DC and MSC1 cells the specific RIG-I ligands poly(I:C) and 5'pppRNA were added to the DC-MSC1 co-cultures and the secretion levels of the pro-inflammatory cytokines IL-6 and TNF- $\alpha$ , the chemokines CXCL8 and CXCL10, as well as the T-lymphocyte polarizing cytokines IL-12 and IL-10 were measured in the supernatants of the activated DC, MSC1 cells and DC-MSC1 co-cultures. We found that both RIG-I ligands could up-regulate the secretion of IL-6 (**Figure 2A**), CXCL8 (**Figure 2C**) and IL-10 (**Figure 2E**) in the DC-MSC1 co-cultures as compared to activated DC and MSC1 cells cultured separately. Although IL-10 was present in the supernatants of both resting and activated DC at low but measurable levels, we could not detect IL-10 in the supernatants of activated MSC1 cells (**Figure 2E**). However, the enhanced level of IL-10 in the supernatant of the co-cultured and stimulated cells suggest that IL-10 could contribute to the anti-inflammatory effects of MSC1 cells presumably via inhibiting the secretion of pro-inflammatory cytokines.

Our results also revealed that the concomitant secretion of TNF- $\alpha$  (**Figure 2B**) and CXCL10 (**Figure 2D**) did not induce a synergistic inflammatory effect but could efficiently be inhibited by MSC1 cells indicating their potent anti-inflammatory activity. In this experimental system the production of IL-12, a key Th1 polarizing cytokine was also dramatically decreased in the presence of MSC1 cells (**Figure 2F**) likely owing to its potential to induce clonal expansion and differentiation of IFN $\gamma$ -producing CD4<sup>+</sup> T-lymphocytes. Moreover, IL-12 was able to stimulate the production of other inflammatory cytokines such as TNF- $\alpha$ , IFN $\gamma$  and the interferon-induced chemokine CXCL10 to multiply the inflammatory response.

In line with these results the production of IL-6 (**Figure 2A**) and CXCL8 (**Figure 2C**), both acting as important mediators of tissue repair and angiogenesis, were detected in the supernatants of DC co-cultured with MSC1 cells at increased levels suggesting that they may contribute to create a local anti-inflammatory milieu in response to inflammatory signals. These observations also imply that beside direct cell-to-cell contacts cytokines and chemokines could also contribute to the regulatory functions of MSC1 cells.

### **MSC1 cells interfere with dendritic cell-mediated allogeneic T-lymphocyte activation and polarization**

To further analyse the outcome of DC-mediated functional activities in the presence of MSC1 cells we sought to measure the secretion of IFN $\gamma$  directly by ELISA and also as a result of DC-mediated allogeneic T-lymphocyte polarization by using ELISPOT assays. Our results demonstrated that IFN $\gamma$  secretion of DC could be induced by both poly(I:C) and 5'pppRNA, however the presence of MSC1 cells decreased its secretion significantly (**Figure 3A**). The potential of poly(I:C) or 5'pppRNA-stimulated DC to drive allogeneic T-lymphocyte polarization confirmed that MSC1 cells are able to interfere with this translational event through exerting potent anti-inflammatory effects on DC activation (**Figure 3B**). These results altogether demonstrate that DC, upon interacting with MSC1 cells, can create an anti-inflammatory local environment.

### **The effect of MSC1 cells on DC migration**

Previous results demonstrated that efficient presentation of peptide antigens to naive T-lymphocytes critically depends on the migration of tissue resident DC to the draining lymph nodes [33-35]. As a novel finding our previous studies demonstrated that the low expression level of the Nav1.7 ion channel, in combination with the high expression of CCR7, is the pre-

requisite of DC migration from the site of inflammation to the lymph nodes [32]. Based on this information we used a transwell system to study the effects of MSC1 cells on the migratory potential of DC in the presence or absence of MSC1 cells. In this experimental setting the upper chamber contained resting DC or DC stimulated either with poly(I:C) or the inflammatory cytokine cocktail in the presence or absence of MSC1 cells. The lower chamber was filled with fresh medium containing 0.5% BSA and the chemokine CCL19/MIP-3 $\beta$ , a strong inducer of activated DC migration guided by the membrane bound chemokine receptor CCR7. The upper and lower chambers were connected by a membrane of 5  $\mu$ m pore size to allow chemokine driven migration of DC towards CCL19 gradients. The migrated cells were collected from the lower chamber and after 24h the number of migrating cells was counted by flow cytometry. In correlation with the level of CCR7 chemokine expression in the presence of MSC1 cells a statistically significant decrease in the number of migratory cells was observed (**Figure 4A, B**).

Considering that the expression of matrix metalloproteinase enzymes (MMP) and their specific inhibitors (TIMP) play important roles in regulating DC migration, we also assessed the contribution of these enzymes to the regulation of MSC1 cell-mediated inhibitory functions. Expression of the MMP9, MMP12, TIMP1 and TIMP2 genes, which could be induced by the interaction of the CCR7 receptor with its specific ligand CCL19, was monitored in both resting and activated DC in the presence or absence of MSC1 cells. We showed that the expression of MMP9 was up-regulated (**Figure 4C**), whereas that of MMP12 was down regulated (**Figure 4D**). The relative gene expression levels of TIMP1 and TIMP2 were higher in CD1a<sup>-</sup> DC than in its inflammatory CD1a<sup>+</sup> counterpart [32] and could be associated with a DC phenotype similar to cells co-cultured with MSC1 cells (**Figure 4E, F**).

Another regulatory mechanism that might be involved in regulating MSC1 cell-related functions could be associated with the activity of the Nav1.7 ion channel. We have previously

shown that functionally active Nav1.7 is predominantly expressed in resting DC but its expression is decreased in parallel with DC activation thus providing a sensitive tool for correlating the state of DC activation to Nav1.7 channel activity [32]. When the expression of Nav1.7 was measured in activated DC without MSC1 cells, the level of Nav1.7 expression decreased, even when the cells were stimulated by the inflammatory cocktail or by poly (I:C), but it was increased in DC suppressed by MSC1 cells (**Figure 4G**). These changes are in good accordance with the low/undetectable numbers of inflammatory CD1a<sup>+</sup> DC detected in these cultures as compared to the tolerogenic CD1a<sup>-</sup> DC subset [36,37].

### **MSC1 cells inhibit the expression of RIG-like receptor family members**

Our results summarized in Figure 1 and 2 show that MSC1 cells are potent inhibitors of DC functionality induced by the specific ligands poly(I:C) and 5'pppRNA of the cytosolic RIG-like receptors RIG-I, MDA5 and LGP-2 recognizing dsRNA, ssRNA or DNA [24]. Taking a step further we hypothesized that co-culturing DC with MSC1 cells might have a direct inhibitory effect on the expression of intracellular RIG-I receptors. Indeed, our Q-PCR results revealed that mRNA expression of the RIG-I, MDA-5 and LGP-2 receptors belonging to the RLR family of cytosolic pattern recognition receptors is up-regulated upon stimulation by specific ligands but in the presence of MSC1 cells this induction does not occur likely due to the inhibited expression of these receptors (**Figure 5A-C**) accompanied by the decreased production of IFN $\beta$  (**Figure 5D**). Based on these data we conclude that in the presence of MSC1 cells the specific ligands of RIG-I could not be recognized by DC due to the down modulation of RLR receptor expression by MSC1 cells.

## **DISCUSSION**

The unique capability of MSC of different origin is to modulate the outcome of allogeneic bone marrow transplantation that has been pioneered by Le Blanc and her group in humans and was confirmed in several mice models [9,38]. These results also revealed the potential of MSC to exert multiple effects on other cell types through regulatory, anti-inflammatory and by stander effects and via targeting injured tissues as described in several diseases, such as grade IV severe acute graft versus host disease (GVHD) of the gut and liver [6] and steroid resistant severe acute GVHD [39]. The utility of MSC also involves the prevention of transplanted skin graft rejection [40], treatment of osteogenesis imperfecta by mismatched allogeneic liver-derived MSC in immune competent fetus [41], autoimmune encephalomyelitis [42], diabetes mellitus [43] and collagen-induced arthritis [44]. Further studies also demonstrated that the anti-inflammatory cytokine environment created by the transplanted MSC at the site of inflammation was able to improve the outcome of acute renal, neural and lung injury [42,45,46]. Thus in the past years MSC became clinically important cell types due to their regenerative potential that can be utilized in cell therapies aimed to treat inflammatory and autoimmune disorders or apply them for tissue engineering. This approach is also supported by the unique capability of MSC to bypass MHC compatibility for inducing immune suppression [47]. It is well established that MSC1 cells can be stimulated by TLRs [26] and upon stimulation they migrate to damaged tissues driven by cytokines, chemokines and secreted extracellular matrix (ECM) proteins [48]. Beside the secretion of soluble factors the direct contact of MSC with immune cells seems to be essential for creating a supportive local milieu [49]. However, application of these cells for therapeutic interventions is limited due to the low number of MSC available. To overcome this restriction several groups explored new strategies for identifying cell lines with immune suppressive properties [50]. In a previous collaborative study we described the phenotypic, some functional as well as the differentiating capability of a cell line of embryonic origin with MSC-like phenotype and

compared its functional activities with BM-MSC and a human foreskin fibroblast cell line (HFF) [17]. This analysis demonstrated similar morphology, phenotype and functional attributes resembling BM-derived MSC and offered us to exploit the further functional characterization of these cells in terms of their immune suppressive potential. The present study provided evidence for the potent immune suppressive nature of the MSC1 cell line characterized by its profound effects on multiple functional properties of resting and activated monocyte-derived DC. In a recent report the immune suppressive effects of human ESC-derived MA07 cells on CD83 expression and IL-12p70 secretion by IFN $\gamma$  and LPS activated DC was observed that was associated with enhanced IL-2 induced expansion of regulatory T-cells [16].

Our results performed with human primary monocyte-derived DC also revealed that MSC1 cells could affect the maturation, activation and a wide array of functional activities in a similar manner as human BM-MSC. As a consequence, the phenotype of DC was modified in the presence of MSC1 cells resulting in decreased expression of the CD1a and CD83 activation markers on the surface of activated DC induced by an inflammatory cytokine cocktail or by specific ligands of the RIG-I receptors. A similar effect was described in LPS-induced activation of mice in the presence of BM-MSCs [51] and TNF- $\alpha$  induced stimulation of umbilical cord blood or bone marrow derived MSC [52]. In contrast, the expression of the CD80 and CD86 co-stimulatory molecules was up-regulated in the human DC-MSC co-cultures, while in mice the expression of these molecules was decreased in the presence of both murine and human BM-MSC [53,54] indicating species related differences in the inhibitory function of MSC.

The secretion of CXCL8 and IL-6 by MSC1 and CXCL10 by MSC2 cells suggested the induction of different cytokine combinations induced by these MSC subsets [26] similar to our results showing increased secretion of IL-6, CXCL8 and CXCL10 by MSC1 cells activated



by poly(I:C). However, no detectable changes could be observed when the RIG-I specific 5'pppRNA ligand was used for DC activation. In contrast, the activation of co-cultured DC and MSC resulted in dramatic changes in the secretion of cytokines and chemokines. Even though changes in cytokine secretion by DC have been demonstrated in the presence of MSC of various origins the mechanistic background of this regulation has not been explored.

It was also shown that MSC, isolated from mouse embryonic fibroblasts could induce the generation of IL-10 dependent regulatory DC via SOCS3 activation leading to increased secretion of anti-inflammatory IL-10 [55]. The cytokines IL-6 and CXCL8 have been considered as important mediators of tissue repair and angiogenesis. In LPS stimulated DC the presence of MSC increased the secretion of IL-6 and CXCL8 [56,57] and in response to poly(I:C) or 5'pppRNA the increased secretion of IL-6, IL-10 and CXCL8 was also detected opposing the significantly decreased levels of IL-12 and TNF- $\alpha$  in the supernatant of the DC-MSCI co-cultures. These results suggest that the contact and/or the communication of cells and soluble factors may trigger anti-inflammatory and/or healing mechanisms that shift the balance of the immune response to tissue regeneration and changes in cytokine and chemokine secretion.

The CCL19 chemokine is expressed in the thymus and lymph nodes and its secretion is essential for the migration of CCR7<sup>high</sup> DC to the draining lymph nodes where they interact with naive T-lymphocytes [33]. However, the expression of CCR7 and the migration of DC derived from the DC-MSCI co-cultures exhibited significantly decreased migration as compared to DC cultured in the absence of MSCI cells. One of the mechanisms involved in this complex regulation was discovered recently [32] showing the contribution of the voltage gated membrane channel Nav1.7, which regulates intracellular Ca<sup>2+</sup> concentration in DC and also acts as a master regulator of the cell cycle. High intracellular Ca<sup>2+</sup> concentration was shown to be essential for DC activation and was associated with down regulated Nav1.7

expression. As anticipated, the expression of the voltage gated sodium channel Nav1.7 was also dramatically down modulated and could be used as a sensitive indicator of the DC activation state [32]. On the contrary, Nav1.7 expression was maintained in the presence of MSC1 cells, which may indicate either dysregulation of the Ca<sup>2+</sup> signal accompanying DC activation or an alternative pathway for the regulation of Nav1.7 expression. Either way, MSC1 cells interfere with DC activation thereby identifying this sodium channel as a novel marker of suppressed DC. Overall, these data indicate that in the presence of MSC1 cells the generation of DC results in 'semi-activated', Nav1.7<sup>high</sup> and CCR7<sup>low</sup> cells, which express high amounts of active MMP9 enzyme and also TIMP with inhibitory potential to induce dramatically reduced migratory potential of DCs.

Up-regulation of CXCL8 chemokine was shown to support the expression of MMP-2 and MMP-9 in trophoblast cells [58]. The balance of MMP and TIMP is also known to have an impact on cell migration, homeostasis, survival and other functional activities of immune cells via regulating the cleavage of extracellular matrix components (ECM) to generate soluble cytokines, chemokines and growth factors together with their matching receptors [59]. Considering that the regulation of the MMP-TIMP axis is also involved in the triggering of signal transduction pathways connected to DC functions, we provoked DC migration with an inflammatory cocktail or with poly(I:C) and could increase the expression level of MMP-9 and MMP-12 in DC accompanied by decreased TIMP-1 and TIMP-2 expression showing a tight control of the inflammatory response [31]. However, in the presence of MSC1 cells, the expression level of TIMP was dramatically up-regulated and resulted in inhibited DC migration via blocking ECM degradation. More importantly, these effects could also modify the regulatory capacity of MMP by the secretion and proteolytic cleavage of cytokines and chemokines. Based on these results we conclude that MSC1 cells are able to modulate the expression of key molecules involved in DC migration. Moreover, the presence of MSC1 cells

may also have an impact on the level and activity of secreted cytokines and chemokines, and can affect the expression of their receptors for fine tuning DC activities in the actual inflammatory environment.

*In vivo* studies have revealed the inhibition of antigen-specific effector T-cell functions in the course of mouse allograft rejection [40], in graft versus host disease [60], autoimmune encephalomyelitis [42] and collagen induced arthritis [44]. In a murine model DC conditioned by allogeneic MSC could inhibit MHC class I and II expression and antigen presentation by DC while decreased CD69 expression on CD8<sup>+</sup> T-cells. In a human system, the presence of activated DC co-cultured with BM-MSc was shown to decrease the secretion of IFN $\gamma$  by T-cells [61]. In the present study, the inhibited secretion of IFN $\gamma$  could be detected by a sensitive ELISPOT assay. In this experimental setting human allogeneic T-lymphocytes, primed by activated DC and co-cultured with MSC1 cells were able to down regulate T-cell polarization to the Th1 direction to a similar extent as DC co-cultured with BM-MSc. In our human *in vitro* studies MSC also could alter DC functions significantly, and together with our previous studies confirmed the potential of MSC1 cells to exert inhibitory signals on antigen-specific T-cell responses.

In this context however, the expression of RIG-like receptors has not been investigated. We hypothesized that the functional changes induced by the RLR ligand poly(I:C) are the consequence of decreased RLR expression. Our results indeed demonstrated that in the presence of MSC1 cells the expression of RIG-I, MDA-5 and LGP-2 in DC was dramatically impaired. Referring to the previously described partially activated DC with tolerogenic and immune suppressive properties [62] MSC1 cells may assist the maintenance of DC in a 'semi-mature' suppressive state [63]. Our results obtained in a human system suggest that MSC1 cells with unlimited proliferating capacity could be harnessed as a reliable model for inhibiting DC activation by MSC-mediated immune suppression.

## **ACKNOWLEDGEMENT**

This research work was supported by the Hungarian Scientific Research Grants OTKA NK 101538, the European Union and the State of Hungary, co-financed by the European Social Fund in the framework of the TÁMOP 4.2.4. A/2-11-1-2012-0001 ‘National Excellence Program’ and the TÁMOP 4.2.2.A-11/1/KONV-2012-0023 ‘DEFENSE-NET’ project.

## **DISCLOSURE STATEMENT**

The authors declare no competing financial interest.

## REFERENCES

1. Beyer Nardi N and da Silva Meirelles L (2006) Mesenchymal stem cells: isolation, in vitro expansion and characterization. *Handb Exp Pharmacol*: 249-282.
2. Kern S, Eichler H, Stoeve J, Kluter H and Bieback K (2006) Comparative analysis of mesenchymal stem cells from bone marrow, umbilical cord blood, or adipose tissue. *Stem Cells* 24: 1294-1301.
3. Branch MJ, Hashmani K, Dhillon P, Jones DR, Dua HS and Hopkinson A (2012) Mesenchymal stem cells in the human corneal limbal stroma. *Invest Ophthalmol Vis Sci* 53: 5109-5116.
4. Friedenstein AJ, Chailakhyan RK, Latsinik NV, Panasyuk AF and Keiliss-Borok IV (1974) Stromal cells responsible for transferring the microenvironment of the hemopoietic tissues. Cloning in vitro and retransplantation in vivo. *Transplantation* 17: 331-340.
5. Pedemonte E, Benvenuto F, Casazza S, Mancardi G, Oksenberg JR, Uccelli A and Baranzini SE (2007) The molecular signature of therapeutic mesenchymal stem cells exposes the architecture of the hematopoietic stem cell niche synapse. *BMC Genomics* 8: 65.
6. Le Blanc K, Rasmusson I, Sundberg B, Gotherstrom C, Hassan M, Uzunel M and Ringden O (2004) Treatment of severe acute graft-versus-host disease with third party haploidentical mesenchymal stem cells. *Lancet* 363: 1439-1441.
7. Le Blanc K and Mougiakakos D (2012) Multipotent mesenchymal stromal cells and the innate immune system. *Nat Rev Immunol* 12: 383-396.
8. English K (2013) Mechanisms of mesenchymal stromal cell immunomodulation. *Immunol Cell Biol* 91: 19-26.
9. Stagg J and Galipeau J (2013) Mechanisms of immune modulation by mesenchymal stromal cells and clinical translation. *Curr Mol Med* 13: 856-867.
10. Glennie S, Soeiro I, Dyson PJ, Lam EW and Dazzi F (2005) Bone marrow mesenchymal stem cells induce division arrest anergy of activated T cells. *Blood* 105: 2821-2827.
11. Corcione A, Benvenuto F, Ferretti E, Giunti D, Cappiello V, Cazzanti F, Risso M, Gualandi F, Mancardi GL, Pistoia V and Uccelli A (2006) Human mesenchymal stem cells modulate B-cell functions. *Blood* 107: 367-372.
12. Ramasamy R, Fazekasova H, Lam EW, Soeiro I, Lombardi G and Dazzi F (2007) Mesenchymal stem cells inhibit dendritic cell differentiation and function by preventing entry into the cell cycle. *Transplantation* 83: 71-76.
13. Philippe B, Luc S, Valerie PB, Jerome R, Alessandra BR and Louis C (2010) Culture and Use of Mesenchymal Stromal Cells in Phase I and II Clinical Trials. *Stem Cells Int* 2010: 503593.
14. Siddappa R, Licht R, van Blitterswijk C and de Boer J (2007) Donor variation and loss of multipotency during in vitro expansion of human mesenchymal stem cells for bone tissue engineering. *J Orthop Res* 25: 1029-1041.
15. Lian Q, Lye E, Suan Yeo K, Khia Way Tan E, Salto-Tellez M, Liu TM, Palanisamy N, El Oakley RM, Lee EH, Lim B and Lim SK (2007) Derivation of clinically compliant MSCs from CD105+, CD24-differentiated human ESCs. *Stem Cells* 25: 425-436.
16. Kimbrel EA, Kouris NA, Yavanian GJ, Chu J, Qin Y, Chan A, Singh RP, McCurdy D, Gordon L, Levinson RD and Lanza R (2014) Mesenchymal stem cell population derived from human pluripotent stem cells displays potent immunomodulatory and therapeutic properties. *Stem Cells Dev* 23: 1611-1624.
17. Varga N, Vereb Z, Rajnavolgyi E, Nemet K, Uher F, Sarkadi B and Apati A (2011) Mesenchymal stem cell like (MSCl) cells generated from human embryonic stem cells support pluripotent cell growth. *Biochem Biophys Res Commun* 414: 474-480.
18. Apati A, Orban TI, Varga N, Nemeth A, Schamberger A, Krizsik V, Erdelyi-Belle B, Homolya L, Varady G, Padanyi R, Karaszi E, Kemna EW, Nemet K and Sarkadi B (2008) High level functional expression of the ABCG2 multidrug transporter in undifferentiated human embryonic stem cells. *Biochim Biophys Acta* 1778: 2700-2709.

19. Liu YP and Hematti P (2009) Generation of mesenchymal stromal cells from HOXB4-expressing human embryonic stem cells. *Cytotherapy* 11: 716-725.
20. Machado Cde V, Telles PD and Nascimento IL (2013) Immunological characteristics of mesenchymal stem cells. *Rev Bras Hematol Hemoter* 35: 62-67.
21. Billingham RE (1948) Dendritic cells. *J Anat* 82: 93-109.
22. Jitschin R, Mougiakakos D, Von Bahr L, Volkl S, Moll G, Ringden O, Kiessling R, Linder S and Le Blanc K (2013) Alterations in the cellular immune compartment of patients treated with third-party mesenchymal stromal cells following allogeneic hematopoietic stem cell transplantation. *Stem Cells* 31: 1715-1725.
23. Wang Q, Sun B, Wang D, Ji Y, Kong Q, Wang G, Wang J, Zhao W, Jin L and Li H (2008) Murine bone marrow mesenchymal stem cells cause mature dendritic cells to promote T-cell tolerance. *Scand J Immunol* 68: 607-615.
24. Szabo A, Bene K, Gogolak P, Rethi B, Lanyi A, Jankovich I, Dezso B and Rajnavolgyi E (2012) RLR-mediated production of interferon-beta by a human dendritic cell subset and its role in virus-specific immunity. *J Leukoc Biol* 92: 159-169.
25. Yang K, Wang J, Xiang AP, Zhan X, Wang Y, Wu M and Huang X (2013) Functional RIG-I-like receptors control the survival of mesenchymal stem cells. *Cell Death Dis* 4: e967.
26. Waterman RS, Tomchuck SL, Henkle SL and Betancourt AM (2010) A new mesenchymal stem cell (MSC) paradigm: polarization into a pro-inflammatory MSC1 or an Immunosuppressive MSC2 phenotype. *PLoS One* 5: e10088.
27. Sallusto F and Lanzavecchia A (1994) Efficient presentation of soluble antigen by cultured human dendritic cells is maintained by granulocyte/macrophage colony-stimulating factor plus interleukin 4 and downregulated by tumor necrosis factor alpha. *J Exp Med* 179: 1109-1118.
28. Steinman RM, Pack M and Inaba K (1997) Dendritic cell development and maturation. *Adv Exp Med Biol* 417: 1-6.
29. Redondo-Munoz J, Jose Terol M, Garcia-Marco JA and Garcia-Pardo A (2008) Matrix metalloproteinase-9 is up-regulated by CCL21/CCR7 interaction via extracellular signal-regulated kinase-1/2 signaling and is involved in CCL21-driven B-cell chronic lymphocytic leukemia cell invasion and migration. *Blood* 111: 383-386.
30. Werb Z (1997) ECM and cell surface proteolysis: regulating cellular ecology. *Cell* 91: 439-442.
31. Kis-Toth K, Bacskai I, Gogolak P, Mazlo A, Szatmari I and Rajnavolgyi E (2013) Monocyte-derived dendritic cell subpopulations use different types of matrix metalloproteinases inhibited by GM6001. *Immunobiology* 218: 1361-1369.
32. Kis-Toth K, Hajdu P, Bacskai I, Szilagyi O, Papp F, Szanto A, Posta E, Gogolak P, Panyi G and Rajnavolgyi E (2011) Voltage-gated sodium channel Nav1.7 maintains the membrane potential and regulates the activation and chemokine-induced migration of a monocyte-derived dendritic cell subset. *J Immunol* 187: 1273-1280.
33. Sallusto F, Schaerli P, Loetscher P, Schaniel C, Lenig D, Mackay CR, Qin S and Lanzavecchia A (1998) Rapid and coordinated switch in chemokine receptor expression during dendritic cell maturation. *Eur J Immunol* 28: 2760-2769.
34. Kellermann SA, Hudak S, Oldham ER, Liu YJ and McEvoy LM (1999) The CC chemokine receptor-7 ligands 6CKine and macrophage inflammatory protein-3 beta are potent chemoattractants for in vitro- and in vivo-derived dendritic cells. *J Immunol* 162: 3859-3864.
35. Sanchez-Sanchez N, Riol-Blanco L, de la Rosa G, Puig-Kroger A, Garcia-Bordas J, Martin D, Longo N, Cuadrado A, Cabanas C, Corbi AL, Sanchez-Mateos P and Rodriguez-Fernandez JL (2004) Chemokine receptor CCR7 induces intracellular signaling that inhibits apoptosis of mature dendritic cells. *Blood* 104: 619-625.
36. Szatmari I, Gogolak P, Im JS, Dezso B, Rajnavolgyi E and Nagy L (2004) Activation of PPARgamma specifies a dendritic cell subtype capable of enhanced induction of iNKT cell expansion. *Immunity* 21: 95-106.

37. Gogolak P, Rethi B, Szatmari I, Lanyi A, Dezso B, Nagy L and Rajnavolgyi E (2007) Differentiation of CD1a- and CD1a+ monocyte-derived dendritic cells is biased by lipid environment and PPARgamma. *Blood* 109: 643-652.
38. Uccelli A, Moretta L and Pistoia V (2008) Mesenchymal stem cells in health and disease. *Nat Rev Immunol* 8: 726-736.
39. Le Blanc K, Frassoni F, Ball L, Locatelli F, Roelofs H, Lewis I, Lanino E, Sundberg B, Bernardo ME, Remberger M, Dini G, Egeler RM, Bacigalupo A, Fibbe W, Ringden O, Developmental Committee of the European Group for B and Marrow T (2008) Mesenchymal stem cells for treatment of steroid-resistant, severe, acute graft-versus-host disease: a phase II study. *Lancet* 371: 1579-1586.
40. Bartholomew A, Sturgeon C, Siatskas M, Ferrer K, McIntosh K, Patil S, Hardy W, Devine S, Ucker D, Deans R, Moseley A and Hoffman R (2002) Mesenchymal stem cells suppress lymphocyte proliferation in vitro and prolong skin graft survival in vivo. *Exp Hematol* 30: 42-48.
41. Le Blanc K, Gotherstrom C, Ringden O, Hassan M, McMahon R, Horwitz E, Anneren G, Axelsson O, Nunn J, Ewald U, Norden-Lindeberg S, Jansson M, Dalton A, Astrom E and Westgren M (2005) Fetal mesenchymal stem-cell engraftment in bone after in utero transplantation in a patient with severe osteogenesis imperfecta. *Transplantation* 79: 1607-1614.
42. Zappia E, Casazza S, Pedemonte E, Benvenuto F, Bonanni I, Gerdoni E, Giunti D, Ceravolo A, Cazzanti F, Frassoni F, Mancardi G and Uccelli A (2005) Mesenchymal stem cells ameliorate experimental autoimmune encephalomyelitis inducing T-cell anergy. *Blood* 106: 1755-1761.
43. Dong QY, Chen L, Gao GQ, Wang L, Song J, Chen B, Xu YX and Sun L (2008) Allogeneic diabetic mesenchymal stem cells transplantation in streptozotocin-induced diabetic rat. *Clin Invest Med* 31: E328-337.
44. Augello A, Tasso R, Negrini SM, Cancedda R and Pennesi G (2007) Cell therapy using allogeneic bone marrow mesenchymal stem cells prevents tissue damage in collagen-induced arthritis. *Arthritis Rheum* 56: 1175-1186.
45. Tsuda H, Yamahara K, Otani K, Okumi M, Yazawa K, Kaimori JY, Taguchi A, Kangawa K, Ikeda T, Takahara S and Isaka Y (2014) Transplantation of allogenic fetal membrane-derived mesenchymal stem cells protects against ischemia/reperfusion-induced acute kidney injury. *Cell Transplant* 23: 889-899.
46. Zhao F, Zhang YF, Liu YG, Zhou JJ, Li ZK, Wu CG and Qi HW (2008) Therapeutic effects of bone marrow-derived mesenchymal stem cells engraftment on bleomycin-induced lung injury in rats. *Transplant Proc* 40: 1700-1705.
47. Le Blanc K, Tammik L, Sundberg B, Haynesworth SE and Ringden O (2003) Mesenchymal stem cells inhibit and stimulate mixed lymphocyte cultures and mitogenic responses independently of the major histocompatibility complex. *Scand J Immunol* 57: 11-20.
48. Aggarwal S and Pittenger MF (2005) Human mesenchymal stem cells modulate allogeneic immune cell responses. *Blood* 105: 1815-1822.
49. Gur-Wahnon D, Borovsky Z, Beyth S, Liebergall M and Rachmilewitz J (2007) Contact-dependent induction of regulatory antigen-presenting cells by human mesenchymal stem cells is mediated via STAT3 signaling. *Exp Hematol* 35: 426-433.
50. Sykova E and Frostyakov S (2013) Stem cells in regenerative medicine. *Laser Ther* 22: 87-92.
51. Spaggiari GM, Abdelrazik H, Becchetti F and Moretta L (2009) MSCs inhibit monocyte-derived DC maturation and function by selectively interfering with the generation of immature DCs: central role of MSC-derived prostaglandin E2. *Blood* 113: 6576-6583.
52. Saeidi M, Masoud A, Shakiba Y, Hadjati J, Mohyeddin Bonab M, Nicknam MH, Latifpour M and Nikbin B (2013) Immunomodulatory effects of human umbilical cord Wharton's jelly-derived mesenchymal stem cells on differentiation, maturation and endocytosis of monocyte-derived dendritic cells. *Iran J Allergy Asthma Immunol* 12: 37-49.
53. Zhang B, Liu R, Shi D, Liu X, Chen Y, Dou X, Zhu X, Lu C, Liang W, Liao L, Zenke M and Zhao RC (2009) Mesenchymal stem cells induce mature dendritic cells into a novel Jagged-2-dependent regulatory dendritic cell population. *Blood* 113: 46-57.

54. Djouad F, Charbonnier LM, Bouffi C, Louis-Pence P, Bony C, Apparailly F, Cantos C, Jorgensen C and Noel D (2007) Mesenchymal stem cells inhibit the differentiation of dendritic cells through an interleukin-6-dependent mechanism. *Stem Cells* 25: 2025-2032.
55. Liu X, Qu X, Chen Y, Liao L, Cheng K, Shao C, Zenke M, Keating A and Zhao RC (2012) Mesenchymal stem/stromal cells induce the generation of novel IL-10-dependent regulatory dendritic cells by SOCS3 activation. *J Immunol* 189: 1182-1192.
56. Lai HY, Yang MJ, Wen KC, Chao KC, Shih CC and Lee OK (2010) Mesenchymal stem cells negatively regulate dendritic lineage commitment of umbilical-cord-blood-derived hematopoietic stem cells: an unappreciated mechanism as immunomodulators. *Tissue Eng Part A* 16: 2987-2997.
57. van den Berk LC, Roelofs H, Huijs T, Siebers-Vermeulen KG, Raymakers RA, Kogler G, Figdor CG and Torensma R (2009) Cord blood mesenchymal stem cells propel human dendritic cells to an intermediate maturation state and boost interleukin-12 production by mature dendritic cells. *Immunology* 128: 564-572.
58. Jovanovic M, Stefanoska I, Radojic L and Vicovac L (2010) Interleukin-8 (CXCL8) stimulates trophoblast cell migration and invasion by increasing levels of matrix metalloproteinase (MMP)2 and MMP9 and integrins alpha5 and beta1. *Reproduction* 139: 789-798.
59. Khokha R, Murthy A and Weiss A (2013) Metalloproteinases and their natural inhibitors in inflammation and immunity. *Nat Rev Immunol* 13: 649-665.
60. Sudres M, Norol F, Trenado A, Gregoire S, Charlotte F, Levacher B, Lataillade JJ, Bourin P, Holy X, Vernant JP, Klatzmann D and Cohen JL (2006) Bone marrow mesenchymal stem cells suppress lymphocyte proliferation in vitro but fail to prevent graft-versus-host disease in mice. *J Immunol* 176: 7761-7767.
61. Nazarov C, Lo Surdo J, Bauer SR and Wei CH (2013) Assessment of immunosuppressive activity of human mesenchymal stem cells using murine antigen specific CD4 and CD8 T cells in vitro. *Stem Cell Res Ther* 4: 128.
62. Lutz MB and Schuler G (2002) Immature, semi-mature and fully mature dendritic cells: which signals induce tolerance or immunity? *Trends Immunol* 23: 445-449.
63. English K, Barry FP and Mahon BP (2008) Murine mesenchymal stem cells suppress dendritic cell migration, maturation and antigen presentation. *Immunol Lett* 115: 50-58.