



**PROTEIN KINASE C ISOFORMS PLAY DIFFERENTIAL ROLES
IN THE REGULATION OF HUMAN SEBOCYTE BIOLOGY**

Journal:	<i>Journal of Investigative Dermatology</i>
Manuscript ID:	JID-2011-0692.R1
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Géczy, Tamás; University of Debrecen, Medical and Health Science Center, Department of Physiology, DE-MTA "Lendület" Cellular Physiology Research Group</p> <p>Oláh, Attila; University of Debrecen, Medical and Health Science Center, Department of Physiology, DE-MTA "Lendület" Cellular Physiology Research Group</p> <p>Tóth, Balázs; University of Debrecen, Medical and Health Science Center, Department of Physiology, DE-MTA "Lendület" Cellular Physiology Research Group</p> <p>Czifra, Gabriella; University of Debrecen, Medical and Health Science Center, Department of Physiology, DE-MTA "Lendület" Cellular Physiology Research Group</p> <p>Szóllósi, Attila; University of Debrecen, Medical and Health Science Center, Department of Physiology, DE-MTA "Lendület" Cellular Physiology Research Group</p> <p>Szabó, Tamás; University of Debrecen, Medical and Health Science Center, Department of Pediatrics</p> <p>Zouboulis, Christos; Dessau Medical Center, Departments of Dermatology and Immunology</p> <p>Paus, Ralf; University of Lübeck, Department of Dermatology; University of Manchester, Epithelial Sciences, School of Translational Medicine</p> <p>Biro, Tamas; University of Debrecen, Medical and Health Science Center, Department of Physiology, DE-MTA "Lendület" Cellular Physiology Research Group</p>
Key Words:	protein kinase C (PKC) isoenzymes, sebaceous gland, SZ95 sebocytes, lipid synthesis, apoptosis

DEBRECENI EGYETEM
ORVOS- ÉS EGÉSZSÉGTUDOMÁNYI CENTRUM
ÉLETTANI INTÉZET

Sejt- és Molekuláris Élettani Laboratórium
Molekuláris Medicina Kutatóközpont
(EU Center of Excellence)

DE-MTA „Lendület” Sejtélettani Kutatócsoport
4032 Debrecen, Nagyerdei krt 98.

Vezető: Dr. Bíró Tamás
egyetemi docens



UNIVERSITY OF DEBRECEN
MEDICAL AND HEALTH SCIENCE CENTER
DEPARTMENT OF PHYSIOLOGY

Laboratory for Cellular and Molecular Physiology
Research Center for Molecular Medicine
(EU Center of Excellence)

DE-MTA „Lendület” Cellular Physiology Research Group
Nagyerdei krt 98, Debrecen, H-4032 Hungary-EU

Chief: Dr. Tamás Bíró, M.D., Ph.D., D.Sc.
Associate Professor

Phone: +36-52-255-575. FAX: +36-52-255-116. Cellular: +36-209-816-326 or +36-20-583-7074
Email: biro.tamas@med.unideb.hu, biro.lcmp@gmail.com

To: Professor Paul. R. Bergstresser
Editor
Journal of Investigative Dermatology
Editorial Office

December 19, 2011

Dear Professor Bergstresser:

Please find enclosed our revised manuscript (**JID-2011-0692R1**) which we now re-submit for publication to the *Journal of Investigative Dermatology*.

During the revision, we have corrected the manuscript according to the very constructive comments and suggestions of both the Editors and the Reviewers. In the below pages you find our point-by-point answers to the Reviewers. As requested, all changes are underlined in the revised text.

In addition, to show novel experimental data, we have supplemented the manuscript with 3 Supplementary Figures and the corresponding Supplementary data file (these files are also uploaded at the JID website).

We hope that these revisions, along with our point-by-point answers to comments of the Reviewers, are sufficient for considering our work for publication in your distinguished journal.

Yours sincerely,

Tamás Bíró, M.D., Ph.D., D.Sc.

Department of Physiology, DE-MTA „Lendület” Cellular Physiology Research Group,
University of Debrecen, Medical and Health Science Center, Research Center for
Molecular Medicine, 4032 Debrecen, Nagyerdei krt. 98. PO Box 22, Hungary

Email: biro.tamas@med.unideb.hu or biro.lcmp@gmail.com

Phone: +36-52-255-575

FAX: +36-52-255-116

Response to the Reviewers

Response to Reviewer 1.

In the paper titled "PROTEIN KINASE C ISOFORMS PLAY DIFFERENTIAL ROLES IN THE REGULATION OF HUMAN SEBOCYTE BIOLOGY" the authors demonstrate differential and antagonistic roles of PKC isoforms in a human sebocyte cell line. The authors show that PKCdelta and PKCzeta are important regulators of lipid synthesis and apoptosis.

Major Revisions:

1. What was the cell viability in Fig 3 after 48 and 72hrs of AA treatment? In figure 5 you demonstrate that there is a 50% decrease in cell viability about 48hrs of AA treatment. So which cells are you measuring PKC levels in if 50% of the cells are dead? Are you just measuring the cells in the population that express less PKCdelta?
--> Very important note, thank you for it. In Figure 5.b., we indeed show that the mitochondrial membrane potential (and early sign of the induction of apoptosis) was significantly decreased following 48-hr AA-treatment. However, we also present that necrosis was not initiated (no change in the intensity of the necrosis marker SYTOX Green) suggesting the lack of sudden changes in the viable cell number.

However, in order to exclude the "PKCdelta-specific selection" mentioned by Reviewer 1, we have performed additional experiments and investigated the effects of prolonged (1-3 days) AA-treatments on the living cell number, using CyQUANT cellular proliferation assay. As now shown in the new Supplementary Figure S2, we found that despite of the induced apoptosis there were no significant changes in the cell count compared to the appropriate vehicle control groups. These results indicate that (i) the early apoptotic processes shown in Figure 5.b. may reduce the cell count only in a longer time scale; (ii) most (if not all) cells were alive at the time when the Western blots were performed; and (iii) hence Western blots show the PKC expression of the whole cell population.

To mention these, we now present in Result section, paragraph 10, new last 2 sentences. In addition, we attach Supplementary data with the corresponding Materials and Methods section and Figure legends.

2. In figures 1, 3, and 4 there are no western blot loading controls such as beta-actin, alpha-tubulin, or GAPDH. I would like to see one of these loading controls included with the isoform western blots.

--> Thank you for the note. Since the goal of the Western blots shown in Figure 1. was only the demonstration of the presence of the PKC isoforms (and not the presentation of the potential changes in their expression level), loading controls could not provide further support for the message of the figure.

1
2
3 However, in the cases of Figures 3. and 4., Reviewer 2 is absolutely right; therefore
4 the following changes were made:

- 5 • For Figure 3b, loading controls are now shown in the new Supplementary
6 Figure S1a (with this modification, the originally shown loading control was
7 omitted from Figure 3b). This is now mentioned as the last sentence in
8 paragraph 6 of the Results section. In addition, a new Figure legend is shown
9 in the Supplementary data.
- 10 • For Figure 4a, loading controls are now shown in the new Supplementary
11 Figure S1b. This is now mentioned as the last sentence in paragraph 7 of the
12 Results section. In addition, a new Figure legend is shown in the
13 Supplementary data.
- 14 • In the modified Figure 4b, we now show the loading controls for those PKCs
15 which were changed upon siRNA transfection; i.e., loading control of
16 cPKCalpha for the upper panel and loading control of aPKCzeta for the lower
17 panel.
18
19
20
21
22
23
24

25 *Minor Revisions:*

26 *3. I would like to see a paragraph in the discussion talking about the prostaglandin*
27 *pathway in response to the addition of AA. I would like to see discussed whether a*
28 *COX inhibitor would also decrease the levels of apoptosis in these cells treated with*
29 *AA.*
30

31 --> Very good suggestion. We now present a new paragraph 4 to the Discussion
32 section. Please note that 3 additional references are now shown in the Reference list.
33
34

35
36 *4. I would also like to see the paragraph of speculation about applications in human*
37 *disease removed. Cell culture is notoriously poor at predicting human responses to*
38 *drugs. If you would like to make this type of speculation I think it would be*
39 *appropriate that there be some animal testing that demonstrates PKC inhibitors as*
40 *good drugs to inhibit sebocyte diseases or cancers.*
41

42 --> As requested, the last paragraph was removed from the Discussion section.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Response to Reviewer 2

Geczy and colleagues investigated the role of PKC family in sebocytes by using human immortalized SZ95 sebocyte cell culture. They examined the expression of PKC family proteins in SZ95 cells with Q-PCR, western and immunostaining, then tested the effects of PKC chemical inhibitors on SZ95 cell lipid synthesis on basal, PMA- and arachidonic acid-induced lipid synthesis and cell survival. Furthermore, isoform-specific knocking down of PKC α , δ , ζ was performed to study isoform-specific effects on SZ95 cell lipid synthesis and cell death. They observed that PKC α and PKC δ are required for PMA-induced lipid synthesis, while PKC δ also involved arachidonic acid-induced lipid synthesis and apoptosis. In contrast, PKC ζ inhibits the basal lipid synthesis and apoptosis. The authors conclude that PKC family play isoform-specific roles in regulation of human sebaceous gland function.

Comments:

The report provides interesting information on the role of PKC in human sebocytes. The manuscript is clearly written with sound data. The study implicates the importance of PKC family proteins in sebocyte biology and provides a potential target pathway for the clinical management of inflammatory SG disorders.

--> Thank you very much for the positive evaluation of our work.

Here are a few specific concerns.

1. The error bars and p value shown in Figure 2-5 appear based on readings from 4 different wells from the same experiment. For transfected cells (transfection is a stochastic process), the results from 3-4 addition independent experiments should be included and analyzed for mean and p value.

--> Very good point. Now we present the mean and p values of the average of 3 independent experiments. In addition, the corresponding figure legends are also changed.

2. The authors need to discuss what the downstream effectors of PKCs in this setting are. Inflammatory cytokines and lipid synthesis pathway enzymes could be examined by Q-PCR as the group has done for other study before. PKC ζ inhibits the fatty acid synthesis in Hela cells through phosphorylates mtF0F1-ATPase β -subunit. Does PKC ζ play a similar role in the SZ95 cells?

--> Very interesting point. Actually, the Reviewer is "knocking on opened doors", since we have already started the investigation of how inflammatory cytokines and other agents involved in innate (and/or adaptive) immunity are involved in the action of AA and how the PKC isoforms may influence their expressions. As a new Supplementary Figure S3, we present our preliminary Q-PCR data indicating that 24-hr AA treatment of SZ95 sebocytes markedly alter the expression of certain interleukins, pro-inflammatory agents, anti-microbial peptides, etc. In addition, we

1
2
3 present a new Discussion paragraph (only in the Supplementary Data section) which
4 discuss these findings.
5

6 However, according to our opinion, the second part of the note of Reviewer 2
7 requires a more detailed answer. Using a human hepatoma cell line (HepG2) Guo et
8 al. (J. Lipid Res . 2011. 52: 908–922.) have demonstrated that the aPKC ζ -driven
9 phosphorylation of the β -subunit of the mtF0F1-ATP-ase suppresses the activity of
10 this enzyme, which in turn leads to the inhibition of the mitochondrial acyl-
11 CoA:glycerol-sn-3-phosphate acyltransferase 1 (mtGPAT1) enzyme in the cells.
12 Since cellular GPAT enzymes control the first and rate-limiting step of TAG
13 synthesis, the activation of this pathway (by anthocyanin, a naturally occurring plant
14 pigment) has been described to play an important role in the reduction of the de novo
15 lipid biosynthesis in liver cells, thereby reducing high-glucose-induced hepatic
16 steatosis. Therefore it would seem plausible that, by activating the same signaling
17 pathway, aPKC ζ would have similar inhibitory effects on sebocyte lipid synthesis as
18 well.
19
20

21
22 However the following factors needs to be taken into consideration before one
23 decides to dissect the same pathway in sebocytes. As mentioned above, the first
24 committed step of TAG biosynthesis is controlled by GPAT enzymes. To date four
25 mammalian GPAT isoforms have been identified, two of which (GPAT1, GPAT2)
26 localizes to the mitochondrial outer membrane, whereas GPAT3 and GPAT4 reside
27 in the ER. GPAT1 expression levels are highest in liver and adipose tissue, where it
28 plays a key role in determining the fate of FFA molecules, by channeling them toward
29 glycerolipid synthesis (and thereby away from β -oxidation). Thus GPAT1 has a
30 pivotal regulatory role in whole-body lipid metabolism, facilitating pathways that
31 enable energy storage. Apart from these two tissues, mitochondrial GPAT activity is
32 also significant in muscle, brain, lung and kidney. However, other tissues possess
33 only a 10% GPAT activity attributable to the mitochondrial GPAT (GPAT1&2)
34 isoforms (Takeuchi and Reue, Am J Physiol Endocrinol Metab 296: E1195–E1209,
35 2009). Although the exact GPAT expression pattern of sebocytes is yet to be
36 described, the most important feature of sebocyte lipid synthesis is that it takes place
37 mainly in the smooth ER of the cells, and therefore does not involve mitochondrial
38 enzymes. Throughout their differentiation sebocytes increase their intracellular sER
39 content, thereby enabling the production and accumulation of an increased level of
40 intracellular lipids. (This process probably does not have any significant effect on
41 whole-body lipid metabolism, unlike the ones described in hepatocytes and
42 adipocytes) Ultimately, the cells disintegrate (at the end stage of their maturation)
43 and release their lipid content into the center of the gland (holocrine secretion).
44
45
46

47 The complex mixture of lipids secreted by the sebaceous gland is called sebum. It
48 consists of a wide variety of lipid molecules. TAG molecules constitute only 30-50%
49 of this lipid mixture, the rest of which consists mainly of FFA, wax esters, squalene
50 and cholesterol (Picardo et al, Dermato-Endocrinology 1:2, 68-71;2009; Smith and
51 Thiboutot, J. Lipid Res. 2008. 49: 271–281.). Since our methods, that analyze the
52 level of sebocyte lipid production, do not distinguish between these individual
53 components, we cannot determine whether or not the changes in overall lipid content
54 of sebocytes are attributable to changes in TAG levels within the cells.
55
56
57
58
59
60

1
2
3 Taken together, we would predict that the mechanism by which aPKC ζ activation
4 inhibits mitochondrial GPAT1-driven TAG synthesis in hepatocytes probably does not
5 have a major impact on the sebaceous lipid production of sER-related enzymatic
6 pathways in sebocytes. Therefore, in our future studies we would like to direct our
7 investigative efforts towards other potential downstream mechanisms.
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

PROTEIN KINASE C ISOFORMS PLAY DIFFERENTIAL ROLES IN THE REGULATION OF HUMAN SEBOCYTE BIOLOGY

**Tamás Géczy¹, Attila Oláh¹, Balázs I. Tóth¹, Gabriella Czifra¹, Attila G. Szöllősi¹,
Tamás Szabó², Christos C. Zouboulis³, Ralf Paus^{4,5}, Tamás Bíró¹**

¹Department of Physiology, DE-MTA “Lendület” Cellular Physiology Research Group
and ²Department of Pediatrics, University of Debrecen, Medical and Health Science
Center, Research Center for Molecular Medicine, Debrecen, Hungary

³Departments of Dermatology and Immunology, Dessau Medical Center, Dessau,
Germany

⁴Department of Dermatology, University of Lübeck, Lübeck, Germany

⁵Epithelial Sciences, School of Translational Medicine, University of Manchester,
Manchester, UK

Running title: PKC isoforms regulate human sebocyte biology

Key words: Protein kinase C, isoenzymes, human SZ95 sebocytes, lipid synthesis, apoptosis

Corresponding author

Tamás Bíró, M.D., Ph.D., Department of Physiology, DE-MTA “Lendület” Cellular Physiology Research Group, University of Debrecen, Medical and Health Science Center, Research Center for Molecular Medicine, 4032 Debrecen, Nagyerdei krt. 98. PO Box 22, Hungary, Email: biro@phys.dote.hu, Phone: +36-52-416-634, FAX: +36-52-432-289

ABSTRACT

Protein kinase C (PKC) isoforms play crucial roles in cutaneous signaling. Interestingly, we lack information about their involvement in human sebaceous gland biology. Therefore, in the current study, we investigated the functions of the PKC system in human immortalized SZ95 sebocytes. Using molecular biological approaches, imaging, and functional assays, we report that SZ95 sebocytes express the conventional cPKC α ; the novel nPKC δ , ϵ , and η ; and the atypical aPKC ζ . Activation of the PKC system by phorbol 12-myristate 13-acetate (PMA) stimulated lipid synthesis (a hallmark of differentiation), and resulted in translocation and then down-regulation of cPKC α and nPKC δ . In good accord with these findings, the effect of PMA was effectively abrogated by inhibitors and siRNA-mediated “silencing” of cPKC α and nPKC δ . Of further importance, molecular or pharmacological inhibition of nPKC δ also prevented the lipogenic and apoptosis-promoting action of arachidonic acid. Finally, we also found that “knock-down” of the endogenous aPKC ζ activity markedly increased basal lipid synthesis and apoptosis, suggesting its constitutive activity in suppressing these processes. Collectively, our findings strongly argue for that certain PKCs play pivotal, isoform-specific, differential, and antagonistic roles in the regulation of human sebaceous gland-derived sebocyte biology.

Key words: protein kinase C (PKC); isoenzymes; human sebaceous gland; human SZ95 sebocytes; lipid synthesis; apoptosis.

INTRODUCTION

Protein kinase C (PKC) comprises a family of serine/threonine kinases (Nishizuka, 1988, 1992) that play crucial roles in the regulation of various biological mechanisms including e.g. cell growth, differentiation, and survival. The PKC family includes the i) calcium- and diacylglycerol (DAG) or phorbol ester-dependent “conventional” isoforms (PKC α , β I, β II, and γ ; cPKCs); ii) calcium-independent “novel” nPKCs (PKC δ , ϵ , η , and θ); and iii) calcium- and phorbol ester-independent “atypical” aPKCs (PKC ζ and λ /I). These isoenzymes possess a characteristic expression pattern in a given tissue and cell type, and differentially regulate numerous cellular processes in an isoenzyme specific fashion (Gould and Newton, 2008; Reyland, 2009; Newton, 2010; Rosse et al, 2010).

Emerging evidence suggest a pivotal role of the PKC system in cutaneous biology (Denning, 2004; Breitkreutz et al, 2007). Indeed, PKC activity has been shown to modulate the proliferation and differentiation of epidermal keratinocytes (Lee et al, 1998; Papp et al, 2003). In addition, isoenzyme-specific roles of PKCs expressed by keratinocytes have also been reported; i.e. cPKC α and nPKC δ were found to be key components in stimulating epidermal differentiation whereas PKC ϵ was shown to promote skin tumor development (Lee et al, 1997; Li et al, 1999a; Jansen et al, 2001; Neill et al, 2003; Papp et al, 2004).

Importantly, PKCs are also involved in controlling hair growth and cycling. Several PKC isoforms (e.g. cPKC α and β , nPKC δ and ϵ) were shown to be expressed in various compartments (e.g. outer root sheath keratinocytes, dermal papilla) of the hair follicle (Eicheler et al, 1997; Li et al, 2003; Neill et al, 2003). Moreover, levels of certain

1
2
3 isoforms appeared to alter with the hair cycle; for example, expression of cPKC α
4
5 showed good correlation with increasing hair growth (anagen) in various mouse models
6
7 (Li et al, 1999b, 2003). Of further importance, modification of PKC activity by either
8
9 phorbol esters or synthetic inhibitors resulted in profound alterations in human hair shaft
10
11 elongation as well as dermal papilla fibroblast growth and survival (Harmon et al, 1995;
12
13 Ferraris et al, 1997).
14
15
16
17

18
19 Intriguingly, we lack data on the functional role of the PKC system in the other
20
21 component of the human pilosebaceous unit, i.e. the sebaceous gland and its
22
23 sebocytes. Sebocytes, engaged in holocrine (sebum) secretion, are major site of
24
25 hormone synthesis and metabolism in the human skin, and possess numerous
26
27 ionotropic, metabotropic, and nuclear receptor-coupled pathways (Zouboulis et al, 2002;
28
29 Zouboulis and Bohm, 2004; Alestas et al, 2006; Zhang et al, 2006; Dobrosi et al, 2008;
30
31 Tóth et al, 2009, 2011) which may all function as potential regulators of PKCs. Hence
32
33 sebocytes provide a highly instructive research tool for exploring PKC functions, which
34
35 promises additional sets of information that may complement those obtainable with
36
37 human hair follicle and with other extrafollicular cell types of the human skin.
38
39
40
41
42
43
44

45
46 Therefore, in the current study, we identified and investigated members of the PKC
47
48 system in SZ95 sebocytes – a human SG-derived immortalized cell line which
49
50 possesses striking functional similarities to those of primary human sebocytes
51
52 (Zouboulis et al, 1999) – and dissected their isoform-specific role in the regulation of
53
54 sebocyte biology.
55
56
57
58
59
60

RESULTS

Human SZ95 sebocytes possess a characteristic PKC isoform pattern

We first investigated the presence of PKC isoforms in SZ95 sebocytes. Using immunocytochemistry, Western blotting, and quantitative “real-time” PCR, we provide the first evidence that SZ95 cells express cPKC α ; nPKC δ , ϵ , and η ; and aPKC ζ (**Figure 1a and b**). Importantly, none of the above techniques identified other members of the family (i.e. cPKC β I, β II, and γ ; nPKC η and θ ; aPKC λ I; and the unique PKC μ -PKD) (data not shown). In addition, Q-PCR also revealed that, among the above isoforms, cPKC α , nPKC δ , and aPKC ζ exhibited the highest expressions whereas the levels of nPKC ϵ and η were rather low (**Figure 1b**).

Pharmacological modulation of PKC activity alters lipid synthesis but does not influence cell viability of SZ95 sebocytes

One of the main hallmarks of sebocyte differentiation is the synthesis of various lipids, among which neutral lipids represent a major component (Rosenfield, 1989; Thody and Shuster, 1989; Doran et al, 1991, Zouboulis et al, 1998). Therefore, we investigated the effects of PKC-acting agents on the lipid content of cultured SZ95 sebocytes. Nile Red staining-based quantitative FLIPR measurement revealed that treatment with the general PKC activator phorbol 12-myristate 13-acetate (PMA) (up to 48 hrs) significantly stimulated lipid synthesis in a dose-dependent fashion (**Figure 2a**). This effect was equally inhibited by GF109203X (inhibitor of cPKCs and nPKCs), Gö6976 (inhibitor of cPKCs), and Rottlerin (inhibitor of nPKC δ) (**Figure 2b**). Of note, none of the inhibitors alone (i.e. when applied without PMA) affected basal lipid content of the cells (**Figure 2b**). Furthermore, none of the above protocols altered cell viability or induced cell death

1
2
3 of any form (apoptosis, necrosis); namely, the PKC activator or the inhibitors did not
4 significantly modify the viable cell number (MTT-based colorimetric proliferation assay,
5 **Figure 2c**), nor did they induce necrotic (SYTOX-Green accumulation assay) or
6 apoptotic cell death (fluorimetric measurement of the decrease of mitochondrial
7 membrane potential – DiIC₁(5) labeling) (**Figure 2d**).

14
15
16
17
18 *The effect of PMA to stimulate lipid synthesis of SZ95 sebocytes is mediated by cPKC α*
19 *and nPKC δ*

20
21
22 The above results suggested the involvement of multiple PKC isoforms in the action of
23 PMA. However, numerous reports indicate that the above inhibitors may often target
24 other signaling molecules (Gschwendt et al, 1994; Alessi, 1997) which question their
25 specificity and/or selectivity for the PKC isoforms. Therefore, in order to define the
26 specific roles of the PKC isoenzymes that were expressed at the highest levels (i.e.
27 cPKC α , nPKC δ , and aPKC ζ) in mediating the lipid synthesis-stimulating effect of PMA,
28 we have carried out a series of cellular imaging and molecular biology experiments.

29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Activation of a given PKC isoform is accompanied by the translocation of the molecule to
another cellular compartment. Therefore, we first investigated the potential alteration in
the subcellular localization of PKCs upon short-term (1 hr) PMA treatment. As revealed
by immunostaining of PKCs followed by confocal microscopy analysis, 10 nM PMA
induced the translocation of cPKC α and nPKC δ . As expected, being a PMA-insensitive
isoform, the subcellular localization of aPKC ζ was not changed upon the application of
the phorbol ester (**Figure 3a**).

1
2
3 Prolonged activation of the PKC isoforms results in the cleavage of the molecule by
4 proteases resulting in a decreased expression of the given PKC (down-regulation).
5
6
7 Therefore, we also assessed the effect of long-term (up to 3 days) PMA administration
8 on the levels of the PKCs. Consistent with the above confocal microscopy data, as
9 revealed by Western blot analysis, the phorbol ester markedly down-regulated cPKC α
10 and nPKC δ whereas the level of aPKC ζ was not altered (**Figure 3b**). Loading controls
11 are presented in Supplementary Figure S1a.
12
13
14
15
16
17
18
19

20
21
22 The involvement of cPKC α and nPKC δ in mediating the cellular effects of PMA was
23 further assessed by siRNA technique. Western blot analysis demonstrated that the
24 expressions of “targeted” PKC isoforms were significantly “knocked-down” by the
25 specific siRNA probes at day 2 after transfection and remained suppressed on day 3 as
26 well (**Figure 4a-c**). Specificity and selectivity of the procedure was demonstrated by the
27 followings (i) siRNA-mediated silencing of a given isoforms did not affect the expression
28 of the other two PKCs investigated (**Figure 4a and b**); (ii) scrambled siRNA probes had
29 no effect on the expression of the PKC isoforms (**Figure 4a-c**). Additional loading
30 controls are presented in Supplementary Figure S1b.
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45
46 Silencing of cPKC α and nPKC δ did not modify the basal lipid synthesis (**Figure 4d**) and
47 did not induce apoptosis/necrosis (**Figure 4e**) in SZ95 sebocytes. However, in good
48 accord with the above data, “knock-down” of either of these isoforms significantly
49 reduced the action of PMA to promote lipid synthesis (**Figure 4d**). Intriguingly, the
50 siRNA-mediated silencing of aPKC ζ dramatically increased the basal lipid synthesis of
51 the sebocytes (**Figure 4d**) and, furthermore, suppressed cellular viability and induced
52
53
54
55
56
57
58
59
60

1
2
3 apoptosis (but not necrosis) (**Figure 4e**). Of further importance, elevation of basal lipid
4 content was so marked in the α PKC ζ knocked-down cells that PMA was unable to
5 further stimulate lipid accumulation (**Figure 4d**).
6
7
8
9

10
11
12 *Effects of arachidonic acid to stimulate lipid synthesis and induce apoptosis of SZ95*
13 *sebocytes are selectively mediated by n PKC δ*
14
15

16
17 The above effects of PMA strikingly resembled the previously described findings with
18 arachidonic acid (AA), one of the most effective inducers of lipid synthesis in SZ95
19 sebocytes (Wróbel et al, 2003; Alestas et al, 2006; Tóth et al, 2009). Therefore, we
20 wished to dissect whether or not the PKC isoforms are also involved in mediating the
21 cellular effect of AA. As expected, AA induced a marked lipid synthesis in SZ95
22 sebocytes (**Figure 5a**). This effect was significantly (yet only partially) abrogated by both
23 GF109203X and Rottlerin but not by Gö6976 (**Figure 5a**) suggesting that cPKC α most
24 probably does not participate in mediating the effect of AA. Furthermore, PMA was able
25 to further increase the lipid synthesis in the AA-treated cells (**Figure 5a**).
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 Our previous studies have also shown that the effect of AA to augment lipid synthesis in
42 SZ95 sebocytes was accompanied by the induction of sebocyte apoptosis (Wróbel et al,
43 2003; Tóth et al, 2009). Therefore, we investigated the involvement of the PKC system
44 in the AA-induced apoptotic process. As expected, AA induced apoptosis (fluorimetric
45 measurement of mitochondrial membrane potential) but not necrosis (SYTOX-Green
46 accumulation assay) in SZ95 sebocytes (**Figure 5b**). This effect of AA was almost
47 completely prevented by GF109203X and Rottlerin but not by Gö6976, similar to the
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 actions of these inhibitors on the AA-induced lipid synthesis. Finally, we found that the
4
5 PKC activator PMA did not modify the effect of AA to promote apoptosis (**Figure 5b**).
6
7

8
9
10 Next, using the above cellular and molecular approaches, we further investigated the
11
12 isoform-selective involvement of certain PKCs in the action of AA. Confocal microscopy
13
14 analysis revealed that a short-term AA treatment (1 hr) of SZ95 sebocytes selectively
15
16 translocated nPKC δ whereas it did not modify the subcellular localizations of cPKC α and
17
18 aPKC ζ (**Figure 3a**). Similarly, long-term (1-3 days) administration of AA selectively
19
20 down-regulated nPKC δ whilst expressions of the other two isoforms were insignificantly
21
22 changed (**Figure 3b**). In addition, in order to assess the effect of prolonged AA
23
24 treatment on cell growth, proliferation was additionally measured for 3 days. As shown in
25
26 Supplementary Figure S2, long-term application of AA did not significantly affect the
27
28 growth rate suggesting that the selective down-regulation of nPKC δ by prolonged AA
29
30 application was not due to the suppression in cell number or selective suppression of
31
32 viability of those SZ95 sebocytes which may express nPKC δ at high levels.
33
34
35
36
37
38
39
40

41 Moreover, the effect of AA to stimulate lipid synthesis and induce apoptosis was
42
43 significantly suppressed in those SZ95 sebocytes in which the expression of nPKC δ
44
45 (but, notably, not that of cPKC α) was silenced by siRNA (**Figure 5c and d**). In aPKC ζ -
46
47 silenced sebocytes, AA was able to further stimulate the already highly elevated (basal)
48
49 lipid synthesis; however, in these cells, the apoptosis was so marked (and hence the
50
51 mitochondrial membrane potential was so reduced) that the effect of AA to induce
52
53 further apoptosis was not measurable (**Figure 5c and d**). Nevertheless, these findings
54
55
56
57
58
59
60

1
2
3 collectively argue for the selective involvement of nPKC δ in mediating the cellular effects
4
5 of AA.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only

DISCUSSION

In this study, we report that human sebaceous gland-derived SZ95 sebocytes possess a characteristic PKC isoform pattern (i.e. the highly expressed cPKC α , nPKC δ and aPKC ζ ; and the poorly expressed nPKC ϵ and η) which is similar to that described on human epidermal keratinocytes (Lee et al, 1997, 1998; Papp et al, 2003). In addition, we provide the first evidence that the highly expressed PKC isoforms play central (yet differential and antagonistic) roles in regulating cellular growth, survival, and lipid synthesis of human sebocytes.

Specifically, we found that the activation of the PKC system by the phorbol ester PMA stimulated lipid synthesis. A series of pharmacological (using various inhibitors) and molecular biology (employing siRNA-mediated “silencing”) studies revealed that this action engaged two isoforms, i.e. cPKC α and nPKC δ . Since these isoenzymes were reported to stimulate differentiation of human epidermal keratinocytes (Lee et al, 1997; Li et al, 1999a; Papp et al, 2004); and, furthermore, since holocrine sebum production is one of the key hallmarks of sebocyte differentiation (Zouboulis et al, 1999, 2002; Alestas et al, 2006), we can conclude that cPKC α and nPKC δ play key roles in promoting differentiation of these cutaneous cell types.

Moreover, nPKC δ was also involved in selectively mediating the lipogenic action of AA, one of the most widely studied inducers of lipid synthesis in SZ95 sebocytes (Wróbel et al, 2003; Alestas et al, 2006; Dobrosi et al, 2008; Tóth et al, 2009), which was previously shown to stimulate skin PKC activity (Reynolds et al, 1993; Lo et al, 1994). Indeed, both the pharmacological and the molecular inhibition of nPKC δ suspended the lipogenic

1
2
3 effect of AA. Of further importance, in SZ95 sebocytes with “silenced” nPKC δ AA was
4
5 unable to induce apoptosis which suggests the pro-apoptotic role of nPKC δ (in addition
6
7 to its pro-lipogenic action). Apparently, cPKC α is not involved in mediating the actions of
8
9 AA since i) the AA-induced lipid synthesis and apoptosis was unaltered on sebocytes
10
11 with “silenced” cPKC α ; and ii) the AA-elevated lipid synthesis (mediated by nPKC δ) was
12
13 further increased by the co-administration of PMA which (besides stimulating nPKC δ)
14
15 additionally activates cPKC α .
16
17
18
19
20
21

22 With respect to the potential link between the prostaglandin pathways and the action of
23 AA, it should be noted that sebocytes were found to produce metabolites of both the
24 lipoxigenase (LOX) and the cyclooxygenase (COX) pathways (Alestas et al, 2006).
25 Moreover, it was also shown that COX inhibitors (indomethacin, diclofenac etc.)
26 enhanced (Iwata et al, 2005) whereas the LOX inhibitor Zileuton dramatically decreased
27 sebaceous lipid production (Zouboulis et al, 2005; Zouboulis, 2009). These data suggest
28 that AA metabolites may play opposite roles in the regulation of the sebaceous lipid
29 synthesis; i.e. LOX products seem to increase, whereas COX products seem to
30 decrease it. Therefore, the facts that increased lipid synthesis and apoptotic processes
31 are usually “walking hand-in-hand” during the sebaceous differentiation, and that COX
32 inhibition enhances lipid production make COX inhibitors very unlikely to decrease AA-
33 induced apoptosis of the sebocytes. (For our preliminary data on the effect of AA on
34 immune profile of SZ95 sebocytes, see **Supplementary Figure S3** and **Supplementary**
35 **Discussion**).
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Interestingly, neither the siRNA-mediated “knock-down”, nor the pharmacological
4 inhibition of cPKC α and nPKC δ resulted in alterations in cellular viability and basal lipid
5 synthesis of the sebocytes. These data implicate that the endogenous activities of the
6 two isoforms are not crucial for regulating basal growth and differentiation of sebocytes.
7
8 However, results obtained with PMA and AA strongly suggest their central involvement
9 in such conditions when the engagement of various signal transduction mechanisms
10 results in production of PKC-activating intracellular secondary messengers such as e.g.
11 DAG, Ca²⁺, etc. Since human sebocytes express a wide-array of metabotropic and
12 nuclear receptors and their coupled “lipogenic” signaling pathways (Zouboulis et al,
13 2002; Zouboulis and Bohm, 2004; Alestas et al, 2006; Zhang et al, 2006; Dobrosi et al,
14 2008; Tóth et al, 2009, 2011), it is proposed that cPKC α and/or nPKC δ may play a role
15 in their actions. (Evidently, the validity of the above hypothesis should be investigated in
16 the future.)
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

36 The siRNA experiments resulted in other intriguing findings. Namely, we found a
37 dramatically increased lipid synthesis and apoptosis rate in cells with “silenced” aPKC ζ
38 levels. These results, on the one hand, suggest that – in contrast to the pro-lipogenic
39 actions of cPKC α and nPKC δ , and the pro-apoptotic effect of nPKC δ – aPKC ζ rather
40 inhibits lipid production and apoptosis of SZ95 sebocytes. On the other hand, our
41 findings also implicate that the endogenous and, even more importantly, *constitutive*
42 aPKC ζ activity is an indispensable regulatory factor for promoting physiological sebocyte
43 proliferation and survival.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Our preclinical data that certain PKC isoforms specifically regulate human sebocyte
4 biology may have clinical implications which invite subsequent studies. Namely, it
5 deserves systematic analysis whether selective α -PKC ζ inhibitors can be employed in the
6 management of sebaceous tumors. Likewise, the data reported here will hopefully
7 encourage one to explore whether the targeted inhibition of cPKC α and/or nPKC δ
8 activity could become a useful treatment strategy for such human dermatoses (e.g. acne
9 vulgaris) which are characterized by pathologically elevated sebum production of the
10 sebaceous glands.
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

MATERIALS AND METHODS

Materials

Phorbol 12-myristate 13-acetate (PMA), GF 109203X and arachidonic acid (AA) were purchased from Sigma-Aldrich (St. Louis, MO). Gö6976 and Rottlerin were from Calbiochem (Nottingham, UK). The primary antibodies against cPKC α , β _I, β _{II}, γ ; nPKC ϵ and η ; and aPKC ζ and PKD-PKC μ were obtained from Sigma-Aldrich, and the ones against nPKC δ and θ ; and aPKC λ / ι were from Santa Cruz Biotechnology (Santa Cruz, CA)

Cell culturing

Immortalized human SZ95 sebocytes (derived from facial sebaceous glands; Zouboulis et al., 1999) were cultured in Sebomed Basal medium (Biochrom, Berlin, Germany) supplemented with 10% fetal bovine serum (Invitrogen, Paisley, UK), 5 ng/ml human epidermal growth factor (Sigma-Aldrich), 1 mM CaCl₂, 50 U/ml penicillin and 50 μ g/ml streptomycin (both from Teva, Debrecen, Hungary) (Dobrosi et al, 2008; Tóth et al, 2009)

Western blotting

Immunoblotting was performed to assess the protein expression of the various PKCs in SZ95 cells. Cell lysates were subjected to SDS-PAGE (7.5% gels were loaded with 40 μ g protein per lane), transferred to BioBond nitrocellulose membranes (Whatman, Maidstone, UK), and probed with the above mentioned rabbit primary antibodies against distinct PKC isoforms (dilution 1:100-1:1000). Horseradish peroxidase-conjugated rabbit

1
2
3 anti-goat IgG antibody (1:1000, Bio-Rad, Hercules CA, USA) was employed as a
4 secondary antibody, and the immunoreactive signal was visualized by a SuperSignal®
5 West Pico Chemiluminescent Substrate enhanced chemiluminescence kit (Pierce
6 Rockford, IL) using LAS-3000 Intelligent Dark Box (Fuji, Tokyo, Japan). To confirm equal
7 loading, membranes were re-probed with an anti cytochrome-C antibody (Cyt-C, 1:50,
8 Santa Cruz). Where indicated, signal intensity was analyzed with densitometry, using
9 the Image Pro Plus 4.5.0 software (Media Cybernetics, Silver Spring, MD) (Gönczi et al,
10 2008; Szegedi et al, 2009).
11
12
13
14
15
16
17
18
19
20
21
22
23

24 *Immunostaining of PKCs, confocal microscopy*

25
26 Identification and determination of the subcellular localization of PKC isozymes in SZ95
27 cells were carried out by confocal microscopy following immunolabeling. Cells were fixed
28 in acetone, permeabilized by 0.1% Triton-X-100 (Sigma-Aldrich) and then incubated with
29 the above rabbit anti-PKC primary antibodies for 60 min (dilution 1:100-1:200).
30 Coverslips were then stained with a FITC-conjugated secondary antibody (Vector
31 Laboratories, Burlingame, CA, USA) (dilution 1:200), and examined in a Zeiss LSM
32 confocal microscopy system (Carl Zeiss Inc., Oberkochen, Germany) (Bodó et al, 2005;
33 Dobrosi et al, 2008; Tóth et al, 2009).
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 *Quantitative “real-time” PCR (Q-PCR)*

49
50 Q-PCR was carried out on an ABI Prism 7000 sequence detection system (Applied
51 Biosystems, Foster City, CA) using the 5' nuclease assay. Total RNA was isolated from
52 cell lysates using TRIzol (Invitrogen). Three micrograms of total RNA was then reverse-
53 transcribed into cDNA by using 15 U of AMV reverse transcriptase (Promega, Madison,
54
55
56
57
58
59
60

1
2
3 Wl) and 0.025 $\mu\text{g}/\mu\text{l}$ oligodT primers (Promega). TaqMan primers and probes (Applied
4 Biosystems) were applied for subsequent PCR amplification (assay IDs,
5 Hs00176973_m1 for cPKC α , Hs00176998_m1 for cPKC β , Hs00177010_m1 for cPKC γ ,
6 Hs00178914_m1 for nPKC δ , Hs00178455_m1 for nPKC ϵ , Hs00178933_m1 for nPKC η ,
7 Hs00234709_m1 for nPKC θ , Hs00177051_m1for aPKC ζ , and Hs00702254_s1 for
8 aPKC λ/i) using the TaqMan universal PCR master mix protocol (Applied Biosystems).
9
10 As internal controls, transcripts of glyceraldehyde 3-phosphate dehydrogenase
11 (GAPDH) were determined (assay ID, Hs99999905_m1 for human) (Dobrosi et al, 2008;
12 Tóth et al, 2009, 2011).
13
14
15
16
17
18
19
20
21
22
23
24
25
26

27 *Assessment of viability*

28
29 Viable cell number was assessed by measuring the ability of cellular dehydrogenases to
30 convert the tetrazolium salt MTT (Sigma-Aldrich) to formazan. Cells were plated onto 96-
31 well plates (Greiner Bio-One, Frickenhausen, Germany) (15,000 cells/well) in
32 quadruplicates. After treatment, cells were incubated with 0.5 mg/ml MTT for 3 hours,
33 and the amount of formazan crystals (generated within the cells) was determined
34 colorimetrically according to our previous reports (Bodó et al, 2005; Kiss et al, 2008;
35 Szegedi et al, 2009).
36
37
38
39
40
41
42
43
44
45
46
47

48 *Assessment of apoptosis*

49
50 Reduced mitochondrial membrane potential serves as an early indicator of the onset of
51 apoptotic processes. SZ95 cells (15,000 cells/well) were plated onto 96-well black-
52 well/clear-bottom plates (Greiner Bio One) in quadruplicates. After treatment, their
53 mitochondrial membrane potential was determined using MitoProbe™ DiIC $_1(5)$ Assay Kit
54
55
56
57
58
59
60

1
2
3 (Invitrogen). The fluorescence intensity (reflecting the level of mitochondrial membrane
4 potential) was measured on FlexStation II³⁸⁴ Fluorescence Image Plate Reader (FLIPR)
5
6 (Molecular Devices, San Francisco, CA), as described in our previous reports (Dobrosi
7
8 et al., 2008; Tóth et al, 2010, 2011).
9
10
11

12 13 14 15 *Assessment of cytotoxicity (necrosis)*

16
17 The cytotoxic effects of the above PKC-acting reagents were determined by SYTOX
18 Green labeling (Invitrogen). Ruptured plasma membranes enable the penetration (and
19 subsequent nucleic acid binding) of the fluorescent dye into necrotic cells, whereas
20 viable cells with intact surface membranes display negligible SYTOX Green uptake.
21
22 SZ95 cells were plated onto 96-well black-well/clear-bottom plates (Greiner Bio One).
23
24 After treatment, supernatants were removed and the cells were incubated with a 1 μ M
25 solution of SYTOX Green. The fluorescence intensity of the dye was measured on
26
27 FLIPR, as described previously (Dobrosi et al., 2008; Tóth et al, 2010, 2011).
28
29
30
31
32
33
34
35
36
37

38 39 *Assessment of intracellular lipid content*

40
41 For a quantitative measurement of the intracytoplasmic lipid content, Nile Red (Sigma-
42 Aldrich) labeling was employed. SZ95 cells (15,000 cells/well) were plated onto 96-well
43 black-well/clear-bottom plates (Greiner Bio One) in quadruplicates. Following treatment,
44
45 supernatants were removed from wells and cells were incubated in 1 μ g/ml Nile Red for
46
47 20 min. The fluorescence intensity of the dye (excitation: 485 nm; emission: 565 nm)
48
49 reflecting the neutral lipid content of intracytoplasmic lipid droplets was measured on
50
51
52
53
54
55
56
57
58
59
60
60 FLIPR (Wróbel et al, 2003; Alestas et al, 2006, Dobrosi et al., 2008).

RNA interference (siRNA)

SZ95 sebocytes were seeded on six-well culture plates in medium lacking antibiotics and were grown until reaching 40-60% confluence. Cells were then transfected with specific Stealth diRNAi oligonucleotides (at 40 nM, all from Santa Cruz) against cPKC α (sc-44227), nPKC δ (sc-36253) and aPKC ζ (sc-29451) using Lipofectamine RNAiMAX transfection reagent (Invitrogen). siRNA Negative Control Duplexes (scrambled siRNA), were used as controls (sc-37007 and sc-44230; Santa Cruz). Immunoblotting was performed daily (for 4 days) to follow the changes in the expression level of the given PKC isoform after transfection. For subsequent cellular assays (assessing lipid content, cell viability, apoptosis, and cytotoxicity) cells were trypsinized 24 h after transfection replated onto 96-well plates and subjected to the above mentioned procedures (Dobrosi et al., 2008; Tóth et al, 2010, 2011).

Statistical analysis

When applicable, data were analyzed using a two-tailed un-paired *t*-test and $P < 0.05$ values were regarded as significant differences.

1
2
3 **CONFLICT OF INTEREST**
4
5
6
7

8 The authors declare no competing financial interests.
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only

ACKNOWLEDGEMENT

This work was supported in part by Hungarian (OTKA NK78398, TÁMOP-4.2.2-08/1/2008-0019, TÁMOP 4.2.1./B-09/1/KONV-2010-0007, ETT 329-07) and EU (FP7-REGPOT-2008-1/22992) research grants. GC is a recipient of the János Bolyai scholarship of the Hungarian Academy of Sciences whereas AO is a recipient of the Richter "Talentum" Fellowship of the Richter Talentum Foundation, Hungary. The SZ95 sebaceous gland cell line is protected by the patents and patent applications EP1151082, DE59913210D, AU200019804, US2002034820, CA2360762, CN1344314T, JP2002535984, IL144683D, PL350191, HU0200048, AT319813T, DK1151082T, and KR31762.

REFERENCES

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- Alessi DR (1997) The protein kinase C inhibitors Ro 318220 and GF 109203X are equally potent inhibitors of MAPKAP kinase-1 β and p70 S6 kinase. *FEBS Letters* 402:121-3
- Alesta T, Ganceviciene R, Fimmel S *et al.* (2006) Enzymes involved in the biosynthesis of leukotriene B4 and prostaglandin E2 are active in sebaceous glands. *J Mol Med* 84:75-84
- Bodó E, Bíró T, Telek A *et al.* (2005) A “hot” new twist to hair biology – Involvement of vanilloid receptor-1 (VR1/TRPV1) signaling in human hair growth control. *Am J Pathol* 166:985-98
- Breitkreutz D, Braiman-Wiksman L, Daum N *et al.* (2007) Protein kinase C family: on the crossroads of cell signaling in skin and tumor epithelium. *J Cancer Res Clin Oncol* 133:793-808
- Denning MF (2004) Epidermal keratinocytes: regulation of multiple cell phenotypes by multiple protein kinase C isoforms. *Int J Biochem Cell Biol* 36:1141-6
- Dobrosi N, Tóth IB, Nagy G *et al.* (2008) Endocannabinoids enhance lipid synthesis and apoptosis of human sebocytes via cannabinoid receptor-2- mediated signaling. *FASEB J* 22:3685-95
- Doran TI, Baff R, Jacobs P, Pacia E (1991) Characterization of human sebaceous cells in vitro. *J Invest Dermatol* 96:341-8
- Eicheler W, Huth A, Happle R, Hoffmann R (1997) Phorbol-myristate-acetate, but not interleukin-1 beta or insulin-like growth factor-I, regulates protein kinase C isoenzymes in human dermal papilla cells. *Acta Derm Venereol* 77:361-4

- 1
2
3 Ferraris C, Cooklis M, Polakowska RR, Haake AR (1997) Induction of apoptosis through
4 the PKC pathway in cultured dermal papilla fibroblasts. *Exp Cell Res* 234:37-46
5
6
7
8 Gould CM, Newton AC. (2008) The life and death of protein kinase C. *Curr Drug Targets*
9
10 9:614-25
11
12
13 Gönczi M, Telek A, Czifra G *et al.* (2008) Altered calcium handling following the
14 recombinant overexpression of protein kinase C isoforms in HaCaT cells. *Exp*
15
16 *Dermatol* 17:584-91
17
18
19
20 Gschwendt M, Müller HJ, Kialbassa K *et al.* (1994) Rottlerin, a novel protein kinase
21 inhibitor. *Biochem Biophys Res Commun* 199:93-8
22
23
24
25 Harmon CS, Nevins TD, Bollag WB. (1995) Protein kinase C inhibits human hair follicle
26 growth and hair fibre production in organ culture. *Br J Dermatol* 133:686-93.
27
28
29 Iwata C, Akimoto N, Sato T *et al.* (2005) Augmentation of lipogenesis by 15-deoxy-
30 $\Delta^{12,14}$ -prostaglandin J2 in hamster sebaceous glands: identification of cytochrome
31 P-450-mediated 15-deoxy- $\Delta^{12,14}$ -prostaglandin J2 production. *J Invest Dermatol*
32 125:865–872
33
34
35
36
37
38
39 Jansen AP, Dreckschmidt NE, Verwiebe ED *et al.* (2001) Regulation of the induction of
40 epidermal ornithine decarboxylase and hyperplasia to the different skin tumor-
41 promotion susceptibilities of protein kinase C alpha, -delta, and -epsilon transgenic
42 mice. *Int J Cancer* 93:635-43
43
44
45
46
47
48 Kiss B, Bíró T, Czifra G *et al.* (2008): Investigation of micronized titanium-dioxide
49 penetration in human skin xenografts and its effect on cellular functions of human
50 skin-derived cells. *Exp Dermatol* 17:659-67
51
52
53
54
55 Lee YS, Dlugosz AA, McKay R *et al.* (1997) Definition by specific antisense
56 oligonucleotides of a role for protein kinase C α in expression of differentiation
57
58
59
60

1
2
3 markers in normal and neoplastic mouse epidermal keratinocytes. *Mol Carcinog*
4
5 18:44-53
6

7
8 Lee YS, Yuspa SH, Dlugosz AA (1998) Differentiation of cultured human epidermal
9
10 keratinocytes at high cell densities is mediated by endogenous activation of the
11
12 protein kinase C pathway. *J Invest Dermatol* 111:762-6
13
14

15 Li L, Lorenzo PS, Bogi K *et al.* (1999a) Protein kinase C δ targets mitochondria, alters
16
17 mitochondrial membrane potential, and induces apoptosis in normal and neoplastic
18
19 keratinocytes when overexpressed by an adenoviral vector. *Mol Cell Biol*
20
21 19:8547-58
22
23

24 Li LF, Fiedler VC, Kumar R (1999b) The potential role of skin protein kinase C isoforms
25
26 alpha and delta in mouse hair growth induced by diphencyprone-allergic contact
27
28 dermatitis. *J Dermatol* 26:98-105
29
30

31 Li LF, Guo J, Gao ZF (2003) Overexpression of skin protein kinase C-alpha in anagen
32
33 hair follicles during induced growth of mouse hair. *Clin Exp Dermatol* 28:429-33
34
35

36 Lo HH, Bartek GA, Fischer SM (1994) In vitro activation of mouse skin protein kinase C
37
38 by fatty acids and their hydroxylated metabolites. *Lipids* 29:547-53
39
40

41 Neill GW, Ghali LR, Green JL *et al.* (2003) Loss of protein kinase C alpha expression
42
43 may enhance the tumorigenic potential of Gli1 basal cell carcinoma. *Cancer Res*
44
45 63:4692-7
46
47

48 Newton AC (2010) Protein kinase C: poised to signal. *Am J Physiol Endocrinol Metab*
49
50 298:E395-402
51
52

53 Nishizuka Y (1988) The molecular heterogeneity of protein kinase C and its implication
54
55 for cellular regulation. *Nature* 334:661-5
56
57
58
59
60

- 1
2
3 Nishizuka Y (1992) Intracellular signaling by hydrolysis of phospholipids and activation
4 of protein kinase C. *Science* 258:607-14
5
6
7
8 Papp H, Czifra G, Bodó E *et al.* (2004) Opposite roles of protein kinase C isoforms in
9 proliferation, differentiation, apoptosis, and tumorigenicity of human HaCaT
10 keratinocytes. *Cell Mol Life Sci* 61:1095-1105
11
12
13
14 Papp H, Czifra G, Lázár J *et al.* (2003) Protein kinase C isozymes regulate proliferation
15 and high cell density-mediated differentiation of HaCaT keratinocytes. *Exp*
16 *Dermatol* 12:811-24
17
18
19
20
21
22 Reyland ME (2009) Protein kinase C isoforms: Multi-functional regulators of cell life and
23 death. *Front Biosci* 14:2386-99
24
25
26
27 Reynolds NJ, Talwar HS, Baldassare JJ *et al.* (1993) Differential induction of
28 phosphatidylcholine hydrolysis, diacylglycerol formation and protein kinase C
29 activation by epidermal growth factor and transforming growth factor-alpha in
30 normal human skin fibroblasts and keratinocytes. *Biochem J* 294:535-44
31
32
33
34
35
36 Rosenfield RL (1989) Relationship of sebaceous cell stage to growth in culture. *J Invest*
37 *Dermatol* 92:751-4
38
39
40
41 Rosse C, Linch M, Kermorgant S *et al.* (2010) PKC and the control of localized signal
42 dynamics. *Nat Rev Mol Cell Biol* 11:103-12
43
44
45
46 Szegedi A, Páyer E, Czifra G *et al.* (2009) Protein kinase C isoenzymes differentially
47 regulate the differentiation-dependent expression of adhesion molecules in human
48 epidermal HaCaT keratinocytes. *Exp Dermatol* 18:122-129
49
50
51
52
53 Thody AJ, Shuster S (1989) Control and function of sebaceous glands. *Physiol Rev*
54 69:383-416
55
56
57
58
59
60

- 1
2
3 Tóth IB, Dobrosi N, Dajnoki A *et al.* (2011) Endocannabinoids modulate human
4
5 epidermal keratinocyte proliferation and survival via the sequential engagement of
6
7 cannabinoid receptor-1 and transient receptor potential vanilloid-1. *J Invest*
8
9 *Dermatol* 131:1095-1104
10
11
12 Tóth IB, Géczy T, Griger Z *et al.* (2009) Transient receptor potential vanilloid-1 signaling
13
14 as a regulator of human sebocyte biology. *J Invest Dermatol* 129:329-39
15
16
17 Wróbel A, Seltmann H, Fimmel S *et al.* (2003). Differentiation and apoptosis in human
18
19 immortalized sebocytes. *J Invest Dermatol* 120:175-81
20
21
22 Zhang Q, Seltmann H, Zouboulis CC, Konger RL (2006) Involvement of PPARgamma in
23
24 oxidative stress-mediated prostaglandin E(2) production in SZ95 human
25
26 sebaceous gland cells. *J Invest Dermatol* 126:42-8
27
28
29 Zouboulis CC. (2009) Zileuton, a new efficient and safe systemic anti-acne drug.
30
31 *Dermatoendocrinol* 1(3):188-192
32
33
34 Zouboulis CC, Bohm M (2004) Neuroendocrine regulation of sebocytes - a pathogenetic
35
36 link between stress and acne. *Exp Dermatol* 13 Suppl 4: 31-5
37
38
39 Zouboulis CC, Saborowski A, Boschnakow A. (2005) Zileuton, an oral 5-lipoxygenase
40
41 inhibitor, directly reduces sebum production. *Dermatology* 210(1):36-38
42
43
44 Zouboulis CC, Seltmann H, Hiroi N *et al.* (2002). Corticotropin-releasing hormone: an
45
46 autocrine hormone that promotes lipogenesis in human sebocytes. *Proc Natl Acad*
47
48 *Sci USA* 99:7148-53
49
50
51 Zouboulis CC, Seltmann H, Neitzel H, Orfanos CE (1999). Establishment and
52
53 characterization of an Immortalized Human Sebaceous Gland Cell Line (SZ95). *J*
54
55 *Invest Dermatol* 113:1011-20
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Zouboulis CC, Xia L, Akamatsu H *et al.* (1998). The human sebocyte culture model provides new insights into development and management of seborrhoea and acne. *Dermatology* 196:21-31

For Review Only

FIGURE LEGENDS

Figure 1. *Human SZ95 sebocytes possess a characteristic PKC isoform pattern*

a) Specific immunoreactivity of various PKC isoforms as determined by immunofluorescence (FITC, green fluorescence). NC, negative control obtained after omitting the primary antibody. Nuclei were counterstained with DAPI (blue fluorescence). **Insets:** Expressions of PKC isoforms were also determined in cell lysates of SZ95 sebocytes by Western blot analysis. **b)** Q-PCR analysis of mRNA transcript expression profile of the PKC isoenzymes. Data (mean±SEM) are expressed as a fraction of the mean value of expression of the housekeeping gene GAPDH (defined as 1). Two additional experiments yielded similar results.

Figure 2. *PMA treatment stimulates lipid synthesis of SZ95 sebocytes*

Cells were treated with vehicle (Control, C); the PKC-activator PMA (at the indicated doses); the PKC inhibitors GF109203X (GF), Gö6986 (Gö), and Rottlerin (Rottl.) at 10 nM; or combinations for 48 hrs. **a, b)** Quantitative measurement of lipids as assessed by Nile Red labeling followed by FLIPR measurement. Neutral lipids indicate *de novo* synthesized intracellular lipids. **c)** Determination of cell viability by colorimetric MTT assay. **d)** Assessment of apoptosis by fluorimetric DilC₁(5) assay reflecting mitochondrial membrane potential; and of necrosis by fluorimetric SYTOX Green assay reflecting ruptured cell membrane. Data (mean±SEM) are expressed as a percentage of the mean value of the vehicle-treated control group (defined as 100 %, solid line). * marks significant ($P<0.05$) differences compared to the control group. In panel **b**, # marks significant ($P<0.05$) differences compared to the PMA-treated group. n=4 in each group. Three-four additional experiments yielded similar results.

1
2
3
4
5
6 **Figure 3.** *PMA and arachidonic acid differentially translocate and down-regulate certain*
7
8 *PKC isoforms in SZ95 sebocytes*

9
10 Cells were treated with vehicle (Control), 10 nM PMA, and 50 μ M arachidonic acid (AA)
11
12 for 1 hr (a) to assess translocation or for 1-3 days (b) to measure down-regulation of
13
14 PKC isoforms. a) Immunofluorescence labeling (FITC, green fluorescence) of PKCs
15
16 followed by confocal microscopy analysis. Arrows indicate translocation. b) Western blot
17
18 analysis. The amounts of the individual PKC isoforms were quantitated by densitometry
19
20 (optical density, OD) and expressed as the percentage of the value of immunoreactive
21
22 bands of daily-matched control group regarded as 100%. Note the translocation and
23
24 down-regulation of cPKC α and nPKC δ by PMA, and of nPKC δ by AA. Three additional
25
26 experiments yielded similar results.
27
28
29
30
31
32
33

34 **Figure 4.** *cPKC α and nPKC δ are involved in mediating the lipogenic action of PMA*
35
36 *whereas siRNA-mediated "silencing" of aPKC ζ induces lipid synthesis and apoptosis in*
37
38 *SZ95 sebocytes*

39
40 Various siRNA probes against PKC isoforms, as well as a scrambled siRNA probe (Scr.
41
42 si), were introduced to cells by transfection. To evaluate the efficacy of this intervention,
43
44 at days 1-4 after transfection, cells were subjected to Western blot analysis. As controls
45
46 (C), the effect of the transfection reagent was determined. As a house-keeping
47
48 molecule, expression of Cytochrome-C (Cyt-C) was assessed. a, b) Representative
49
50 Western blot data at days 1-4 (a) and at day 2 (b) after transfection. c) Statistical
51
52 analysis of Western blot data. Optical density (OD) values of specific immunosignals
53
54 were determined at day 2 after transfection in 3 independent experiments. Normalized
55
56
57
58
59
60

1
2
3 OD values (to Cyt-C) in each group were then averaged and expressed as mean±SEM
4 as the percentage of the averaged values of the control groups regarded as 100%. *
5 marks significant ($P<0.05$) differences compared to the Scr. si groups. At day 2 after
6 transfection, cells were treated with 10 nM PMA for 48 hrs. **d)** Determination of neutral
7 lipids as assessed by Nile Red labeling followed by FLIPR measurement. **e)**
8 Determination of cell viability by colorimetric MTT assay; of apoptosis by fluorimetric
9 DiIC₁(5) assay; and of necrosis by fluorimetric SYTOX Green assay. Data (mean±SEM)
10 of 3 independent experiments are expressed as a percentage of the mean value of the
11 Scr. si group (defined as 100 %, solid line). * marks significant ($P<0.05$) differences
12 compared to the Scr. si group whereas # marks significant ($P<0.05$) differences
13 compared to the PMA-treated group. n=4 in each group. The Scr. si group displayed no
14 significant difference compared to the non-transfected SZ95 sebocyte population
15 subjected to identical treatment protocols (data not shown). ~~Two additional experiments~~
16 ~~yielded similar results.~~

17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39 **Figure 5.** *The lipogenic and apoptosis-inducing actions of arachidonic acid are mediated*
40 *by nPKCδ in SZ95 sebocytes*

41 Control (**a, b**) or siRNA-transfected (**c, d**, 2 days after transfection) SZ95 sebocytes
42 were treated with vehicle (Control, C); 10 nM PMA; the PKC inhibitors GF109203X (GF),
43 Gö6986 (Gö), and Rottlerin (Rottl.) at 10 nM; 50 μM arachidonic acid (AA); or
44 combinations for 48 hrs. **a, c)** Quantitative measurement of neutral lipids as assessed by
45 Nile Red labeling followed by FLIPR measurement. **b, d)** Assessment of apoptosis by
46 fluorimetric DiIC₁(5) assay (**b, d**) and of necrosis by fluorimetric SYTOX Green assay
47 (**b**). Data (mean±SEM) of 3 independent experiments are expressed as a percentage of
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 the mean value of either the control (**a, b**) or the scrambled siRNA probe (Scr. si; **c, d**)
4
5 group (defined as 100 %, solid line). * marks significant ($P<0.05$) differences compared
6
7 to the control or Scr. si group whereas # marks significant ($P<0.05$) differences
8
9 compared to the AA-treated group. n=4 in each group. The Scr. si group displayed no
10
11 significant difference compared to the non-transfected SZ95 population subjected to
12
13 identical treatment protocols (data not shown). ~~Two additional experiments yielded~~
14
15 ~~similar results.~~
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only







Figure 1

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


a



b



c



d





Figure 2



b






Figure 3

a


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




b



c



d



e






Figure 4

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


a



b



c



d




Figure 5

SUPPLEMENTARY DATA

Supplementary Materials and methods

Quantitative “real-time” PCR (Q-PCR) of immune genes

Q-PCR was carried out on an ABI Prism 7000 sequence detection system (Applied Biosystems, Foster City, CA) using the 5' nuclease assay. Total RNA was isolated from cell lysates using TRIzol (Invitrogen). Three micrograms of total RNA was then reverse-transcribed into cDNA by using 15 U of AMV reverse transcriptase (Promega, Madison, WI) and 0.025 µg/µl oligodT primers (Promega). TaqMan primers and probes (Applied Biosystems) were applied for subsequent PCR amplification (assay IDs, Hs00174092_m1 for interleukin-1α (IL1α), Hs00174097_m1 for IL1β, Hs00985639_m1 for IL6, Hs00174103_m1 for IL8, Hs00174086_m1 for IL10, Hs01038788_m1 for IL18, Hs00174128_m1 for tumor necrosis factor α (TNFα), Hs00189038_m1 for LL37 cathelicidin (CAMP), Hs00914334_m1 for lactotransferrin (LTF), Hs00608345_m1 for β-defensin-1 (bDEF1) and Hs00173615_m1 for pentraxin-3 (PTX3) using the TaqMan universal PCR master mix protocol (Applied Biosystems). As internal controls, transcripts of glyceraldehyde 3-phosphate dehydrogenase (GAPDH) were determined (assay ID, Hs99999905_m1 for human) (Dobrosi et al, 2008; Tóth et al, 2009, 2011).

Assessment of cellular proliferation

The degree of cellular growth (reflecting number of viable cells) was determined by measuring the DNA content of cells using CyQUANT Cell Proliferation Assay Kit (Invitrogen). SZ95 sebocytes (5,000 cells per well) were cultured in 96-well black-well/clear-bottom plates (Greiner Bio-One) in quadruplicates and were treated with 50

1
2
3 μM AA for 24, 48 and 72 hrs. Supernatants were then removed by blotting on paper
4
5 towels, and the plates were subsequently frozen at -70°C . The plates were then
6
7 thawed at room temperature, and 200 μl of CyQUANT dye/cell lysis buffer mixture
8
9 was added to each well. After 5 minutes of incubation, fluorescence was measured at
10
11 490 nm excitation and 520 nm emission wavelengths using FLIPR (Molecular
12
13 Devices).
14
15

16 17 18 **Supplementary Discussion**

19
20 In our ongoing studies, we are currently investigating how AA treatment affects the
21
22 expression of immune and inflammatory processes of SZ95 sebocytes. In addition,
23
24 we are also assessing the involvement of certain PKCs in these processes. Our
25
26 preliminary Q-PCR data suggest that AA treatment markedly alter expressions of
27
28 certain interleukins, pro-inflammatory agents, anti-microbial peptides, etc. which are
29
30 involved in innate (and/or adaptive) immune mechanisms of these cells. Specifically,
31
32 we found that expressions of the well known pro-inflammatory interleukins (IL6 and
33
34 IL18) (Gauldi et al, 1992; Tsutsui et al, 2011]) as well as the anti-microbial peptide β -
35
36 defensin-1 (Prado and da Oca, 2010) and pentraxin-3 (PTX3), a central regulator of
37
38 the relationship between the innate and adaptive immune system (Deban et al,
39
40 2011), were significantly elevated by AA. Evidently, further (ongoing) studies are to
41
42 define the roles of the PKCs in these processes; yet, our preliminary data imply that
43
44 the above members of the sebocyte immune response machinery may function as
45
46 putative AA and PKC downstream targets.
47
48
49
50

51 52 53 **Supplementary References**

54
55
56
57
58
59
60

1
2
3 Deban L, Jaillon S, Garlanda C *et al.* (2001) Pentraxins in innate immunity: lessons
4
5 from PTX3. *Cell Tissue Res* 343:237–249
6

7 Gauldie J, Richards C, Baumann H (1992) IL6 and the acute phase reaction. *Res*
8
9 *Immunol* 143(7):755-759
10

11 Prado E and de Oca M (2010) Human β -defensin 1: A restless warrior against
12
13 allergies, infections and cancer. *Int J Biochem Cell B* 42:800–804
14

15
16 Tsutsui H, Mizutani H, Nakanishi K (2011) Contribution of interleukin 18 to the
17
18 development of infection-associated atopic dermatitis. *Curr Probl Dermatol*
19
20 41:93-103
21
22

23 24 25 **Supplementary Figure Legends**

26
27 **Supplementary Figure S1.** *Demonstration of equal protein loading during Western*
28
29 *blot experiments*
30

31
32 (a) Western blot analysis. To assess equal loading, membranes shown at **Figure 3b**
33
34 were re-probed using an antibody against cytochrome-C (Cyt-C). (b) Western blot
35
36 analysis. To assess equal loading, membranes shown at **Figure 4a** were re-probed
37
38 using an antibody against cytochrome-C (Cyt-C).
39
40

41
42
43 **Supplementary Figure S2.** *Long-term AA treatment does not influence proliferation*
44
45 *of SZ95 sebocytes*
46

47 Proliferation (CyQUANT) assay. Results are expressed in the percentage of the 24-hr
48
49 vehicle control (100%, solid line) as mean \pm SEM of four independent determinations.
50


51
52 Two additional experiments yielded similar results.
53
54
55
56
57
58
59
60

1
2
3 **Supplementary Figure S3. AA regulates expression of genes involved in the innate**
4 *immunity*


5
6
7 Q-PCR analysis of mRNA transcript expression profile of different selected genes
8 involved in the innate immunity. Data are presented by using $\Delta\Delta CT$ method
9 regarding glyceraldehyde-3-phosphate dehydrogenase (GAPDH) normalized mRNA
10 expressions of the vehicle control as 1 (solid line). Data are expressed as mean \pm SD
11 of three independent determinations. Two additional experiments yielded similar
12 results. * mark significant ($P<0.05$) differences compared to the vehicle control.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


a



b




Supplementary Figure S1



Supplementary Figure S2

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Supplementary Figure S3