



1 RESEARCH LETTER

2 **D-Galactose uptake is nonfunctional in the conidiospores of**
 3 ***Aspergillus niger***

4
 5
 6 Erzsébet Fekete¹, Ronald P.de Vries^{2,3}, Bernhard Seiboth⁴, Patricia A. vanKuyk^{2,5}, Erzsébet Sándor⁶,
 7 Éva Fekete¹, Benjamin Metz^{4,7}, Christian P. Kubicek⁴ & Levente Karaffa¹

8
 9 ¹Department of Biochemical Engineering, Faculty of Science and Technology, University of Debrecen, Debrecen, Hungary; ²CBS-KNAW Fungal
 10 Biodiversity Centre, Utrecht, The Netherlands; ³Microbiology and Klyuyver Centre for Genomics of Industrial Fermentation, Utrecht University,
 11 Utrecht, The Netherlands; ⁴Research Area Biotechnology and Microbiology, Institute of Chemical Engineering, TU Wien, Wien, Austria;
 12 ⁵Molecular Microbiology, Institute of Biology Leiden, Leiden University, Leiden, The Netherlands; and ⁶Institute of Food Processing, Quality
 13 Assurance and Microbiology, Faculty of Agricultural and Food Sciences and Environmental Management, University of Debrecen, Debrecen,
 14 Hungary

15
 16
 17
 18 **Correspondence:** Levente Karaffa,
 19 Department of Biochemical Engineering,
 20 Faculty of Science and Technology, University
 21 of Debrecen, H-4010, PO Box 64, Debrecen,
 22 Hungary. Tel.: +36 52 512 900 ext. 62488;
 23 fax: +36 52 533 677; e-mail:
 24 karaffa.levente@science.unideb.hu

25
 26 **Present address:** Benjamin Metz,
 27 Department of Biotechnology, Delft
 28 University of Technology and Klyuyver Centre
 29 for Genomics of Industrial Fermentation,
 30 Julianalaan 67, 2628 BC, Delft, The
 31 Netherlands

32 Received 15 October 2011; revised 2
 33 February 2012; accepted 3 February 2012.

34 DOI: 10.1111/j.1574-6968.2012.02524.x

35 Editor: Matthias Brock

36
 37 **Keywords**
 38 *Aspergillus niger*; D-galactose; Leloir pathway;
 39 uptake; germination.

40
 41
 42 **Introduction**

43 Plant cell wall polysaccharides – the most abundant
 44 organic compounds in nature – can be divided into three
 45 groups: cellulose, hemicellulose and pectin (de Vries &
 46 Visser, 2001). Three main hemicelluloses can be recog-
 47 nized: xylan, mannan and xyloglucan. Xylan contains a
 48 backbone of β-linked D-xylose residues that can be deco-
 49 rated with acetyl-, L-arabinose, D-galactose, (4-O-methyl-)
 50 D-glucuronic acid and ferulic acid. Mannan contains a
 51 β-linked D-mannose backbone that can be decorated with
 52 α- and β-linked D-galactose and, depending on the origin,

36 **Abstract**

The majority of black *Aspergilli* (*Aspergillus* section *Nigri*), including *Aspergillus niger*, as well as many other *Ascomycetes* fail to germinate on D-galactose as a sole carbon source. Here, we provide evidence that the ability of *A. niger* to transport D-galactose is growth stage dependent, being absent in the conidiospores but present in the mycelia. Despite earlier claims, we could identify galactokinase activity in growing cells and all genes of the Leloir pathway (responsible for channelling D-galactose into the EMP pathway) are well induced on D-galactose (and also on lactose, D-xylose and L-arabinose) in the mycelial stage. Expression of all Leloir pathway genes was also detectable in conidiospores, although *galE* (encoding a galactokinase) and *galD* (encoding a galactose-1-phosphate uridylyl transferase) were expressed poorly. These results suggest that the D-galactose-negative phenotype of *A. niger* conidiospores may be due to the lack of inducer uptake.

can contain single D-glucose residues interrupting the mannose main chain (referred to as glucomannan). Xyloglucan contains a β-linked D-glucose backbone that is decorated with α-linked D-xylose residues. Attached to these residues are D-galactose, L-arabinose and/or L-fucose residues. D-Galactose is the only component common to all three hemicelluloses and is also found in pectin (Pauly & Keegstra, 2010). The enzymatic hydrolysis of these polysaccharides is subject to significant industrial interest, both in the food and feed as well as the wood-manufacturing sector (Bhat, 2000). Amongst microorganisms with an ability to produce plant cell wall degrading enzymes,

fungi are by far the most interesting group. Besides certain *Trichoderma* species, black Aspergilli such as *Aspergillus niger* are the most important organisms because of their high protein secretion capacity and wide range of cell wall degrading enzyme activity (de Vries & Visser, 2001).

In recent years, considerable knowledge has been accumulated on the enzyme systems and genes involved in degrading hemicelluloses to their monomers and also about the further metabolism of the hemicellulose monomers in fungi (Flippin *et al.*, 2009). With respect to D-galactose, information has been obtained in *Trichoderma reesei* (Seiboth *et al.*, 2002, 2003, 2004; Karaffa *et al.*, 2006) and *Aspergillus nidulans* (Fekete *et al.*, 2004; Christensen *et al.*, 2011). In addition to the Leloir pathway, these fungi possess a second pathway for D-galactose catabolism, which, in analogy to the L-arabinose catabolic pathway, uses reductive and oxidative reactions to convert D-galactose into D-fructose-6-phosphate (Seiboth & Metz, 2011). Although genome information from *A. niger* has shown the presence of all genes/enzymes needed to degrade D-galactose (Flippin *et al.*, 2009), only few experimental data are available on its metabolism (Mojzita *et al.*, 2011; Koivistoinen *et al.*, 2012). This may be due to the fact that with the exception of *Aspergillus brasiliensis*, D-galactose is considered a very poor carbon source for black Aspergilli including *A. niger* (Meijer *et al.*, 2011), which hampers efforts to cultivate it on D-galactose. Growth on D-galactose containing complex carbohydrates may also be affected, depending on which other carbon sources are present and the ratio of these and galactose in the carbohydrate. The aim of this study was to analyse and understand the physiological background of this phenomenon in *A. niger*.

Materials and methods

Fungal strain and cultivation conditions

Aspergillus niger N402 (FGSC A733; *cspA1*) was used in this study (Bos *et al.*, 1988). The conditions for strain maintenance have been described earlier as has the MM medium used for shake-flask cultivations (de Vries *et al.*, 2004). Carbon sources were used in 1% (w/v) final concentrations and are given at the respective results.

Batch cultures were incubated on a rotary shaker (INFORS HT Multitron; 250 r.p.m.) at 30 °C in 500 mL Erlenmeyer flasks containing 100 mL of medium. Mycelia were pregrown in MM containing glycerol as a carbon source, harvested after 24 h by filtration on a sintered glass funnel, washed with cold sterile tap water and then transferred into fresh MM without glycerol, but supple-

mented with other carbon sources. For transcript analysis, samples were taken after 6 h of further incubation.

Aspergillus niger conidiospores are not formed on D-galactose containing solid medium. As a consequence, except where noted otherwise, we used glycerol as a sole carbon source to conidiate *A. niger* in the experiments aimed at investigating conidial stage events.

Nucleic acid isolation and hybridization

Fungal mycelia or conidia were harvested by filtration, washed with distilled water, frozen and ground under liquid nitrogen. For nucleic acid extraction, the Wizard Genomic DNA Purification Kit and SV Total RNA Isolation System (Promega) were used. Standard methods were used for electrophoresis, blotting and hybridization of nucleic acids (Sambrook *et al.*, 1989). Northern analysis was performed with the PCR DIG Probe Synthesis kit (Roche). 5.5 µg of RNA, respectively, was loaded into each lane. Primers for probe amplifications are given in Table 1.

Analytical methods

Mycelial dry mass was determined by withdrawing 2 × 5 mL aliquots from the culture, suction filtration through a preweighted glass wool filter and drying in an oven at 80 °C until constant weight. Data were averaged and deviated by not more than 14%.

The concentration of D-galactose in the growth medium was determined by HPLC analysis, using an H⁺ exchange column (Bio-Rad Aminex HPX-H⁺), employing 10 mM H₂SO₄ at 55 °C as mobile phase with isocratic elution and a refractive index detection.

To determine the galactokinase activity, an HPLC method was used (Fekete *et al.*, 2002). Specific galactokinase activities are reported as mg protein, which was determined by means of a modification of the method of Lowry (Peterson, 1983) using BSA for calibration.

Measurement of D-galactose uptake

Mycelia were pregrown for 18 h on glycerol as a carbon source, harvested by gentle filtration and resuspended in 20 mL of carbon-free medium (MM) to give a final density of 1 mg mL⁻¹. MM was inoculated with 10⁶, 10⁷ and 10⁹ spores mL⁻¹, respectively, when the D-galactose uptake of conidiospores were tested. After incubation at 30 °C for 60 min, 13.63 µL (0.2 mCi mL⁻¹) of D-galactose-1-¹⁴C (G3143-14C; Sigma) was added to give 100 000–150 000 dpm per mL culture, and a further amount of cold D-galactose was added to give a final concentration of 1 mM. The cultures were incubated for

Table 1. Primers used for the amplification of the genes of the Leloir pathway in *Aspergillus niger*

Gene	Activity function	Locus	Oligonucleotide sequence	Amplicon size (bp)
<i>galE</i>	Galactokinase	An16g04160	galEforw: 5'-TCTTGATTGGCAGTAGTGTC-3' galErev: 5'-CCTCAACCTGGGCAATAG-3'	1085
<i>galD</i>	Galactose-1-phosphate uridylyltransferase	An02g03590	galDforw: 5'-ACCCATTAGAGAGGAGC-3' galDrev: 5'-TTGCTCGGGTGAATGTC-3'	1151
<i>galF</i>	UTP-hexose-1-phosphate uridylyltransferase	An12g00820	galFforw: 5'-ATAACTTCTTCGCCCTCTCC-3' galFrev: 5'-CCTCACCTTCTTGTGAGC-3'	906
<i>galG</i>	UDP-galactose 4-epimerase	An14g03820	galGforw: 5'-ATTGACAGCGTTATCCAC-3' galGrev: 5'-GGGTCATTATTTCTGTC-3'	1053
<i>gpmB</i>	Phosphoglucomutase	An02g07650	gpmBforw: 5'-ACCACCATCGGCTCCAAGAC-3' gpmBrev: 5'-TAGGTGCGGACATCAGGCTC-3'	1175

further 6 h, and 1.0 mL of samples withdrawn in intervals of 30 or 60 min by immediately pipetting them into 1 mL of 1 M D-galactose and vigorous shaking. This treatment stopped further uptake of labelled D-galactose and also released ^{14}C -D-galactose adsorbed to the cells. The suspensions were then spun down in an Eppendorf centrifuge, and the radioactivity in the supernatant measured in a liquid scintillation counter (Wallac, Model 1409). OptiPhase HiSafe 3 (PerkinElmer) was used as a liquid scintillation cocktail.

Chemicals

All chemicals were of analytical grade, and, except where noted otherwise, were purchased from Sigma-Aldrich Kft., Budapest, Hungary.

Reproducibility

All the results presented here are means of 3–5 independent experiments. The data were analysed and visualized by Sigmaplot (Jandel Scientific), and standard deviations (SDs) for each procedure were determined. The SD values were always < 14% of the means.

Results

Growth of *A. niger* on D-galactose

Conidiospores of *A. niger* were unable to germinate in submerged minimal medium with 1% D-galactose as a sole carbon source even after a prolonged incubation. Essentially similar results were obtained on solid medium. However, mycelia of *A. niger* pregrown on glycerol (or on any other carbon source tested such as D-glucose, peptone, L-arabinose, D-xylose) and transferred to fresh medium containing D-galactose as a sole carbon source were able to grow, although at a rate lower than other fungi such as *A. nidulans* (Fekete *et al.*, 2004) or *T. reesei* (Seiboth *et al.*, 2004) (Fig. 1).




Fig. 1. Time-profile of biomass formation (○) and D-galactose consumption (●) of *Aspergillus niger* mycelia preformed on glycerol and transferred to minimal medium with D-galactose as a sole carbon source.

Uptake of D-galactose in *A. niger*

The above results suggested that *A. niger* can grow on D-galactose once the spores have germinated but its conidiospores fail to do so. This suggested to us that transport of D-galactose into the conidia may be nonfunctional. To investigate this hypothesis in more detail, we incubated mycelia and conidiospores, respectively, with ^{14}C -labelled D-galactose, and followed its uptake into the cells. Uptake by mycelia was related to dry weight. As it was practically impossible to determine biomass data for conidiospores in a reproducible way, we could not specify ^{14}C -labelled D-galactose uptake on the same basis in these two sets of experiments. Instead, we employed three different concentrations of conidia, namely 10^6 , 10^7 and 10^9 spores mL^{-1} , respectively, under identical experimental conditions. Any D-galactose uptake was therefore expected to be proportional to the number of conidiospores present in the medium. Data obtained indeed showed that the mycelia preformed on glycerol were able to transport D-galactose (Fig. 2a). On the other hand, there was no ^{14}C -labelled D-galactose uptake by the conidiospores irrespective of their concentration (Fig. 2b), indicating the absence of D-galactose transport at this stage of growth in *A. niger*.




Fig. 2. (a) Time-course of extracellular ^{14}C -labelled D-galactose levels in *Aspergillus niger* culture supernatant inoculated with mycelium at a density of 1 mg mL^{-1} . Mycelium was pregrown on glycerol and biomass stayed constant during the 6 h of incubation. (b) Time-course of ^{14}C -labelled D-galactose uptake of *A. niger* conidiospores. (●), 10^6 spores mL^{-1} ; (○), 10^7 spores mL^{-1} ; (▲), 10^9 spores mL^{-1} .

Aspergillus niger is able to phosphorylate D-galactose

The D-galactose-negative phenotype of *A. niger* was earlier speculated to be the consequence of a lack of galactokinase activity (Elshafei & Abdel-Fatah, 2001). In contrast, cell-free extracts of *A. niger* mycelia prepared by us were able to phosphorylate D-galactose, resulting in a specific galactokinase activity similar in value to that of *A. nidulans* (Ilyés *et al.*, 2004). Galactokinase activity showed a basically constitutive nature, although it was higher during growth on pentoses (notably on D-xylose) and D-galactose than on D-glucose (Table 2). This finding is corroborated by the fact that the genome of *A. niger* contains a locus (An16g04160; *galE*) with obvious similarity to other fungal galactokinases (Flipphi *et al.*, 2009). Northern analysis performed with the respective gene as a probe showed that the gene was transcribed on all carbon sources investigated. Expression on D-galactose was higher than on D-glucose or glycerol, however, lower than on L-arabinose or D-xylose (Fig. 3a).

Aspergillus niger has an intact Leloir pathway

The finding that galactokinase was active prompted us to study whether a full Leloir pathway is operating in *A. niger*. *In silico* data revealed that the *A. niger* genome contains orthologs for each gene of this pathway (Flipphi *et al.*, 2009). Expression studies showed that they are all

expressed in a fashion similar to galactokinase, for example, transcripts were formed on all carbon sources studied, but their transcript levels were higher on pentoses (L-arabinose, D-xylose) and on D-galactose (Fig. 3a). The reason for the higher expression of Leloir pathway genes on L-arabinose and D-xylose than on D-galactose remains unclear at this point and will require further study. Most notably, however, results obtained from conidiospores formed on glycerol or D-glucose showed that while all transcripts of the Leloir pathway genes were also present in conidiospores, *galE* (encoding a galactokinase) and *galD* (encoding an UTP-galactose-1-phosphate uridylyltransferase) were very poorly expressed (Fig. 3b), indicating that the potential to convert D-galactose into an intermediate of the EMP pathway may be dependent on the growth stage of the fungus.

Discussion

Aspergillus niger has a prominent position amongst microorganisms employed in industrial biotechnology, thus it is not surprising that numerous studies have been devoted to its biology (Andersen *et al.*, 2011). However, its metabolic relationship with D-galactose remained obscure; although this hexose is a major component of hemicelluloses and pectin, whose enzymatic hydrolysis is subject to considerable industrial interest. **3**

In this article, we have provided evidence that the D-galactose-negative phenotype of *A. niger* is growth stage dependent, being complete in the conidiospores but only partial in mycelia germinated on any other carbon source. This result required that a D-galactose transporter system needs to be present in *A. niger*. In the yeast *Kluyveromyces lactis*, D-galactose and lactose transport are mediated by the same protein (Baruffini *et al.*, 2006), while in the related species *A. nidulans*, transport of these two sugars are independent (Fekete *et al.*, unpublished data). Galactose permeases from *A. niger* have not yet been identified but BLAST analysis with the *Saccharomyces cerevisiae* *GAL2* gene as a query identified at least four unknown hexose permeases that shared high similarity to Gal2p ($< e^{-120}$), that were contained in neighbouring phylogenetic clades in a phylogenetic tree, with the *S. cerevisiae* Gal2p being the basal protein (Fekete *et al.*, unpublished). Their function is currently investigated by us. In any case, it is clear from our experiments, however, that the transport of **4**

Table 2. Intracellular galactokinase activity as a function of carbon source in *Aspergillus niger*

	Carbon source					
	D-galactose	Lactose	L-arabinose	Glycerol	D-glucose	D-xylose
Galactokinase activity (U per mg protein)	0.28 ± 0.014	0.19 ± 0.012	0.20 ± 0.01	0.11 ± 0.006	0.08 ± 0.05	0.31 ± 0.015




Fig. 3. Expression analysis of the genes comprising the Leloir pathway in *Aspergillus niger* on different carbon sources. (a) Expression in mycelia and (b) expression in conidiospores. Gly, glycerol; Glu, D-glucose; Gal, D-galactose; Ara, L-arabinose; Xyl, D-xylose; Lac, lactose. 18S rRNA was used as control.

D-galactose is not functional in the conidiospores of *A. niger*. While the reason for this unknown, our data suggest that D-galactose uptake in *A. niger* is growth stage dependent; for example, it is expressed in mycelia but not in resting conidia, resembling the behaviour of certain permeases from *T. reesei* (Metz *et al.*, 2011) and *A. nidulans* (Tazebay *et al.*, 1997; Amillis *et al.*, 2004; Pantazopoulou *et al.*, 2007).

D-Galactose metabolism via the Leloir pathway is a ubiquitous trait in pro- and eukaryotic cells (Frey, 1996). It involves an ATP-dependent galactokinase (EC 2.7.1.6) to form D-galactose 1-phosphate, which is subsequently transferred to UDP-glucose in exchange with D-glucose 1-phosphate by D-galactose 1-phosphate uridylyltransferase (EC 2.7.7.12). The resulting UDP-galactose is a substrate for the reaction catalysed by UDP-galactose 4-epimerase (EC 5.1.3.2), resulting in UDP-glucose. While we did not determine specific enzyme activities apart from that of galactokinase, gene expression data strongly suggest that the Leloir pathway is readily available to convert D-galactose once this sugar is inside of the cell, which occurs only in the mycelial stage of *A. niger*. In the conidiospore stage, however, expression of the genes encoding the first two enzymes of the Leloir pathway was hardly detected, and weak expression was observed for the other three genes of the pathway as well. As we demonstrated that the conidia are unable to transport D-galactose, we conclude that the D-galactose-negative phenotype of the *A. niger* is unlikely to be caused by a lack of D-galactose

catabolism. Rather, the phenomenon seems to be mainly uptake related in conidiospores. Therefore, the reduced expression observed for the Leloir genes in conidiospores may be due to the lack of inducer (D-galactose) uptake and appears to be a secondary effect rather than the cause of the nongrowth phenotype. Future studies will address this in more detail.

Acknowledgements

The project was carried out in the framework of an Austrian-Hungarian Intergovernmental Science & Technology Cooperation Programme (AT-18/2007). Research at the University of Debrecen was supported by the Hungarian Scientific Research Fund (OTKA; K67667 and K1006600) and the National Office for Research and Technology (NKTH; A2-2006-0017). E.F. is supported by a Bolyai János Research Scholarship (BO/00519/09/8). B.S. was supported by the Austrian Science Foundation (P19421).

References

- Amillis S, Cecchetto G, Sophianopoulou V, Koukaki M, Scazzocchio C & Djalilinas G (2004) Transcription of purine transporter genes is activated during the isotropic growth phase of *Aspergillus nidulans* conidia. *Mol Microbiol* **52**: 205–216.
- Andersen MR, Salazar MP, Schaap PJ *et al.* (2011) Comparative genomics of citric-acid-producing *Aspergillus*

- niger* ATCC 1015 versus enzyme-producing CBS 513.88. *Genome Res* **21**: 885–897.
- Baruffini E, Goffrini P, Donnini C & Lodi T (2006) Galactose transport in *Kluyveromyces lactis*: major role of the glucose permease Hgt1. *FEMS Yeast Res* **6**: 1235–1242.
- Bhat MK (2000) Cellulases and related enzymes in biotechnology. *Biotechnol Adv* **18**: 355–383.
- Christensen U, Gruben BS, Madrid S, Mulder H, Nikolaev I & de Vries RP (2011) Unique regulatory mechanism for D-galactose utilization in *Aspergillus nidulans*. *Appl Environ Microbiol* **77**: 7084–7087.
- de Vries RP & Visser J (2001) *Aspergillus* enzymes involved in degradation of plant cell wall polysaccharides. *Microbiol Mol Biol Rev* **65**: 497–522.
- de Vries HP, Burgers K, van de Vondervoort PJI, Frisvad JC, Samson RA & Visser J (2004) A new black *Aspergillus* species, *A. vadensis*, is a promising host for homologous and heterologous protein production. *Appl Environ Microbiol* **70**: 3954–3959.
- Elshafei AM & Abdel-Fatah OM (2001) Evidence for a non-phosphorylated route of galactose breakdown in cell-free extracts of *Aspergillus niger*. *Enzyme Microb Technol* **29**: 76–83.
- Fekete E, Karaffa L, Sándor E, Seiboth B, Biró S, Szentirmai A & Kubicek CP (2002) Regulation of formation of the intracellular beta-galactosidase activity of *Aspergillus nidulans*. *Arch Microbiol* **179**: 7–14.
- Fekete E, Karaffa L, Sándor E, Bányai I, Seiboth B, Gyémánt G, Sepsi A, Szentirmai A & Kubicek CP (2004) The alternative D-galactose degrading pathway of *Aspergillus nidulans* proceeds via L-sorbose. *Arch Microbiol* **181**: 35–44.
- Flipphi M, Sun J, Robellet X, Karaffa L, Fekete E, Zeng AP & Kubicek CP (2009) Biodiversity and evolution of primary carbon metabolism in *Aspergillus nidulans* and other *Aspergillus* spp. *Fungal Genet Biol*, **46**(suppl 1): 19–44.
- Frey PA (1996) The Leloir pathway: a mechanistic imperative for three enzymes to change the stereochemical configuration of a single carbon in galactose. *FASEB J* **10**: 461–470.
- Ilyés H, Fekete E, Karaffa L, Fekete E, Sándor E, Szentirmai A & Kubicek CP (2004) CreA-mediated carbon catabolite repression of beta-galactosidase formation in *Aspergillus nidulans* is growth rate dependent. *FEMS Microbiol Lett* **235**: 147–151.
- Karaffa L, Fekete E, Gamauf C, Szentirmai A, Kubicek CP & Seiboth B (2006) D-Galactose induces cellulase gene expression in *Hypocrea jecorina* at low growth rates. *Microbiology* **152**: 1507–1514.
- Koivistoinen OM, Richard P, Penttilä M, Ruohonen L & Mojzita D (2012) Sorbitol dehydrogenase of *Aspergillus niger*, SdhA, is part of the oxido-reductive D-galactose pathway and essential for D-sorbitol catabolism. *FEBS Lett* **6**: ???–???. doi:10.1016/j.febslet.2012.01.004.
- Meijer M, Houbraeken JA, Dalhuijsen S, Samson RA & de Vries RP (2011) Growth and hydrolase profiles can be used as characteristics to distinguish *Aspergillus niger* and other black *aspergilli*. *Stud Mycol* **69**: 19–30.
- Metz B, Seidl-Seiboth V, Haarmann T, Kopchinskiy A, Lorenz P, Seiboth B & Kubicek CP (2011) Expression of biomass degrading enzymes is a major event during conidia development in *Trichoderma reesei*. *Eukaryot Cell* **10**: 1527–1535.
- Mojzita D, Koivistoinen OM, Maaheimo H, Penttilä M, Ruohonen L & Richard P (2011) Identification of the galactitol dehydrogenase, LadB, that is part of the oxido-reductive D-galactose catabolic pathway in *Aspergillus niger*. *Fungal Genet Biol* **7**: ???–???. doi:10.1016/j.fgb.2011.11.005.
- Pauly M & Keegstra K (2010) Plant cell wall polymers as precursors for biofuels. *Curr Opin Plant Biol* **13**: 305–312.
- Peterson GL (1983) Determination of total protein. *Methods Enzymol* **91**: 86–105.
- Sambrook J, Fritsch EF & Maniatis T (1989) *Molecular Cloning: A Laboratory Manual*, 2nd edn. Cold Spring Harbor Laboratory Press, Plainview, NY.
- Seiboth B & Metz B (2011) Fungal arabinan and L-arabinose metabolism. *Appl Microbiol Biotechnol* **89**: 1665–1673.
- Seiboth B, Hofmann G & Kubicek CP (2002) Lactose metabolism and cellulase production in *Hypocrea jecorina*: the *gal7* gene, encoding galactose-1-phosphate uridylyltransferase, is essential for growth on galactose but not for cellulase induction. *Mol Genet Genomics* **267**: 124–132.
- Seiboth B, Hartl L, Pail M & Kubicek CP (2003) D-xylose metabolism in *Hypocrea jecorina*: loss of the xylitol dehydrogenase step can be partially compensated for by *lad1*-encoded L-arabinitol-4-dehydrogenase. *Eukaryot Cell* **2**: 867–875.
- Seiboth B, Hartl L, Pail M, Fekete E, Karaffa L & Kubicek CP (2004) The galactokinase of *Hypocrea jecorina* is essential for cellulase induction by lactose but dispensable for growth on D-galactose. *Mol Microbiol* **51**: 1015–1025.
- Tazebay UH, Sophianopoulou V, Scazzocchio C & Diallinas G (1997) The gene encoding the major proline transporter of *Aspergillus nidulans* is upregulated during conidiospore germination and in response to proline induction and amino acid starvation. *Mol Microbiol* **24**: 105–117.

Author Query Form

Journal: FEMSLE

Article: 2524

Dear Author,

During the copy-editing of your paper, the following queries arose. Please respond to these by marking up your proofs with the necessary changes/additions. Please write your answers on the query sheet if there is insufficient space on the page proofs. Please write clearly and follow the conventions shown on the attached corrections sheet. If returning the proof by fax do not write too close to the paper's edge. Please remember that illegible mark-ups may delay publication.

Many thanks for your assistance.


Query reference	Query	Remarks
1	AUTHOR: Seiboth and Metz, 2010 has been changed to Seiboth and Metz, 2011 so that this citation matches the Reference List. Please confirm that this is correct.	
2	AUTHOR: Bos <i>et al.</i> , 1988 has not been included in the Reference List, please supply full publication details.	
3	AUTHOR: Please check and confirm that whether the edit made in the sentence " However, its metabolic relationship...considerable industrial interest." retains the intended meaning.	
4	AUTHOR: Please provide all author names along with their initials for unpublished data.	
5	AUTHOR: Pantazopoulou <i>et al.</i> , 2007 has not been included in the Reference List, please supply full publication details.	
6	AUTHOR: Please provide the volume number, page range for reference Koivistoinen <i>et al.</i> (2012).	
7	AUTHOR: Please provide the volume number, page range for reference Mojzita <i>et al.</i> (2011).	

USING e-ANNOTATION TOOLS FOR ELECTRONIC PROOF CORRECTION


Required software to e-Annotate PDFs: Adobe Acrobat Professional or Adobe Reader (version 8.0 or above). (Note that this document uses screenshots from Adobe Reader X)

The latest version of Acrobat Reader can be downloaded for free at: <http://get.adobe.com/reader/>

Once you have Acrobat Reader open on your computer, click on the [Comment](#) tab at the right of the toolbar:



This will open up a panel down the right side of the document. The majority of tools you will use for annotating your proof will be in the [Annotations](#) section, pictured opposite. We've picked out some of these tools below:



1. Replace (Ins) Tool – for replacing text.




Strikes a line through text and opens up a text box where replacement text can be entered.

How to use it

- Highlight a word or sentence.
- Click on the [Replace \(Ins\)](#) icon in the Annotations section.
- Type the replacement text into the blue box that appears.

standard framework for the analysis of microeconomics. Nevertheless, it also led to the emergence of a new paradigm of strategic behaviour. The number of competitors in the industry is that the structure of the industry is a key component of the main components of the industry. At the micro level, are exogenous variables important works on entry by Shiraz (M henceforth) we open the 'black b



2. Strikethrough (Del) Tool – for deleting text.



Strikes a red line through text that is to be deleted.

How to use it

- Highlight a word or sentence.
- Click on the [Strikethrough \(Del\)](#) icon in the Annotations section.

there is no room for extra profits and the number of competitors are zero and the number of competitors (net) values are not determined by the number of competitors. Blanchard and ~~Kiyotaki~~ (1987), perfect competition in general equilibrium. The effects of aggregate demand and supply in the classical framework assuming monopoly power are an exogenous number of firms

3. Add note to text Tool – for highlighting a section to be changed to bold or italic.



Highlights text in yellow and opens up a text box where comments can be entered.

How to use it

- Highlight the relevant section of text.
- Click on the [Add note to text](#) icon in the Annotations section.
- Type instruction on what should be changed regarding the text into the yellow box that appears.

dynamic responses of mark ups consistent with the **VAR** evidence

sation... y Ma... and... on n... to a... on... stent also with the demand-



4. Add sticky note Tool – for making notes at specific points in the text.




Marks a point in the proof where a comment needs to be highlighted.

How to use it

- Click on the [Add sticky note](#) icon in the Annotations section.
- Click at the point in the proof where the comment should be inserted.
- Type the comment into the yellow box that appears.

and supply shocks. Most of the... a... number... dard fr... cy. Nev... ole of st... ber of competitors and the imp... is that the structure of the secto



USING e-ANNOTATION TOOLS FOR ELECTRONIC PROOF CORRECTION


5. Attach File Tool – for inserting large amounts of text or replacement figures.



Inserts an icon linking to the attached file in the appropriate place in the text.

How to use it

- Click on the [Attach File](#) icon in the Annotations section.
- Click on the proof to where you'd like the attached file to be linked.
- Select the file to be attached from your computer or network.
- Select the colour and type of icon that will appear in the proof. Click OK.



6. Add stamp Tool – for approving a proof if no corrections are required.




Inserts a selected stamp onto an appropriate place in the proof.

How to use it

- Click on the [Add stamp](#) icon in the Annotations section.
- Select the stamp you want to use. (The [Approved](#) stamp is usually available directly in the menu that appears).
- Click on the proof where you'd like the stamp to appear. (Where a proof is to be approved as it is, this would normally be on the first page).

of the business cycle, starting with the
 on perfect competition, constant re
 production. In this environment goods
 extra profits and the market for marke
 he market for goods is determined by the model. The New-Key
 otaki (1987), has introduced produc
 general equilibrium models with nomin
 and... Most of this literature




7. Drawing Markups Tools – for drawing shapes, lines and freeform annotations on proofs and commenting on these marks.

Allows shapes, lines and freeform annotations to be drawn on proofs and for comment to be made on these marks..

How to use it

- Click on one of the shapes in the [Drawing Markups](#) section.
- Click on the proof at the relevant point and draw the selected shape with the cursor.
- To add a comment to the drawn shape, move the cursor over the shape until an arrowhead appears.
- Double click on the shape and type any text in the red box that appears.



For further information on how to annotate proofs, click on the [Help](#) menu to reveal a list of further options:

