



Characterization of the ionic liquid obtained by chlorosulfonation of 1-methylimidazole: 1-methyl-3-sulfonic acid imidazolium chloride, 1-methylimidazolium chlorosulfate or a zwitterionic salt?

Béla Urbán^a, Gábor Szalontai^b, Máté Papp^{a,1}, Csaba Fehér^a, Attila C. Bényei^c, Rita Skoda-Földes^{a,*}

^a Institute of Chemistry, Department of Organic Chemistry, University of Pannonia, Egyetem u. 10. (P.O.Box 158), H-8200 Veszprém, Hungary

^b NMR Laboratory, University of Pannonia, Egyetem u. 10, P.O.Box 158, H-8200 Veszprém, Hungary

^c Laboratory for X-ray Diffraction, University of Debrecen, 4002 Debrecen, P.O.Box 400, Hungary

ARTICLE INFO

Article history:

Received 1 December 2020

Accepted 3 January 2021

Available online 5 January 2021

Keywords:

Brønsted acid

Chlorosulfate

Ionic liquids

NMR spectroscopy

X-ray diffraction

ABSTRACT

A great number of organic reactions catalyzed by the ionic liquid product of chlorosulfonation of 1-methylimidazole have been reported recently. At the same time controversial assumptions have appeared on the real structure of the catalyst. In the present report the primarily formed chlorosulfonation product is proved to be 1-methylimidazolium chlorosulfate ([HMim]⁺[SO₃Cl][−]) instead of 1-methyl-3-sulfonic acid imidazolium chloride, reported previously. The former structure is confirmed by X-ray crystallography and NMR spectroscopy, including ¹H-, ¹³C-, ¹⁷O- and ¹⁵N-¹H HSQC measurements. ¹H and ¹⁷O NMR experiments support fast hydrolysis of [HMim][SO₃Cl] resulting in the formation of [HMim][HSO₄] in the presence of traces of water.

© 2021 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Acid-catalyzed reactions involve one of the most important technologies applied in the chemical industry. The most common catalysts are mineral acids, used in homogeneous phase reactions. Although they usually show high catalytic activity, they often suffer from some drawbacks, such as the problems of side reactions, large amounts of acidic wastes and corrosion of the equipment. Isolation of the products can also be tedious, which can cause environmental problems. As a result, great efforts have been devoted to the development of new catalysts. One possibility is the application of solid acids which, as non-volatile materials, are less harmful than traditional liquid acids [1]. However, these heterogeneous systems often show an inferior activity when compared to their homogeneous counterpart. As another approach, increasing attention has been paid to the application of task specific ionic liquids as catalysts [2]. Especially acidic ionic liquids were found to be useful and even recyclable alternatives to conventional Brønsted or Lewis acid catalysts [3]. Among them, ionic liquids (ILs) incorporating imidazolium cations with *N*-alkyl sulfonic groups serve as efficient Brønsted acids in a great variety of organic transformations [4]. Furthermore, they can be converted into -SO₃H functionalized dual Brønsted-Lewis acidic ILs possessing complex

metal halide anions as the Lewis acidic site [5]. The first example for a Brønsted acidic IL with an SO₃H moiety attached directly to imidazolium nitrogen, 1-methyl-3-sulfonic acid imidazolium chloride ([MSim]Cl, **1**, Fig. 1), was reported by Zolfigol et al. in 2010 [6]. Since then, the catalytic activity of ILs incorporating the [MSim]⁺ cation (**1**) and different anions [7] was tested in a wide range of organic reactions. At the same time, no detailed structural data are available for these compounds. Only ¹H and ¹³C NMR spectra were reported [6] that do not give sufficient information about the position of the SO₃ moiety. Recently, Shiflett et al. found that chlorosulfonation of 1-methylimidazole led to a mixture of two compounds, an overall neutral zwitterion (**2**) as the main product and another derivative that was claimed to be **1** [8]. The formation of the former was proved by crystallography but beyond that only ¹H and ¹³C spectra of the mixture of the two compounds were presented. Based on ¹H NMR data, it was supposed that compound **2** could be converted to **1** when an equimolar amount of HCl was added [8]. It should also be mentioned that Zolfigol described the product as a viscous oil [6] while the other group obtained a solid material and they recorded a melting point of 325.8 °C for the zwitterionic component **2** [8].

As part of our ongoing interest in the application of acidic IL catalysts in ring opening of epoxides [9] as well as oligomerization of alkenes [10], we decided to explore the activity of ILs with *N*-SO₃H moieties. Before that the clarification of the somewhat contradictory results seemed to be necessary that prompted us to carry out a more detailed investigation of the chlorosulfonation product of 1-methylimidazole.

* Corresponding author.

E-mail address: skodane@almos.uni-pannon.hu (R. Skoda-Földes).

¹ Present address: Institute of Chemistry ELTE Eötvös Loránd University, Budapest, Hungary.




Fig. 1. The structure of the product of chlorosulfonation of 1-methylimidazole, suggested by Zolfigol et al. [6] ([MSim]Cl, **1**) and the zwitterionic structure **2** reported by Shiflett et al. [8].

2. Experimental

2.1. Computational methods

Theoretical calculations were performed with the Gaussian 09 software package [11]. Molecular geometries were optimized using the long-range corrected hybrid density functional CAM-B3LYP [12] and the standard 6-311++G** basis set. The optimizations were performed without any symmetry constraint. The vibrational analysis yielded no imaginary frequencies verifying that each of the optimized structures is a true minimum on the potential energy surface. The Gibbs free energies were refined using the CPCM solvent model (with Solvent=dichloromethane) built in the Gaussian 09 suite.

2.2. Synthetic methods

Chlorosulfonation of 1-methylimidazole was carried out based on the procedure reported by Zolfigol [6].

Method A: Chlorosulfonic acid (1.05 ml, 15.6 mmol) was added dropwise into a room temperature solution of 1-methylimidazole (1.2 ml, 15 mmol) in 150 ml dry CH₂Cl₂. The mixture was stirred overnight and then left to stand for one hour. After that the CH₂Cl₂ was decanted. The residue was washed with dry CH₂Cl₂ (3 × 50 ml) and dried in vacuum to obtain a viscous oil that solidified upon standing two weeks under strictly inert conditions.

Method B: Chlorosulfonic acid (1.05 ml, 15.6 mmol) was added dropwise into a cooled solution of 1-methylimidazole (1.2 ml, 15 mmol) in 150 ml dry CH₂Cl₂. The mixture was stirred for 20 min and left to stand a further 20 min to obtain a colorless solid precipitate. After that the CH₂Cl₂ was decanted. The residue was washed with dry CH₂Cl₂ (3 × 50 ml) and dried in vacuum.

[HMim][HSO₄] (**4**) was prepared by the reaction of 1-methylimidazole and H₂SO₄. Concentrated sulfuric acid (1.34 ml, 25.1 mmol) was added dropwise to 2 ml (25.1 mmol) 1-methylimidazole under vigorous stirring in an ice bath. After warming to room temperature it was stirred for 48 h and then was washed with 5 × 5 ml diethyl ether and was dried in vacuum at 70 °C for 10 h.

[HMim]Cl was prepared by the reaction of 1-methylimidazole and HCl. HCl was liberated from the reaction of NaCl and H₂SO₄ in a separated flask connected with the vessel containing 2 ml (25.1 mmol) 1-methylimidazole. The reaction mixture was stirred at room temperature overnight, then it was washed with 5 × 10 ml diethyl ether and was dried in vacuum at 60 °C for 10 h.

2.3. X-ray diffraction analysis

X-ray quality crystals of **3** were grown from the neat sample obtained by Method A upon standing. A suitable crystal was fixed onto a Mitegen loop using high density oil. Diffraction intensity data collection was carried out using a Bruker-D8 Venture diffractometer equipped with INCOATEC IμS 3.0 dual (Cu and Mo) sealed tube micro sources and Photon II Charge-Integrating Pixel Array detector using Mo Kα (λ = 0.71073 Å) radiation. Low temperature data collection (150 K) was applied to protect the sample from traces of moisture and also because of the low melting point. High multiplicity data collection and

integration was performed using the APEX3 (Ver. 2017.3-0, Bruker AXS Inc., 2017.) software. Data reduction and multi-scan absorption correction was performed using SAINT (Ver. 8.38A, Bruker AXS Inc., 2017). The structure could be solved using direct methods and refined on F² using the SHELXL program [13] incorporated into APEX3 suite. Refinement was performed anisotropically for all non-hydrogen atoms. Hydrogen atoms were placed into geometric positions. The structure is stabilized by strong intermolecular N-H...O and weak C-H...O hydrogen bonds. The CIF file was manually edited using the publCIF software [14]. Results of X-ray diffraction structure determinations were very good according to the Checkcif of PLATON software [15] and structural parameters such as bond length and angle data are in the expected range. Further crystallographic and refinement details can be seen in the SI. Powder diffraction data were collected using the same instrument applying Cu Kα (λ = 1.54178 Å) radiation. Integration and background correction was performed with Bruker's DIFFRAC.EVA 4.2.2.3 software.

CCDC entry 2,023,770 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

2.4. NMR investigations





All NMR investigations were carried out on a Bruker Avance II 400 MHz spectrometer using a 5 mm BBO probe with temperature regulation. The relaxation delays applied were 1, 4 and 10 s for the ¹H, ¹³C, and ¹⁵N nuclei, respectively. The standard Bruker pulse sequences were used. 2–3000 scans were sufficient to record ¹⁷O NMR spectra.

3. Results and discussion

3.1. Quantum chemical calculations

Acidities of some well-known acids and different sulfonated imidazolium ions were compared based on their dissociation free energies (Table 1). Sulfonation of imidazole derivatives is reported to take place in position four or five under usual conditions for electrophilic substitutions [16]. 2-Sulfonated imidazoles were prepared by oxidation of 2-substituted thiols [17]. These sulfonic acids were shown to exist mainly as zwitterionic compounds (Fig. 2) according to the crystallographic data [16,17]. In the present case however, the formation of a cationic species is supposed so data for the proposed [6] *N*-sulfonated

Table 1
Relative dissociation free energies for different acids in CH₂Cl₂ solvent.

Entry	AH	ΔG _{diss} ΔG _{diss} [kJ/mol] ^a
1	HCl	0
2	H ₂ SO ₄	−34.2 (128.9) ^b
3	CF ₃ SO ₃ H	−59.4
4		−4.3
5		−80.6
6		−104.1
7		−126.7

^a ΔG_{diss} − ΔG_{diss}(HCl).

^b 2nd H⁺ in parenthesis.





Fig. 2. Zwitterionic structures of sulfonated imidazoles [16,17].

cation ([MSim]) (entry 7) as well as those for cationic C-4 (entry 5) and C-2 sulfonated derivatives (entry 6) were determined.

The dissociation free energies defined by eq. (1) were calculated using the cam-B3LYP density functional with the 6-311++G** basis set. The geometry optimisations and free energy calculations were performed using the conductor-like polarisable continuum model (CPCM) with CH₂Cl₂ solvent. The dissociation energy data, relative to HCl are summarized in Table 1. For the sulfonated imidazolium ions, the calculations gave surprisingly high energies (Table 1 entries 5–7) compared to the data of common acids. The value for the [MSim]⁺ cation indicates much higher acidity of this structure than that of CF₃SO₃H.



$$\Delta G_{\text{diss}} = G_{\text{A}^-} + G_{\text{H}^+} - G_{\text{AH}} \quad (1)$$

G_{A^-} : Calculated Gibbs free energy for A[−].

G_{AH} : Calculated Gibbs free energy for AH.

G_{H^+} : Gibbs free energy of H⁺ ($G_{\text{H}^+} = -26.28$ kJ/mol) [18]

Geometry optimizations were also performed for the *N*-sulfonated cation ([MSim]⁺) in the presence of chloride, hydrogensulfate or triflate anions as well as for the deprotonated form (2) with the corresponding acid (Scheme 1, Table 2). In each case, the zwitterion 2 had much lower Gibbs free energy.

These results support the assumption that the equilibrium depicted in Scheme 1 is shifted considerably to the right side for Cl[−], HSO₄[−] or OTf[−] anions. That means that the *N*-sulfonated derivative may exist mainly in the zwitterionic form 2 similarly to other sulfonated derivatives (Fig. 2). In other words, hydrogen chloride is not a sufficiently strong acid to protonate zwitterion 2, especially when they are present in equimolar amounts. It should be mentioned however that sulfuryl imidazolium salts, the ester derivatives of the proposed ILs [MSim]A, are known compounds and their synthesis and applications were reported before [19].

The presence of zwitterion 2 in the two component mixture was proved by Shiflett [8] but we felt that some more evidence was necessary to be certain about the structure of the other component. Accordingly, a more detailed investigation of the chlorosulfonation product of 1-methylimidazole was carried out. Besides the formation of the zwitterionic form 2, a simple acid-base reaction leading to the [HMim]⁺ cation (Scheme 2) was also taken into consideration. [HMim][SO₃Cl] (3) had been obtained before by the reaction of 1-methylimidazolium

Table 2

Gibbs free energy differences between 2 + HA and [MSim]A structures in dichloromethane^a.

Entry	Anion	ΔG [kJ/mol]
1	chloride	−85.8
2	hydrogensulfate	−75.4
3	triflate	−71.9

^a $G(2 + \text{HA}) - G([\text{MSim}] + \text{anion})$

hydrogensulfate ([HMim][HSO₄], 4) with SOCl₂ and was characterised by HRMS and ¹H NMR spectroscopy [20].

3.2. Preparation of ionic liquids

Chlorosulfonation of 1-methylimidazole was carried out based on the report of Zolfigol [6]. A careful addition of chlorosulfonic acid to the room temperature solution of 1-methylimidazole and stirring the mixture overnight (Method A) led to the formation of a highly viscous oil forming a separate phase. After the removal of the dichloromethane phase, the material solidified upon standing two weeks under strictly anhydrous conditions and even crystals suitable for X-ray investigations could be isolated. In another procedure, chlorosulfonic acid was added dropwise to a cooled solution of 1-methylimidazole in dichloromethane (Method B). In this case, a colorless solid precipitated almost immediately that was separated after stirring the mixture for 20 min.

3.3. X-ray measurements


Crystals degraded rapidly upon warming so diffraction data had to be collected at low temperature (150 K). The compound crystallized in the P2₁/c monoclinic space group. Crystallographic data are summarized in Table 3. The results revealed the formation of an imidazolium salt with a ClSO₃[−] anion: ([HMim][ClSO₃] (3) Scheme 2, Fig. 3). It should be mentioned that to the best of our knowledge, X-ray data for only two structures incorporating the ClSO₃[−] anion were reported before [21].

Comparison of the PXRD of the bulk sample shows that all the bulk crystalline material is the same as the single crystal salt structure 3 (Fig. S3).


3.4. NMR investigations

Next, detailed NMR investigations [22] of the materials obtained by the two different methods (A and B) were carried out.

NMR data of a freshly prepared neat sample of the ionic liquid obtained by Method A supported the X-ray structure and proved the formation of 3. It is noteworthy that because of the peculiar properties, such as the extreme viscosity or high polarity or low melting point of the ionic liquids, recording and interpretation of spectra require special attention [23]. In the ¹H NMR spectrum of the neat sample (Fig. 4) a one-proton signal could be found at 12.7 ppm and three more with 1: 2: 3 integral ratios at 8.41 (2-H), 7.11 (4-H, 5-H), and 3.57 (N-CH₃)



Scheme 1. Possible deprotonation the *N*-sulfonated cation ([MSim]⁺) in the presence of anions



Scheme 2. Possible products of the reaction of 1-methylimidazole with chlorosulfonic acid

ppm, respectively. Unlike the 4-H, 5-H proton signals, the corresponding carbons are clearly resolved in the $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum (Fig. 5).

The high chemical shift of the NH proton (12.7 ppm) can be explained by the existence of a hydrogen bond between this proton and one of the oxygen atoms of the SO_3Cl^- anion in the neat sample, similarly to the solid structure (Fig. S2).

It should also be mentioned that another set of signals can be seen in the ^1H and $^{13}\text{C}\{^1\text{H}\}$ NMR spectra with a relative integral ratio of about 1/10 (Fig. 4) that shows the presence of some minor component(s).

The two nitrogens of the imidazole ring gave completely overlapping signals at 170.7 ppm in the inverse-gated $^{15}\text{N}\{^1\text{H}\}$ NMR spectrum (Fig. 6). It is important to notice that this experiment is not directly influenced by a protonation – deprotonation exchange [22].

pH-dependence of ^{15}N NMR shifts of 1-methylimidazole had been reported earlier [24]. While an approximately 80 ppm difference was observed between the $^{15}\text{N}_1$ and $^{15}\text{N}_3$ signals under basic conditions (corresponding to 1-methylimidazole), a single peak had been detected for the two nitrogens in acidic solutions at $\text{pH} < 5$, where the 1-methylimidazolium cation is in excess. So the overlap of the signals in the present case, though somewhat unexpected, may provide another proof for the formation of the $[\text{HMim}]^+$ cation.

Table 3
Summary of crystallographic data for compound 3.

Crystal data	
Chemical formula	$\text{C}_4\text{H}_7\text{N}_2 \cdot \text{ClO}_3\text{S}$
M_r	198.63
Crystal system, space group	Monoclinic, $P2_1/c$
Temperature (K)	150
a, b, c (Å)	5.9038 (3), 18.0587 (11), 7.4778 (5)
β (°)	95.048 (2)
V (Å ³)	794.15 (8)
Z	4
Radiation type	Mo $K\alpha$
μ (mm ⁻¹)	0.70
Crystal size (mm)	0.35 × 0.26 × 0.21
Data collection	
Diffractometer	Bruker D8 VENTURE
Absorption correction	Numerical <i>SADABS2016/2</i> - Bruker AXS area detector scaling and absorption correction
$T_{\text{min}}, T_{\text{max}}$	0.78, 0.87
No. of measured, independent and observed [$I > 2\sigma(I)$] reflections	8179, 1487, 1427
R_{int}	0.019
$(\sin \theta/\lambda)_{\text{max}}$ (Å ⁻¹)	0.610
Refinement	
$R[F^2 > 2\sigma(F^2)], wR(F^2), \chi^2$	0.041, 0.139, 1.35
S	
No. of reflections	1487
No. of parameters	102
H-atom treatment	H-atom parameters constrained
$\Delta_{\text{max}}, \Delta_{\text{min}}$ (e Å ⁻³)	0.96, -0.91




Fig. 3. X-ray structure of ionic liquid ([HMim][SO_3Cl]) (3) with partial numbering scheme and selected bond distances. S–O distances are in the range of 1.433–1.439 Å. Thermal ellipsoids drawn at 50% probability.




Fig. 4. ^1H -400 MHz NMR spectrum of a neat sample of compound 3 (obtained by Method A).




Fig. 5. $^{13}\text{C}\{^1\text{H}\}$ - 100.3 MHz NMR spectrum of a neat sample of compound **3** (obtained by Method A).




Fig. 6. $^{15}\text{N}\{^1\text{H}\}$ inverse-gated- spectrum of a neat sample of compound **3** (obtained by Method A).

The proton-detected ^{15}N - ^1H through-bond (HSQC) experiment using a typical one-bond coupling value ($[1J(^1\text{H}-^{15}\text{N}) = 100 \text{ Hz}]$ [17] exhibited an extremely strong cross peak (Fig. 7) between the 12.7 ppm proton signal and the 170.7 ppm N_3 nitrogen signal, confirming that they are directly connected. However, this correlation




Fig. 7. Proton-detected ^{15}N - ^1H HSQC spectrum of a neat sample of compound **3** ($J = 110 \text{ Hz}$) (obtained by Method A).




Fig. 8. Proton-detected ^{15}N - ^1H HSQC spectrum of a neat sample of compound **3** ($J = 11 \text{ Hz}$) (obtained by Method A).

disappears, as soon as the NH protons take part in an exchange process fast on the ^1H time scale. Note that the weaker correlation seen on Fig. 8 points to the N_1 nitrogen coupled also to the 12.7 proton by a three bond coupling. These observations correspond to the presence of the $[\text{HMim}]^+$ cation.

The highly concentrated neat sample made possible also the recording of spectra of rare nuclei such as ^{17}O within reasonable accumulation time. Thus, to identify the structure of the anion, the ^{17}O NMR spectrum was also attempted at natural abundance. At room temperature it presented a broadened peak at 240.8 ppm (Fig. 9). This chemical shift can be indicative of the presence of a Cl-S-O moiety [20]. Incorporation of chlorine results in a considerable downfield shift of the ^{17}O signal. Some literature examples are as follows: $\text{CH}_3\text{SO}_2\text{OCH}_3$: 170 ppm [25], $\text{CH}_3\text{SO}_2\text{Cl}$: 238 ppm [25], $\text{CH}_3\text{OSO}_2\text{OCH}_3$: 140 ppm [26], $\text{CH}_3\text{OSO}_2\text{Cl}$: 219 ppm [26]. At the same time, δ_{O} of sulfonic acid derivatives remains far below 200 ppm (e.g. 140 ppm for p-toluenesulfonic acid monohydrate [27], 171 ppm for Cys- SO_3H) [28].) Also, an increase in the negative charge on oxygens around sulfur was found to lead to deshielding (e.g. $\delta_{\text{O}}(\text{H}_2\text{SO}_4) = 140 \text{ ppm}$, while $\delta_{\text{O}}(\text{SO}_4^{2-}) = 167 \text{ ppm}$) [25a].

The sample, stored for a period of two months in a closed bottle, displayed a bit different ^1H NMR spectrum compared to the fresh material (Fig. 10). Although the signals of the $[\text{HMim}]^+$ cation remained unaltered, some increase in the integral value of the peak at 10.8 ppm was observed. Another marked feature is the disappearance of the signals of




Fig. 9. ^{17}O -direct detection (zg) not-proton decoupled (linewidth $\sim 300 \text{ Hz}$) NMR spectrum of a neat sample of compound **3** (obtained by Method A).




Fig. 10. a) 400 MHz ^1H NMR spectra of a freshly made neat sample prepared by Method B (300 K); b) ^1H NMR spectrum of the same sample after a two-month storage (304 K).

the 1-methylimidazole moiety of the minor component. This proves that the proton with the peak at 10.8 ppm does not belong to the latter.

It can be assumed that ionic liquid **3** is extremely sensitive to moisture, similarly to chlorosulfonic acid [29] and the changes in the ^1H NMR spectrum (Fig. 10) can be attributed to partial hydrolysis. The deliberate addition of $\text{DMSO-}d_6$ containing traces of water to the fresh sample resulted in a considerable change in the integral value of the signal around 11 ppm (Fig. 11b). In a sample with higher dilution (Fig. 11c), signals corresponding to protons of the methyl group and the imidazole ring became sharper. Besides, two broad signals could clearly be distinguished from the narrow ones: one above 14 ppm, corresponding to approximately one proton, and another one at 7.5 ppm with a larger integral value (Fig. 11c). NOESY spectra indicate slow exchange between the two (Fig. S4).

Essentially the same spectra were obtained in $\text{DMSO-}d_6$ for the material prepared by Method B (Fig. S5) showing that the two methods led to the formation of the same major product. The integral value of the peak around 11 ppm decreased in the spectrum of a more concentrated sample (probably due to the lower water content) together with some downfield shift of the signal (Fig. S6). The broad singlet with the highest chemical shift moved to the other direction, in

accordance with the behaviour of the other sample (Fig. 11). Signals corresponding to a minor compound incorporating the 1-methylimidazole moiety could be distinguished again in all of these spectra, similarly to the sample prepared by Method A.

A broadening of the most de-shielded two signals (~ 14 and ~ 12 ppm) was observed when the temperature was increased to 323 K and they collapsed into a single broad peak upon further heating (Fig. S7), supporting an exchange between them. These signals disappeared when D_2O was added to the mixture and a new broad H_2O peak was detected at 4.4 ppm indicating a deuterium-proton exchange (Fig. 12). It is noteworthy that the singlets corresponding to the minor component were absent from the spectrum obtained by heating this sample to 323 K. Only four signals with the ratio of 1:1:1:3 could be distinguished at 8.63 ppm, 7.40 ppm, 7.38 ppm and 3.79 ppm, respectively, supporting the presence of a single compound.

Fast hydrolysis of the chlorosulfate anion [29] of $[\text{HMim}][\text{SO}_3\text{Cl}]$ (**3**) may lead to the formation of protons and different anions (Scheme 3).

To prove the hypothesis of the formation of HSO_4^- (or SO_4^{2-}) anions in the partially hydrolysed samples, the spectra were compared with those of $[\text{HMim}][\text{HSO}_4]$ (**4**), prepared by the reaction of 1-methylimidazole and H_2SO_4 (Figs. S8–S10).

The signals in the ^1H NMR spectrum of a neat sample of $[\text{HMim}][\text{HSO}_4]$ (**4**) (Fig. S8) is very similar to those of the main components of the partially hydrolysed samples of substances obtained by Methods A (Fig. 11b) and B (Fig. S5). An exchange between the two protons exhibiting the broad signals at 13.2 ppm and 11.0 ppm was proved by a change in the temperature of the ^1H NMR experiment (Fig. S10). This behaviour is also similar to that of the hydrolysed material (Fig. S7).

The presence of HSO_4^- (or SO_4^{2-}) anions in a hydrolysed sample could also be proved by ^{17}O NMR measurements. The chemical shift at 165.5 ppm observed for the substance obtained by Method B in the presence of water (and D_2O) (Fig. 13b) corresponded well to the ^{17}O signal of $[\text{Hmim}][\text{HSO}_4]$ (**4**) (Fig. 13c), as well as to that of KHSO_4 (163 ppm) [30], supporting the hydrolysis of the anion of **3** to form HSO_4^- ions. A similar behaviour could be noticed on comparing the ^{17}O NMR spectra of chlorosulfonic acid and a chlorosulfonic acid / water mixture. The ClSO_3^- ions immediately and fiercely form HSO_4^- (and HCl) and practically remove water from the mixture (see spectrum at Fig. S11).

It should also be mentioned that spectra depicted in Fig. 11a and Fig. S5 are very similar to that reported by Shiflett [8], except that a greater difference (major / minor ratio = 10 and 8, respectively) was observed between the amount of the major and minor components.




Fig. 11. a) 400 MHz ^1H NMR spectra of a neat sample prepared by Method A (300 K); b) ^1H NMR spectrum of the same sample after addition of 30% $\text{DMSO-}d_6$ (300 K); c) ^1H NMR spectrum of a diluted sample (70 mg / 0.4 ml $\text{DMSO-}d_6$ (300 K) (The shift of the methyl singlet in spectrum a) is adjusted to the same signals of the spectra obtained in $\text{DMSO-}d_6$ b).)





Fig. 12. 400 MHz ^1H NMR spectra of the material obtained by Method B in a mixture of $\text{DMSO-}d_6$ (0.3 ml) and D_2O (0.1 ml) a) 313 K; b) 323 K.



Scheme 3. Hydrolysis of [HMim][SO₃Cl] (**3**) in the presence of added water

Shiflett proved the formation of zwitterion **2** as one of the two products by X-ray investigations [8]. The other derivative was supposed to be [MSim]Cl (**1**), although the explanation for the appearance of two different peaks in the region of acidic protons in the ¹H NMR spectrum remained somewhat vague.

In our case, the integral values of the high-frequency shifted two protons (~13–14 ppm and ~11–12 ppm) suggests that they do not belong to the minor product.

Based on the above results, it can be supposed that the minor component in the substances prepared by Methods A or B is zwitterion **2**. For this compound, only signals corresponding to the methylimidazole core can be distinguished either in the ¹H- or in the ¹³C NMR spectra. It was prepared before [31] by the reaction of 1-methylimidazole and SO₃ and was found to lose SO₃ in the presence of water. It was also shown that it could convert alcohols to alkyl sulfates. Consequently, its hydrolysis may result in the formation of [HMim][HSO₄] (**4**), similarly to that of the main component of the material obtained by Method A, [HMim][ClSO₃] (**3**) (Scheme 3). This may give an explanation for the presence of one set of signals in the region of imidazolium protons in samples contaminated with water, corresponding to the [HMim]⁺

cation. The signal around 14 ppm can be attributed to the NH proton of the same cation, while the peak with altering intensity around 11 ppm can be assigned to the HSO₄[−] anion. That means that wet samples contain mixtures of [HMim][ClSO₃] (**3**), [HMim][HSO₄] (**4**) and, eventually, zwitterion **2**. NMR investigations of the substance obtained by Method B (Figs. S5–S7) strongly suggest that it is also a mixture of zwitterion **2**, and ionic liquids **3** and **4**.

It should be mentioned that the experiments discussed above prove that the zwitterion **2** is much less prone to hydrolysis than **3** and is converted to **4** only on heating (Fig. 12), or on prolonged interaction with water (Fig. 10).

It should be emphasized that high chemical shift of the NH proton of the [HMim]⁺ cation in neat samples can be due to the existence of hydrogen bonds, e.g. between N³H of the cation and O¹ of the anion in case of **3**, as indicated by the X-ray structure (Fig. S2). Also, the chemical shifts of NH protons of **3**, **4** and [HMim]Cl in the neat samples (Fig. S12) change in a reversed order to the acidity of the conjugate acid of the anion. This phenomenon was observed before for diethylmethylamine based protic ionic liquids with the chemical shift of NH protons ranging from 14.35 ppm to 7 ppm with increasing acid strength of the acid used during the protonation of the amine. This tendency was explained by the increase in the strength of hydrogen bond formed between the anion and the protonated base, causing a deshielding of the exchangeable proton [32].

4. Conclusions

The results presented above show that the reaction of chlorosulfonic acid and 1-methylimidazole can lead to a mixture of [HMim][SO₃Cl] (**3**) and a minor component **2** at room temperature. The primarily formed ionic liquid is very sensitive to moisture and readily hydrolyses to [HMim][HSO₄] (**4**). A slower reaction of zwitterion **2** with water leads to the same product (**4**). The formation of [MSim]Cl as a stable product of the present reaction remains to be proved. ¹H and ¹³C NMR data are not sufficient for this as NH chemical shifts (and even ¹³C and ¹⁵N NMR chemical shifts [24]) of 1-methylimidazole are subject to changes depending on the conditions (dilution, pH, etc.). Also, it can be seen that the chlorosulfonation reaction is extremely sensitive not only to moisture but also to other experimental details. The original report describes the formation of a single product [6], Shiflett obtained a mixture with two components of comparable amounts [8]. The present reactions led to the same two products but in a different ratio. Chlorosulfonic acid is prepared by the reaction of SO₃ and HCl. The reaction is reversible [33], so chlorosulfonic acid may contain some SO₃ that can lead to the formation of zwitterion **2** in an amount corresponding to the SO₃ content of the reagent.

It should be mentioned that the formation of **3** cannot be due to the lack of proper inertness of our procedure as it was shown to decompose in the presence of moisture soon. At the same time, under sufficiently inert conditions it can survive a two-month long storage.

Good catalytic activity of the chlorosulfonation product is also not a proper proof for the existence of [MSim]Cl and can also be explained by the formation of ionic liquid **3**. Bao et al. investigated the acidity of [HMim][SO₃Cl] (**3**), obtained by the reaction of SO₂Cl and [Hmim][HSO₄], and showed that the ionic liquid **3** had dual Brønsted–Lewis acidity [20]. Moreover, much superior catalytic activity of the latter salt compared to that of [Hmim][HSO₄] was observed in esterification




Fig. 13. 54.2 MHz ¹⁷O NMR spectra without proton decoupling. a) neat [HMim][SO₃Cl] (**3**) δ = 240.8 ppm, b) substance obtained by Method B in the presence of DMSO-*d*₆ and D₂O (50 μL D₂O, 298 K) δ = 165.5 ppm (HSO₄[−]), δ = 13.1 ppm (DMSO), and δ = 0.0 ppm (excess H₂O was added as reference) and c) [HMim][HSO₄] (**4**) in the presence of DMSO-*d*₆ (50 μL) δ = 163.6 ppm.

reactions. It should be mentioned, that the great majority of the catalytic processes carried out with [MSim]⁺ I⁻s [6,7] are condensations resulting in the formation of water, so hydrolysis or partial hydrolysis may take place even under strictly inert conditions in the course of the reactions. The hydrolysis product (Scheme 3) can be considered a mixture of a protic ionic liquid and an inorganic acid. Such mixtures were shown to exert improved catalytic activity over the single components [34], that could be awaited in the present case, too.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgement

The authors acknowledge the financial support of the Economic Development and Innovation Operative Program of Hungary (GINOP-2.3.2-15-2016-00053) and the National Research, Development and Innovation Office (K120014). A. C. Bényei thanks for the support of projects GINOP-2.3.2-15-2016-00008 and GINOP-2.3.3-15-2016-00004 financed by the EU and co-financed by the European Regional Development Fund.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.molliq.2021.115276>.

References

- [1] F. Liu, K. Huang, A. Zheng, F.-S. Xiao, S. Dai, Hydrophobic solid acids and their catalytic applications in green and sustainable chemistry, *ACS Catal.* 8 (2018) 372–391, <https://doi.org/10.1021/acscatal.7b03369>;
- [2] K. Wilson, J.H. Clark, Solid acids and their use as environmentally friendly catalysts in organic synthesis, *Pure Appl. Chem.* 72 (2000) 1313–1319, <https://doi.org/10.1351/pac200072071313>;
- [3] T. Okuhara, Water-tolerant solid acid catalysts, *Chem. Rev.* 102 (2002) 3641–3666, <https://doi.org/10.1021/cr0103569>.
- [4] (a) H. Olivier-Bourbigou, L. Magna, D. Morvan, Ionic liquids and catalysis: recent progress from knowledge to applications, *Appl. Catal. A* 373 (2010) 1–56, <https://doi.org/10.1016/j.apcata.2009.10.008>;
- (b) R. Ratti, Ionic liquids: synthesis and applications in catalysis, *Adv. Chem.* (2014) <https://doi.org/10.1155/2014/729842> Article ID 729842.
- [5] (a) L.C. Brown, J.M. Hogg, M. Swadźba-Kwaśny, Lewis acidic ionic liquids, *Top. Curr. Chem.* (Z) 375 (2017) 78, <https://doi.org/10.1007/s41061-017-0166-z>;
- (b) A.S. Amarasekara, Acidic ionic liquids, *Chem. Rev.* 116 (2016) 6133–6183, <https://doi.org/10.1021/acs.chemrev.5b00763>;
- (c) M. Vafaezadeh, H. Alinezhad, Brønsted acidic ionic liquids: green catalysts for essential organic reactions, *J. Mol. Liq.* 218 (2016) 95–105, <https://doi.org/10.1016/j.molliq.2016.02.017>;
- (d) W. Hui, Y. Zhou, Y. Dong, Z.J. Cao, F.Q. He, M.Z. Cai, D.J. Tao, Efficient hydrolysis of hemicellulose to furfural by novel superacid SO₃H functionalized ionic liquids, *Green Energy Environ.* 4 (2019) 49–55, <https://doi.org/10.1016/j.gjee.2018.06.002>.
- [4] P. Sarma, A.K. Dutta, R. Borah, Design and exploration of –SO₃H group functionalized Brønsted acidic ionic liquids (BAILs) as task-specific catalytic systems for organic reactions: a review of literature, *Catal. Surv. Asia* 21 (2017) 70–93, <https://doi.org/10.1007/s10563-017-9227-0>.
- [5] C. Chiappe, S. Rajamani, Structural effects on the physico-chemical and catalytic properties of acidic ionic liquids: an overview, *Eur. J. Org. Chem.* (2011) 5517–5539, <https://doi.org/10.1002/ejoc.201100432>.
- [6] M.A. Zolfigol, A. Khazaei, A.R. Moosavi-Zare, A. Zare, 3-Methyl-1-sulfonic acid imidazolium chloride as a new, efficient and recyclable catalyst and solvent for the preparation of N-sulfonyl imines at room temperature, *J. Iran. Chem. Soc.* 7 (2010) 646–651, <https://doi.org/10.1007/BF03246053>.
- [7] (a) M.A. Zolfigol, A. Khazaei, A.R. Moosavi-Zare, Abdolkarim Zare, Ionic liquid 3-methyl-1-sulfonic acid imidazolium chloride as a novel and highly efficient catalyst for the very rapid synthesis of bis(indolyl)methanes under solvent-free conditions, *Org. Prep. Proc. Int.* 42 (2010) 95–102, <https://doi.org/10.1080/00304940903585495>;
- (b) M.A. Zolfigol, A. Khazaei, A.R. Moosavi-Zare, A. Zare, V. Khakyzadeh, Rapid synthesis of 1-amidoalkyl-2-naphthols over sulfonic acid functionalized imidazolium salts, *Appl. Catal. A* 400 (2011) 70–81, <https://doi.org/10.1016/j.apcata.2011.04.013>;
- (c) A. Khazaei, M.A. Zolfigol, A.R. Moosavi-Zare, A. Zare, E. Ghaemian, V. Khakyzadeh, Z. Asgari, A. Hasaninejad, Sulfonic acid functionalized imidazolium salts/FeCl₃ as novel and highly efficient catalytic systems for the synthesis of benzimidazoles at room temperature, *Sci. Iran. C* 18 (2011) 1365–1371;
- (d) M.A. Zolfigol, H. Vahedi, S. Azimi, A.R. Moosavi-Zare, Benzoylation of aromatic compounds catalyzed by 3-methyl-1-sulfonic acid imidazolium tetrachloroaluminate and silica sulfuric acid under mild conditions, *Synlett* 24 (2013) 1113–1116, <https://doi.org/10.1055/s-0033-1338386>;
- (e) P. Gogoi, A.K. Dutta, P. Sarma, R. Borah, Development of Brønsted–Lewis acidic solid catalytic system of 3-methyl-1-sulfonic acid imidazolium transition metal chlorides for the preparation of bis(indolyl)methanes, *Appl. Catal. A* 492 (2015) 133–139, <https://doi.org/10.1016/j.apcata.2014.12.013>;
- (f) A. Khazaei, A. R. Moosavi-Zare, S. Firoozmand, M. R. Khodadadian, Synthesis, characterization and application of 3-methyl-1-sulfonic acid imidazolium tetrachloroferrate as nanostructured catalyst for the tandem reaction of β-naphthol with aromatic aldehydes and amide derivatives, *Appl. Organometal Chem.* 2017, e4058, <https://doi.org/10.1002/aoc.4058>;
- (g) N.G. Khaligh, Preparation, characterization and use of 3-methyl-1-sulfonic acid imidazolium hydrogen sulfate as an eco-benign, efficient and reusable ionic liquid catalyst for the chemoselective trimethylsilyl protection of hydroxyl groups, *J. Mol. Catal. A* 349 (2011) 63–70, <https://doi.org/10.1016/j.molcata.2011.08.021>;
- (h) N.G. Khaligh, Synthesis of coumarins via Pechmann reaction catalyzed by 3-methyl-1-sulfonic acid imidazolium hydrogen sulfate as an efficient, halogen-free and reusable acidic ionic liquid, *Catal. Sci. Technol.* 2 (2012) 1633–1636, <https://doi.org/10.1039/C2CY20196H>;
- (i) N.G. Khaligh, Four-component one-pot synthesis of unsymmetrical polyhydroquinoline derivatives using 3-methyl-1-sulfonic acid imidazolium hydrogen sulfate as a catalyst, *Chin. J. Catal.* 35 (2014) 1036–1042, [https://doi.org/10.1016/S1872-2067\(14\)60038-3](https://doi.org/10.1016/S1872-2067(14)60038-3);
- (j) M.A. Zolfigol, A. Khazaei, A.R. Moosavi-Zare, A. Zare, H.G. Kruger, Z. Asgari, V. Khakyzadeh, M. Kazem-Rostami, Design of ionic liquid 3-methyl-1-sulfonic acid imidazolium nitrate as reagent for the nitration of aromatic compounds by in situ generation of NO₂ in acidic media, *J. Org. Chem.* 77 (2012) 3640–3645, <https://doi.org/10.1021/jo300137w>.
- [8] R. Kore, V. Day, M.B. Shiflett, Structural identification for the reaction of chlorosulfonic acid with tertiary N-donor ligand – ionic liquid or zwitterionic compound? *ACS Sustain. Chem. Eng.* 7 (2019) 4631–4636, <https://doi.org/10.1021/acssuschemeng.8b06573>.
- [9] (a) R. Horváth, S. Skoda-Földes, Z. Mahó, L. Berente, Kollár, facile ring opening of 2,3-epoxy-steroids with aromatic amines in ionic liquids, *Steroids* 71 (2006) 706–711, <https://doi.org/10.1016/j.steroids.2006.04.006>;
- (b) A. Horváth, D. Frigyes, S. Mahó, Z. Berente, L. Kollár, R. Skoda-Földes, facile synthesis of steroidal vicinal hydroxy-sulfides via the reaction of steroidal epoxides with thiols in the presence of an ionic liquid, *Synthesis* (2009) 4037–4041, <https://doi.org/10.1055/s-0029-1217031>;
- (c) A. Horváth, Á. Szájlí, R. Kiss, J. Kóti, S. Mahó, R. Skoda-Földes, Ionic liquid promoted Wagner-Meerwein rearrangement of 16α,17α-epoxy androstanes and estranes, *J. Org. Chem.* 76 (2011) 6048–6056, <https://doi.org/10.1021/jo2006285>.
- [10] (a) E. Fehér, J. Kriván, R. Hancsók, Skoda-Földes, Oligomerisation of isobutene with silica supported ionic liquid catalysts, *Green Chem.* 14 (2012) 403–409, <https://doi.org/10.1039/C1GC15989E>;
- (b) C. Fehér, E. Kriván, J. Kovács, J. Hancsók, R. Skoda-Földes, Support effect on the catalytic activity and selectivity of SILP catalysts in isobutene trimerization, *J. Mol. Catal. A* 372 (2013) 51–57, <https://doi.org/10.1016/j.molcata.2013.02.008>;
- (c) C. Fehér, S. Tomasek, J. Hancsók, R. Skoda-Földes, Oligomerization of light olefins in the presence of a supported Brønsted acidic ionic liquid catalyst, *Appl. Catal. B* 239 (2018) 52–60, <https://doi.org/10.1016/j.apcatb.2018.08.013>.
- [11] M.J. Frisch, G.W. Trucks, H.B. Schlegel, G.E. Scuseria, M.A. Robb, J.R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G.A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H.P. Hratchian, A.F. Izmaylov, J. Bloino, G. Zheng, J.L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J.E. Peralta, Montgomery Jr., F. Ogliaro, M. Bearpark, J.J. Heyd, E. Brothers, K.N. Kudin, V.N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J.C. Burant, J. Tomasi, Iyengar, M. Cossi, N. Rega, N.J. Millam, M. Klene, J.E. Knox, J.B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R.E. Stratmann, O. Yazyev, A.J. Austin, R. Cammi, C. Pomelli, J.W. Ochterski, R.L. Martin, K. Morokuma, V.G. Zakrzewski, G.A. Voth, P. Salvador, J.J. Dannenberg, S. Dapprich, A.D. Daniels, Ö. Farkas, J.B. Foresman, J.V. Ortiz, J. Cioslowski, D.J. Fox, Gaussian 09, Revision A.02, Gaussian, Inc., Wallingford, CT, 2009.
- [12] T. Yanai, D.P. Tew, N.C. Handy, A new hybrid exchange–correlation functional using the coulomb-attenuating method (CAM-B3LYP), *Chem. Phys. Lett.* 393 (2004) 51–57, <https://doi.org/10.1016/j.cplett.2004.06.011>.
- [13] G. Sheldrick, Crystal structure refinement with SHELXL, *Acta Crystallogr. C* 71 (2015) 3–8, <https://doi.org/10.1107/S2053229614024218>.
- [14] S.P. Westrip, publCIF: software for editing, validating and formatting crystallographic information files, *J. Appl. Crystallogr.* 43 (2010) 920–925, <https://doi.org/10.1107/S0021889810022120>.
- [15] A.L. Spek, Structure validation in chemical crystallography, *Acta Crystallogr. D* 65 (2009) 148–155, <https://doi.org/10.1107/S090744490804362X>.
- [16] A.P. Purdy, R. Gilardi, J. Luther, R.J. Butcher, Synthesis, crystal structure, and reactivity of alkali and silver salts of sulfonated imidazoles, *Polyhedron* 26 (2007) 3930–3938, <https://doi.org/10.1016/j.poly.2007.04.024>.

- [17] O.M. Lezina, S.A. Rubtsova, D.V. Belykh, P.A. Slepukhin, A.V. Kutchin, Oxidation of 1-methyl-1H-imidazole-2-thiol with chlorine dioxide, *Russ. J. Org. Chem.* 49 (2013) 112–118, <https://doi.org/10.1134/S1070428013010193>.
- [18] J.A. Mejías, S. Hamad, S. Lago, Calculation of the free energy of proton transfer from an aqueous phase to liquid acetonitrile, *J. Phys. Chem. B* 105 (2001) 9872–9878, <https://doi.org/10.1021/jp003433x>.
- [19] Y. Desoky, J. Hendel, L. Ingram, S.D. Taylor, Preparation of trifluoroethyl- and phenyl-protected sulfates using sulfuryl imidazolium salts, *Tetrahedron* 67 (2011) 1281–1287, <https://doi.org/10.1016/j.tet.2010.11.085>.
- [20] Q. Bao, K. Qiao, D. Tomida, C. Yokoyama, 1-Methylimidazolium chlorosulfate ([HmIm] SO_3Cl): a novel ionic liquid with dual Brønsted–Lewis acidity *Chem. Lett.* 39 (2010) 728–729, <https://doi.org/10.1246/cl.2010.728>.
- [21] a) T. Höhle, T.F.C. Mijlhoff, The crystal structure of $NOSO_3Cl$, *Recl. Trav. Chim. Pays-Bas* 86 (1967) 1153–1158, <https://doi.org/10.1002/recl.19670861015>;
b) T. Laube, X-ray crystal structures of a benzonorbornenyl cation and of a protonated benzonorbornenol, *J. Am. Chem. Soc.* 126 (2004) 10904–10912, <https://doi.org/10.1021/ja040115t>.
- [22] S. Berger, S. Braun, 200 and More NMR Experiments: A Practical Course, Wiley-VCH, 2004.
- [23] (a) D. Bankmann, R. Giernoth, Magnetic resonance spectroscopy in ionic liquids, *Prog. Nucl. Mag. Res. Sp.* 51 (2007) 63–90, <https://doi.org/10.1016/j.pnmrs.2007.02.007>;
(b) K. Damodaran, Recent NMR studies of ionic liquids, in *Annual Reports on NMR Spectroscopy* Vol. 88 (Ed.: G. Webb), Elsevier, 2016 216–244.
- [24] M. Alei Jr., L.O. Morgan, W.E. Wageman, T.W. Whaley, pH dependence of ^{15}N NMR shifts and coupling constants in aqueous imidazole and 1-methylimidazole. Comments on estimation of tautomeric equilibrium constants for aqueous histidine, *J. Am. Chem. Soc.* 102 (1980) 2881–2887, <https://doi.org/10.1021/ja00529a002>.
- [25] (a) P. Gerothanassis, Oxygen-17 NMR spectroscopy: basic principles and applications (part I), *Prog. Nucl. Mag. Res. Sp.* 56 (2010) 95–197, <https://doi.org/10.1016/j.pnmrs.2009.09.002>;
(b) B.A. Suvorov, Nature of “chlorine effect” in the ^{17}O NMR spectra of sulfur compounds XSO_2Y Russ, *J. Gen. Chem.* 82 (2012) 1098–1100, <https://doi.org/10.1134/S1070363212060096>.
- [26] G. Barbarella, C. Chatgililoglu, S. Rossini, V. Tugnoli, ^{33}S and ^{17}O NMR of compounds containing the SO_2 moiety. The chlorine effect, *J. Magn. Res.* 70 (1986) 204–212, [https://doi.org/10.1016/0022-2364\(86\)90002-8](https://doi.org/10.1016/0022-2364(86)90002-8).
- [27] V. Lemaître, M.E. Smith, A. Watts, A review of oxygen-17 solid-state NMR of organic materials—towards biological applications, *Solid State Nucl. Magn. Reson.* 26 (2004) 215–235, <https://doi.org/10.1016/j.ssnmr.2004.04.004>.
- [28] N. Fujiwara, D. Yoshihara, H. Sakiyama, H. Eguchi, K. Suzuki, Solution oxygen-17 NMR application for observing a peroxidized cysteine residue in oxidized human SOD1, *Hyperfine Interact.* 237 (2016) 114, <https://doi.org/10.1007/s10751-016-1320-7>.
- [29] Boudin S. Yiin, Dale W. Margerum, Kinetics of hydrolysis of the chlorosulfate ion, *Inorg. Chem.* 27 (1988) 1670–1672, <https://doi.org/10.1021/ic00283a002>.
- [30] M. Eriksen, R. Fehrmann, G. Hatem, M. Gaune-Escard, O.B. Lapina, V.M. Mastikhin, Conductivity, NMR, thermal measurements, and phase diagram of the $K_2S_2O_7$ – $KHSO_4$ system, *J. Phys. Chem.* 100 (1996) 10771–10778, <https://doi.org/10.1021/jp953744i>.
- [31] T.P. Bochkareva, B.V. Passet, K.R. Popov, N.V. Platonova, T.I. Kovalchuk, Sulfonation of substituted azoles with sulfur trioxide in dichloroethane, *Chem. Heterocycl. Compd.* 23 (1987) 1084–1089, <https://doi.org/10.1007/BF00476538>.
- [32] S.K. Davidowski, F. Thompson, W. Huang, M. Hasani, S.A. Amin, C.A. Angell, J.L. Yarger, NMR characterization of ionicity and transport properties for a series of diethylmethylamine based protic ionic liquids, *J. Phys. Chem. B* 120 (18) (2016) 4279–4285, <https://doi.org/10.1021/acs.jpcc.6b01203>.
- [33] G. A. Olah, G. K. S. Prakash, Á. Molnár, J. Sommer, *Superacid Chemistry*, Wiley, 2009, pp. 36–37.
- [34] (a) P. Wasserscheid, M. Sessing, W. Korth, W. Hydrogensulfate and tetrakis (hydrogensulfato)borate ionic liquids: synthesis and catalytic application in highly Brønsted-acidic systems for Friedel–Crafts alkylation, *Green Chem* 4 (2002) 134–138, <https://doi.org/10.1039/B109845B>;
(b) S. Tang, A.M. Scurto, B. Subramaniam, Improved 1-butene/isobutane alkylation with acidic ionic liquids and tunable acid/ionic liquid mixtures, *J. Catal* 268 (2009) 243–250, <https://doi.org/10.1016/j.jcat.2009.09.022>;
(c) K. Matuszek, A. Brzeczek-Szafran, D. Kobus, D.R. MacFarlane, M. Swadźba-Kwaśny, A. Chrobok, Protic ionic liquids based on oligomeric anions $[(HSO_4)_x]^-$ ($x = 0, 1, \text{ or } 2$) for a clean ϵ -caprolactam synthesis, *Aust. J. Chem.* 72 (2019) 130–138, <https://doi.org/10.1071/CH18384>.