

# Early hemorheological changes in a porcine model of intravenously given *E. coli* induced fulminant sepsis

Norbert Nemeth<sup>a,\*</sup>, Mariann Berhes<sup>b</sup>, Ferenc Kiss<sup>a</sup>, Endre Hajdu<sup>b</sup>, Adam Deak<sup>a</sup>, Abel Molnar<sup>a</sup>, Judit Szabo<sup>c</sup> and Bela Fulesdi<sup>b</sup>

<sup>a</sup>*Department of Operative Techniques and Surgical Research, Institute of Surgery, University of Debrecen, Hungary*

<sup>b</sup>*Department of Anesthesiology and Intensive Therapy and, University of Debrecen, Hungary*

<sup>c</sup>*Department of Medical Microbiology, Faculty of Medicine, University of Debrecen, Hungary*

**Abstract.** The pathophysiology of hemorheological and microcirculatory disturbances in septic process -mostly during the early hours- still not clarified in all the details, yet. In anesthetized pigs living *E. coli* (ATCC 25922 strain) was administered intravenously with an increasing concentration and the animals were observed for 8 hours. Before the intervention and in every 2 hours arterial (cannulated femoral artery) and venous (cannulated external jugular vein) blood samples were collected for hemorheological laboratory tests: blood and plasma viscosity, ESR, leukocyte anti-sedimentation rate, erythrocyte deformability (together with osmoscan parameters) and erythrocyte aggregation (using light-transmission and laser back-scattering methods). Control animals were stable over the 8-hour anesthesia, while septic animals died by the 6th hours in a fulminant sepsis. Over the experimental period, the tendency of impairment in erythrocyte deformability (together with osmotic gradient ektacytometry parameters) and the controversial decreasing of erythrocyte aggregation values (declining all aggregation index values, elongating  $t_{1/2}$ ) were well detected in this porcine model during the early hours (4–6) of fulminant sepsis. The *in vitro* effect of these bacteria on erythrocytes' micro-rheological parameter was similar: decreasing red blood cell deformability and lowering aggregation. Further studies are needed to clarify the early micro-rheological changes of bacteremia and the developing sepsis.

**Keywords:** Red blood cell deformability, red blood cell aggregation, *E. coli*, sepsis, animal model

## 1. Introduction

Sepsis still means a serious challenge in the clinical practice. It is estimated that 13 million people become septic each year, the mortality of severe sepsis is about 30–50% and of the septic shock it is over 50% [23, 40, 47]. The pathophysiology of the septic process (colonization – infection – bacteremia and with systemic inflammatory response syndrome turning towards sepsis – severe sepsis – septic shock and multiorgan failure) is very complex and not completely clarified, yet. The sepsis pathophysiology includes numerous links to the rheology of blood since sepsis is considered to be a disorder of the microcirculation [24, 42, 45].

\*Corresponding author: Norbert Nemeth, M.D., Ph.D., Department of Operative Techniques and Surgical Research, Faculty of Medicine, University of Debrecen, H-4032 Debrecen, Nagyerdei krt. 98., Hungary. Tel./Fax: +36 52 416 915; E-mail: nemeth@med.unideb.hu.

30 The disturbed microcirculation in sepsis can be resulted by numerous factors including blood and  
31 plasma viscosity alterations, disturbed micro-vascular blood flow and vascular resistance, decreased red  
32 blood cell deformability, increased red blood cell aggregation among others [4, 16, 24, 42]. It is known  
33 that fibrinogen concentration and plasma viscosity increases, and red blood cell deformability impairs and  
34 their aggregation is enhanced significantly both in experimental sepsis and in clinical manifestations [4,  
35 7, 8, 34, 38]. Furthermore, filterability of neutrophils also significantly decreases, while their aggregation  
36 markedly increases in severe sepsis and septic shock [11, 48, 47].

37 However, the magnitude of changes and the dynamics of micro-rheological alterations as well as  
38 the 'mystified border' between reversibility and irreversibility are still unclear. Numerous experimental  
39 models are known using various animal species and different methods to induce the sepsis [17, 18, 37].  
40 Besides the limitations of the experimental model and sometimes the difficult comparability, these *in vivo*  
41 studies are highly important to reveal and understand the pathomechanism. Intravenously administered  
42 live bacteria method is one of the ways, by which an increasing magnitude of bacteremia can be produced,  
43 being comparable with some clinical situation [3, 12, 18, 37, 44]. However, the complete hemorheological  
44 characteristics of the very early hours of induced bacteremia and of the response reaction is not known  
45 completely.

46 In our study we aimed to investigate whether *E. coli* bacteremia of increasing concentration results in  
47 sepsis and whether micro-rheological parameters do alter during the early hours of the septic process in  
48 a porcine model.

## 49 2. Materials and methods

### 50 2.1. Experimental animals and protocol

51 The experiments were carried in accordance with the European Community guidelines and State  
52 Regulations with the approval of the University of Debrecen Committee of Animal Welfare (reg. Nr.:  
53 21/2013. DEMAB). Nine juvenile female Hungahib pigs (bodyweight:  $17.8 \pm 2$  kg) were subjected into  
54 Control ( $n = 4$ ) or Septic Group ( $n = 5$ ). Under general anesthesia (15 mg/kg ketamin, i.m.+1 mg/kg xylazin  
55 i.m.) tracheostomy was performed for assisted ventilation, and the left external jugular vein and femoral  
56 artery have been cannulated for sampling and hemodynamic measurements. The animals did not receive  
57 anticoagulant.

58 In *Sepsis group Escherichia coli* suspension ( $2.5 \times 10^5$ /ml; strain: ATCC 25922, Department of Medi-  
59 cal Microbiology, University of Debrecen) was intravenously administrated in a continuously increasing  
60 manner as the followings: 2 ml in the first 30 min, then 4 ml in 30 min and afterwards 16 ml/h for 2 hours (so  
61 a total of  $9.5 \times 10^6$  *E. coli* within 3 hours) (Fig. 1). In the *Control Group* the anesthesia was maintained  
62 for 8 hours, infusion was administered as a similar volume of isotonic saline solution and no other  
63 intervention was made. At the end of the experimental period the animals have been over-anaesthetized.

64 At the beginning of the procedure (prior to *E. coli* administration) blood samples were collected in  
65 parallel from the cannulated external jugular vein and the femoral artery (base samples) directly into tubes  
66 containing K<sub>3</sub>-EDTA (1.5 mg/ml) for all the laboratory tests, except for sedimentation measurements of  
67 venous blood, where 0.109 M sodium-citrate was used as anticoagulant. Further arterial and venous blood  
68 samples were collected at 2, 4, 6 and 8 hours after the time point when the infusion started. At the same  
69 times, body temperature (central venous, T [°C]), mean arterial pressure (MAP [mmHg]), heart rate (HR  
[1/min]) and oxygen saturation (Sat [%]) were recorded.




Fig. 1. Protocol of the intravenous administration of living *E. coli* suspension ( $2.5 \times 10^5/\text{ml}$ ). Dashed lined plot shows the step-wisely increased volume of the suspension [ml], the continuous black line represents the sum amount of the administered *E. coli* bacteria ( $\times 10^5$ ) during the first 3 hours of the experiment.

## 2.2. *In vitro* study (effect of the presence of living bacteria on blood rheology)

Venous blood samples taken from two animals at the beginning of the experiment (base or normal blood sample) were divided into two sub-samples. Those were mixed with suspension of *E. coli* bacteria at half and total dose that was presented *in vivo* (according to the totally given amount of bacteria described above and the circulating blood volume, as 60 ml/kg).

Within 15–20 minutes after the mixing and after 2 hours of incubation at room temperature micro-rheological tests (red blood cell deformability, osmotic gradient ektacytometry, red blood cell aggregation) were performed.

## 2.3. Laboratory measurements

### 2.3.1. Hematological parameters

A Sysmex F-800 semi-automated microcell counter (TOA Medical Electronics Co., Japan) was used to determine hematological parameters. In this paper white blood cell count (WBC [ $\times 10^3/\mu\text{l}$ ]), red blood cell count (RBC [ $\times 10^6/\mu\text{l}$ ]), hematocrit (Hct [%]), mean corpuscular volume (MCV [fl]), mean corpuscular hemoglobin content (MCH [pg]), and platelet count (Plt [ $\times 10^3/\mu\text{l}$ ]) are shown for comparisons.

### 2.3.2. Erythrocyte sedimentation rate and leukocyte anti-sedimentation rate

The tube containing the sodium-citrate-anticoagulated venous blood samples were taken in vertical positions and left for gravity sedimentation. After 1 hour the regular erythrocyte sedimentation rate (ESR [mm/h]) was registered.

To determine the leukocyte antisedimentation rate (LAR [%]) the upper and the lower part of the sedimentation blood column were gently separated and the white blood cell count was measured in both subsamples. The LAR was calculated using the Bogar's formula [9]:

$$LAR = 100 \times (WBC_{\text{upper}} - WBC_{\text{lower}}) / (WBC_{\text{upper}} + WBC_{\text{lower}}).$$

This parameter can provide information about the distribution of light and heavy polymorphonuclear leukocytes [9].

### 2.3.3. Blood and plasma viscosity

Viscosity measurements were carried out at 37°C using Hevimet-40 capillary viscosimeter (Hemorex Ltd., Hungary) within 1 hour after sampling. Plasma was prepared by centrifuging at 800 g for 10 minutes. Whole blood viscosity values (WBV [mPas]) at 90 s<sup>-1</sup> shear are used for the comparisons, and according to the Matrai formula, correction for 40% hematocrit has been also performed:  $WBV_{40\%/PV} = (WBV_{Hct}/PV)^{40\%/Hct}$ , where WBV<sub>40%</sub>: corrected for 40% Hct; WBV<sub>Hct</sub>: whole blood viscosity measured at the native Hct; PV: plasma viscosity; Hct: actual hematocrit value [%] of the sample [26].

### 2.3.4. Red blood cell deformability

Red blood cell deformability was determined by a LoRRca MaxSis Osmoscan device (Mechatronics BV, The Netherlands). Blood sample of 5 µl was taken into 1 ml of isotonic polyvinyl-pyrrolidone solution (360 kDa PVP in normal phosphate buffered saline; viscosity = 27 mPas, osmolality = 290–300 mOsm/kg; pH 7.3) and mixed gently. Based upon the laser diffraction pattern changes the elongation index (EI) values were determined in the function of shear stress (SS) in a range of 0.3–30 Pa. The EI is equal to  $(L-W) / (L+W)$ , where L is the length and W is the width of the diffractogram [19]. EI increases with red blood cell deformability. The measurements were carried out at 37 °C. Comparing EI-SS curves Lineweaver-Burk analysis was performed, calculating the maximal elongation index (EI<sub>max</sub>) and the shear stress values at half EI<sub>max</sub> (SS<sub>1/2</sub> [Pa]):  $1/EI = SS_{1/2} / EI_{max} \times 1/SS + 1/EI_{max}$ . Furthermore, EI<sub>max</sub> / SS<sub>1/2</sub> ratio was also calculated [6].

For the osmotic gradient ektacytometry (osmoscan) test [15] 250 µl of blood was gently mixed in 5 ml PVP solution. At constant shear stress of 30 Pa the elongation index values were continuously determined while the osmolality was changing (rising gradually from 0 to 500 mOsmol/kg). Among the device-given parameters we analyzed the minimal elongation index values measured at low-osmotic environment (minimal EI), the maximal elongation index values (maximal EI), the belonging osmolality values (minO and maxO as 'optimal' osmolality), and the area under the individual elongation index-osmolality curves (AUC). Additionally, we calculated further parameters describing the phase between minEI and maxEI in the function of osmolality: ΔEI, as the difference between maximal and minimal EI values; ΔO, as the difference between osmolality values at maximal and minimal EI; and ΔEI / ΔO [31].

### 2.3.5. Red blood cell aggregation

Based on light-transmittance method, a Myrenne MA-1 erythrocyte aggregometer (Myrenne GmbH, Germany) was used for determining aggregation index values: M (at shear rate of 0 s<sup>-1</sup>) and M1 (at shear rate of 3 s<sup>-1</sup>) 5 or 10 seconds after disaggregation. The indices (M 5 s, M1 5 s, M 10 s, M1 10 s) increase with enhanced red blood cell aggregation [19]. The LoRRca device was also used to measure red blood cell aggregation parameters by syllectometry determining the following parameters: amplitude (Amp [au]), aggregation index (AI [%]) and the aggregation half-time (t<sub>1/2</sub> [s]) [19].

## 2.4. Statistical analyses

Data are presented as means ± standard deviation (S.D.). According to the normality of data distribution, for inter-group (Control vs. Sepsis) comparison Student *t*-test or Mann-Whitney RS test, for

intra-group comparison (vs. base values) one-way ANOVA tests (Dunn's or Bonferroni's method) were used. A  $p$  value less than 0.05 was considered as statistically significant.

### 3. Results

#### 3.1. Hemodynamical changes and mortality rate

In the Septic group, as expected, all animals developed fulminant sepsis and died within by 3-7 hours (2 animals in 3–4 hours, and 3 in 6–7 hour). In the septic animals the heart rate rose and mean arterial pressure dropped, their ratio increased significantly compared both the base values (at the 6th hour:  $p < 0.001$ ) and the Control group ( $p = 0.004$ ). The control animals showed stable condition over the 8-hour anesthesia (Table 1).

#### 3.2. Hematological parameters

Table 2 summarizes selected hematological parameters. Total leukocyte count showed a continuously decreasing tendency in the Sepsis group, both in arterial and venous samples. By the 4th and 6th hours the values were significantly lower than the base values and that of the Control group ( $p < 0.001$ ). The Control values were constant showing only an elevation by the end of the experimental period. The monocyte-granulocyte% showed slight increasing in both groups, but without significant differences.

Red blood cell count was almost constant in both groups, however, by the 6th hour it was increased together with the hematocrit in the septic animals. Intergroup difference reached a significant level at the 4th and 6th hours, both in arterial and venous blood. MCV values were slightly higher in Septic group compared to the Control.

Platelet count dominantly decreased in septic animals, reaching the significance level by the 6th hours compared both the base and control values ( $p < 0.001$ ).

#### 3.3. Erythrocyte sedimentation rate and leukocyte anti-sedimentation rate

Erythrocyte sedimentation rate (ESR) remained constant in the control animals during the entire experimental period. In septic animals the ESR values increased stepwise till the 4th hours, and then decreased (Table 3).

Table 1

Body temperature (Temp), heart rate (HR) / mean arterial pressure (MAP) ratio and oxygen saturation values of the Control and the Sepsis groups during the experimental period

Variable	Group	Base	2 h	4 h	6 h	8 h
Temp [°C]	Control	36.73 ± 0.47	37.43 ± 0.41	37.83 ± 0.7	38.76 ± 1.35	39 ± 1.47
	Sepsis	37 ± 1.21	37.85 ± 0.97	38.2 ± 0.42	38.5 ± 0.28	–
HR / MAP [1/mmHg.min]	Control	0.86 ± 0.21	1.09 ± 0.34	0.98 ± 0.17	0.98 ± 0.21	1 ± 0.2
	Sepsis	0.82 ± 0.18	1.48 ± 1.24	1.31 ± 0.24 *	3.09 ± 0.82 *#	–
Saturation [%]	Control	97 ± 1.8	96.7 ± 0.9	94 ± 2.7	95.2 ± 1.5	95 ± 1.4
	Sepsis	94.2 ± 3.1	91 ± 4.35	93.6 ± 6.1	90 ± 15.5	–

means ± S.D., \* $p < 0.05$  vs. base, # $p < 0.05$  vs. Control; at 4 h and at 6 h  $n = 3$  in Sepsis Group (used test: Bonferroni's method of one-way ANOVA on rank test, Mann-Whitney rank sum test).

Table 2

Changes of selected hematological parameters in arterial and venous blood samples of the Control and the Sepsis groups

Variable	Group	Sample	Base	2 h	4 h	6 h	8 h
WBC [ $\times 10^3/\mu\text{l}$ ]	Control	A	15.3 $\pm$ 2.39	16.61 $\pm$ 0.79	17.63 $\pm$ 2.05	17.97 $\pm$ 0.61	20.45 $\pm$ 2.64
		V	15.21 $\pm$ 1.97	17.36 $\pm$ 1.17	16.95 $\pm$ 2.69	16.16 $\pm$ 2.68	17.45 $\pm$ 7.05
	Sepsis	A	14.88 $\pm$ 3.21	13.94 $\pm$ 4.98	8.88 $\pm$ 3.77 <sup>##</sup>	7.05 $\pm$ 3.58 <sup>##</sup>	–
		V	14.53 $\pm$ 2.64	14.49 $\pm$ 5.08	8.85 $\pm$ 4.09 <sup>##</sup>	7.88 $\pm$ 3.64 <sup>##</sup>	–
Mo+Gr% [%]	Control	A	43.25 $\pm$ 9.54	41.88 $\pm$ 8.84	53.2 $\pm$ 10.39	60.21 $\pm$ 13.39	68.25 $\pm$ 12.3
		V	46.47 $\pm$ 10.7	41.42 $\pm$ 10.02	52.7 $\pm$ 11.26	58.35 $\pm$ 10.16	67.31 $\pm$ 8.22
	Sepsis	A	48.81 $\pm$ 6.77	46.14 $\pm$ 11.27	45.73 $\pm$ 16.42	59.14 $\pm$ 20.49	–
		V	45.85 $\pm$ 7.03	45.49 $\pm$ 11.7	47.15 $\pm$ 20.2	53.15 $\pm$ 18.42	–
RBC [ $\times 10^6/\mu\text{l}$ ]	Control	A	6.21 $\pm$ 0.51	6.48 $\pm$ 0.47	5.97 $\pm$ 0.85	5.99 $\pm$ 0.84	6.21 $\pm$ 0.71
		V	6.27 $\pm$ 0.43	6.45 $\pm$ 0.31	5.85 $\pm$ 0.83	5.53 $\pm$ 1.14	5.78 $\pm$ 0.53
	Sepsis	A	6.65 $\pm$ 0.44	6.17 $\pm$ 0.32	6.7 $\pm$ 0.7	6.98 $\pm$ 0.89	–
		V	6.44 $\pm$ 0.64	6.49 $\pm$ 0.33	6.85 $\pm$ 0.8	6.63 $\pm$ 0.93	–
MCV [fl]	Control	A	59.83 $\pm$ 4.54	62.76 $\pm$ 6.3	60.17 $\pm$ 2.82	58.71 $\pm$ 1.23	60.5 $\pm$ 3.41
		V	57.65 $\pm$ 2.8	60.3 $\pm$ 3.88	58.26 $\pm$ 1.47	60.42 $\pm$ 5.72	60.11 $\pm$ 1.62
	Sepsis	A	61.99 $\pm$ 2.62	65.23 $\pm$ 7.88	65.93 $\pm$ 4.4	64.26 $\pm$ 3.05	–
		V	63.88 $\pm$ 4.58	64.81 $\pm$ 6.44	63.63 $\pm$ 2.07	64.86 $\pm$ 3.26	–
MCH [pg]	Control	A	15.21 $\pm$ 0.87	14.76 $\pm$ 0.11	15.42 $\pm$ 0.89	15.48 $\pm$ 1.4	14.58 $\pm$ 0.99
		V	14.91 $\pm$ 0.24	15.12 $\pm$ 0.39	15.07 $\pm$ 0.31	15.03 $\pm$ 1.21	15.37 $\pm$ 0.33
	Sepsis	A	15.66 $\pm$ 0.64	16.08 $\pm$ 0.83	17.08 $\pm$ 0.85	16.43 $\pm$ 1.76	–
		V	16.35 $\pm$ 1.39	16 $\pm$ 0.86	16.73 $\pm$ 1.3	16.81 $\pm$ 0.92	–
Plt [ $\times 10^3/\mu\text{l}$ ]	Control	A	608.5 $\pm$ 40.2	612.1 $\pm$ 127.4	565 $\pm$ 152.1	630.6 $\pm$ 105.8	555 $\pm$ 131.1
		V	562.1 $\pm$ 119.3	562.3 $\pm$ 145.1	517.6 $\pm$ 114.8	578 $\pm$ 83.7	477.6 $\pm$ 88.7
	Sepsis	A	470.5 $\pm$ 55.1	401.75 $\pm$ 125.4	382.3 $\pm$ 126.2	221.5 $\pm$ 32.5 <sup>##</sup>	–
		V	421.6 $\pm$ 88.9	384.6 $\pm$ 111.6	346.1 $\pm$ 108	204.5 $\pm$ 42.09 <sup>##</sup>	–

means  $\pm$  S.D., \* $p$  < 0.05 vs. base, # $p$  < 0.05 vs. Control. A = arterial blood sample, V = venous blood sample at 4 h and at 6 h  $n$  = 3 in Sepsis Group (used test: Bonferroni's method of one-way ANOVA on rank test, Mann-Whitney rank sum test).

Leukocyte anti-sedimentation rate showed a declining tendency in the septic animals: almost every 2 hours it decreased by 15–20%. The values differed significantly from the Control group (at 2 h:  $p$  = 0.031, at 4 h:  $p$  = 0.05, at 6 h:  $p$  = 0.049) (Table 3).

### 3.4. Blood and plasma viscosity

Whole blood viscosity moderately decreased over the experimental period in the control animals, thanking to the blood samplings and infusion. However, plasma viscosity did not changed importantly. The WBV values corrected for 40% hematocrit were almost constant. In the Sepsis group the whole blood viscosity showed an elevation by the 4th hours while plasma viscosity continuously decreased (Table 3).

### 3.5. Red blood cell deformability

Red blood cell deformability of the Sepsis group showed worsening; in every 2 hours the  $SS_{1/2}$  increased and the  $EI_{\max}/SS_{1/2}$  ratio decreased both in arterial and venous blood. However, the differences did not reach a statistically significant level. There was no obvious arterio-venous difference (Fig. 2, A-F).

Table 3

Changes of whole blood viscosity (WBV), plasma viscosity (PV), corrected WBV at 40% hematocrit (WBV<sub>40%</sub>), erythrocyte sedimentation rate (ESR) and leukocyte anti-sedimentation rate (LAR) of the Control and the Sepsis groups.

Variable	Group	Base	2 h	4 h	6 h	8 h
WBV [mPas] at shear rate of 90 s <sup>-1</sup>	<i>Control</i>	2.87 ± 0.6	2.74 ± 0.62	2.31 ± 0.89	2.46 ± 0.39	2.28 ± 0.34
	<i>Sepsis</i>	3.03 ± 0.51	2.66 ± 0.65	3.02 ± 0.71	2.77 ± 0.91	–
PV [mPas]	<i>Control</i>	0.98 ± 0.09	0.98 ± 0.03	1 ± 0.11	1 ± 0.21	0.9 ± 0.12
	<i>Sepsis</i>	1.03 ± 0.15	0.95 ± 0.02	0.94 ± 0.02	0.91 ± 0.03	–
WBV <sub>40%</sub> [mPas]	<i>Control</i>	3.41 ± 1.29	2.98 ± 0.85	2.58 ± 0.81	3.19 ± 1.09	2.57 ± 0.31
	<i>Sepsis</i>	2.95 ± 0.46	2.57 ± 0.65	2.72 ± 0.38	2.54 ± 0.59	–
Htc / WBV [%/mPas]	<i>Control</i>	12.96 ± 3.4	14.8 ± 4.08	15.66 ± 3.38	13.77 ± 3.25	12.81 ± 5.13
	<i>Sepsis</i>	13.8 ± 2.5	16.47 ± 4.01	14.73 ± 2.43	16.25 ± 4.26	–
ESR [mm/h]	<i>Control</i>	3.5 ± 1	3.5 ± 1	3 ± 1.15	3.75 ± 1.25	3.75 ± 1.25
	<i>Sepsis</i>	3.75 ± 0.5	4.25 ± 0.5	4.66 ± 1.15	3.66 ± 0.57	–
LAR [%]	<i>Control</i>	21.98 ± 14.33	27.09 ± 7.33	21.11 ± 6.01	22.02 ± 6.74	19.41 ± 7.62
	<i>Sepsis</i>	17.68 ± 4.9	12.95 ± 8.12 <sup>#</sup>	10.48 ± 4.59 <sup>#</sup>	6.91 ± 6.52 <sup>#</sup>	–

means ± S.D., \**p* < 0.05 vs. base, <sup>#</sup>*p* < 0.05 vs. Control; at 4 h and at 6 h *n* = 3 in Sepsis Group (used test: Bonferroni's method of one-way ANOVA on rank test, Mann-Whitney rank sum test).

Table 4

Changes of selected osmotic gradient ektacytometry (osmoscan) parameters in arterial and venous blood samples of the Control and the Sepsis groups.

Variable	Group	Sample	Base	2 h	4 h	6 h	8 h
minEI	<i>Control</i>	A	0.111 ± 0.007	0.115 ± 0.011	0.111 ± 0.009	0.118 ± 0.011	0.126 ± 0.017
		V	0.120 ± 0.011	0.117 ± 0.017	0.121 ± 0.009	0.127 ± 0.008	0.122 ± 0.021
	<i>Sepsis</i>	A	0.105 ± 0.014	0.116 ± 0.023	0.130 ± 0.03	0.141 ± 0.016 *	–
		V	0.115 ± 0.024	0.104 ± 0.019	0.131 ± 0.009	0.121 ± 0.02	–
maxEI	<i>Control</i>	A	0.502 ± 0.01	0.499 ± 0.01	0.491 ± 0.007	0.493 ± 0.004	0.490 ± 0.004
		V	0.501 ± 0.011	0.500 ± 0.009	0.488 ± 0.01	0.487 ± 0.011	0.484 ± 0.01
	<i>Sepsis</i>	A	0.491 ± 0.012	0.487 ± 0.009	0.491 ± 0.011	0.491 ± 0.016	–
		V	0.490 ± 0.01	0.488 ± 0.011	0.493 ± 0.013	0.490 ± 0.014	–
minO [mOsm/kg]	<i>Control</i>	A	187.7 ± 5.8	188.5 ± 8.5	192.5 ± 7.6	191.2 ± 3.6	190.5 ± 3.8
		V	189.7 ± 5.9	191.7 ± 6.8	191.5 ± 4.1	192.7 ± 7	192.2 ± 4.5
	<i>Sepsis</i>	A	194 ± 7.3	200.6 ± 10.4	195.3 ± 3.5	191 ± 3.6	–
		V	199.4 ± 10.7	198.2 ± 10.6	197.3 ± 0.5	181 ± 23.5	–
maxO [mOsm/kg]	<i>Control</i>	A	379.2 ± 13.8	380 ± 12.7	380 ± 8.7	369.5 ± 11.2	382.2 ± 22.7
		V	377.7 ± 13.4	388 ± 13.2	383 ± 17	390.7 ± 14.4	391 ± 21.1
	<i>Sepsis</i>	A	379 ± 20.9	390.4 ± 28.1	392.6 ± 8.3	394.6 ± 14.3 <sup>#</sup>	–
		V	387.8 ± 15.9	391.2 ± 20.4	398.3 ± 12.8	386.3 ± 9.45	–
AUC	<i>Control</i>	A	117.7 ± 5.4	112.9 ± 5.1	109.3 ± 6.7	107.5 ± 8.9	109.9 ± 9.2
		V	112 ± 5.3	110.9 ± 5.4	108.3 ± 9.7	105.9 ± 5.9	104.3 ± 7.5
	<i>Sepsis</i>	A	115.3 ± 4.3	106.7 ± 15.3	108.1 ± 7.4	107 ± 11.5	–
		V	105 ± 12.5	104.9 ± 10.6	108 ± 2.6	107.7 ± 11.4	–

means ± S.D., \**p* < 0.05 vs. base, <sup>#</sup>*p* < 0.05 vs. Control. A = arterial blood sample, V = venous blood sample at 4 h and at 6 h *n* = 3 in Sepsis Group (used test: Bonferroni's method of one-way ANOVA on rank test, Mann-Whitney rank sum test).




Fig. 2. EI<sub>max</sub> (A,B), SS<sub>1/2</sub> [Pa] (C,D), EI<sub>max</sub>/SS<sub>1/2</sub> [1/Pa] (E,F) values in arterial and venous blood of Control and Sepsis Groups. means ± S.D., \**p* < 0.05 vs. base, #*p* < 0.05 vs. Control (at 4 h and at 6 h *n* = 3 in Sepsis Group; used test: Bonferroni's method of one-way ANOVA on rank test, Mann-Whitney rank sum test).

165 Investigating the osmoscan parameters, the minEI values continuously increased mostly in the arterial  
 166 blood of the septic animals (at 6 h: *p* = 0.017 vs. base). The belonging osmolality values showed a  
 167 decreasing tendency both in arterial and venous blood. The maximal EI lowered, the belonging osmolality  
 168 slightly increased in the septic animals (at 6 h: *p* = 0.047 vs. Control). The AUC showed larger differences:






Fig. 3.  $\Delta EI$  (A,B),  $\Delta O$  (C,D) and  $\Delta EI/\Delta O$  (E,F) values in arterial and venous blood of Control and Sepsis Groups. means  $\pm$  S.D., \* $p < 0.05$  vs. base, # $p < 0.05$  vs. Control (at 4 h and at 6 h  $n = 3$  in Sepsis Group; used test: Bonferroni's method of one-way ANOVA on rank test, Mann-Whitney rank sum test).

it decreased by the 2nd hour and showed further lowering in the septic animals (Table 4).  $\Delta EI$  values gradually decreased mostly in the arterial blood of the septic animals, while  $\Delta O$  increased stepwise till the 6th hour. Their ratio ( $\Delta EI/\Delta O$ ) showed more expressed alteration: it gradually decreased mostly in the arterial blood (Fig. 3, A-F).

### 3.6. Red blood cell aggregation

In Sepsis group the red blood cell aggregation index (AI%) remained almost unchanged till the end of the 4th hour, then declined (at 6 h:  $p < 0.001$  vs. base and  $p = 0.003$  vs. Control in the arterial blood;  $p = 0.011$  vs. base and  $p = 0.002$  vs. Control in the venous blood). This alteration was more prominent in the venous blood. The amplitude values showed marked arterio-venous differences in both the Control and Sepsis group. The declination, which was more expressed in venous blood samples, reached the significance level from the 2nd hour (at 2 h:  $p = 0.001$  vs. base, at 4 h:  $p < 0.001$  vs. base in the venous blood; at 6 h:  $p = 0.02$  vs. base and  $p = 0.043$  vs. Control in the arterial, and  $p < 0.001$  vs. base and  $p = 0.029$  vs. Control in venous blood). The  $t_{1/2}$  was elongated and rose by the 6th hours in septic animals ( $p < 0.001$  vs. base and  $p = 0.003$  vs. Control in the venous,  $p = 0.02$  vs. base and  $p = 0.002$  vs. Control in the venous blood samples) (Table 5).

Parameters tested by the Myrenne aggregometer showed again a decreasing tendency in the Septic group at the 4th and 6th hour. The difference was significant by the 6th hours in case of the '5 sec M' (arterial blood:  $p < 0.001$  vs. base and Control; venous blood:  $p = 0.001$  vs. base and  $p = 0.002$  vs. Control; arterio-venous difference:  $p = 0.003$ ), '5 sec M1' (arterial blood:  $p = 0.027$  vs. base; venous blood:  $p < 0.001$  vs. base and  $p = 0.023$  vs. Control), '10 sec M' (arterial blood:  $p = 0.022$  vs. base,  $p = 0.028$  vs. Control; venous blood:  $p < 0.001$  vs. base and Control) as well as in '10 sec M1' (only in venous blood:  $p = 0.027$  vs. base; arterio-venous difference:  $p = 0.053$ ) index values (Fig. 4, A-H).

Table 5

Changes of red blood cell aggregation parameters (aggregation index, AI; amplitude, Amp and aggregation half time,  $t_{1/2}$ ) in arterial and venous blood samples of the Control and the Sepsis groups during the experimental period

Variable	Group	Sample	Base	2 h	4 h	6 h	8 h
AI [%]	Control	A	67.39 ± 7.75	66.96 ± 3.48	64.65 ± 3.17	64.18 ± 5.4	61.09 ± 7.33
		V	67.28 ± 6.54	68.35 ± 2.93	61.36 ± 4.79	67.57 ± 2.34	66.77 ± 4.39
	Sepsis	A	63.64 ± 3.39	61.35 ± 6.24	62.21 ± 6.39	55.31 ± 2.85 <sup>#</sup>	–
		V	63.69 ± 2.78	62.75 ± 5.23	63.48 ± 5.12	51.43 ± 8.8 <sup>#</sup>	–
Amp	Control	A	19.16 ± 1.67	20.61 ± 2.05	18.83 ± 3.43	20.52 ± 3.22	18.73 ± 2.63
		V	15.37 ± 0.67 <sup>+</sup>	15.4 ± 1.78 <sup>+</sup>	12.81 ± 2.84 <sup>+</sup>	12.96 ± 3.71 <sup>+</sup>	12.16 ± 2.45 <sup>+</sup>
	Sepsis	A	20.85 ± 1.08	19.63 ± 2.84	19.96 ± 3.78	16.39 ± 5.31 <sup>#</sup>	–
		V	17.76 ± 2.39 <sup>+</sup>	13.08 ± 3.03 <sup>++</sup>	10.93 ± 2.74 <sup>++</sup>	8.82 ± 1.44 <sup>++#</sup>	–
$t_{1/2}$ [s]	Control	A	1.88 ± 0.81	1.8 ± 0.33	2 ± 0.35	2.11 ± 0.59	2.51 ± 0.93
		V	1.85 ± 0.67	1.68 ± 0.27	2.37 ± 0.57	1.75 ± 0.28	1.83 ± 0.43
	Sepsis	A	2.17 ± 0.35	2.49 ± 0.8	2.28 ± 0.64	3.19 ± 0.44 <sup>#</sup>	–
		V	2.16 ± 0.3	2.29 ± 0.61	2.23 ± 0.51	3.85 ± 1.48 <sup>#</sup>	–

means ± S.D., \* $p < 0.05$  vs. base, <sup>#</sup> $p < 0.05$  vs. Control; <sup>+</sup> $p < 0.05$  vs. A. A = arterial blood sample, V = venous blood sample at 4 h and at 6 h  $n = 3$  in Sepsis Group (used test: Bonferroni's method of one-way ANOVA on rank test, Mann-Whitney rank sum test).

Table 6

Changes of selected micro-rheological parameters in normal venous blood samples ( $n = 2$ ), and subsamples mixed with half or total dose of living *E. coli* bacteria just after the mixing and over 2 hours of incubation. Results of the supplementary *in vitro* study part

Variable	Base values	With half dose of bacteria after mixing	With total dose of bacteria after mixing	With half dose of bacteria after 2-hour incubation	With total dose of bacteria after 2-hour incubation
EI at 3 Pa	0.362 ± 0.003	0.350 ± 0.012	0.352 ± 0.001	0.342 ± 0.011	0.349 ± 0.001
EI <sub>max</sub>	0.509 ± 0.004	0.517 ± 0.004	0.510 ± 0.009	0.490 ± 0.008	0.485 ± 0.015
SS <sub>1/2</sub> [Pa]	1.38 ± 0.06	1.64 ± 0.17	1.55 ± 0.17	1.41 ± 0.16	1.37 ± 0.07
EI <sub>max</sub> / SS <sub>1/2</sub> [Pa <sup>-1</sup> ]	0.36 ± 0.01	0.31 ± 0.03	0.33 ± 0.04	0.35 ± 0.04	0.35 ± 0.02
minEI	0.112 ± 0.005	0.106 ± 0.014	0.096 ± 0.001	0.128 ± 0.001	0.103 ± 0.015
maxEI	0.498 ± 0.001	0.496 ± 0.003	0.486 ± 0.014	0.494 ± 0.005	0.482 ± 0.005
minO	192.5 ± 2.1	185.5 ± 2.1	185.5 ± 2	192.5 ± 0.7	189 ± 2.8
maxO	379.5 ± 14.8	361 ± 2.8	356.5 ± 7.7	371.5 ± 10.6	352.5 ± 4.9
AUC	115.9 ± 5.3	121.6 ± 3.8	113.2 ± 10.9	111.7 ± 9.5	109.7 ± 10.1
ΔEI	0.386 ± 0.015	0.390 ± 0.017	0.389 ± 0.023	0.365 ± 0.005	0.379 ± 0.02
ΔO	187 ± 16.9	175.5 ± 0.7	171 ± 5.65	179 ± 11.3	163.5 ± 7.7
ΔEI / ΔO *100	0.207 ± 0.027	0.222 ± 0.011	0.227 ± 0.006	0.204 ± 0.015	0.232 ± 0.023
AI%	59.31 ± 3.46	55.8 ± 2.9	46.12 ± 7.74	59.46 ± 8.45	47.77 ± 4.56
Amp	21.01 ± 0.29	15.87 ± 3.66	10.04 ± 6.87	15.8 ± 10.99	9.5 ± 7.28
t <sub>1/2</sub> [sec]	2.7 ± 0.44	3.17 ± 0.4	4.76 ± 1.54	2.69 ± 1.07	4.5 ± 0.83
5 sec M	3.06 ± 0.69	2.48 ± 0.65	1.4 ± 1.02	3.7 ± 0.97	1.68 ± 1.28
5 sec M1	6.17 ± 1.38	3.65 ± 1.14	2.7 ± 0.3	2.65 ± 0.78	1.82 ± 1.47
10 sec M	6.78 ± 1.72	7.68 ± 0.75	3.76 ± 1.88	10.5 ± 2.21	5.36 ± 4.68
10 sec M1	14.46 ± 7.63	11.38 ± 2.07	5.78 ± 1.87	10.63 ± 0.79	5.28 ± 4.59

means ± S.D.9.

### 3.7. Results of the *in vitro* study

Half dose of bacteria resulted in 5–6% decrease of red blood cell count, while it was about 13–15% when using the total dose. The values were slightly increased after the 2-hour incubation. The red blood cell distribution width (RDW-CV%) did not change just after mixing, while after the incubation period it increased by 4–5% and 9–10% when half or total dose of bacteria was used. The change of MCV did not exceed 2–4%.

Table 6 summarizes the red blood cell deformability (normal and osmotic gradient ektacytometry) and aggregation results. At half dose of the bacteria the aggregation parameters slightly decreased. At total dose of the *E. coli*, all the aggregation parameters were deteriorated: the index values (AI% and all the Myrenne index values) markedly decreased, which declination became more expressed after 2 hours of incubation. At the same time, red blood cell deformability impaired moderately.

## 4. Discussion

There are numerous animal models of sepsis using various methods to induce bacteremia, or using endotoxins among others [17, 18, 37, 43]. Our current model was based upon recent papers using living *E. coli* colony and administered intravenously in an increasing concentration within a relatively short period




Fig. 4. Aggregation index 5 s M (A,B), 5 s M1 (C,D), 10 s M (E,F) and 10 s M1 (G,H) values in arterial and venous blood of Control and Sepsis Groups. means  $\pm$  S.D., \* $p < 0.05$  vs. base, # $p < 0.05$  vs. Control (at 4 h and at 6 h  $n = 3$  in Sepsis Group; used test: Bonferroni's method of one-way ANOVA on rank test, Mann-Whitney rank sum test).

206 [3, 12, 44]. We concentrated on the early effects of the developing fulminant sepsis. By our knowledge  
207 there is no other paper dealing with the early hemorheological changes during i.v. administered *E. coli*  
208 induced sepsis in the pig.

209 Chvojka et al. used a peritonitis-induced septic porcine model [14]. The measurements were completed  
210 12, 18 and 22 hours after induction of the peritonitis. The hemodynamic parameters reached significant  
211 difference (decrease of MAP, increased of MPAP and CO, decrease in SVR, increase of CVP and PAOP)  
212 by 12 hours together with decrease in pH and rise in IL-6, TNF- $\alpha$  level and so reflecting the hyperdynamic  
213 circulation, the inflammatory response with oxidative and metabolic stress [14]. This was a peritonitis-  
214 induced model with a relatively quick development of the sepsis. In our model we used intravenous  
215 administration of living bacteria resulting in a rapid, fulminant sepsis. Wester et al. [46] used inactivated  
216 *N. meningitidis* for inducing sepsis in pigs. When skin and tongue microcirculation was investigated  
217 200 minutes after sepsis induction, the clinical signs of the sepsis have been obviously developed. This  
218 period matches well with our experimental phase: the time period of 2–4 hours after starting the i.v.  
219 administration of the living bacteria.

220 Numerous papers reported important results on microcirculatory and rheological disturbances in sepsis  
221 [1, 2, 8, 13, 14, 16, 21, 25, 28, 30, 34, 40, 42, 45]. Most of them concluded significant worsening of  
222 the micro-rheological parameters: impaired red blood cell deformability and enhanced red blood cell  
223 aggregation is a common finding [e.g., 8, 28, 34, 42, 45].

224 Red blood cell deformability can be altered by numerous factors in septic process, including 2,3 diphos-  
225 phoglycerate (by increased internal viscosity), nitric oxide (bound by hemoglobin, modulating membrane  
226 properties), increased intracellular concentration (by decreased activity of Ca<sup>2+</sup> - ATPase pump), sialic  
227 acid (signal recognition for capturing by the reticulo-endothelial system, modifying cell shape and increas-  
228 ing aggregation), white blood cells (producing reactive oxygen species, increase aggregation with red  
229 blood cells), reactive oxygen species themselves (lipid peroxidation, sulfhydryl-crosslinking in proteins,  
230 methemoglobin formation) [4, 34].

231 Baskurt and Mat [5] in a rat model of experimental sepsis clearly demonstrated that the temperature  
232 of the measurement (in the laboratory device) has a great importance in detecting the changes of micro-  
233 rheological parameters, such as red blood cell aggregation and deformability. While decreased elongation  
234 index of septic group has not been detected at 20 °C, 25 °C or even at 30 °C, the significant decrease of red  
235 blood cell deformability were found at 37 °C. Also red blood cell aggregation index values were obviously  
236 higher at body temperature [5]. Jagger et al. found significant alteration at 25 °C [20]. Piagnerelli et al.  
237 found a loss of biconcavity (the red blood cell shape was determined by flow cytometry) in septic patients  
238 compared to healthy volunteer. The difference was not depending on the temperature of sample [36].

239 Reggiori et al. investigated red blood cell deformability and aggregation values in intensive care unit  
240 patients, reporting early alterations of both micro-rheological parameters in septic patients. The elongation  
241 index decreased almost at all shear stresses. Aggregation index values increased and  $t_{1/2}$  shortened than  
242 that of the non-septic patients [39].

243 Piagnerelli et al. [35] analyzed red blood cell membrane proteins in healthy volunteers, non-septic  
244 intensive care unit patients and septic patients (severe sepsis or septic shock). They found that red blood  
245 cell membrane skeletal protein content was modified in critical ill patients without additional changes in  
246 sepsis [35]. Other cell components have also great importance. Also Piagnerelli et al. [33] investigated  
247 erythrocyte shape and sialic acid content of the cell membrane in septic patients (blood samples were  
248 collected during the first 24 hours of the sepsis). They found by flow cytometry analyses membrane  
249 sialic acid content significantly lowered in septic patients, the glycophorine A content was higher than  
250 in the healthy volunteers. They found that spherical index significantly decreased in sepsis. Erythrocyte

251 of septic patients had a decreased capacity of sphericity in hypoosmolar conditions. It has been shown  
252 that red blood cells' membrane sialic acid content significantly decreases in sepsis related to increased  
253 neuraminidase activity [33].

254 Moutzouri et al. reported the evidence about decreased red blood cell deformability as a prognostic  
255 marker in sepsis, and also being useful for monitoring the severity of sepsis in the clinical practice [28].  
256 The patients who developed nosocomial sepsis were entered into the study within 24 hours of the diagno-  
257 sis; consequently the early hours of the septic procedures have not been investigated from the beginning.  
258 The impairment of erythrocyte deformability was obvious and expressed a gradual deterioration in sep-  
259 sis, severe sepsis, ARDS, and septic shock, in this sequence. Red blood cell rigidity showed positive  
260 correlation with the projected mortality and the simplified acute physiology score (SAPS) II [28].

261 Leukocytes play a determining role in the pathophysiology of sepsis, having impact on the microcircu-  
262 lation, too [11, 32, 40, 47, 48]. Their margination is strongly influenced by the rheology: by the magnitude  
263 of the axial migration of the red blood cells. The erythrocyte aggregation is known to enhance it [29].  
264 Kirschenbaun et al. demonstrated not only impaired red blood cell deformability and enhanced aggre-  
265 gation but also an increased platelet-neutrophil interaction with decreased leukocyte deformability [22].  
266 Yodice et al. reported increased leuko-aggregation and decreased neutrophil deformability in patients  
267 with severe sepsis and septic shock [48].

268 Micro-rheological abnormalities are characteristic not solely for sepsis and septic shock. Sordia et  
269 al. in a rat model of traumatic shock also found significant increased of erythrocyte aggregability and  
270 decreased deformability [41].

271 In our study we found a continuously decreasing red blood cell deformability of Sepsis Group during the  
272 experimental period. The changing in osmoscan parameters also occurred. The phase of the osmoscan  
273 curve that is related to the cell deformability and membrane stability (the curve part between min EI  
274 and maxEI together with the belonging osmolality values) showed the mostly expressed alterations.  
275 The difference between these points clearly showed a gradually worsening tendency. It might reflect  
276 not only the decreasing red cell deformability but the susceptibility of the cells to damage by altered  
277 micro-environmental osmotic conditions.

278 Interestingly, in our study we observed decreasing values of the aggregation-related parameters and it  
279 seemed that the aggregation process becomes slower and lower in its magnitude within 6th hour after  
280 starting the intravenous administration of the living bacteria. However, red blood cell deformability  
281 impaired. The systemic inflammatory response reaction was obvious, therefore the presence of sepsis  
282 occurred and mortal septic shock has been developed quickly. It is interesting to deal with the data of  
283 decreased aggregation of the erythrocytes in this experiment. It is obscure why the aggregation parameters  
284 showed an opposite behavior than in other studies. Speculating on this controversial finding it is important  
285 to note that our investigations were performed in parallel with the developing septic shock in a very early  
286 period of a fulminant sepsis. Furthermore the direct effect of the existing bacteremia (with an increasing  
287 concentration) might also have an effect on the aggregation. Therefore, we tested the effect *in vitro* in  
288 two normal blood sample that has been mixed with suspension of *E. coli* bacteria approximately in the  
289 same concentration as it was presented *in vivo* (according to the totally given amount of bacteria and  
290 the circulating blood volume). It was found that the impairment of red blood cell deformability and the  
291 decrease of aggregation also occurred *in vitro* and changed further after the 2-hour incubation period. It  
292 is important to note that this *E. coli* strain has hemolysin causing  $\beta$ -hemolysis. In the *in vivo* part of the  
293 study we did not observe any decrease in red blood cell count or changes in MCV (Table 2). In the *in*  
294 *vitro* study both the red blood cell count and the MCV decreased after mixing the blood with the bacteria  
295 suspension.

It is well-known that *E. coli* hemolysin jeopardizes the red blood cell membrane by creating pores with ring-shaped heptamer complexes, thus causing irreversible osmotic swelling. The presence of complement is also important. Furthermore, another way of red cell damage is linked to the endotoxin effect. Pöschl et al. concluded that endotoxin binding may directly affect the mechanical properties of the erythrocytes. The elongation index - shear stress curves were very similar to each other -and being significantly worsened compared to healthy control- in septic blood and blood samples *in vitro* incubated with endotoxin (LPS, *E. coli* O111:B4) [38]. The deterioration of red blood cell deformation (decrease of the elongation index) was associated with an increasing  $\beta$ -hydroxymyristic acid (HMA) concentration in cell membrane. The LPS in washed red blood cells resulted in much lower changes compared to the protocol when whole blood was mixed with LPS. They also suggested that endotoxin might have a maximum effect on red blood cell deformability: as of 25  $\mu$ g LPS (corresponding to 0.66  $\mu$ g/ml HMA) bound to red blood cell membrane [38].

Since the late 1990s, the role of nitric oxide (NO) has been widely investigated. The increase of NO in septic shock -but with an initial deficit- is known to contribute to the early microvascular deterioration. Morel et al. [27] used a rat CLP (caecal ligation and puncture) model of sepsis, and found an increased NO concentration in the first 3 hours of the septic shock (peaked at 2 hours). In parallel, microcirculatory flow (tested by laser Doppler flowmetry) gradually decreased over the experiment till 3 hours after the septic shock onset. According to their results, NO deficit does not occur to cause adverse changes in the microcirculation during the very early phase of the septic process [27].

It is also known that NO has an effect on red blood cells improving their deformability and aggregation [4, 7, 10]. The effect of NO in the very early hours of the septic process might have an influence on the micro-rheological parameters during septic process. Speculating with this issue, we cannot exclude the NO effect on the red blood cell aggregation when interpret the results: why we found sustained erythrocyte aggregation during the first 2–6 hours of the sepsis. However, this issue should be further investigated with other strains of bacteria, too.

## 5. Conclusion

Due to the complexity of the findings and also in respect of the case number of this study, we cannot state that these micro-rheological changes are characteristic. However, the tendency of impairment in red blood cell deformability (together with osmotic gradient ektacytometry parameters) and the controversial decreasing of red blood cell aggregation values (declining all aggregation index values and amplitude, elongating  $t_{1/2}$ ) were well detected in this porcine model during the early hours (4–6) of fulminant sepsis that was induced by i.v. administration of living *E. coli*. The *in vitro* effect of these bacteria on erythrocytes' micro-rheological parameter was similar: decreasing red blood cell deformability and lowering aggregation. Further studies are needed for clarifying the early micro-rheological changes of bacteremia and the developing sepsis.

## Acknowledgments

Authors are grateful to the technical and laboratory staff of the Department of Operative Techniques and Surgical Research at University of Debrecen. The project was supported by the Hungarian Brain Research Program - Grant No. KTIA\_13\_NAP-A-II/5.

The authors comply with the Ethical Guidelines for Publication in *Clinical Hemorheology and Microcirculation* as published on the IOS Press website and in Volume 44, 2010, pp. 1-2 of this journal.

There is no conflict of interest.

## References

- [1] E. Alt, B.R. Amann-Vesti, C. Madl, G. Funk and R. Koppensteiner, Platelet aggregation and blood rheology in severe sepsis/septic shock: Relation to the Sepsis-related Organ Failure Assessment (SOFA) score, *Clin Hemorheol Microcirc* **30** (2004), 107–115.
- [2] M.E. Astiz, G.E. DeGent, R.Y. Lin and E.C. Rackow, Microvascular function and rheologic changes in hyperdynamic sepsis, *Crit Care Med* **23** (1995), 265–271.
- [3] A. Barratt-Due, A. Sokolov, A. Gustavsen, B.C. Hellerud, K. Egge, S.E. Pischke, J.K. Lindstad, A. Pharo, A. Castellheim, E.B. Thorgersen and T.E. Mollnes, Polyvalent immunoglobulin significantly attenuated the formation of IL-1beta in Escherichia coli-induced sepsis in pigs, *Immunobiology* **218** (2013), 683–689.
- [4] O.K. Baskurt, Mechanisms of blood rheology alterations, in: Handbook of Hemorheology and Hemodynamics, O.K. Baskurt, M.R. Hardeman, M.W. Rampling and H.J. Meiselman, eds., IOS Press, Amsterdam, The Netherlands, 2007, pp. 170-190.
- [5] O.K. Baskurt and F. Mat, Importance of measurement temperature in detecting the alterations of red blood cell aggregation and deformability studied by ektacytometry: A study on experimental sepsis in rats, *Clin Hemorheol Microcirc* **23** (2000), 43–49.
- [6] O.K. Baskurt and H.J. Meiselman, Data reduction methods for ektacytometry in clinical hemorheology, *Clin Hemorheol Microcirc* **54** (2013), 99–107.
- [7] O.K. Baskurt, B. Neu and H.J. Meiselman, Red Blood Cell Aggregation, CRC Press, Boca Raton, USA, 2012, pp. 223-267.
- [8] O.K. Baskurt, A. Temiz and H.J. Meiselman, Red blood cell aggregation in experimental sepsis, *J Lab Clin Med* **130** (1997), 183–190.
- [9] L. Bogar and M. Tekeres, Leukocyte flotation during gravity sedimentation of the whole blood, *Clin Hemorheol Microcirc* **22** (2000), 29–33.
- [10] M. Bor-Kucukatay, R.B. Wenby, H.J. Meiselman and O.K. Baskurt, Effects of nitric oxide on red blood cell deformability, *Am J Physiol Heart Circ Physiol* **284** (2003), H1577–H1584.
- [11] K.A. Brown, S.D. Brain, J.D. Pearson, J.D. Edgeworth, S.M. Lewis and D.F. Treacher, Neutrophils in development of multiple organ failure in sepsis, *The Lancet* **368** (2006), 157–169.
- [12] A. Castellheim, E.B. Thorgersen, B.C. Hellerud, A. Pharo, H.T. Johansen, F. Brosstad, P. Gaustad, H. Brun, E. Fosse, T.I. Tonnessen, E.W. Nielsen and T.E. Mollnes, New biomarkers in an acute model of live Escherichia coli-induced sepsis in pigs, *Scand J Immunol* **68** (2008), 75–84.
- [13] V.J. Castro, M.E. Astiz and E.C. Rackow, Effect of crystalloid and colloid solutions on blood rheology in sepsis, *Shock* **8** (1997), 104–107.
- [14] J. Chvojka, R. Sykora, A. Krouzecky, J. Radej, V. Varnerova, T. Karvunidis, O. Hes, I. Novak, P. Radermacher and M. Matejovic, Renal haemodynamic, microcirculatory, metabolic and histopathological responses to peritonitis-induced septic shock in pigs, *Crit Care* **12** (2008), R164.
- [15] M.R. Clark, N. Mohandas and S.B. Shohet, Osmotic gradient ektacytometry: Comprehensive characterization of red cell volume and surface maintenance, *Blood* **61** (1983), 899–910.
- [16] S. Chien, S. Usami, J. Dellenback and V. Magazinovic, Blood rheology after hemorrhage and endotoxin, *Adv Exp Med Biol* **33** (1972), 75–93.
- [17] C.T. Esmon, Why do animal models (sometimes) fail to mimic human sepsis? *Crit Care Med* **32** (2004), S219–S222.
- [18] M.P. Fink, Animal models of sepsis, *Virulence* **5** (2014), 143–153.
- [19] M.R. Hardeman, P.T. Goedhart and S. Shin, Methods in hemorheology, in: Handbook of Hemorheology and Hemodynamics, O.K. Baskurt, M.R. Hardeman, M.W. Rampling and H.J. Meiselman, eds., IOS Press, Amsterdam, The Netherlands, 2007, pp. 242-266.
- [20] J.E. Jagger, C.G. Ellis, W.J. Sibbald and O. Eichelbrönnner, Measurement temperature plays a pivotal role in the distribution of erythrocyte deformability after LPS, *Biorheology* **38** (2001), 439–448.



- 383 [21] T. Kato, Relation between blood viscosity and hematocrit in rabbits subjected to repeated treatment with endotoxin,  
384 *Resuscitation* **21** (1991), 61–66.
- 385 [22] L.A. Kirschenbaum, M. Aziz, M.E. Astiz, D.C. Saha and E.C. Rackow, Influence of rheologic changes and platelet-  
386 neutrophil interactions on cell filtration in sepsis, *Am J Respir Crit Care Med* **161** (2000), 1602–1607.
- 387 [23] M.M. Levy, M.P. Fink, J.C. Marshall, E. Abraham, D. Angus, D. Cook, J. Cohen, S.M. Opal, J.L. Vincent and G.  
388 Ramsay, 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference, *Crit Care Med* **31** (2003),  
389 1250–1256.
- 390 [24] D.J. Lundy and S. Trzeciak, Microcirculatory dysfunction in sepsis, *Crit Care Clin* **25** (2009), 721–731.
- 391 [25] C. Madl, R. Koppensteiner, B. Wendelin, K. Lenz, L. Kramer, G. Grimm, A. Kranz, B. Schneeweiss and H. Ehringer,  
392 Effect of immunoglobulin administration on blood rheology in patients with septic shock, *Circ Shock* **40** (1993),  
393 264–267.
- 394 [26] A. Matrai, R.B. Whittington and E. Ernst, A simple method of estimating whole blood viscosity at standardized hematocrit,  
395 *Clin Hemorheol* **7** (1987), 261–265.
- 396 [27] J. Morel, J.Y. Li, P. Eyenga, A. Meiller, M.P. Gustin, G. Bricca, S. Molliex and J.P. Viale, Early adverse changes in liver  
397 microvascular circulation during experimental septic shock are not linked to an absolute nitric oxide deficit, *Microvasc*  
398 *Res* **90** (2013), 187–191.
- 399 [28] A.G. Moutzouri, A.T. Skoutelis, C.A. Gogos, Y.F. Missirlis and G.M. Athanassiou, Red blood cell deformability in patients  
400 with sepsis: A marker for prognosis and monitoring of severity, *Clin Hemorheol Microcirc* **36** (2007), 291–299.
- 401 [29] G.B. Nash, T. Watts, C. Thornton and M. Barigou, Red cell aggregation as a factor influencing margination and adhesion  
402 of leukocytes and platelets, *Clin Hemorheol Microcirc* **39** (2008), 303–310.
- 403 [30] O.O. Nduka and J.E. Parrillo, The pathophysiology of septic shock, *Crit Care Nurs Clin North Am* **23** (2011),  
404 41–66.
- 405 [31] N. Nemeth, F. Kiss and K. Miszti-Blasius, Interpretation of osmotic gradient ektacytometry (osmoscan) data: A comparative  
406 study for methodological standards and data reduction – submitted to *Scand. J Clin Lab Invest* (2014).
- 407 [32] M. Nishino, H. Tanaka, H. Ogura, Y. Inoue, T. Koh, K. Fujita and H. Sugimoto, Serial changes in leukocyte deformability  
408 and whole blood rheology in patients with sepsis or trauma, *J Trauma* **59** (2005), 1425–1431.
- 409 [33] M. Piagnerelli, K.Z. Boudjeltia, A. Rapotec, T. Richard, D. Brohee, S. Babar, V. Bouckaert, A.C. Simon, J.P. Toko, T.  
410 Walravens, J.L. Vincent and M. Vanhaeverbeek, Neuraminidase alters red blood cells in sepsis, *Crit Care Med* **37** (2009),  
411 1244–1250.
- 412 [34] M. Piagnerelli, K.Z. Boudjeltia, M. Vanhaeverbeek and J.L. Vincent, Red blood cell rheology in sepsis, *Intensive Care*  
413 *Med* **29** (2003), 1052–1061.
- 414 [35] M. Piagnerelli, F. Cotton, M. Van Nuffelen, J.L. Vincent and B. Gulbis, Modifications in erythrocyte membrane protein  
415 content are not responsible for the alterations in rheology seen in sepsis, *Shock* **37** (2012), 17–21.
- 416 [36] M. Piagnerelli, K. Zouaoui Boudjeltia, P. Piro, D. Brohee, M. Vanhaeverbeek and J.L. Vincent, Effects of sample temperature  
417 on red blood cell shape in septic patients, *Clin Hemorheol Microcirc* **30** (2004), 463–466.
- 418 [37] L.F. Poli-de-Figueiredo, A.G. Garrido, N. Nakagawa and P. Sannomiya, Experimental models of sepsis and their clinical  
419 relevance *Shock* **30** Suppl 1 (2008), 53–59.
- 420 [38] J.M.N. Pöschl, C. Leray, P. Ruef, J.P. Cazenave and O. Linderkamp, Endotoxin binding to erythrocyte membrane and  
421 erythrocyte deformability in human sepsis and *in vitro*, *Crit Care Med* **31** (2003), 924–928.
- 422 [39] G. Reggiori, G. Occhipinti, A. De Gasperi, J.L. Vincent and M. Piagnerelli, Early alterations of red blood cell rheology in  
423 critically ill patients, *Crit Care Med* **37** (2009), 3041–3046.
- 424 [40] D.G. Remick, Pathophysiology of sepsis, *Am J Pathol* **170** (2007), 1435–1444.
- 425 [41] T. Sordia, J. Tatarishvili and G. Mchedlishvili, Hemorheological disorders in the microcirculation during septic shock in  
426 rats, *Clin Hemorheol Microcirc* **35** (2006), 223–226.
- 427 [42] P.E. Spronk, D.F. Zandstra and C. Ince, Bench-to-bedside review: Sepsis is a disease of the microcirculation, *Crit Care* **8**  
428 (2004), 462–468.
- 429 [43] M.E. Starr and H. Saito, Sepsis in old age: Review of human and animal studies, *Aging Dis* **5** (2014), 126–136.
- 430 [44] E.B. Thorgersen, B.C. Hellerud, E.W. Nielsen, A. Barratt-Due, H. Fure, J.K. Lindstad, A. Pharo, E. Fosse, T.I. Tonnessen,  
431 H.T. Johansen, A. Castellheim and T.E. Mollnes, CD14 inhibition efficiently attenuates early inflammatory and hemostatic  
432 responses in *Escherichia coli* sepsis in pigs, *FASEB J* **24** (2010), 712–722.
- 433 [45] A. Tyagi, A.K. Sethi, G. Girotra and M. Mohta, The microcirculation in sepsis, *Indian J Anaesth* **53** (2009),  
434 281–293.

- 435 [46] T. Wester, E. Haggblad, Z.A. Awan, A. Barratt-Due, M. Kvernebo, P.S. Halvorsen, T.E. Mollnes and K. Kvernebo, Assess-  
436 ments of skin and tongue microcirculation reveals major changes in porcine sepsis, *Clin Physiol Funct Imaging* **31** (2011),  
437 151–158.
- 438 [47] J.L. Wynn and H.R. Wong, Pathophysiology and treatment of septic shock in neonates, *Clin Perinatol* **37** (2010),  
439 439–479.
- 440 [48] P.C. Yodice, M.A. Astiz, B.M. Kurian, R.Y. Lim and E.C. Rackow, Neutrophil rheological changes in septic shock, *Am J*  
441 *Respir Crit Care Med* **155** (1997), 38–42.

Uncorrected Author Proof